



ISSN: 2348-1900

Plant Science Today

<http://www.plantsciencetoday.online>



Research Article

Allelopathic inhibition of germination, seedling growth and cell division of selected plant species by *Calotropis procera* (Ait.) Ait.

Farrukh Hussain¹, Aneela Rasool², Khalid Aziz¹, Sapna Raisham², Sajid Aziz¹, Lal Badshah² & Wahid Hussain³

¹Institute of Biological Sciences, Sarhad University of Science & Information Technology, Peshawar, Pakistan

²Department of Botany, University of Peshawar, Pakistan

³Department of Botany, GPGC Parachinar, Kurram, Pakistan

Article history

Received: 12 July 2019

Accepted: 08 September 2019

Published: 01 January 2020

Abstract

Calotropis procera (Ait.) Ait. is perennial medicinal obnoxious shrub growing in Pakistan up to 1500 m altitude. Hot and water aqueous extracts from leaves and young stems of *C. procera* were used against *Pennisetum glaucum* (Linn.) R. Br., *Setaria italica* (Linn.) P. Beauv., *Brassica campestris* Linn. and *Lactuca sativa* L. under laboratory condition. It was seen that germination, seedling growth, fresh and dry biomass reduced in concentration dependent manner. It was observed that the allelopathic effects depended upon the tested species, growth parameter measured, soaking duration and concentration of the donor plant material. The *C. procera* litter incorporated into the growth medium inhibited the test species used. The *C. procera* extracts from leaves were more inhibitory than stem extracts. The tendency of inhibition was radical growth > germination > plumule growth suggesting radicle growth to be a better measure of allelopathy. Leaf extracts significantly reduced division and size of cells. It is suggested that aqueous extract from *C. procera* can be further assessed against microbes and weed under laboratory and field condition.

Publisher

Horizon e-Publishing Group

Keywords: allelopathy; *Calotropis procera*; aqueous extracts; seedling growth; inhibited cell division and expansion

Citation: Hussain F, Rasool A, Aziz K, Raisham S, Aziz S, Badshah L, Hussain W. Allelopathic inhibition of germination, seedling growth and cell division of selected plant species by *Calotropis procera* (Ait.) Ait. Plant Science Today 2020;7(1):1-8. <https://doi.org/10.14719/pst.2020.7.1.606>

Copyright: © Hussain et al. (2020). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>).

*Correspondence

Farrukh Hussain

✉ farrukh.biotech@suit.edu.pk

Indexing: Plant Science Today is covered by Scopus, Web of Science, BIOSIS Previews, ESCI, CAS, AGRIS, UGC-CARE, CABI, Google Scholar, etc. Full list at <http://www.plantsciencetoday.online>

Introduction

Calotropis procera (Ait.) Ait. (Family Asclepiadaceae) is an erect perennial obnoxious, non-palatable medicinal shrub found up to 1500 m

altitude in drier parts of Pakistan. The non-palatability to livestock, insects and wild animals facilitates its natural spread. Some evidences regarding the allelopathy of *Calotropis* species are

Samreen *et al.* (1), Abdul-Farid *et al.* (2), Akinyede *et al.* (3), Umar *et al.* (4), Shetta *et al.* (5) and Hilal-ul-Zaman & Ahmad (6), which reported retarded cell division and seedling growth of various tested species.

Allelopathy affects the diversity, functioning and productivity of natural and agroecosystems (7, 8). *Lantana* (9) and *Celtis* (10) allelopathically reduced the productivity by affecting various growth parameters of susceptible plants. Raihan *et al.* (11) reported decreased germination and radical growth of lettuce by extracts from leaves and barks of 145 plant species. Anwar *et al.* (12) reported the allelopathic nature of papaya leaf extract on various parameters of some weeds. Similarly, Qasem (13) screened 135 plants for their allelopathic capability against the germination and radical growth of wheat. Furthermore, some plants exhibited strong to weak inhibition, while others were stimulatory. *Amaranthus retroflexus* significantly reduced the stomatal opening, photosynthetic pigments and reduced other growth parameters (14). Li *et al.*, (15) reported that *Veronica persica* had herbicidal effects against weeds due to presence of 21 compounds in ethanolic extracts. Wu *et al.* (16) concluded that *Eucalyptus* significantly retarded germination and early growth of *Physalis* that can be used as herbicide. The allelopathic effects of leaf, stem and bark of *Pistacia* were tested against crops and weeds by Tahir *et al.* (17). They identified 45 compounds from tested parts, which caused inhibition. Similarly, *Melia azedarach* exhibited strong allelopathy against chick pea and black gram (18). The allelopathic substances responsible for allelopathy were identified.

All these studies suggested that germination, early seedling growth, biochemical and physiological processes were differentially affected. The allelopathic effects usually depended upon donor plant species, test species, parts assayed, concentration and duration of soaking of plant materials. It is also reported that cytological changes and abnormalities in cell division, expansion and chromosomal aberrations occur due to extracts from the donor species.

Since some studies (1-6) on the allelopathy of *Calotropis procera* are available, therefore the present study further envisaged its allelopathic stress against germination and seedling growth of some common crop species, and its effect on the division and development of cell as a possible allelopathic mechanism. This endeavor can be extended to identify its allelochemicals and phytochemicals responsible for the allelopathy.

Materials and Methods

Leaves and young stems of mature *Calotropis procera* were collected at flowering stage from Peshawar University Campus during November,

2017. The plant material was air dried in shade at room temperature (25-30 °C) and powdered. Glassware, thoroughly washed with tap water, was sterilized at 170 °C for at least 4 h. *Pennisetum glaucum* (Linn.) R.Br., *Setaria italica* (Linn.) P. Beauv., *Brassica campestris* Linn., *Lactuca sativa* Linn. and *Allium cepa* Linn. were used as the test species in the following experiments. The results were statistically analyzed using *t*-test.

1. Aqueous extract bioassay

Five gm dried powdered leaves and young stems were separately soaked in 100 ml distilled water at 25 °C for 24 and 48 h without stirring; and filtered through Whatman filter paper. These different aqueous extracts were tested against seeds of above mentioned test species on 2-folds of Whatman filter papers in petri dishes. Tests were made by soaking the filter papers with 15 ml each of the extracts. In control extract was replaced with same volume of distilled water. Each treatment was replicated 10 times, each with 10 seeds of test species. Germination, growth of plumule and radicle were measured after 72 h incubation at 25 °C. Twenty seedlings from test and control treatments were randomly selected for determining fresh and dry weight. Seedlings were dried at 65 °C for 72 h. Moisture contents were calculated on oven dry basis (19).

For the preparation of hot water extract, 5g leaves and stems were separately boiled in 100 ml distilled water for 5 minutes and filtered. These hot-water extracts were cooled to room temperature and tested against the same test species as described above. Although, obtaining hot water extract is an unnatural process because the temperature in nature never rises to boiling point, yet many studies (7, 8, 20, 21) have suggested that not only phytochemicals were resistant but were easily released.

2. Effect of litter

Litter bed bioassay

Five gm powdered leaf and stem litter of *C. procera* were uniformly spread in a petri dish and topped with a single sheet of filter paper. In control, litter was replaced with 5 gm fine pieces of filter papers. All the dishes were provided with 20 ml distilled water, which were considered optimal moisture. After 6 hr, seeds of the test species were placed on the top of filter paper. There were 10 replicates, each with 10 seeds in each of the treatments. The petri dishes were incubated as before and the same parameters were measured.

Pot experiment

The effect of litter was further tested by placing 5 gm crushed leaf litter in small plastic pots (measuring 6x9 cm tapering down to 6 cm) containing equal volume of coarse sand, which had been sterilized at 170 °C for 4 h. For each

treatment five replicates, each with 10 seeds were made. Control treatment consisted of same weight of fine pieces of filter papers. These pots were incubated at 25 °C in dark. All the pots were irrigated with 25 ml Hoagland nutrient solution (22). Germination was recorded after 72h and thereafter, the pots were shifted to partial shady condition (veranda). Seedling growth was noted after 15 days, which was sufficient time to show the differences in the test and control treatments.

3. Rain leachate bioassay

Natural rain leachates were collected from crushed leaves following Hussain *et al.* (8) during spring rains. The original rain leachates were dilute; therefore they were concentrated to 50% (highly concentrated) and 25% (less concentrated) of original concentration in rotavapor. These three concentrations (original, 50%, 25%) of rain leachates were tested against seeds and seedling growth of *L. sativa* and *A. cepa* in aqueous extract bioassay as above.

4. Effect on cell development (Cell size and cell area)

Plant extract from leaves and stem-barks, prepared by soaking 5 g materials in 100 ml distilled water for 48 h, were tested as before against seeds of *P. glaucum*, *S. italica*, *L. sativa* and

A. cepa in standard filter paper bioassay using distilled water as control. After 72 h incubation at 25 °C, the radicle tips were excised and placed in concentrated solution of chloral hydrate. After 10-12 hr, the tips were randomly taken out, placed on plain microscopic glass slide under cover slip and gently pressed to spread the material in a thin layer. The size of cells was measured between 3rd to 5th cortical layers in a row over fixed distance. Ten tips, each with 5 counts, were observed.

5. Effect on cell division

Radicle-tips from the preceding aqueous extract bioassay were saved; and fixed in Carnoy's fixer (mixture of absolute alcohol: glacial acetic acid, 3:1) for 12 h. These tips were then thoroughly washed with distilled water and saved in 70% alcohol at 5-10 °C. The meristematic tips were hydrolyzed in *IN* HCl at 60 °C for 10 to 15 min. HCl was thoroughly washed with distilled water and fixed in basic fuchsin in airtight vials in dark. Within 30 min the tips turned violet. A single root-tip was placed in 2-3 drops of aceto-carmin on a slide and slightly warmed. The cover slip was gently pressed to disperse the tissue. This temporary slide was used for counting dividing/non-dividing cells under microscope with 100x magnification (Olympus, XC-401 A, Shinjuku Monolith, Nish Shinjuku, Tokyo, Japan).

Table 1. Effect of aqueous extracts of *Calotropis procera* on the germination and various growth parameters of test species.

Test species	Control	Soaking time (h)	Extracts									
			Leaves	% of control	Stems	% of control	Control	Soaking Time (h)	Leaves	% of control	Stems	% of control
			Germination (%)				Fresh weight (mg)					
<i>Pennisetum glaucum</i>	67	24	12*	18.4	22	32.8	690	24	370	53.6*	330	47.8*
		48	15*	22.4	33	49.3	48	310	44.9*	320	46.4*	
<i>Setaria italica</i>	85	24	61*	71.8	74	87.0	960	24	400	41.7*	400	41.7*
		48	63*	74	68	80.0	48	400	41.7*	800	83.3*	
<i>Brassica campestris</i>	78	24	45*	57.7	60	76.9	450	24	350	77.8*	270	60.0*
		48	34*	43.6	50	64.0	48	270	60.0*	250	55.6*	
			Radicle Growth (mm)				Dry weight (mg)					
<i>Pennisetum glaucum</i>	23.2	24	12.2*	53.6	10.1*	43.5	180	24	170	94.4	170	94.4
		48	9.0**	38.8	12.0	51.7	48	130	72.2*	130	72.2*	
<i>Setaria italica</i>	10.7	24	6.4*	59.8	5.30	50	400	24	110	27.5	110	27.5
		48	4**	37.4	7.4**	69.2	48	100	25.0*	300	75.0	
<i>Brassica campestris</i>	13.9	24	6.4**	46.0	8.8	63.3	80	24	70	87.5	70	87.5
		48	2.1**	15.1	9.1	65.5	48	70	87.5	75	93.8	
			Plumule Growth (mm)				Moisture contents (%)					
<i>Pennisetum glaucum</i>	6.7	24	2.3**	34.3	2.3	34.3	283	24	117	41.3*	94	33.2*
		48	1.8**	26.9	1.34	19.4	48	146	51.6*	138	48.8*	
<i>Setaria italica</i>	8.5	24	1.80**	21.2	3.60**	42.4	425	24	207	48.7*	72	17*
		48	1.40**	16.5	2.10**	24.7	48	290	68.2*	33	7.8*	
<i>Brassica campestris</i>	7.8	24	2.60**	33.3	2.60**	33.3	462	24	337	72.9*	285	61.7*
		48	1.90**	24.4	2.00**	25.6	48	285	61.7*	212	45.9*	

Each value is a mean of 10 replicates, each with 10 seeds. *and ** Significantly different from control at $P = 0.05$ and at $P = 0.01$

Table 2. Effect of Hot water extracts on the germination (%) and various growth parameters of test species.

Test species	Control	Extracts								
		Leaves	% of control	Stems	% of control	Control	Leaves	% of control	Stems	% of control
Germination						Fresh weight (mg)				
<i>Pennisetum glaucum</i>	67	5**	75	13**	19.4	690	240	34.8*	310	44.9*
<i>Setaria italica</i>	85	62*	72.9	62*	72.9	900	480	53.3*	500	55.6*
<i>Brassica campestris</i>	78	43*	55.1	58*	74.4	450	330	73.3*	280	62.2*
<i>Lactuca sativa</i>	98	50*	51.0	62*	63.3	550	307	55.8*	377	68.6
Radicle Growth (mm)						Dry Weight (mg)				
<i>Pennisetum glaucum</i>	23.2	8.1*	34.9	9.0*	38.8	180	150	83.3	150	83.3
<i>Setaria italica</i>	10.7	5.5**	51.4	6.2**	57.9	400	170	42.5	180	45.0
<i>Brassica campestris</i>	13.9	4.4**	31.7	9.2**	66.2	80	70	87.5	70	87.5
<i>Lactuca sativa</i>	7.0	3.0	42.2	3.0	42.7	319	195	61.2	200	52.7
Plumule Growth (mm)						Moisture contents (%)				
<i>Pennisetum glaucum</i>	5.6	1.8**	32.1	1.5	26.7	283	60	21.5*	106*	37.5
<i>Setaria italica</i>	3.8	2**	52.6	2.2**	57.9	125	182	146*	117	93.6
<i>Brassica campestris</i>	3.1	2.3**	74.92	2.0**	64.5	462	371	80.3	300	64.9
<i>Lactuca sativa</i>	8.5	5.0	58.2	6.0	70.0	72.4	57.4	79.3	88.5	122.2

Each value is a mean of 10 replicates, each with 10 seeds. * and ** Significantly different from control at $P = 0.05$ and at $P = 0.01$

Results and Discussion

1. Aqueous extract bioassay

The aqueous extracts from leaves and stems of *C. procer* significantly inhibited the germination, radicle and plumule growth of all the test species in various treatments (Table 1). Extracts from leaves in this case were more inhibitory than stems. Higher soaking duration (48 h) was found to have a greater inhibitory effect than shorter duration (24 h). The fresh weight of all the test species declined under test condition, especially in leaves (Table 1). The dry weight of *S. italica* significantly declined in leaf and stem extracts, that of *B. campestris* slightly reduced in test condition. *P. glaucum* remained unaffected in 24 h extract (Table 1). Moisture contents of all the tested species declined in all the treatments. The 48 h extracts from leaves were strongly inhibitory than stem extracts. The present findings are in line with those of Raihan *et al.* (11), Bakhshayeshan-Agdam *et al.* (14) and Thakur *et al.* (18), who also observed similar allelopathic effects in their studies. Hussain & Ilahi (7) and Hussain *et al.* (8) reported that leaf extracts of *Cenchrus* and *Bothriochloa* strongly inhibited the germination than stem and root extracts. The findings regarding differential phytotoxicity agree with contemporary workers. For example, Raihan *et al.* (11) observed that extracts from bark and leaves of donor species differentially inhibited the test species. The response of wheat and cucumber towards extracts from *Amaranthus* varied (14). Li

et al. (15) concluded that various weed species responded differently to *Veronica* extracts. Similarly, the allelopathic inhibition depended upon the test species, parts of plant used for extraction, concentration and parameters measured as reported in many studies (12, 13, 17, 23-25). All these studies are in line with the present findings. Hot water extracts diminished the germination, radicle and plumule growth (Table 2), and fresh weight (Table 2) of tested species in all the treatments. The moisture contents of *P. glaucum* and *B. campestris* in leaf and stem extracts and that of *L. sativa* in leaf extract declined. The moisture contents of *S. italica* growing in leaf extract enhanced; but in stem extract there was no effect. The stem extract enhanced the moisture contents of *L. italica*. Hot water extracts from leaves exhibited strong inhibition than stem extracts. Lodhi & Nickell (20) also stated that hot water extracts from *Celtis* reduced gas exchange capacity and moisture contents of test seedlings. The results are also in line with those of Hussain & Ilahi (7) and Hussain *et al.*, (8, 21) who reported that hot water extracts from shoots of *Cenchrus* and *Bothriochloa* decreased germination and various growth parameters of tested seedlings.

2. Effect of litter

Addition of plant litter generally improves fertility and soil condition, but, litter from many plants intoxicates the immediate habitat by releasing water soluble phytotoxins. In the present case, in litter bed bioassay the added *Calotropis* litter

Table 3. Effect of *Calotropis procera* litter on the germination and growth parameters of test species.

Test species	Control	Litter								
		Leaves	% of control	Stems	% of control	Control	Leaves	% of control	Stems	% of control
Germination						Fresh weight (mg)				
<i>Pennisetum glaucum</i>	35	2*	5.7	7*	20	100	20	20*	30	30*
<i>Setaria italica</i>	31	16*	51.6	20*	64.5	100	40	40*	50	50*
<i>Brassica campestris</i>	43	30*	69.7	31.8*	74	100	51	51*	55	55*
<i>Lactuca sativa</i>	85	53*	62.4	57*	67	495	379	76.6	361	72.9
Radicle Growth (mm)						Dry Weight (mg)				
<i>Pennisetum glaucum</i>	21.4	7*	32.7	6.2*	29	50	10	20*	10	20*
<i>Setaria italica</i>	48.2	2.8*	5.8	9.7*	20.1	70	30	42.9*	40	57*
<i>Brassica campestris</i>	22	4.2*	19.0	6.2**	28.2	65	31	47.7*	43	66.2*
<i>Lactuca sativa</i>	5.9	2.1	35.8	3.0	47.5	392	350	89.3	310	79.8
Plumule Growth (mm)						Moisture contents (%)				
<i>Pennisetum glaucum</i>	4.7	1.5*	31.9	2.1*	44.7	100	100	100	200	200*
<i>Setaria italica</i>	23.8	7.7*	32.4	8.1*	34.0	42	33*	78.6	25	59.5*
<i>Brassica campestris</i>	17.2	1.2*	7.0	2.5**	14.5	53	64	120	27	50.9*
<i>Lactuca sativa</i>	6.8	5.6	82.8	5.7	84.6	26.3	8.3	31.6	16.5	62.6

Each value is a mean of 5 replicates, each with 10 seeds. * and ** Significantly different from control at $P = 0.05$ and at $P = 0.01$

suppressed the germination, radicle and plumule growth of various test species (Table 3). The fresh and dry weight of *P. glaucum*, *B. campestris*, *S. italica* and *L. sativa* reduced significantly in all the treatments (Table 3). The moisture contents of *S. italica* and *L. sativa* reduced in both the extracts and that of *B. campestris* declined in stem extracts; but gained moisture in leaf extracts. Moisture contents of *P. glaucum* remained unaffected by leaf extract but were enhanced by stem extract (Table 6). In pot experiment, the incorporated litter reduced the germination, height, fresh and dry weight of all the tested species (Table 4). The moisture contents of *S. italica* and *L. sativa* decreased in both the treatments; while the moisture contents of *P. glaucum* and *B. campestris* increased. However, Premathilake *et al.* (26) and Thakur *et al.* (27) observed that incorporated litter was not harmful to the test species in pot experiments. Kluthe *et al.* (28) reported phytotoxicity of litter that supports the present findings.

3. Rain leachate bioassay

The results of preceding experiments were further confirmed by the phytotoxicity of natural rain leachates against *L. italica* and *A. sativa* (Table 5). The original concentration of rain leachates was low to inhibit germination, plumule and radical growth; but 50% and 25% concentrated extracts significantly diminished germination and radical growth of both the test species. The order of inhibition was: 50% > 25% > original rain leachate concentration. The inhibitory effects were due to release of some phytochemicals from *Calotropis*. Although, we did not isolated phytochemicals in this study, yet many other studies have identified

phytochemicals responsible for allelopathy by *Calotropis*. Alkaloids, tannins, flavonoides, phenolic compounds, saponins, glycosides and many other phytochemicals have been reported (4, 6, 29-31). We expect that allelopathy is operative through these phytochemicals in the present case. In nature, litter soaked by rain or moist soil releases various phytochemicals similar to the present effort thereby intoxicating the immediate soil. The phytotoxicity of rain leachates was concentration dependent, species tested and parameters measured. The released phytotoxins accumulate to physiologically active concentration in the habitat that adversely affects the associated species, thereby supporting the present findings.

4. Effect on division and growth of cell

Allelochemicals cause death, blistering and growth inhibition of cell. Adverse biochemical changes and decreased chlorophyll contents within the plant body decreases uptake of water and nutrients. This leads to limited growth, biological, physiological and cytological functions of susceptible plants. It was demonstrated that leaf extracts not only significantly reduced cell division (Table 6), but also lessened the cell size and development. The dividing cells were 38%, 35.8%, 24.5% and 25.1% respectively in *P. glaucum*, *S. italica*, *B. campestris* and *A. cepa*. Similarly, the size of cell decreased in test condition. This is one of mechanisms for limiting radicle and shoot growth. The findings concur with those of Santosa *et al.* (32), Hussain *et al.* (33, 34) and Hussain & Ilahi (7), who affirmed that aqueous extracts of *Kielemeyra*, *Eragrostis* and *Cenchrus* reduced both the division and development of cells. Cheng *et al.* (35) stated that low concentration of

Table 4. Effect of added litter (Mulch) of *Calotropis procera* on the germination (%) and various growth parameters seedling of test species in pot experiment.

Test species	Control	Mulch	% of control
Germination (%)			
<i>Pennisetum glaucum</i>	26	11*	42.3
<i>Setaria italica</i>	13	2*	15.4
<i>Brassica campestris</i>	34	15*	44.1
<i>Lactuca sativa</i>	80	60*	75.0
Seedling height (mm)			
<i>Pennisetum glaucum</i>	111.5	77.3*	69.3
<i>Setaria italica</i>	87.8	59.8*	68.1
<i>Brassica campestris</i>	95.6	55**	57.5
<i>Lactuca sativa</i>	5	4.2	84.0
Fresh weight (mg)			
<i>Pennisetum glaucum</i>	700	500*	71.4
<i>Setaria italica</i>	200	100*	50
<i>Brassica campestris</i>	700	100*	14.3
<i>Lactuca sativa</i>	550	350	63.6
Dry weight (mg)			
<i>Pennisetum glaucum</i>	500	300	60
<i>Setaria italica</i>	100	55	55
<i>Brassica campestris</i>	590	70	11.86
<i>Lactuca sativa</i>	290	233	80.3
Moisture contents (%)			
<i>Pennisetum glaucum</i>	40	66.67*	151.5
<i>Setaria italica</i>	100	81.82*	81.8
<i>Brassica campestris</i>	18.6	42.86*	230.6
<i>Lactuca sativa</i>	89.7	50.2*	56.0

Germination is a mean of 5 replicates, each with 10 seeds; others mean of 5 replicates each with 2 seedlings.

*Significantly different from control at $P = 0.05$

allelochemicals from garlic promoted the cell length but higher concentration reduced these activities. Likewise, Chaudhuri *et al.* (36) observed reduced nuclear volume and increased interphase chromatin material. Raoof & Siddiqui (37), Santosa *et al.*, (32) and Irum *et al.* (38) stated that various phytochemicals causing allelopathic stress reduce growth and productivity. The decreased cell division and growth by allelopathic stress is reported in many studies (23, 39-43). However, Fonseca *et al.* (44) reported no effect on mitotic

index and cell division by *Schinus* leaf extracts. The presence of various phytochemicals (3, 6, 45) in *Calotropis procera* can be responsible for its observed allelopathy. The findings of Lubini *et al.* (46), Sharma *et al.* (47) and Abdelmigid & Morsi (48) support the present results with respect to inhibition of cell division.

The present findings conclude that *Calotropis procera* is strongly allelopathic at least against the tested species. The allelopathic effects depended upon the parts of *Calotropis* used in

Table 5. Effect of rain leachates from *Calotropis procera* leaves on the germination and radicle growth of *Lactuca sativa* and *Allium cepa*.

Species	Parameter	Control	Original low concentration rain leachates	25% concentrated rain leachates	50% (Strongly) concentrated rain leachates
<i>Lactuca sativa</i>	Germination	100	90 (90%)	75 (75%)	45 (45%)
	Radicle Growth	25.9	20.2 (77.99%)	15.35 (59.27%)	10.92 (42.16)
<i>Allium cepa</i>	Germination	86	79 (91.86%)	62 (72.09%)	49 (55.06%)
	Radicle Growth	5.38	3.94 (73.23)	2.01 (37.36%)	1.28 (23.79%)

Table 6. Effect of *Calotropis procera* leaf extract on the division and size of cells of test species.

Species	No. of Dividing cells in Control	No. of Dividing cells in Lest extract	% of Control
<i>Pennisetum glaucum</i>	250	95*	38.0
<i>Setaria italica</i>	279	100*	35.8
<i>Brassica campestris</i>	200	49*	24.5
<i>Allium cepa</i>	311	78*	25.1
Cell Sizes, μ			
<i>Pennisetum glaucum</i>	26.2	16.92*	64.58
<i>Setaria italica</i>	17.5	9.22*	52.69
<i>Brassica campestris</i>	22.9	8.25*	36.03
<i>Allium cepa</i>	30.0	12.92*	43.07

bioassays, the response of receptor (test) species and the growth parameter determined. It was observed that germination, seedling growth, fresh and dry mass and moisture contents of various test species responded independently. Further studies are needed to identify the possible allelochemicals and search for their use as alternative herbicides and insecticides.

Acknowledgements

The authors are grateful to the Lab Staff of Department of Botany, University of Peshawar for their assistance.

Authors' contributions

All the authors contributed equally to the work presented in this paper.

Competing Interests

The authors declared that they have no conflict of interest.

References

- Samreen U, Hussain F, Sher Z. Allelopathic potential of *Calotropis procera* (Ait.) Ait. Pak. J Pl Sci 2009;15(1):7-14
- Abdul-Farid I, Magdi El-Syed, Eman M. Allelopathic potential of *Calotropis procera* and *Morettia philaeana*. Int J Agric & Biol 2013;15(1):130
- Akindede PO, Fatunla OA, Ibrahim KA, Afolayan CO. Antibacterial and phytochemical screening of *Calotropis procera* leaf extracts against vancomycin and methicillin resistant bacteria isolated from wound samples in hospital patients. Jour Complem & Alternative Med Res 2017; 2(1):1-14. <https://doi.org/10.9734/JOCAMR/2017/30975>
- Umar S, Shuaib NH, Dangore II, Hassan KY. Evaluation of allelopathic effects of *Calotropis procera* on sorghum and maize. Standard Sci Res Essays 2014; 2(8):345-49.
- Shetta ND, Alshahrani TS, Aref IM, Nasser R.A. Allelopathic potential of *Calotropis procera* and *Eucalyptus* species on germination and growth of some timber trees. Allelopathy Journal 2017; 40(1):81-94. <https://doi.org/10.26651/2017-40-1-1068>
- Hilal-Ul-Zaman, Ahmad S. Antibacterial activity and phytochemical analysis of leaf extracts of *Calotropis procera*. Acta Scientific Pharmaceu Sci 2017; 1.5 (2017):19-21.
- Hussain F, Ilahi I. Allelopathic potential of *Cenchrus ciliaris* L. and *Bothriochloa pertusa* (L) A. Camus. J Sci & Technol 2009; 33(1&2): 47-55
- Hussain F, Ilahi I, Malik SA, Dasti AA, Ahmad B. Allelopathic effects of rain leachates and root exudates of *Cenchrus ciliaris* L and *Bothriochloa pertusa* (L) A. Camus. Pak J Bot 2011; 43(1): 341-50.
- Saxena MK. Aqueous leachate of *Lantana camara* kills water hyacinth. J. Chem. Ecol 2000; 26(10):2435-48. <https://doi.org/10.1023/A:1005539230307>
- Ahmad I, Hussain F, Barkatullah, Ahmad B. Phytotoxic potential of *Celtis australis* L. (Family Ulmaceae) against four crop species. Pak J Bot 2014;46(6):2063-67.
- Raihan I, Miiyaural R, Baki BB, Fujii Y. Assessment of allelopathic potential of goniothalamine allelochemicals from Malaysian plant *Goniothalamus andersonii* J. Snclair by sandwich method. Allelopathy Journal 2019;46(1):25-40. <https://doi.org/10.26651/allelo.j/2019-46-1-1196>
- Anwar T, Ilyas N, Qureshi R, Malik M. A. Allelopathic potential of *Carica papaya* against selected weeds of wheat crop. Pak J Bot 2019; 51(1):279-87. [https://doi.org/10.30848/PJB2019-1\(37\)](https://doi.org/10.30848/PJB2019-1(37))
- Qasem JR. A survey of the phytotoxicity of common weeds, wild grown species and medicinal plants on wheat. Allelopathy Journal 2017;42(2):179-94 <https://doi.org/10.26651/allelo.j/2017-42-2-1115>
- Bakhshayeshan-Agdam H, Salehi-Lisar SY, Motafakkerazad R. Allelopathic effects of redroot pigweed (*Amaranthus retroflexus* L) aqueous extract on cucumber and wheat. Allelopathy Journal 2019; 46(1):55-72. <https://doi.org/10.26651/allelo.j/2019-46-1-1198>
- Li ZR, LiuYB, Zhou XM, Li XG, Bai LY. Allelopathic herbicidal effects of crude ethanolic extracts of *Veronica persica* (Lour) Merr. on weeds. Allelopathy Journal 2019; 46(1):85-96. <https://doi.org/10.26651/allelo.j/2019-46-1-1200>
- Wu H, Lin JM, Zhang JB. Allelopathic effects of *Eucalyptus salubris* F. Muell. and *E. brockwayii* C.A. Gardner on germination and seedling growth of prairie ground cherry (*Physalis hederifolia* A. Gray). Allelopathy Journal 2019; 46(1):109-20. <https://doi.org/10.26651/allelo.j/2019-46-1-1202>
- Tahir NA, Azeez HA, Hama-Amin HH, Rashid JS, Omer DA. Antibacterial activity and allelopathic effects of extracts from leaf, stem and bark of Mt. Atlas mastic tree (*Pistacia atlantica* subsp *kurdica*) on crops and weeds. Allelopathy Journal 2019; 46(1):121-32. <https://doi.org/10.26651/allelo.j/2019-46-1-1203>
- Thakur NS, Kumar D, Chauhan RS, Hedge HT, Gunaga RP. Allelopathic effects of *Melia azedarach* L on germination, growth and yield of black gram and chickpea. Allelopathy Journal 2019; 46(1):133-44. <https://doi.org/10.26651/allelo.j/2019-46-1-1204>
- Hussain F. Field and Laboratory Manual of Plant Ecology. 1989;University Grants Commission, Islamabad.
- Lodhi MAK, Nickell GI. Effects of leaf litter of *Celtis laevigata* on growth, water contents, and carbon dioxide exchange rates of three grass species. Bull Torrey Bot Club 1973; 100:59-165. <https://doi.org/10.2307/2484627>
- Hussain F, Ahmad B, Illahi I. Allelopathic effects of *Cenchrus ciliaris* L and *Bothriochloa pertusa* (L) A. Camus. Pak. J. Bot 2010; 42(5): 3587-3604.
- Hoagland DR, Arnon DI. The water-culture method of growing plants without soil. Calif Agr Expt Sta Circ 347. 1950.
- Bassey CS, Okoi EP. Comparative evaluation of the allelopathic effects of the leaf extracts of three Asteraceae species (*Ageratum conyzoides*, *Vernonia amygdalina*, *Artemisia annua*). Archives of Current Research International 2018; 15(3):1-8. <https://doi.org/10.9734/ACRI/2018/45397>
- Gulzar A, Siddiqui MB. Allelopathic effects of *Calotropis procera* (Ait.) R. Br. on growth and antioxidant activity of *Brassica oleracea* var *botrytis*. Jour Saudi Soc Agric Science 2017; 16:375-82. <https://doi.org/10.1016/j.jssas.2015.12.003>
- Manzoor SA, Zubair M, Hussain SB, Akhtar SNR, Nouman W. Evaluation of allelopathic effects of *Calotropis procera* against wheat (*Triticum aestivum*) using seedling vigor and germination indices. J Biodiver & Environ Sci 2013;3(5):25-30.
- Premathilake UGAT, Wathugalal DL, Dharmadasa RM. Phytotoxic potential of lemon grass (*Cymbopogon citrates* (DC) Stapf. Allelopathy Journal 2018; 45(2):261-68. <https://doi.org/10.26651/allelo.j/2018-44-2-1168>
- Thakur NS, Parmar AG, Gunaga RP, Patel DP. Allelopathic influence of leaf and leaf litter of white cedar (*Melia azedarach* L.) on eggplant and okra. Allelopathy Journal 2017; 42(2):295-306. <https://doi.org/10.26651/allelo.j/2017-42-2-1124>

28. Kluthe BG, Ali MBHB, Stephenson SL. Allelopathic influence of *Eucalyptus* on common Kenyan agricultural crops. *J Agron Agric Sci* 2018;1(1):2-6.
29. Chandrawat P, Sharma RA. The genus *Calotropis*: An overview on bioactive principles and their biosafety. *Res J Recent Sci* 2016;5(1):61-70.
30. Gulzar A, Siddiqui MB, Bi S. Phenolic acid allelochemicals induced morphological, ultrastructural, and cytological modification on *Cassia sophora* L. and *Allium cepa* L. *Protoplasma* 2016;253(5):1211-21. <https://doi.org/10.1007/s00709-015-0862-x>
31. Waheed N, Jabeen K, Iqbal S, Javid A. Biopesticidal activity of *Calotropis procera* against *Macrophomina phaseolina*. *African J Tradit Complement & Altern Med* 2016;13(6):163-167. <https://doi.org/10.21010/ajtcam.v13i6.23>
32. Santos VHMde, Daneluzzi GS, Silva LP, da Silva RMG. Evaluation of allelopathic potential of leaf extracts of *Kielmeyera coriacea* on *Lactuca sativa* L. *Biosci J*.2015; 31(1): 259-67. <https://doi.org/10.14393/BJ-v31n1a2015-17662>
33. Hussain F, Zaidi MI, Chughtai SR. Allelopathic effects of Pakistani Weeds: *Eragrostis poaeoides* P. Beauv. *Pak J Sci Ind Res* 1984;27:159-64.
34. Hussain F, Abidi N, Malik ZH. *Imperata cylindrica* affects germination, early growth and cell division and development in some crop species. *Pak J Sci Ind Res* 1994; 37:100-03.
35. Cheng F, Cheng Z, Meng H, Tang X. The garlic allelochemicals diallyl disulfide affects tomato root growth by influencing cell division, phytohormone balance and expansion gene expression. *Front Plant Sci* 2016;7:1199. <https://doi.org/10.3389/fpls.2016.01199>
36. Chaudhuri A, Kundu LM, Datta S, Chatterjee S, Goswami S, Roy GC et al. Allelopathic effects of aerial parts aqueous extract of *Amelocissus latifolia* (Roxb) Planch. in apical meristem cells. *Asian J. Plant Sci. Res* 2015;5(3):11-16.
37. Raoof KM, Siddiqui MB. Allelotoxic effect of parthenin on cytomorphology of broad bean (*Vicia faba* L.). *Jour Saudi Soc Agric Sci* 2013; 12:143-46. <https://doi.org/10.1016/j.jssas.2012.11.001>
38. Irum A, Tabassum S, Qureshi R, Gulfraz M, Anwar P. Phytochemical analysis of medicinally important constituents of *Teucrium stocksianum* Boiss. *Pak J Bot* 2019; 51(3): 1067-74. [https://doi.org/10.30848/PJB2019-3\(17\)](https://doi.org/10.30848/PJB2019-3(17))
39. Cruz-Ortega R, Anaya GAL, Ramos L. Effects of allelopathic compounds of corn pollen on respiration and cell division of water melon. *J ChemEcol*1988; 14(1):71-86. <https://doi.org/10.1007/BF01022532>
40. Rai PD, Paudel N, Shakya S. R. Cytological effects of leaf extract of *Artemisia vulgaris* L. on meristematic cells of *Allium cepa* L. *Our Nature* 2012;10: 242-48. <https://doi.org/10.3126/on.v10i1.7787>
41. Mohammad FI, El-Ashry ZM. Cytogenetic effect of allelochemicals *Brassica nigra* L. extracts on *Pisum sativum* L. *World Appl Sci. Journal* 2012;20(3):344-53.
42. Talukdar D. Allelopathic effects of *Lantana camara* L. on *Lathyrus sativus* L.: Oxidative imbalance and cytogenetic consequences. *Allelopathy Journal* 2013;31 (1):71-90.
43. FrescuraVDal-Souto, Kuhn AW, Laughinghouse HD, Nicoloso FT, Lopes SJ, Tedesco SB. Evaluation of the allelopathic, genotoxic, and antiproliferative effect of the medicinal species *Psychotria brachypoda* and *Psychotria birotula* (Rubiaceae) on the germination and cell division of *Eruca sativa* (Brassicaceae). *Caryologia* 2013; 66:138-44. <https://doi.org/10.1080/00087114.2013.821832>
44. Fonseca VB, Tavares VRdeS, Goncalves VdM, Fritag RA, Bobrowski VL. Allelopathic potential of leaves and flowers extracts of *Schinus terebinthifolius* Raddi. *Ceintifica* 2016; 44(1):35-39. <https://doi.org/10.15361/1984-5529.2016v44n1p35-39>
45. Ayeni MJ, Akinyede OA. Effects of *Calotropis procera* (Ait.) R.Br. leaves on the germination and early growth of soybeans (*Glycine max* L.). *J Agric & Vet Sci* 2014; 7(4): 5-9. <https://doi.org/10.9790/2380-07420509>
46. Lubini G, Fachinnetto JM, Laughinghouse HD, Paranhos JT, Silva ACF, Tedesco SB. Extracts affecting mitotic division in root-tip meristematic cells. *Biologia*2008; 63(5): 647-651. <https://doi.org/10.2478/s11756-008-0108-x>
47. Sharma M, Tandon S, Agarwal V, Bhat KG, Kappadi D, Chandrashekhar P et al. Evaluation of antibacterial activity of *Calotropis gigantea* against *Streptococcus mutans* and *Lactobacillus acidophilus*: an in vitro comparative study. *J Conserv Dentistry* 2015; 18(1):457-460. <https://doi.org/10.4103/0972-0707.168809>
48. Abdelmigid HA, Morsi MM. Cytotoxic and molecular aspects of allelopathic effects of leaf residues of *Eucalyptus globulus* on soyabean (*Glycine max*). *J. Eng & Biotechnol* 2017; 15:297-302. <https://doi.org/10.1016/j.jgeb.2017.08.005>

