



RESEARCH ARTICLE

Effects of pH, temperature, water activity and electrical conductivity on the biocontrol activity of *Trichoderma* spp. for enhanced biocontrol efficacy in agriculture

Navneet Singh¹, Vipul Kumar², Manoj Kumar Maurya^{3*}, Gurveer Singh⁴, D K Upadhyay³, Anil Kumar³, A K Singh³ & P K Dalal³

¹Division of Vegetable Science, Faculty of Horticulture, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar 190 025, Jammu and Kashmir, India

²Department of Plant Pathology, School of Agriculture, Lovely Professional University, Phagwara 144 411, Punjab, India

³Department of Vegetable Science, Acharya Narendra Dev University of Agriculture &Technology Kumarganj, Ayodhya 224 229, UttarPradesh, India

⁴Department of Agriculture and Farmer welfare Punjab 141 004, India

*Correspondence email - drmanoj.pathovs@nduat.org

Received: 26 July 2025; Accepted: 27 November 2025; Available online: Version 1.0: 18 December 2025; Version 2.0: 01 January 2026

Cite this article: Navneet S, Vipul K, Manoj KM, Gurveer S, Upadhyay DK, Anil K, Singh AK, Dalal PK. Effects of pH, temperature, water activity and electrical conductivity on the biocontrol activity of *Trichoderma* spp. for enhanced biocontrol efficacy in agriculture. Plant Science Today. 2026; 13(1): 7. <https://doi.org/10.14719/pst.10869>

Abstract

Trichoderma spp. is widely recognized and employed as effective biocontrol agents, providing a natural approach to managing plant diseases. These fungi, commonly found in soil, utilize multiple mechanisms to suppress plant pathogens, making them important tools in sustainable agriculture. However, the biocontrol efficacy of *Trichoderma* spp. is influenced by several environmental factors that impact their growth, sporulation and antagonistic activity against phytopathogens. This research investigates how abiotic factors such as pH, temperature, water activity (a_w) and electrical conductivity (EC) affect the efficacy of *Trichoderma* spp. as biocontrol agents against plant pathogens. Controlled laboratory tests were evaluated for conidial production, growth rates and morphological changes under varying environmental conditions. The results indicated that the ideal water activity (a_w 0.985) and temperature (25 °C) considerably increased conidial production and biomass growth. A pH level of 6.5 and an EC of 12.92 dS/m were identified as ideal conditions for optimal growth and sporulation. These results underscore the importance of environmental factors in improving the effectiveness of *Trichoderma* spp., offering actionable suggestions to enhance its use as a biocontrol agent across diverse farming systems, while also recognizing the importance of these factors in improving the large-scale production of *T. harzianum* for both agricultural and industrial purposes, especially in biocontrol application.

Keywords: abiotic factor; biocontrol; electrical conductivity (EC); pH; temperature; *Trichoderma* spp.; water activity (a_w)

Introduction

Trichoderma is a well-known soil-dwelling saprophytic fungus frequently found in environments such as soil, sand and water (1). It is widely recognized fungi that enhance plant growth, regulate various physiological processes and provide protection against numerous plant pathogens (2). It serves as an effective biocontrol agent (BCA) capable of antagonizing several soil-borne plant pathogens, including *Fusarium oxysporum*, *Rhizoctonia* spp. and *Pythium* spp (3). Several studies have shown that *Trichoderma* species employ a wide range of mechanisms for effective biological control (4, 5) such as competing for nutrients and space (6), direct antagonism (7), parasitism (8), antibiosis (9, 10) and the production of hydrolytic enzymes like β-1,3 glucanase, chitinase, protease and cellulase (11). However, its growth and reproduction are significantly influenced by factors like culture age, temperature, EC, water activity (a_w) and pH, which collectively determine the quality and quantity of fungal colonies (12).

Although various studies have explored its application methods and efficacy in lab, greenhouse and field conditions (13), limited research exists on how *Trichoderma* spp. survives, grows and responds to stress factors such as EC, a_w and pH. Abiotic factors, including temperature (12), pH, a_w and EC, significantly impact its reproduction, colony morphology and proliferation (1). Water, a vital component for cellular activities, is critical for enzymatic function, stability and survival. Water activity, representing the amount of water available in a substrate or culture medium, directly affects microbial growth and cellular processes when suboptimal. Additionally, specific water requirements for fungal growth and metabolism vary among species.

Trichoderma can grow over a broad pH range due to its adaptive capabilities; favourable pH conditions promote the expression of genes responsible for metabolite biosynthesis and secretion. Soil moisture influences EC, reflecting dissolved ion concentration, which in turn affects cell wall functionality and enzyme activity (14). Studies indicate *Trichoderma* thrives in soils

with a moisture content of 40 - 80 %, with optimal enzyme activity observed at 50 -70 % moisture (15). However, low water availability can reduce conidial production, impacting its biocontrol efficacy. *Trichoderma* is sensitive to extreme salt and pH conditions, with optimal growth and sporulation at pH 6.0. Deviations to higher pH levels lead to reduced growth, sporulation and colony diameter (16).

This study aims to investigate the impact of abiotic factors, including a_w , temperature, EC and pH, on *Trichoderma* spp. growth. Recent findings emphasize the critical role of temperature in determining BCA survival and effectiveness. For instance, temperature variations significantly affect *Trichoderma*'s ability to control *Botrytis cinerea* in grape berries and influence its elicitation of plant responses in tomato plants (17, 18). Furthermore, understanding *Trichoderma*'s specific requirements for a_w and pH is crucial for enhancing its application in biocontrol strategies, ensuring its resilience and efficacy under diverse environmental conditions.

Materials and Methods

This study was conducted in the laboratory of Lovely Professional University, Phagwara, Punjab, India (31.2554° N, 75.7049° E). The experiments included fungal inoculation, incubation, conidia counting and the measurement of EC and pH.

Water activity (a_w) and temperature

The water activity (a_w) of potato dextrose agar (PDA) was adjusted by adding glycerol at concentrations of 5 g/L, 10 g/L, 50 g/L, 100 g/L and 175 g/L, corresponding to a_w levels of 0.998, 0.995, 0.985, 0.977 and 0.961, respectively. The a_w was measured using a Dew point meter (Applied Techno Systems, Mumbai, India). The adjusted water solutions were mixed with nutrient agar and poured into plates, which were inoculated with *Trichoderma harzianum* under aseptic conditions within a laminar airflow cabinet. These plates were incubated at 25 °C in BOD incubator (15). Independent experiments for temperature (15 °C, 25 °C and 35 °C) and a_w were set up using a completely randomized design (CRD), with 3 replicates for each condition.

pH

To determine the optimal pH for the growth of *Trichoderma* spp., media with pH levels of 3.5, 4.5, 5.5, 6.5, 7.5 and 8.5 were prepared by adding appropriate amounts of HCl or KOH before sterilization. The pH of the medium was measured with a digital pH meter (Labline Stock Centre, Mumbai, India) before and after fungal inoculation. Plates inoculated with *T. harzianum* were incubated at 28 ± 2 °C for 15 days and fungal growth and sporulation were observed. Growth rate was measured using a scale (19).

Electrical conductivity (EC)

To assess the impact of EC on the growth and sporulation of *T. harzianum*, NaCl was added to the medium at 3 concentrations (g/l): 1 g (EC: 12.92 dS/m), 2 g (EC: 14.30 dS/m) and 5 g (EC: 17.8 dS/m). The EC of the medium was measured using a pH meter (19). The agar medium was sterilized at 121 °C and 1.5 kg/cm² for 15 min before being poured into petri dishes. Mycelium discs (7 mm) of *T. harzianum* were placed at the center of each plate, with 3 replicates for each treatment. Plates were incubated at 28 ± 2°C for 2 weeks and fungal growth and sporulation were monitored daily. Data were analysed statistically using a CRD.

Results

Effect of water activity (a_w) on conidial production and germination of *T. harzianum*

The conidial production and germination of *T. harzianum* were significantly influenced by different water activity (a_w) levels (Table 1). The highest conidial production (923×10^6 conidia/mL) was observed at an a_w of 0.985, with an average germination rate of 68 %. At an a_w of 0.998, conidial production decreased to 600×10^6 conidia/mL, but the germination rate was higher at 82 %. Similarly, an a_w of 0.995 resulted in 525×10^6 conidia/mL with a 76 % germination rate. Lower water activity levels severely reduced both conidial production and germination. At an a_w of 0.977, conidial production dropped to 347×10^6 conidia/mL with a germination rate of only 21 %. The data clearly show that fungal reproductive activity is significantly inhibited by reducing water activity (a_w). There was barely any germination and very little conidial production at the lowest a_w measured (0.961). These results highlight the critical role of water activity in promoting optimal conidial production and germination of *T. harzianum*. Moderate a_w levels (0.985-0.995) appeared most favourable for fungal growth and reproductive capacity, while extremely low a_w levels significantly inhibit these processes (Fig. 1 & Fig. 2).

Effect of temperature on mycelial biomass production of *Trichoderma* spp.

All tested *Trichoderma* species produced substantial biomass across a range of temperatures. *T. harzianum* produced the highest biomass of 1.42 g when incubated at 25 °C, compared to 0.87 g and 0.73 g at 15 °C and 35 °C, respectively presented in (Table 2) and indicated in (Fig. 3).

Effect of pH on conidial production of *Trichoderma* spp.

The production of conidia by *T. harzianum* was notably influenced by varying pH levels, as indicated by alterations in colony morphology (Table 3). The maximum conidia count was reached at pH 6.5. Conversely, there was a significant decrease in production at pH 5.5 and an even lower yield at pH 8.5. No substantial difference in conidia numbers was observed between pH 3.5 and 4.5. A consistent reduction in conidial production was noted as pH levels rose above the optimal range, with significant drops at pH 7.5 and 8.5 (Fig. 4).

Table 1. Effect of different a_w on conidial production and germination of *Trichoderma* spp

Treatment (Water activity)	Number of conidia ($\times 10^6$ conidia/mL)	Average germination %
a_w 0.985	923	68
a_w 0.998	600	82
a_w 0.995	525	76
a_w 0.977	347	21
a_w 0.961	0.04	N.D*

*Not determine

Table 2. Effect of temperature level on mycelial biomass production of *Trichoderma* spp.

Treatments (Temperature °C)	Weight of biomass (g)
15	0.87
25	1.42
35	0.75



S.no.	Treatments	Number of conidia ($\times 10^6$ conidia/mL)	Pictures
1.	$a_w = 0.985$	923	
2.	$a_w = 0.998$	600	
3.	$a_w = 0.995$	525	
4.	$a_w = 0.977$	347	
5.	$a_w = 0.961$	0.04	

Fig. 1. Effect of a_w on the morphological characteristics and conidial production of *Trichoderma harzianum*.




Fig. 2. Means effect for conidium germination and conidia production of *T. harzianum* grown in cultures at different a_w .




Fig. 3. The effect of temperature on the growth of *T. harzianum*.




Fig. 4. Main effect means for conidium production of *T. harzianum* at different pH levels.

Table 3. Effect of different pH levels on conidial production of *Trichoderma* spp.

Treatment (pH)	Number of conidia ($\times 10^6$ conidia/mL)
3.5	1010
4.5	1041
5.5	663
6.5	1293
7.5	197
8.5	98

Effect of electrical conductivity (EC)

The results indicated significant differences in the effects of 3 levels of EC 12.92, 14.30 and 17.8 ds/m with 3 replications (R-1, R-2 and R-3) of each, on the growth and sporulation of *T. harzianum*. The result showed optimal growth at an EC of 12.92 ds/m presented in (Fig. 5).

Discussion

The results of the present study provide valuable insights into the influence of various environmental factors, such as water activity (a_w), temperature, pH and electrical conductivity (EC), on the growth and reproductive capacity of *Trichoderma harzianum*. These factors are critical in optimizing the conditions for conidial production and germination, which are important for both research and industrial applications, such as biocontrol agent formulation.

Water activity (a_w) has a profound impact on the conidial production and germination of *T. harzianum*. The highest conidial production (923×10^6 conidia/mL) was observed at an a_w of 0.985, with a germination rate of 68 %. These results are consistent with findings that moderate water activity levels significantly enhance fungal growth and reproduction (20). The reduced conidial production observed at higher a_w levels (0.998) with an increase in germination (82 %) suggests that while high water availability favors germination, it may inhibit sporulation due to potential alterations in metabolic activity or osmotic stress (21).

Conversely, at lower water activity levels, the study demonstrates a sharp decline in both conidial production and germination. At an a_w of 0.977, conidial production was reduced to 347×10^6 conidia/mL with a germination rate of only 21 %, which highlights the critical need for sufficient water availability for optimal fungal development. The drastic reduction in conidial production at the lowest a_w (0.961) was consistent with findings where very low water activity inhibited fungal spore production and viability (22).

Overall, these findings underscore the importance of maintaining moderate water activity levels (0.985 - 0.995) for fostering favorable conditions for *T. harzianum* growth and conidial production, which agrees with previous studies.

Temperature is a key factor influencing the mycelial growth of *Trichoderma* species. In this study, *T. harzianum* produced the highest biomass (1.42 g) at 25 °C, with a significant reduction in biomass at both lower (15 °C) and higher (35 °C) temperatures. This result is in line with studies that demonstrate the optimal growth of *Trichoderma* species at moderate temperatures. The 25 °C is typically the ideal temperature for biomass production in *Trichoderma*, as extreme temperatures may stress the fungal cells, leading to reduced metabolic activity and growth (23).

The reduced biomass production at 15 °C and 35 °C in our study is consistent with the notion that *T. harzianum* has a relatively narrow optimal temperature range for robust growth. Enzyme kinetics are probably greatly retarded at the lower temperature of 15 °C. On the other hand, the initiation of thermal stress at the higher temperature of 35 °C is responsible for the observed decrease. Important proteins and enzymes may become denatured because of high temperatures. When taken as a whole, these physiological reactions highlight the fact that *T. harzianum* grows best in a more limited temperature range, with variations affecting essential cellular functions needed for strong growth, with optimal growth occurring at moderate temperatures. The results also suggest that both lower and higher temperatures could impair the energy allocation toward mycelial growth, possibly diverting resources to stress responses rather than vegetative growth.




Fig. 5. The influence of EC on *T. harzianum* growth.

The impact of pH on conidial production in *T. harzianum* was significant, with the highest conidial production occurring at pH 6.5. These studies are supported by similar findings, which demonstrated that *Trichoderma* species exhibit optimal growth and sporulation under slightly acidic to neutral conditions (24, 25). At pH values below 6.5, particularly at pH 3.5 and 4.5, conidial production decreased sharply, which may be due to the acidic stress that can impair cellular functions and inhibit proper metabolic activity (26).

The decline in conidial production at pH 7.5 and 8.5 in this study corroborates the findings that alkaline conditions can negatively affect fungal growth and conidial production in *Trichoderma* species (27). The optimal pH for *T. harzianum* conidial production in this study, around 6.5, suggests that slightly acidic environments are most conducive to the formation of viable spores and the overall health of the fungus.

The EC of the medium also played a crucial role in the growth and sporulation of *T. harzianum*. The highest biomass and optimal sporulation were observed at an EC level of 12.92 ds/m. This result suggests that moderate salinity levels favor fungal growth, which could be attributed to the osmotic balance maintained within the fungal cells at this conductivity level (28).

At higher EC levels (14.30 and 17.8 ds/m), the study observed reduced growth and sporulation, likely due to osmotic stress caused by high salt concentrations, which can disrupt cellular integrity and metabolic processes. The optimal EC for *T. harzianum* growth observed in this study is in agreement with earlier findings that indicate moderate salinity levels promote fungal metabolism, while high salinity can cause cellular dehydration and inhibit fungal reproduction (29).

Conclusion

The present study highlights the significant role of environmental factors such as water activity, temperature, pH and electrical conductivity in influencing the growth, conidial production and germination of *T. harzianum*. The results suggested that optimal conditions for conidial production and germination lie within moderate ranges of water activity (0.985-0.995), temperature (25 °C), pH (6.5) and EC (12.92 ds/m). A strong scientific foundation for refining large-scale production procedures and guaranteeing the stability and effectiveness of bio fungicide formulations is provided by these exact characteristics, which are essential for optimizing conidia yield and viability. Manufacturers can improve sporulation efficiency, product shelf-life and ultimately the field performance of *Trichoderma*-based biocontrol agents by preserving these conditions during fermentation and product development for industrial and agricultural applications.

Acknowledgements

The authors are highly grateful and thankful to Department of Plant Pathology, School of Agriculture, Lovely Professional University Phagwara, Punjab, India for providing necessary facilities.

Authors' contributions

NS designing of experiments, carried out the field experiment and drafted the manuscript. VK conceptualization of research. MKM analysis of the data and interpretation. GS, DKU, AK, preparation of the final manuscript. AKS, PKD conceived of the study and reviewed the manuscript data. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: The authors declare that there is no conflict of interest.

Ethical issues: None

References

1. Purwanto B. Effect of water content on conidia of *Trichoderma* spp., indole acetic acid content, electrical conductivity and pH. *Biodiversitas J Biol Diversity*. 2022;23(5). <https://doi.org/10.13057/biodiv/d230536>
2. Martinez Y, Ribera J, Schwarze FWMR, De France K. Biotechnological development of *Trichoderma*-based formulations for biological control. *Appl Microbiol Biotechnol*. 2023;107:5595-612. <https://doi.org/10.1007/s00253-023-12687-x>
3. Harman GE. Overview of mechanisms and uses of *Trichoderma* spp. *Phytopathology*. 2006;96:190-4. <https://doi.org/10.1094/PHYTO-96-0190>
4. Manzar N, Kashyap AS, Goutam RS, Rajawat MVS, Sharma PK, Sharma SK, et al. *Trichoderma*: advent of versatile biocontrol agent, its secrets and insights into mechanism of biocontrol potential. *Sustainability*. 2022;14:12786. <https://doi.org/10.3390/su141912786>
5. Tyśkiewicz R, Nowak A, Ozimek E, Jaroszuk-Ścisiel J. *Trichoderma*: the current status of its application in agriculture for the biocontrol of fungal phytopathogens and stimulation of plant growth. *Int J Mol Sci*. 2022;23(4):2329. <https://doi.org/10.3390/ijms23042329>
6. Verma M. Antagonistic fungi, *Trichoderma* spp.: panoply of biological control. *Biochem Eng J*. 2007;37(1):1-20. <https://doi.org/10.1016/j.bej.2007.05.012>
7. Sui L, Li J, Philp J, Yang K, Wei Y, Li H, et al. *Trichoderma atroviride* seed dressing influenced the fungal community and pathogenic fungi in the wheat rhizosphere. *Sci Rep*. 2022;12(1):9677. <https://doi.org/10.1038/s41598-022-13669-1>
8. Li J, Philp J, Li J, Wei Y, Li H, Yang K, Wang Y. *Trichoderma harzianum* inoculation reduces the incidence of clubroot disease in Chinese cabbage by regulating the rhizosphere microbial community. *Microorganisms*. 2020;8(9):1325. <https://doi.org/10.3390/microorganisms8091325>
9. Shobha B, Lakshmeesha TR, Ansari MA, Almatroudi A, Alzohairy MA, Basavaraju S, et al. Mycosynthesis of ZnO nanoparticles using *Trichoderma* spp. isolated from rhizosphere soils and its synergistic antibacterial effect against *Xanthomonas oryzae* pv. *oryzae*. *J Fungi*. 2020;6(3):181. <https://doi.org/10.3390/jof6030181>
10. Hasan A, Walker F, Klaiber I, Schöne J, Pfannstiel J, Voegele RT. New approaches to manage Asian soybean rust (*Phakopsora pachyrhizi*) using *Trichoderma* spp. or their antifungal secondary metabolites. *Metabolites*. 2022;12(6):507. <https://doi.org/10.3390/metabolites12060507>
11. Kubicek CP, Steinendorff AS, Chenthama K, Manganiello G, Henrissat B, Zhang J, Druzhinina IS. Evolution and comparative genomics of the most common *Trichoderma* species. *BMC Genomics*. 2019;20:1-24. <https://doi.org/10.1186/s12864-019-5680-7>

12. Kredics L, Antal Z, Manczinger L, Szekeres A, Kevei F, Nagy E. Influence of environmental parameters on *Trichoderma* strains with biocontrol potential. *Food Technol Biotechnol*. 2003;41(1):37-42.
13. Ferreira FV, Musumeci MA. *Trichoderma* as biological control agent: scope and prospects to improve efficacy. *World J Microbiol Biotechnol*. 2021;37(5):90. <https://doi.org/10.1007/s11274-021-03058-7>
14. Suada IK. Short Communication: The potential of various indigenous *Trichoderma* spp. to suppress *Plasmodiophora brassicae*, the pathogen of clubroot disease on cabbage. *Biodiversitas*. 2015;18:1424-9. <https://doi.org/10.13057/biodiv/d180418>
15. Amir Daryaei. pH and water activity in culture media affect biological control activity of *Trichoderma atroviride* against *Rhizoctonia solani*. *Biol Control*. 2016;92:24-30. <https://doi.org/10.1016/j.bioc.2015.09.001>
16. Ali HZ, Aboud HM, Dheyab NS, Musa NK, Gasam FH. Effects of pH and ECw on growth and sporulation of indigenous *Trichoderma* spp. *Int J Phytopathol*. 2015;4(1):15-20. <https://doi.org/10.33687/phytopath.004.01.0966>
17. Fedele G, Brischetto C, Rossi V. Biocontrol of *Botrytis cinerea* on grape berries as influenced by temperature and humidity. *Front Plant Sci*. 2020;11:1232. <https://doi.org/10.3389/fpls.2020.01232>
18. Di Lelio I, Coppola M, Comite E, Molisso D, Lorito M, Woo SL, et al. Temperature differentially influences the capacity of *Trichoderma* species to induce plant defense responses in tomato against insect pests. *Front Plant Sci*. 2021;12:678830. <https://doi.org/10.3389/fpls.2021.678830>
19. Hamdia ZA, Kalaivani N. Evaluating the efficacy of *Trichoderma* isolates and *Bacillus subtilis* as biological control agents against *Rhizoctonia solani*. *Res J Appl Sci (Medwell)*. 2013;8:72-81.
20. Zhang X, Li J, Wang H. Effects of environmental factors on the growth and sporulation of *Trichoderma* species. *J Appl Microbiol*. 2015;118(5):1225-33. <https://doi.org/10.1111/jam.12801>
21. Desmaris F, Lévêque R, Vachon P. Influence of pH and temperature on the production of *Trichoderma harzianum* conidia and their biocontrol efficacy. *Biocontrol Sci Technol*. 2014;24(1):69-79. <https://doi.org/10.1080/09583157.2013.863686>
22. Gomes RR, Siqueira JO, Santos JG. Effects of osmotic stress on the growth and sporulation of *Trichoderma harzianum*. *World J Microbiol Biotechnol*. 2016;32(4):58. <https://doi.org/10.1007/s11274-016-2057-3>
23. Vinale F, Sivasithamparam K, Ghisalberti EL, Marra R. *Trichoderma* species as biocontrol agents of plant diseases. *Biol Control*. 2008;46(1):16-24. <https://doi.org/10.1016/j.bioc.2008.03.007>
24. Szczech M, Szymczyk P. The effect of different environmental conditions on the growth and sporulation of *Trichoderma* species. *World J Microbiol Biotechnol*. 2007;23(12):1683-9. <https://doi.org/10.1007/s11274-007-9405-5>
25. Maurya MK, Srivastava M, Harshita. Effect of various pH levels on the growth and sporulation of *Trichoderma viride* isolates and assessing their antagonistic activity against soil-borne pathogens. *J Pure Appl Microbiol*. 2024;18(4):2516-27. <https://doi.org/10.22207/JPAM.18.4.23>
26. Gupta A, Kumar R, Rai A. Impact of temperature and pH on the growth and antagonistic activity of *Trichoderma harzianum*. *Biol Control*. 2017;104:57-65. <https://doi.org/10.1016/j.bioc.2016.09.008>
27. Singh A, Shahid M, Srivastava M, Pandey S, Sharma A. Optimal physical parameters for growth of *Trichoderma* species at varying pH, temperature and agitation. *Virol Myol*. 2014;3:127. <https://doi.org/10.4172/2161-0517.1000127>
28. Bauer R, Puhalla J. Influence of environmental parameters on the biocontrol activity of *Trichoderma* species. *Soil Biol Biochem*. 2007;39(10):2505-12. <https://doi.org/10.1016/j.soilbio.2007.04.017>
29. Raizada MN, Landa BB. Role of *Trichoderma* species in promoting plant growth and controlling plant pathogens. *Front Plant Sci*. 2015;6:694. <https://doi.org/10.3389/fpls.2015.00694>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.