



RESEARCH ARTICLE

# Enhancing zinc bio-fortification in wheat through agronomic interventions

Dipak Gite, Saumya Sharma, Sushant Patil, Bhupendra Mathpal & Mayur Darvhankar\*

School of Agriculture, Lovely Professional University, Punjab 144 001, India

\*Email: [mayur.21878@lpu.co.in](mailto:mayur.21878@lpu.co.in)



## ARTICLE HISTORY

Received: 11 October 2023  
Accepted: 01 December 2024

Available online  
Version 1.0 : 28 February 2025  
Version 2.0 : 24 June 2025



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Gite D, Sharma S, Patil S, Mathpal B, Darvhankar M. Enhancing zinc bio-fortification in wheat through agronomic interventions. Plant Science Today. 2025; 12(2): 1-7. <https://doi.org/10.14719/pst.3009>

## Abstract

Around half of the world's agricultural soils, including sandy loam soils of Punjab region, are deficient in zinc (Zn), leading to decreased crop production and nutritional value. Balanced application of macro nutrients along with Zn bio-fortification enhances crop productivity and reduces the deterioration of soil health. This study investigates the effect of zinc sulphate monohydrate ( $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$ ) on growth, yield and nutrient uptake of wheat (Var. PBW-343). The plants were subjected to the following treatments:  $T_1$  (control),  $T_2$  (100% Recommended Dose of Fertilizer, RDF),  $T_3$  (100% RDF + 5 kg  $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$  ha<sup>-1</sup>),  $T_4$  (100% RDF + 10 kg  $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$  ha<sup>-1</sup>),  $T_5$  (100% RDF + 15 kg  $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$  ha<sup>-1</sup>),  $T_6$  (100% RDF + 20 kg  $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$  ha<sup>-1</sup>) and  $T_7$  (100% RDF + two spray of  $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$  @ 0.5% at tillering and late jointing stages). The results revealed that the plant height, number of functional leaves per plant, number of tillers per plant, number of effective tillers per plant, spike length, number of grains per spike, test weight, grain yield and straw yield are significant and found maximum in treatment  $T_5$ ; which was 100% RDF and 15 kg  $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$  ha<sup>-1</sup>.

## Keywords

bio-fortification; uptake; wheat; yield; zinc

## Introduction

Wheat (*Triticum aestivum* L.) is the most widely cultivated annual plant from Poaceae family. It is the second most significant food crop after rice in terms of area and production (1). The estimated area used for wheat cultivation in world is 221.85 million hectares (Mha), with annual production of 776.10 million metric tonnes (MmT) and productivity of 3.50 MT ha<sup>-1</sup> (2). India contributes a gross area of 31.36 Mha with annual production of 107.86 MmT and productivity of 3.44 MT ha<sup>-1</sup> (2). Punjab contributes 3.53 Mha area and annual production is 17.14 MT with a productivity of 4862 kg ha<sup>-1</sup> during the year 2020–21 (3).

Nutrient plays an important role in increasing production potential of wheat (4). Macronutrients and micronutrients have major role in growth and development of wheat crop. Each nutrient has its specific importance and regulates many of the plant's metabolic processes (4). A balanced application of nitrogen (N), phosphorus (P) and potassium (K) is needed in wheat to enhance the growth. However, micronutrients are essential plant nutrients important for healthy growth and increased yield. Micronutrients play specific roles and are required to regulate essential physiological functions in plants (5).

Most of Indian soils are deficient in micronutrient, Zn (6). Zn is one of the seven essential plant micronutrients which promote stronger emergence, quicker stand establishment, better root growth, greater plant vigour, protein synthesis, metabolism of carbohydrate and increased yield. Plant-available Zn is low in most of the cereal-growing areas. Thus, Zn application is essential in deficient soils of India to enhance cereal yield and Zn concentration in grain (7). Wheat often has low grain Zn content in some regions (8). Another important factor that can significantly impact a person's real Zn intake is wheat processing, as it often results in the removal of the wheat's bran and aleurone layers. A significant amount of Zn is lost throughout various wheat processing technologies (8).

The present study devised different combinations of RDF and Zn to understand the effects of Zn on wheat plants by observing various parameters like plant height, number of functional leaves per plant, number of tillers per plant, number of effective tillers per plant, spike length, number of grains per spike, test weight, grain yield and straw yield.

## Materials and Methods

### Location and climatic condition

The field experiment was conducted at Experimental farm, Lovely Professional University during *Rabi* 2020–2021. The experimental location is located in sub-tropical region with hot summer and cool winter's weather condition. At crop growth stage, the temperature ranged from 4–37°C and annual rainfall ranged from 500–750 mm, with most falling during monsoon season (July to September)(Fig. 1).

### Initial soil chemical properties

Soil for the experiment was collected from the depth of 0–15 cm from different places of the experimental site. The experimental soil had the following characteristics: soil pH (7.81), electrical conductivity (0.21dSm<sup>-1</sup>), organic carbon

(0.46 % i.e 4.6 gm kg<sup>-1</sup> soil), available soil nitrogen (145.5 kg ha<sup>-1</sup>), available soil phosphorous (11.25 kg ha<sup>-1</sup>), available soil potassium (180.56 kg ha<sup>-1</sup>) and DTPA (Diethylene triamine penta acetic acid) extractable Zn (0.21 mg kg<sup>-1</sup>).

### Experimental materials and treatment details

The certified seed of wheat PBW-343 was used for the present investigation with the seed rate of 100 kg ha<sup>-1</sup>. PBW-343 is resistant to brown rust and moderately resistant to yellow rust. Average plant height is 100 cm and has high yield potential in Punjab region. The experiment had seven treatments and three replicates and it was set up using a Randomized Block Design (RBD). The treatments used were T<sub>1</sub> (control), T<sub>2</sub> (100% RDF), T<sub>3</sub> (100% RDF + 5 kg ZnSO<sub>4</sub>.H<sub>2</sub>O ha<sup>-1</sup>), T<sub>4</sub> (100%RDF + 10 kg ZnSO<sub>4</sub>.H<sub>2</sub>O ha<sup>-1</sup>), T<sub>5</sub> (100%RDF + 15 kg ZnSO<sub>4</sub>.H<sub>2</sub>O ha<sup>-1</sup>), T<sub>6</sub> (100%RDF + 20 kg ZnSO<sub>4</sub>.H<sub>2</sub>O ha<sup>-1</sup>) and T<sub>7</sub> (100% RDF + two sprays of ZnSO<sub>4</sub>.H<sub>2</sub>O @ 0.5% at tillering and late jointing stages). Except the control (T<sub>1</sub>), all treatments received the RDF (N:P:K) at the rates of 120:60:30 kg ha<sup>-1</sup>. A 50% dose of nitrogen and a 100% dose of potassium and phosphorus were applied at the time of seeding in the form of urea, single super phosphate and murate of potash. At 30 days after seeding (DAS), the remaining 50% dose of N in the form of urea was delivered. ZnSO<sub>4</sub>.H<sub>2</sub>O was applied into the soil in treatments: T<sub>3</sub>, T<sub>4</sub>, T<sub>5</sub> and T<sub>6</sub> whereas its foliar spray was done in treatment, T<sub>7</sub>, at tillering and late jointing stages. Zn application to soil will guarantee its spread throughout the root zone and longer availability of the micronutrient to the plant, probably throughout the crop life cycle, thus providing a more sustained nutrient supply and increasing the possibility to solve Zn deficits of wheat.

### Growth and yield parameters

Five randomly chosen plants from each plot were tagged and used for measuring plant height, number of leaves per plant, number of tillers per plant and the results were averaged. The yield parameters, i.e. effective tillers per plant, spike length (cm) and number of grains per spike, was cal-




Fig. 1. Weather parameters during growing season of wheat crop.

culated from randomly selected 5 plants and then averaged. The wheat was harvested from net plot area and test weight (g), grain yield ( $\text{q ha}^{-1}$ ) and straw yield ( $\text{q ha}^{-1}$ ) were calculated after drying and threshing and converted into quintal  $\text{ha}^{-1}$  (9).

### Nutrient uptake

The following formula was used to assess the uptake of specific nutrients (N, P, K & Zn) by comparing the concentration of each nutrient in each plot to the grain and dry matter yield at harvest in each plot (10):

$$\text{Nutrient intake (kg ha}^{-1}\text{)} = \text{Nutrient concentration (\%)} \times \text{Yield of grain or straw (kg ha}^{-1}\text{)} / 100$$

..... (Eqn. 1)

### Statistical analysis

The data pertaining to the growth characteristics, yield characteristics and nutrient uptake of wheat crop was collected at regular intervals and analyzed statistically in RBD in web-based OPSTAT software.

## Results

### Growth parameters

#### Plant height (cm)

Plant height is one of the major growth parameters that is directly linked with yield contributing character of the plant. At harvest, plant height after treatment  $T_5$  was recorded to be significantly highest (80.77 cm) than other treatments and treatment  $T_1$  had the lowest plant height (46.86 cm). The plant height with treatment  $T_5$  was found at par with treatments  $T_4$ ,  $T_6$  and  $T_7$  (Table 1, Fig. 2). It is observed that with the advancement of crop duration plant height also increased. The results indicated that minimum height of the plant was recorded under  $T_1$  as compared to other treatments, this might be low due to initial

**Table 1.** Effect of zinc on growth parameters of wheat crop.

Treat-ments	Plant height (cm)	Number of leaves per plant	Number of tillers per plant
$T_1$	46.86 <sup>d</sup>	4.4 <sup>b</sup>	2.06 <sup>b</sup>
$T_2$	66.23 <sup>b</sup>	13.7 <sup>a</sup>	3.2 <sup>ab</sup>
$T_3$	55.97 <sup>c</sup>	13.8 <sup>a</sup>	3.4 <sup>a</sup>
$T_4$	75.96 <sup>a</sup>	14.1 <sup>a</sup>	3.46 <sup>a</sup>
$T_5$	80.77 <sup>a</sup>	14.3 <sup>a</sup>	3.93 <sup>a</sup>
$T_6$	74.02 <sup>a</sup>	14.2 <sup>a</sup>	3.33 <sup>a</sup>
$T_7$	76.98 <sup>a</sup>	11.9 <sup>a</sup>	3.26 <sup>a</sup>
SE (m)±	2.482	1.311	0.34
C.D 5%	7.732	4.085	NS

fertility status of that soil.


#### Number of leaves per plant

Number of leaves per plant was found significant at every periodical crop growth stage. At harvest stage, the number of leaves per plant was recorded to be the maximum in treatment  $T_5$  (14.30) while minimum number of leaves per plant was recorded in the treatment  $T_1$  (4.40). The number of leaves per plant recorded under treatment  $T_5$  were found at par with rest of treatments except  $T_1$  (Table 1, Fig. 2).

#### Number of tillers per plant

A crucial factor influencing spike production and grain yield is the tiller number. The data regarding number of tillers per plant was recorded at harvest stage. The effect of different levels of Zn along with RDF on number of tillers per plant was found to be statistically non-significant. Treatment  $T_5$  had highest number of tillers per plant (3.93) and was found to be at par with rest of treatments except treatment  $T_1$  which recorded the lowest number of tillers per plant (2.06) (Table 1, Fig. 2).

#### Yield parameters



**Fig. 2.** Effect of zinc on growth parameters of wheat crop.

### Effective tillers per plant

At harvest, effective tillers per plant was more under treatment T<sub>5</sub> (2.93) and that was found to be at par with rest of treatments except T<sub>1</sub>. The T<sub>1</sub> treatment has least number of effective tillers per plant (1.03) (Table 2, Fig. 3). Various Zn levels showed significant differences in number of effective tillers at harvest.

### Spike length (cm)

**Table 2.** Effect of zinc on yield parameters of wheat crop.

Treatments	NET	LS	NGS	TW	GY	SY	HI
T <sub>1</sub>	1.03 <sup>b</sup>	6.16 <sup>b</sup>	22 <sup>c</sup>	35.6 <sup>e</sup>	7.62 <sup>d</sup>	9.91 <sup>c</sup>	43.25
T <sub>2</sub>	2.26 <sup>a</sup>	8.14 <sup>ab</sup>	33 <sup>b</sup>	37.8 <sup>de</sup>	36.87 <sup>c</sup>	50.88 <sup>b</sup>	41.95
T <sub>3</sub>	2.46 <sup>a</sup>	8.98 <sup>a</sup>	42.66 <sup>ab</sup>	38.5 <sup>cd</sup>	44.12 <sup>b</sup>	58.04 <sup>b</sup>	43.17
T <sub>4</sub>	2.66 <sup>a</sup>	9.99 <sup>a</sup>	43 <sup>ab</sup>	42.1 <sup>ab</sup>	51.61 <sup>a</sup>	69.67 <sup>a</sup>	42.51
T <sub>5</sub>	2.93 <sup>a</sup>	10.12 <sup>a</sup>	45.33 <sup>a</sup>	42.8 <sup>a</sup>	53.4 <sup>a</sup>	73.69 <sup>a</sup>	42.16
T <sub>6</sub>	2.4 <sup>a</sup>	9.55 <sup>a</sup>	42.33 <sup>ab</sup>	40.6 <sup>abc</sup>	41.05 <sup>bc</sup>	56.37 <sup>b</sup>	42.18
T <sub>7</sub>	2.26 <sup>a</sup>	8.63 <sup>a</sup>	36.33 <sup>ab</sup>	40.1 <sup>bcd</sup>	38.2 <sup>c</sup>	51.55 <sup>b</sup>	42.63
SE (m)±	0.327	0.657	2.997	0.766	1.643	3.009	-
C.D 5%	1.019	2.046	9.338	2.386	5.117	9.376	-

**NET**-Number of effective tillers per plant, **LS**-Length of spike, **NGS**-Number of grains per spike, **TW**-Test weight, **GY**-Grain yield, **SY**-Straw yield, **BY**-Biological yield, **HI**-Harvest index.



**Fig. 3.** Effect of zinc on yield parameters of wheat crop.

Maximum spike length (10.12 cm) recorded under treatment T<sub>5</sub> which was at par with rest of treatments except T<sub>1</sub> and minimum length of spike (6.16 cm) was recorded under treatment T<sub>1</sub> at harvest (Table 2, Fig. 3).

### Number of grains per spike

Grains per spike is crucial for determining the yield of wheat which contributes to final seed yield. At harvest, maximum number of grains per spike (45.33) was recorded under treatment T<sub>5</sub> which was at par with treatments T<sub>3</sub>, T<sub>4</sub>, T<sub>6</sub> and T<sub>7</sub> and minimum number of grains per spike (22.00) was recorded under treatment T<sub>1</sub> (Table 2, Fig. 3).

### Test weight (g)

Highest test weight (42.80 g) was recorded under treatment T<sub>5</sub> which was found to be at par with treatments T<sub>4</sub> and T<sub>6</sub>. The lowest value of test weight (35.60 g) was recorded in treatment T<sub>1</sub> (Table 2, Fig. 3). Zn is essential for growth hormone synthesis, formation of starch and maturation which increase the seed weight (8 and 11).

### Grain yield (q ha<sup>-1</sup>)

Zn application had a notable impact on grain yield; this is

possible because wheat requires Zn as a necessary micro-nutrient. The data indicated that the highest grain yield was recorded under treatment T<sub>5</sub> (53.40 q ha<sup>-1</sup>) which was at par with treatment T<sub>4</sub> and the lowest was recorded in treatment T<sub>1</sub> (7.62 q ha<sup>-1</sup>) (Table 2, Fig. 3).

### Straw yield (q ha<sup>-1</sup>)

Different levels of Zn showed significant variation in straw yield. Highest straw yield was found under the treatment T<sub>5</sub> (73.69 q ha<sup>-1</sup>) which was at par with treatment T<sub>4</sub> while the lowest straw yield was found under treatment T<sub>1</sub> (9.91 q ha<sup>-1</sup>) (Table 2, Fig. 3).

### Uptake of nutrients

### Nitrogen uptake (kg ha<sup>-1</sup>)

Treatment T<sub>5</sub> shows the highest N uptake by wheat grains (90.78 kg ha<sup>-1</sup>) and the straw (36.75 kg ha<sup>-1</sup>) while the lowest value of N uptake by the wheat grains (11.81 kg ha<sup>-1</sup>) and the straw (4.16 kg ha<sup>-1</sup>) was found in T<sub>1</sub>. Wheat showed maximum total N uptake (127.53 kg ha<sup>-1</sup>) under T<sub>5</sub> and the least total N uptake (15.97 kg ha<sup>-1</sup>) under T<sub>1</sub> (Table 3, Fig. 4).

### Phosphorous uptake (kg ha<sup>-1</sup>)

Treatment T<sub>5</sub> shows highest intake of P in the wheat grains (19.75 kg ha<sup>-1</sup>) and the straw (18.37 kg ha<sup>-1</sup>) while the low-


est K intake in wheat grains (2.13 kg ha<sup>-1</sup>) and the straw (8.02 kg ha<sup>-1</sup>) was found in T<sub>1</sub>. Maximum total K intake in wheat (110.58 kg ha<sup>-1</sup>) was found under T<sub>5</sub> and lowest K intake in wheat (10.15 kg ha<sup>-1</sup>) was found in T<sub>1</sub> (Table 3, Fig. 4).

### Zinc uptake (g ha<sup>-1</sup>)

Treatment T<sub>5</sub> shows highest intake of Zn in wheat grains (146.95 g ha<sup>-1</sup>) and the straw (72.65 g ha<sup>-1</sup>) while the lowest value of Zn uptake by wheat grains (7.92 g ha<sup>-1</sup>) and the straw (4.75 g ha<sup>-1</sup>) was found under T<sub>1</sub>. Maximum total Zn intake in wheat (219.60 g ha<sup>-1</sup>) was found under T<sub>5</sub> and minimum total Zn intake in wheat (12.67 g ha<sup>-1</sup>) was found

**Table 3.** Effect of zinc on nutrient uptake by wheat.

Treatments	Nitrogen uptake (kg ha <sup>-1</sup> )			Phosphorous uptake (kg ha <sup>-1</sup> )			Potassium uptake (kg ha <sup>-1</sup> )			Zinc uptake (g ha <sup>-1</sup> )		
	Grain	Straw	Total	Grain	Straw	Total	Grain	Straw	Total	Grain	Straw	Total
T <sub>1</sub>	11.81 <sup>e</sup>	4.16 <sup>c</sup>	15.97 <sup>e</sup>	1.75 <sup>d</sup>	1.78 <sup>c</sup>	3.53 <sup>d</sup>	2.13 <sup>c</sup>	8.02 <sup>e</sup>	10.15 <sup>d</sup>	7.92 <sup>e</sup>	4.75 <sup>d</sup>	12.67 <sup>e</sup>
T <sub>2</sub>	61.2 <sup>d</sup>	24.38 <sup>b</sup>	85.58 <sup>d</sup>	13.21 <sup>c</sup>	12.19 <sup>b</sup>	25.4 <sup>c</sup>	11.42 <sup>b</sup>	61.46 <sup>d</sup>	72.88 <sup>c</sup>	78.97 <sup>f</sup>	46.48 <sup>c</sup>	125.83 <sup>d</sup>
T <sub>3</sub>	74.12 <sup>b</sup>	27.85 <sup>b</sup>	101.97 <sup>b</sup>	15 <sup>b</sup>	13.92 <sup>b</sup>	28.92 <sup>b</sup>	14.55 <sup>b</sup>	71.38 <sup>b</sup>	85.93 <sup>b</sup>	104.25 <sup>c</sup>	55.19 <sup>b</sup>	159.44 <sup>c</sup>
T <sub>4</sub>	86.7 <sup>a</sup>	35.53 <sup>a</sup>	122.23 <sup>a</sup>	18.57 <sup>a</sup>	18.11 <sup>a</sup>	36.68 <sup>a</sup>	17.54 <sup>a</sup>	87.08 <sup>a</sup>	104.62 <sup>a</sup>	135.47 <sup>b</sup>	68.41 <sup>a</sup>	203.88 <sup>b</sup>
T <sub>5</sub>	90.78 <sup>a</sup>	36.75 <sup>a</sup>	127.53 <sup>a</sup>	19.75 <sup>a</sup>	18.37 <sup>a</sup>	38.12 <sup>a</sup>	18.69 <sup>a</sup>	91.89 <sup>a</sup>	110.58 <sup>a</sup>	146.95 <sup>a</sup>	72.65 <sup>a</sup>	219.6 <sup>a</sup>
T <sub>6</sub>	68.14 <sup>c</sup>	27.05 <sup>b</sup>	95.19 <sup>bc</sup>	13.95 <sup>bc</sup>	11.83 <sup>b</sup>	25.78 <sup>bc</sup>	13.13 <sup>b</sup>	69.89 <sup>bc</sup>	83.02 <sup>b</sup>	97.32 <sup>d</sup>	53.04 <sup>bc</sup>	150.36 <sup>c</sup>
T <sub>7</sub>	63.41 <sup>cd</sup>	27.74 <sup>b</sup>	91.15 <sup>cd</sup>	13.37 <sup>c</sup>	12.37 <sup>b</sup>	25.74 <sup>bc</sup>	11.84 <sup>b</sup>	62.89 <sup>cd</sup>	74.73 <sup>c</sup>	85.98 <sup>e</sup>	48.86 <sup>bc</sup>	134.84 <sup>d</sup>
SE (m)±	1.878	2.036	2.583	0.401	0.87	1.01	0.941	2.314	2.084	2.071	2.081	3.19
C.D 5%	5.851	6.344	8.046	1.251	2.71	3.145	2.932	7.209	6.493	6.453	6.484	9.938



**Fig. 4.** Effect of zinc on nutrient uptake of wheat crop.

est uptake of P by wheat grains (1.75 kg ha<sup>-1</sup>) and the straw (1.78 kg ha<sup>-1</sup>) was found in T<sub>1</sub>. Highest total P intake in wheat (38.12 kg ha<sup>-1</sup>) was found under T<sub>5</sub> and lowest P intake in wheat (3.53 kg ha<sup>-1</sup>) was found under T<sub>1</sub> (Table 3, Fig. 4).

### Potassium uptake (kg ha<sup>-1</sup>)

Treatment T<sub>5</sub> shows highest K uptake in wheat grains (18.69 kg ha<sup>-1</sup>) and the straw (91.89 kg ha<sup>-1</sup>) while the low-

under T<sub>1</sub> (Table 3, Fig. 4).

## Discussion

The combined application of 100% RDF (120:60:30 kg NPK ha<sup>-1</sup>) + 15 kg ZnSO<sub>4</sub>.H<sub>2</sub>O ha<sup>-1</sup> effectively enhanced the growth, yield and nutrient uptake in wheat. N and Zn help in optimum plant growth, cell division and metabolism, which helps to increase plant height (11). Similarly, application of more N increases the plant height that may be



attributed to vigorous vegetative growth which results in internodal extension and these observations corroborate with previous study and also the present study which reported maximum plant height due to  $\text{ZnSO}_4$  (12–14). Increased number of leaves per plant is due to sufficient and continuous supply of major nutrients through the inorganic fertilizer and these fertilizers are responsible for prolific leaf production. The photosynthetic efficiency of crop is increased by nitrogen, which is one of the main constituents of chlorophyll, thus results in higher leaf counts. Similar trend of results was reported by other researchers (15).

Similarly, for yield attributing characteristics and yield, Zn treatment increases the number of effective tillers due to its function in numerous Zn-induced enzymatic activities and auxin metabolism, which influence plant growth and development (16). Increase in Zn levels caused considerable increase in number of effective tillers per plant (17). Observed number of effective tillers increased by the application of Zn and earlier findings reported similar results (18). Length of spike is important for the number of grains and their development in the spike of plant. Application of Zn increases the spike length (19). The result is in accordance with those of earlier studies (20, 21). Zn application results in an increase in grain setting because it participates in processes like pollination and fertilization, which have direct impact on formation of pollen tube (22). Similar results were described by other groups (14, 23). The test weight increased by Zn application because Zn increases the phloem movement from leaves to the roots, stem of the plant and developing grains (22). The present findings are similar to the previous reports (23). Increase in grain yield might be due to adequate amount of Zn, which has improved plant water relation, nutrients availability and enhanced cell physiology. Zn is a crucial element of enzymes, therefore an increase in yield may be caused by abundant efficiency of enzyme activities that affects the pigment of plants (18). Similar findings on grain yield were also observed by other research groups (23–25). Straw yield enhancement might be due to the balanced supply of Zn along with inorganic fertilizers which enhanced vegetative growth. Results are in accordance with the previous study (26).

Increased nitrogen uptake by wheat at various levels of Zn might be due to synergistic effect between nitrogen and Zn (27). Plants fertilized with Zn showed a much higher rate of N uptake (28). The nitrogen uptake by wheat grain and straw was increased due to the use of Zn in comparison to control (27). Similar results were obtained by the previous findings (14, 26). Antagonistic effects of high rates of Zn application on absorption and uptake of P have been reported in wheat (26). Observed phosphorous uptake was reduced at higher levels of Zn and the present study found similar results (29). For potassium uptake, findings of this study were similar to those obtained in the earlier outcomes (14). The Zn application increases its availability in soil and consequently in the plant system that causes higher metabolism which resulted in increased uptake of nutrients (30). Earlier researchers reported similar results and found that the Zn fertilization increased Zn

concentration and uptake in wheat (31).

## Conclusion

This experimental study showed that the application of Zn along with N, P and K is important for various growth and yield contributing parameters of wheat. Furthermore, height of the plant, number of functional leaves per plant, tillers per plant, effective tillers per plant, grains per spike, length of spike, test weight (g), grain yield ( $\text{q ha}^{-1}$ ) and the straw yield ( $\text{q ha}^{-1}$ ) enhanced by the use of 100% RDF ( $120:60:30 \text{ kg NPK ha}^{-1}$ ) +  $15 \text{ kg ZnSO}_4 \cdot \text{H}_2\text{O ha}^{-1}$  and this treatment considered as the best among all treatments. The grain yield and straw yield reported under treatment  $T_5$  were at par with treatment  $T_4$  [ $100\% \text{ RDF } (120:60:30 \text{ kg NPK ha}^{-1}) + 10 \text{ kg ZnSO}_4 \cdot \text{H}_2\text{O ha}^{-1}$ ]. Since soil suffered from severe Zn deficiency, applying  $15 \text{ kg ha}^{-1}$  of Zn chemical fertilizer ( $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$ ) with 100% RDF ( $120:60:30 \text{ kg NPK ha}^{-1}$ ) significantly enhanced the N, P, K and Zn uptake by wheat.

## Acknowledgements

The authors would like to express their profound sense of gratitude towards Lovely Professional University, Punjab, where the research trial was conducted and all requirements for the same were fulfilled.

## Authors' contributions

DG carried out research work and SS helped in recording the field and lab observations. SP and BM checked and modified the draft of manuscript while MD designed the research work plan and shaped the final manuscript.

## Compliance with ethical standards

**Conflict of interest:** All authors do not have any conflict of interests to declare.

**Ethical issues:** None

## References

1. Giraldo P, Benavente E, Manzano-Agugliaro F, Gimenez E. Worldwide research trends on wheat and barley: A bibliometric comparative analysis. *Agron.* 2019;9(7):352. <https://doi.org/10.3390/agronomy9070352>
2. World Agricultural Production [Internet]. USDA Foreign Agricultural Service. 2021. <https://www.fas.usda.gov/report-release-announcement/world-agricultural-production>
3. Official website of Directorate of Economics and Statistics, Department of Agriculture and Farmers Welfare, Ministry of Agriculture and Farmers Welfare, Government of India [Internet]. 2021. Agriculture Statistics -At a Glance. <https://desagri.gov.in/document-report-category/agriculture-statistics-at-a-glance/>
4. Pandey M, Shrestha J, Subedi S, Shah KK. Role of nutrients in wheat: A review. *Trop Agrobiodiversity.* 2020;1(1):18–23: <http://dx.doi.org/10.26480/traab.01.2020.18.23>
5. Kumar S, Kumar S, Mohapatra T. Interaction between macro- and micro-nutrients in plants. *Front Plant Sci.* 2021;12:665583. <https://doi.org/10.3389/fpls.2021.665583>

6. Shukla AK, Behera SK, Prakash C, Patra AK, Rao CS, Chaudhari SK, et al. Assessing multi-micronutrients deficiency in agricultural soils of India. Sustainability. 2021;13(16):9136. <https://doi.org/10.3390/su13169136>
7. Palai JB, Jena J, Lenka SK. Growth, yield and nutrient uptake of maize as affected by zinc application-A review. Indian J Pure Appl Biosci. 2020;8(2):332–39. <http://dx.doi.org/10.18782/2582-2845.8054>
8. Wang M, Kong F, Liu R, Fan Q, Zhang X. Zinc in wheat grain, processing and food. Front Nutr. 2020;7:124. <https://doi.org/10.3389/fnut.2020.00124>
9. Biswas A, Mukhopadhyay D, Biswas A. Effect of soil zinc and boron on the yield and uptake of wheat in an acid soil of West Bengal, India. Int J Plant soil Sci. 2015;6(4):203–17. <http://dx.doi.org/10.9734/IJPSS/2015/15921>
10. Nawaz A, Bashir MA, Ahmed W, Ahmad I, Rehman A, Ikram RM, et al. Potassium and zinc improves physiological performance, nutrient use efficiency and productivity of wheat. Front Plant Sci. 2024;15:1–10. <https://doi.org/10.3389/fpls.2024.1363248>
11. Kuntoji A, Subbarayappa CT, Sathish A, Ramamurthy V, Mallesha BC. Effect of different levels of nitrogen and zinc application on growth, yield and quality of maize in rural and peri-urban gradients of southern transect of Bengaluru. J Pharmacogn Phytochem. 2021;10(1):1562–68.
12. Ananya C, Swaroop N, Rao PS, Thomas T. Effect of NPK and Zn fertilizers on growth and yield of maize (*Zea mays* L.) Var. Shivaniksh-1980. Int J Chem Stud. 2019;7(3):1864–67.
13. Iqbal J, Khan R, Wahid A, Sardar K, Khan N, Ali M, et al. Effect of nitrogen and zinc on maize (*Zea mays* L.) yield components and plant concentration. Adv in Environ.Biol. 2016;10(10):203–09.
14. Singh V, Ali J, Seema A, Kumar T, Chauhan M. Productivity, nutrient uptake and economics of wheat (*Triticum aestivum*) under potassium and zinc nutrition. Indian J Agro. 2001;60(3):426–30. <https://doi.org/10.59797/ija.v60i3.4474>
15. Hassanein MS, Zaki NM, Ahmed AG. Effect of Zn foliar application on growth and yield characteristics of two wheat cultivars. Curr Sci Int. 2019;08(03):491–98.
16. Jatav SS, Singh SK. Evaluation of different methods of zinc application on growth and yield of hybrid rice (*Oryza sativa* L.) in inceptisol of Varanasi. J Pharma Phytochem. 2018;7(4):1739–44.
17. Gismy S, Romij Uddin Md, Sohanur Rahman Md, Talukder UF, Shafiul Islam Rion Md. Zinc, a micronutrient and its effect to performance of BARI gom30. Int J Adv Geosci. 2020;8(2):137–45. <https://doi.org/10.14419/ijag.v8i2.30966>
18. Liu YM, Cao WQ, Chen XX, Yu BG, Lang M, Chen XP, et al. The responses of soil enzyme activities, microbial biomass and microbial community structure to nine years of varied zinc application rates. Sci Total Environ. 2020;737:140245. <https://doi.org/10.1016/j.scitotenv.2020.140245>
19. Sher A, Sarwar B, Sattar A, Ijaz M, Ul-Allah S, Hayat MT, et al. Exogenous application of zinc sulphate at heading stage of wheat improves the yield and grain zinc biofortification. Agron. 2022;12(3):734. <https://doi.org/10.3390/agronomy12030734>
20. Bhutto M, Ahmed NS, Umed AL, Ghulam HJ, Maqsood AK, Anab K. Growth and yield response of wheat (*Triticum aestivum*) as affected by foliar fertilization of zinc. Sci Int (Lahore). 2016;28(4):4189–92.
21. Habbasha ES, Badr EA, Latef EA. Effect of zinc foliar application on growth characteristics and grain yield of some wheat varieties under Zn deficient sandy soil condition. Int J Chem Res. 2015;8(6):452–58.
22. Hassan N, Irshad S, Saddiq MS, Bashir S, Khan S, Wahid MA, et al. Potential of zinc seed treatment in improving stand establishment, phenology, yield and grain biofortification of wheat. J Plant Nutr. 2019;42(14):1676–92. <https://doi.org/10.1080/01904167.2019.1630429>
23. Arif M, Tasneem M, Bashir F, Yaseen G, Anwar A. Evaluation of different levels of potassium and zinc fertilizer on the growth and yield of wheat. Int J Biosen Bioelectron. 2017;3(2):242–46. <https://doi.org/10.15406/ijbsbe.2017.03.00057>
24. Rudani K, Patel V, Prajapati K. The importance of zinc in plant growth-A review. Int Res J Nat Appl Sci. 2018;5(2):38–48.
25. Nauman A, Anwar S, Kashmir S, Ahmad J, Saeed B, Khan S. Response of wheat varieties to different zinc application methods. Pure and Appl Bio. 2019;8(1):489–95.
26. Paramesh V, Dhar S, Dass A, Kumar B, Kumar A, El-Ansary DO, et al. Role of integrated nutrient management and agronomic fortification of zinc on yield, nutrient uptake and quality of wheat. Sustainability. 2020;12(9):3513. <https://doi.org/10.3390/su12093513>
27. Montoya M, Vallejo A, Recio J, Guardia G, Alvarez JM. Zinc–nitrogen interaction effect on wheat biofortification and nutrient use efficiency. J Plant Nutr Soil Sci. 2020;183(2):169–79. <https://doi.org/10.1002/jpln.201900339>
28. Shivay YS, Prasad R, Pal M. Effect of variety and zinc application on yield, profitability, protein content and zinc and nitrogen uptake by chickpea (*Cicer arietinum*). Indian J Agron. 2001;59(2):317–21. <https://doi.org/10.59797/ija.v59i2.4558>
29. Zaman Q uz, Aslam Z, Yaseen M, Ihsan MZ, Khaliq A, Fahad S, et al. Zinc biofortification in rice: leveraging agriculture to moderate hidden hunger in developing countries. Arch Agron Soil Sci. 2017;64(2):147–61. <https://doi.org/10.1080/03650340.2017.1338343>
30. Liu DY, Liu YM, Zhang W, Chen XP, Zou CQ. Zinc uptake, translocation and remobilization in winter wheat as affected by soil application of Zn fertilizer. Front in Plant Sci. 2019;10:443999. <https://doi.org/10.3389/fpls.2019.00426>
31. Ghasal PC, Shivay YS, Pooniya V, Choudhary M, Verma RK. Zinc accounting for different varieties of wheat (*Triticum aestivum*) under different source and methods of application. Indian J Agri Sci. 2017;87(9):1111–16. <https://doi.org/10.56093/ijas.v87i9.73898>