



REVIEW ARTICLE

Agricultural carbon credits: A pathway to environmental sustainability

K Adhisankaran¹, V Manivannan^{1*}, N Sakthivel², D Balachandar³, K P Ragunath⁴ & V Vasumathi¹

¹Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore - 641003, India

²Department of Agronomy, Agricultural Research Station, Bhavanisagar - 638451, India

³Department of Agricultural Microbiology, Tamil Nadu Agricultural University, Coimbatore - 641003, India

⁴Centre for Water and Geospatial Studies Tamil Nadu Agricultural University, Coimbatore - 641003, India

*Email: manivannanv@tnau.ac.in



ARTICLE HISTORY

Received: 08 March 2024

Accepted: 13 September 2024

Available online

Version 1.0 : 01 November 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Adhisankaran K, Manivannan V, Sakthivel N, Balachandar D, Ragunath KP, Vasumathi V. Agricultural carbon credits: A pathway to environmental sustainability. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.3504>

Abstract

Climate change and ensuring food security for a rapidly growing global population are two of the biggest challenges in agriculture. To meet the commitments made in the Paris Climate Agreement, it is important to use effective methods for managing soil that can help sequester and stabilise carbon. Conservation agriculture has a huge potential to sequester carbon in plants and soil, making it a viable option for carbon trading despite its significant contribution to global greenhouse gas emissions. Carbon sequestration can be achieved through sustainable practices such as adopting conservation agriculture, crop rotation, cover crop cultivation, crop residue incorporation or mulching, effective management of nutrient supply to crops, and transforming towards organic agriculture and agroforestry. These practices promote food security and environmental improvement and help mitigate global warming. Carbon pricing mechanisms are policies that impose a cost on carbon pollution, encouraging people and organisations to choose low-carbon options and reduce their emissions. Agricultural producers can benefit from carbon trading by earning extra revenue by selling their excess carbon credits to those who emit higher amounts of greenhouse gases. Carbon credit systems in agriculture are still in the early stages, so farmers may have more opportunities to participate in future carbon trading.

Keywords

conservation agriculture; carbon pricing; carbon sequestration; GHG emissions; organic carbon

Introduction

Global climate change has led to the exploration of innovative solutions to reduce greenhouse gas (GHG) emissions in all sectors. Atmospheric carbon dioxide (CO₂) levels have surged dramatically from about 277 ppm in 1750 to 419.3 ppm in 2023 which owes to a 51% increase (1). Historically, developed nations have been the primary contributors to GHG emissions, but the past two decades have witnessed a significant rise in emissions from rapidly developing countries like India, despite their lower per-capita emissions (2). This trend highlights the urgent need for effective climate action. Efforts are now being made to reduce carbon footprints in all areas, including agriculture and food production. The critical role of GHGs in climate change has led to international negotiations, elevating carbon to a tradable commodity and fostering the growth of the global carbon market (3). Agriculture and land use change are pivotal in achieving net-zero emissions,

functioning both as sources and sinks of GHGs (4). However, current policy measures have been deemed inadequate, underscoring the need for stable, well-crafted policies to reduce emissions effectively.

Carbon credits have emerged as a promising voluntary market mechanism to address this challenge. They incentivise sustainable agricultural practices, commonly known as carbon farming, which enhance the soil's capacity to sequester carbon and reduce GHG emissions (5). Farmers adopting practices like reduced tillage or optimised fertiliser use can generate carbon credits and trade them in the market, providing a sustainable pathway to transform agricultural systems while preserving farmers' economic viability (6).

Moreover, carbon farming offers a framework for guiding the global economy towards net-zero emissions and advancing climate action in line with the '4 per 1000 Initiative,' particularly the UN Sustainable Development Goals (SDGs) (5). This article explores the potential of carbon credits within the agricultural sector as a strategy to mitigate climate change, highlighting agriculture's dual role in both contributing to and combating GHG emissions. In addition, it assesses carbon trading mechanisms, pricing strategies, and policy frameworks that support global emission reduction targets and promote agriculture practices resilient to climate change.

Carbon credit

A carbon credit denotes reducing or eliminating one ton of CO₂ or its equivalent and other greenhouse gases (GHG) from the atmosphere (7). Carbon credits are important in carbon trading because they serve as a means to mitigate the negative impacts of greenhouse gas emissions on the environment (6). Carbon credits are earned by entities that emit less than their allocated limit in cap-and-trade systems. These credits can be exchanged or sold to other agencies that exceed their emission limits, providing economic incentives for businesses and individuals to reduce their carbon footprint.

Carbon credits provide a mechanism for countries and businesses to work towards meeting emission reduction targets outlined in international agreements like the Kyoto Protocol and the Paris Agreement. The demand for carbon credit is rising, resulting in a shift towards renewable energy alternatives and a low-carbon economy. The maximum carbon credit limit is determined by the rate of carbon emissions, which stands at 50 Gt CO₂-e/year (8). Carbon credit transactions entail the acquisition of emission licenses from individuals or entities capable of reducing greenhouse gas emissions or storing additional carbon. The cost per carbon credit is determined by market forces as buyers and sellers engage in trading activities (6, 8).

Emissions from agriculture

Intensive agricultural practices in developing countries significantly impact climate change due to the emission of greenhouse gases. Agriculture contributes 10–12% of global greenhouse gas emissions out of which, 19% are from India (2). The major activities that result in the

emission of greenhouse gases are rice cultivation, the use of synthetic fertilisers, and the burning of crop residues. Specifically, in India, methane emissions from rice cultivation account for approximately 138.68 kilotons, while N₂O emissions result from the use of synthetic fertilisers, and the burning of crop residues emit around 405.77 kilotons and 3.59 kilotons, respectively (9). Furthermore, drained agricultural peatlands, which comprise only 1% of agricultural land, contribute to about 33% of global cropland greenhouse gas emissions (10). Emissions occur at every production stage, making non-resilient agricultural practices a key concern to climate change. Fig. 1 highlights the significant impact of different categories on greenhouse gas emissions. "Agrifood systems" contribute the most to the emissions, followed by "Emission on agricultural land" and "Farm gate". Other sources have relatively lower emissions. It suggests that to reduce overall emissions, efforts could be focused on areas like managing agrifood systems and emissions on agricultural land (9).




Fig. 1. Emission share of CO₂ equivalent of different categories [Data sourced from FAOSTAT for the year 2021] (9)

Chemical amendments

The increasing demand for food has resulted in a rise in the use of chemical fertilisers, which could harm ecosystems and contribute to greenhouse gas emissions. Nitrogen fertilisers are the major source of anthropogenic N₂O emissions, while their production alone accounts for nearly 1% of global GHG emissions (11). According to FAO predictions, the usage of synthetic N fertilisers is likely to rise globally by 50% from 2012 levels by 2050, which might threaten the climate aim outlined in the Paris Agreement. Phosphorus is essential for plant growth but is often lost to waterways, causing eutrophication (12). Potassium is an important nutrient for plant growth and its primary source is potassium chloride, which contributes to greenhouse gas emissions during production (13). Fig. 2 shows the major components of greenhouse gas emissions in the agrifood system worldwide. Emissions from livestock, synthetic fertilisers and fires in humid tropical forests have the highest impact. Managing livestock emissions and reducing synthetic fertiliser use can help reduce overall emissions (9).

Organic amendment

Organic farming has seen growth across countries, with approximately 70 million hectares of organic land managed by nearly 2.9 million farmers. Nonetheless, the




Fig. 2. Agrifood system emissions by component in India [Data sourced from FAOSTAT for the year 2021] (9)

application of amendments comes with drawbacks, including issues like nutrient eutrophication and the emission of greenhouse gases (3). Animal manure is widely used in agriculture, but it raises CO₂, CH₄ and N₂O emissions. Composting helps to transform organic waste into humus-like substances, but the process must be correctly handled to reduce GHG emissions (14).

Crops

Cereals are an important source of the human diet and account for a significant portion of global N fertiliser consumption. Rice is a major source of greenhouse gas emissions, contributing to around 30% of worldwide agricultural CH₄ emissions (15). According to FAOSTAT (9), rice has the highest N₂O emissions among all the crops in India's agricultural and food system. It contributes to 47.97% of the total emissions, followed by wheat with 27.98% (Fig. 3). This highlights the importance of reducing N₂O emissions from rice and wheat to mitigate the environmental impact of India's agrifood system. Maize output is increasing every year and nitrogen fertilisers are being largely used to increase production, which leads to an increase in emissions (16). Oil crops cultivated to extract oil emit around 2.8-19.7 kg CO₂ eq per kg of palm oil (17).




Fig. 3. Crop-wise emission of N₂O from India [Data sourced from FAOSTAT for the year 2021] (9)

Agricultural carbon sequestration

Carbon sequestration is a process that aims to minimise carbon emissions by stabilising photosynthesis and altering land use. It involves two main approaches: protecting and maintaining ecosystems and managing ecosystems to


enhance carbon sequestration by increasing carbon content in soil and plant biomass (18). Soil can mitigate climate change by acting as the largest terrestrial carbon reservoir. Soil stores about 1526 Pg C of Soil Organic Carbon (SOC) and 940 Pg C of soil inorganic carbon. Soil and vegetation at a depth of one metre hold three times more carbon than the atmosphere, which currently stands at 880 Pg C (4). Approximately 10% of atmospheric CO₂ is cycled by the soil each year. The quantity of carbon in soil is determined by the equilibrium between the inputs of SOC and the rates at which it mineralises. Land use and soil management changes influence organic carbon sequestration rates in soil. It offers further benefits like food security, biodiversity, maintaining water quality and elemental recycling (5). Soil quality can be enhanced by improving the concentration of SOC by implementing sustainable agricultural practices (Table 1). Retention of soil carbon can be done through direct and indirect mechanisms. Direct mechanisms account for 50-80% of stable SOC and involve the synthesis and accumulation of plant-based carbon in microbial biomass. Indirect mechanisms enhance plant growth and increase biomass entering the soil. Carbon sequestration plays a role in the field of agriculture as it helps offset amounts of methane and nitrous oxide emissions caused by agricultural activities (19). Fig. 4 presents the estimated amount of carbon (Pg C yr⁻¹) that can be stored through soil organic carbon sequestration (7). Adopting Sustainable Soil Management (SSM) practices can mitigate between 0.14-0.56 Pg CO₂ eq yr⁻¹ annually. This estimation is based on three scenarios: SSM1 with a 5% increase in carbon input, SSM2 with a 10% increase in carbon input and SSM3 with a 20% increase in carbon input.

Conservation agriculture and carbon sequestration

Intensive agricultural practices like tillage, over-fertilisation and excessive cultivation are used to fulfil the growing demand for food. Due to the degradation of soil characteristics and increased soil erosion caused by these activities, there is a global loss of 25-75% of SOC (20). Diversified cropping systems like intercropping, cover crops and crop rotation can mitigate this and facilitate SOC sequestration (21, 22). Legumes can fix atmospheric nitrogen and increase nitrogen content when integrated into cropping systems (20).

Table 1. Overview of sustainable agricultural practices and its key benefits

| Sustainable agricultural practices | Key benefits | Reference |
|---|---|----------------------------------|
| <ul style="list-style-type: none"> Conservation tillage <ul style="list-style-type: none"> No-tillage Cover crops Crop residue | Enhance SOC stock and improves SOC stabilization | (22, 61, 83, 84, 49, 87, 90, 91) |
| <ul style="list-style-type: none"> Organic farming <ul style="list-style-type: none"> Crop rotation Conservation tillage <ul style="list-style-type: none"> No-tillage Organic farming <ul style="list-style-type: none"> No-tillage | Reduce GHG emission | (21, 72, 84) |
| <ul style="list-style-type: none"> Crop residue Crop rotation | Improve soil quality by improving soil aggregation and soil structure | (44, 84, 85,86,88) |
| <ul style="list-style-type: none"> N fertilisers Cover crops | Enhancing biomass production by increasing carbon and nitrogen inputs | (46, 89) |
| <ul style="list-style-type: none"> Crop residue | Increase soil labile organic carbon fractions | (38) |

**Fig. 4.** Sequestration potential of various systems [Data sourced from FAO] (7)

Conservation Agriculture (CA) refers to a type of farming system involving minimal soil disturbance, permanent soil cover and diversification of crops. The FAO promotes CA as an effective approach to achieving sustainable land management, environmental protection and climate change adaptation and mitigation (23). Numerous global initiatives have identified CA as a significant contributor to the adaptation and mitigation of climate change in the field of agriculture (22). Conservation agriculture is an effective way to maintain soil health and increase SOC storage. It also provides other benefits that promote agricultural sustainability. Conservation agriculture can reduce CO₂ emissions by increasing carbon stored in soil and limiting vehicle use, lowering direct emissions (21). Conservation agriculture entails practically implementing three interrelated concepts and supplementary good agricultural practices, which play an important role in soil carbon sequestration. Further discussion on this topic can be found below.

Effects of tillage on carbon sequestration: Tillage practices significantly affect carbon sequestration and GHG production (24). Soil organic carbon and aggregate stability are interrelated and No-tillage (NT) methods can promote the development of soil aggregates, enhance their resilience and evenly distribute carbon among soil aggregates of various sizes (25). No-tillage increases top-layer carbon storage more than conventional tillage (26, 27) by enhancing soil organic carbon, microbial biomass carbon, dehydrogenase, alkaline phosphatase and β -glucosidase and notably reduces global warming potential (24, 28). According to a previous study, NT methods improved soil total organic carbon stock by 9.0% compared to CT in 0-40 cm soil depth. The carbon pool management index was also improved by 16.5% (29). NT effectively stores both labile and recalcitrant carbon, favouring diverse carbon conversion. NT reduces the alkyl C to O-alkyl C conversion and lignin to polysaccharides ratio, indicating lower organic carbon decomposition (30).

No-tillage resulted in higher SOC stock in the topsoil (0-20 cm) than conventional tillage. After 23 years, SOC was 21% greater under NT in the topsoil, indicating an effective system to recover lost SOC (26). Conservation tillage practices increase SOC accumulation and tillage with straw mulch is more effective in promoting SOC accumulation than other tillage methods (28). Deep tillage increased SOC stock significantly by (7.36%) over conventional tillage. Subsoiling increased SOC (8.76%) than deep ploughing (5.85%). Subsoiling increased SOC in the top 40 cm layer, with the most significant improvement at the 0-10 cm layer (11.41%). Deep ploughing increased SOC stock in the 10-40 cm layer but had no effect on the SOC of the 0-10 cm layer (31). The application of straw to soil increases easily oxidisable and particulate organic carbon. When combined with NT, residue return leads to improved soil carbon sequestration (32).

Effects of crop residue on carbon sequestration: Crop residues play a significant role in increasing soil organic matter, which in turn affects nutrient availability and carbon storage (11). Which further promotes humus formation, improves soil quality and adds carbon and nutrients to the soil (32-34). Many studies conducted in developing regions have reported positive impacts of crop residue on soil quality, including increased organic matter, carbon stock, water holding capacity, nutrient recycling and reduced soil loss (32, 35, 36). Additionally, crop residue provides the necessary energy source for soil organisms, which contributes to the stability of soil aggregates, enhances the permeability of the soil and favours the movement of soil air, water and heat (37). Adding crop residues into the soil can affect both passive and labile soil carbon pools, leading to emissions, whereas the resistant proportion aids in carbon sequestration. The amount of carbon added depends on the method of residue management, as surface retention and incorporation can improve SOC at different depths (38).

Crop residues are the primary source of SOC, containing approximately 45% carbon and their management significantly impacts carbon dynamics (37). Globally, crop residues generate 2.4 Pg C annually, accounting for 25% of global carbon emissions from fossil fuels and cement production in 2021. China, the US, and India are the leading producers of crop residues. Cereal crops contribute the most at 1.62 Pg C year⁻¹, followed by oilseed/pulse crops at 0.45 Pg C year⁻¹. Maize has the highest potential for crop residue production at 0.58 Pg C year⁻¹, followed by rice at 0.49 Pg C year⁻¹. The European Union has a crop residue production potential of 0.18 Pg C, which is equivalent to 16% of its fossil carbon emissions for 2021 (39).

Effects of crop rotation on carbon sequestration: Crop rotation is a practice where different crops are grown sequentially on the same land. Recently, there's renewed interest in this practice for sustainable food production and minimal environmental impact (27, 40). Crop rotation benefits agriculture by improving soil structure, fertility, and nutrient use efficiency as well as controlling pests,

weeds and diseases (27, 41, 42). Diverse crop rotation with cover crops and perennials can improve ecosystem services. In addition, crop residue can increase SOC and total carbon input, due to increased living soil cover.

Leguminous crops increase soil microbial activity, soil organic carbon, soil moisture and total porosity, enhancing the yield of subsequent crops (43, 44). Legume forage crops, used in rotations, provide nitrogen for subsequent non-legume crops and maintain soil nitrogen levels (41). Furthermore, leguminous crops also increase soil stability, porosity, available water, and the retention of soil carbon and nitrogen. Legume crop residue is a major carbon source in agricultural soils, favouring the accumulation of humus and the soil's carbon pool as legume residues decompose faster than non-legumes. Thus, Legume-based cropping is a key option for soil carbon sequestration, maintaining a favourable soil C/N ratio (43).

Grain legumes, with their deep and robust root system, contribute to SOC through leftover shoots and root biomass. The root of the pigeon pea crop, extending more than 1.5 m, is recognised for deep soil carbon sequestration. Carbon sequestration in deep soil is less prone to oxidation and has longer stability. Cultivating legumes in alternation with cereal crops can enhance the storage of carbon in soil by 9-45 Mg C ha⁻¹, while decreasing the release of greenhouse gases by 5-7 times when compared to non-leguminous plants (45). Biological nitrogen fixation indirectly reduces the use of chemical fertilisers, potentially reducing CO₂ emissions (43, 44).

Effect of cover crops on carbon sequestration: Cover crops are a type of crops that are planted between main crops to provide a range of benefits such as increasing soil organic carbon, improving soil structure, reducing erosion and nutrient leaching and promoting carbon sequestration and nitrogen retention (46, 47). Additionally, these crops help in mitigating climate change and reducing farming operations and fossil fuel consumption. Cover crops are also grown as green manure, which produce large biomass and help in enlarging the soil carbon sink (48). Cover crops have the potential to capture carbon in soils without affecting the yield of the primary crop (49). The presence of cover crops led to a 16% increase in SOC on a global scale when compared to not using them and resulted in a carbon sequestration rate of 167 kg C ha⁻¹ yr⁻¹. Cover crop mixtures are more effective at promoting SOC buildup than single species (47). However, the carbon sequestration rate depends on the type of cover crop used, with non-legumes performing better than legumes due to their larger carbon-to-nitrogen ratios (50).

Effect of nutrient management on carbon sequestration

Fertilising with N, P, and K can affect soil carbon stocks by impacting plant productivity, litter composition and microbial metabolism. These nutrients boost plant growth, increase biomass and soil fertility, leading to a positive effect on SOC stocks (51). Organic amendments can help in carbon sequestration, improve soil fertility and combat climate change (52, 53). However, their effectiveness

depends on local environmental conditions (54). Green manure can increase the carbon pool index (CPI) and reduce the carbon management index (CMI). Organic amendments can affect the rate of SOC decomposition, known as the priming effect (55, 56). It's crucial to consider carbon management indices, crop yield and GHG emissions when evaluating agricultural practices (57). Soil organic carbon stocks vary with the balance between carbon capture and decomposition (54). Manures combined with mineral fertiliser improved soil organic carbon storage and sequestration efficiency (58). This combination also enhances soil structure, fertility and microbial biomass, resulting in stable SOC in both macro and micro-aggregates (59, 60).

Nitrogen input impacts soil organic carbon stocks due to changes in fresh particulate organic carbon and decomposed organic carbon output, increased soil lignin content and decreased CO₂ emissions (55, 61-63). Nitrogen bioavailability is critical for organic carbon cycling and sequestration (59). Increased nitrogen effectiveness promotes soil organic carbon sequestration by promoting plant growth and litter formation, reducing microbial activity and promoting aggregate formation (51, 62, 64, 65). It also enhances soil carbon stocks, including soil inorganic carbon, which store more carbon in deep soil than SOC and contribute to 30% of the world's total carbon in the top 1m of soil (66). This enhances soil carbon accumulation by stimulating plant growth and biomass production (67). However, nutrient addition can also accelerate organic matter decomposition, affecting carbon cycling and storage.

Effect of organic farming on carbon sequestration

Organic farming is a sustainable adaptation technique to overcome climate change. It is not just a specific agricultural production system but a systemic approach to sustainable livelihoods, considering physical, economic and socio-cultural factors (68). The Codex Alimentarius Commission has defined organic farming as a holistic food production management system that promotes agro-ecosystem health, biodiversity and soil biological activity while emphasising the use of management practices over off-farm inputs and synthetic materials. It assists in climate change adaptation and mitigation through nutrient management as it avoids using synthetic fertilisers, which play a key role in agricultural GHG emissions (69).

Compared to conventional farming, organic farming has greater potential in soil carbon sequestration (70, 71). This is accomplished by reducing N₂O emissions due to reduced nitrogen input, reduced CO₂ emissions from farming system inputs, and soil carbon sequestration, which is one of the potential mitigation strategies (72, 73). The soil carbon sequestration rate ranges from 200 kg to 2000 kg C ha⁻¹ yr⁻¹, depending on the organic soil management practices that favour sequestering of carbon in the soil ecosystem (71).

Effect of agroforestry on carbon sequestration

Agroforestry is a sustainable land use system that combines agriculture and forestry to maximise benefits and resource efficiency (74). It is highly valued for its potential to support climate change adaptation and mitigation, with the greatest potential for carbon sequestration among all land uses, which can result in both environmental and economic benefits (11, 75). Agroforestry provides a multitude of advantages, including economic, socio-cultural and environmental benefits. These benefits include reduced erosion, enhanced soil fertility and water quality, improved aesthetics, greater biodiversity and carbon sequestration (76-78). Carbon sequestration in agroforestry occurs through the accumulation of aboveground and belowground biomass and soil, with trees playing a crucial role in adaptation and mitigation due to their significant aboveground biomass volume and deep, dense root systems.

Agroforestry is an essential component of developing countries nationally determined contributions, which could be a significant contributor in global climate goals (76). The IPCC has highlighted agroforestry as the most promising method for carbon sequestration by 2040. It estimates that agroforestry can mitigate between 1.1 to 2.2 Pg C through biomass production and soil organic carbon pool (74).

Agroforestry practices can intentionally increase the SOC pool (79). The integration of trees leads to enhanced CO₂ sequestration from the atmosphere through photosynthesis, resulting in enhanced carbon storage in tree components for long-term storage and its deposition in the soil as a stable organic matter pool (carbon storage) (80). In small-scale tropical agroforestry systems, the C sequestration rates range between 1.5-3.5 mg C ha⁻¹ yr⁻¹ (81). Between the ages of 10 and 20-30, trees have the highest capacity to sequester carbon. By the time forests reach the age of 30, they can sequester approximately 200-520 tonnes of CO₂ ha⁻¹ (82).

Carbon trading

Carbon trading, also known as emissions trading, is a mechanism established under Kyoto Protocol (Article 17). It allows countries with unused emission units, i.e. emissions permitted but not utilised, to sell their excess unused units to other countries that have exceeded their emission targets (92). The carbon trading system significantly impacts the economy and society, affecting industries that emit high levels of carbon (93). The pressure to reduce emissions is increasing due to worsening climate conditions. Agricultural production must consider both ecological impact and direct emission reduction pressure during carbon trading. Carbon trading occurs in markets both within and outside the Kyoto Protocol. These markets are used to distribute and trade carbon credits (Fig. 5) through two types of transactions: allowance-based and project-based. In allowance-based transactions, buyers obtain emission allowances that are generated and allocated by regulators operating under cap-and-trade systems, including Assigned Amount Units (AAUs) and European Union (EU) allowances (94). Countries that join the Kyoto Protocol and accept national limits are




Fig. 5. Carbon trading

granted AAUs and assigned quotas to specific emitting activities or industrial units, such as power plants or paper manufacturers, to implement their emission targets. Allowance-based markets feature EU ETS, Australia's New South Wales Greenhouse Gas Abatement Scheme and the CCX, a voluntary but legally enforceable commitment (95). Regarding project-based transactions, the costumer buys emission credits from a project that demonstrates a reduction in GHG emissions over a period, compared to the emissions in the absence of that project. Many of these deals involve the forward sale of credits, which means that buyers are acquiring future reductions in emissions. The Clean Development Mechanism (CDM) and joint implementation plans of the Kyoto Protocol are the best-known instances of these transactions. Carbon credit mechanisms are in the development phase in 6 countries and have already been implemented in 17 countries. (Table 2) (96).

Carbon and its pricing

Carbon, in terms of Carbon credits is usually priced based on the demand and supply of carbon credits by various firms or countries. With the increase in demand, the price of carbon credits increases, and vice-versa.

According to a report (97), maintaining temperatures below 2°C (the upper limit agreed upon in the Paris Agreement) requires carbon prices to be set between USD 40/tons of CO₂ and USD 80/tons of CO₂ by 2020. Additionally, the report suggests that these prices should be further increased to between USD 50/tons of CO₂ and USD 100/tons of CO₂ by 2030.

The policy for greenhouse gas emissions affects the market conditions, which can also be impacted by changes in weather, energy prices and technical improvements. These factors are known to shift unpredictably, and the costs could vary significantly with the development of new technologies. The carbon credit market is a new and rising institution that may have more buyers or sellers at different times with varying prices. Soon, calculating the price of carbon credits will be challenging, even with models that reflect market features, due to the lack of understanding about the exact impact of supply and demand factors on pricing. As the carbon credit market evolves, more data on prices and quantities will incentivise greenhouse gas reductions (8).

Carbon credit systems allow farmers to earn additional income by selling their excess carbon credits to producers who emit higher levels of GHGs (7, 98). While these agricultural carbon credit systems are still in their

Table 2. Global overview of carbon crediting mechanism initiatives

| Country | Mechanism | Implementation status |
|----------------|--|-----------------------|
| Ecuador | Ecuador Crediting Mechanism | Under development |
| Egypt | Egypt Crediting Mechanism | Under development |
| India | India Crediting Mechanism | Under development |
| Indonesia | Indonesia Crediting Mechanism | Under development |
| Vietnam | Vietnam Crediting Mechanism | Under development |
| Mexico | Mexico Crediting Mechanism | Under development |
| United Kingdom | UK Woodland Carbon Code | Implemented in 2011 |
| Spain | Spain FES-CO ₂ Program | Implemented in 2011 |
| Switzerland | Switzerland CO ₂ Attestations Crediting Mechanism | Implemented in 2012 |
| Japan | J-Credit Scheme | Implemented in 2013 |
| Kazakhstan | Kazakhstan Crediting Mechanism | Implemented in 2013 |
| Thailand | Thailand Voluntary Emission Reduction Program | Implemented in 2014 |
| China | China GHG Voluntary Emission Reduction Program | Implemented in 2014 |
| Korea | Republic of Korea Offset Credit Mechanism | Implemented in 2015 |
| United Kingdom | Peatland Code | Implemented in 2015 |
| Sri Lanka | Sri Lanka Carbon Crediting Mechanism | Implemented in 2016 |
| Colombia | Colombia Crediting Mechanism | Implemented in 2017 |
| South Africa | South Africa Crediting Mechanism | Implemented in 2019 |
| Canada | Canada Federal GHG Offset System | Implemented in 2022 |
| Australia | Australian Carbon Credit Unit Scheme | Implemented in 2024 |
| Chile | Chile Crediting Mechanism | Implemented in 2024 |
| Saudi Arabia | Greenhouse Gas Crediting and Offsetting Mechanism | Implemented in 2024 |
| Portugal | Portuguese Voluntary Carbon Market | Implemented in 2024 |

early stages, the carbon credit market is expanding. This growth means that farmers may have additional opportunities to participate in the future. Additionally, farmers can be encouraged to participate in carbon-crediting mechanisms by stressing the benefits and accompanying economic incentives rather than focusing on the possible cash gain.

Policy framework

Agricultural carbon credits are actively traded within the Voluntary Carbon Markets (VCM), where companies either reduce emissions or acquire carbon credits to offset their environmental impact. In January 2024, India introduced the Carbon Credit Trading Scheme (CCTS) to promote sustainable agricultural practices that generate tradable carbon credits for farmers. India is already a major supplier of carbon offsets in the global voluntary market, but ensuring transparency within the ecosystem remains a critical challenge. Green washing, where entities falsely present themselves as environmentally friendly without making real environmental improvements, is a key concern. It can occur when carbon credits are misused by purchasing them without reducing emissions, providing misleading information, or relying on low-quality credits. To address green washing, stakeholders involved in agricultural carbon markets must prioritise high-quality, verified carbon credits, maintain transparent communication and comply with legal standards. Effective checking mechanisms include third-party verification, regulatory oversight and a comprehensive strategy that focuses on direct emission reductions alongside the responsible use of carbon credits.

To further strengthen the policies, particularly in the agriculture sector where small and marginal farmers are predominant, it is crucial to focus on capacity building of farmers and investing in research to develop innovative techniques that improve soil health and enhance crop resilience to combat climate change. Incentives should be provided for adopting low emission technologies such as precision agriculture, organic farming and agroforestry. Government should focus on reducing GHG emissions, improve resource use efficiency, promote strategies to

adapt climate change, and prioritise agricultural productivity. It requires the creation of credible databases or enhancing available databases (such as FAOSTAT and World Bank open knowledge repository) that contain reliable data and scientific research to improve policy decisions. Financial arrangements should be made for smooth implementation, while flexibility for convergence with other national/international activities, institutional arrangements and stakeholder knowledge should also be emphasised. The policies should prioritise sustainability and livelihood, resource inventory and all existing climate-smart practices while ensuring profitability for growers and emissions from farming activities.

Conclusion

The relationship between agriculture and carbon dynamics highlights the need for sustainable practices in the era of climate change. As the global community aims to reduce emissions, strategies like organic farming and conservation agriculture can promote soil health and carbon sequestration. Carbon trading systems offer a viable means of incentivising sustainable practices, with agriculture presenting significant potential for generating carbon credits despite its current limited contribution. Despite agriculture-related emissions mitigating projects accounting for just more than 1% of total carbon credits granted, it is crucial to recognise that land use, including forestry, is the most extensive source of carbon credits, awarding approximately half of its units issued to date (Fig. 6). This highlights the untapped potential of agriculture, which is another form of land use, in this industry. By concentrating on sustainable and carbon-sequestering agricultural practices, the agricultural sector has the potential to significantly increase its share of carbon credits issued. This benefits the environment and provides additional revenue streams for farmers and agricultural businesses. The future of agriculture in carbon trading is promising, with plenty of room for growth and innovation.




Fig. 6. Sector-Wise Share of Carbon credit issues [Data sourced from AgFunderNews] (99)

Acknowledgements

The authors acknowledge the support of Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu in successful completion of review work.

Authors' contributions

KA conceptualized the study, performed the original draft preparation, and contributed to its review and editing. VM conceptualized and supervised the study and contributed to the review and editing of the manuscript. NS, DB and KPR contributed to the review and editing of the manuscript. VV contributed to the literature collection. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None.

References

- Friedlingstein P, O'Sullivan M, Jones MW, Andrew RM, Bakker DCE, Hauck J, et al. Global carbon budget 2023. *Earth Syst Sci Data*. 2023;15(12):5301-69. <https://doi.org/10.5194/essd-15-5301-2023>
- Cariappa AA, Konath NC, Sapkota TB, Krishna VV. Evaluating the potential and eligibility of conservation agriculture practices for carbon credits. *Sci Rep*. 2024;14(1):9193. <https://doi.org/10.1038/s41598-024-59262-6>
- Singh S, Kiran BR, Mohan SV. Carbon farming: a circular framework to augment CO₂ sinks and to combat climate change. *Env Sci Adv*. 2024;3(4):522-42. <https://doi.org/10.1039/D3VA00296A>
- Rodrigues CID, Brito LM, Nunes LJR. Soil carbon sequestration in the context of climate change mitigation: A review. *Soil Systems*. 2023;7(3):64. <https://doi.org/10.3390/soilsystems7030064>
- Phelan L, Chapman PJ, Ziv G. The emerging global agricultural soil carbon market: the case for reconciling farmers' expectations with the demands of the market. *Environmental Development*. 2024;49:100941. <https://doi.org/10.1016/j.envdev.2023.100941>
- Lokuge N, Anders S. Carbon-credit systems in agriculture: a review of literature. The School of Public Policy Publications. 2022;15. <https://doi.org/10.55016/ojs/sppp.v15i1.74591>
- Peralta G, Di Paolo L, Luotto I, Omuto C, Mainka M, Viatkin K, et al. Global soil organic carbon sequestration potential map (GSOCseq v1. 1)–Technical manual [Internet]. Food and Agriculture Org. 2022 [updated 2022 Jan 18; cited 2024 Feb 14]. Available from: <https://doi.org/10.4060/cb2642en>
- Mathews JA. How carbon credits could drive the emergence of renewable energies. *Energy Policy*. 2008;36(10):3633-39. <https://doi.org/10.1016/j.enpol.2008.05.033>
- FAOSTAT. Food and agriculture data [Internet]. FAO; 2024 [updated 2024 Feb 14; cited 2024 Feb 14]. Available from: <https://www.fao.org/faostat/en/#data/>.
- Anthony TL, Silver WL. Hot spots and hot moments of greenhouse gas emissions in agricultural peatlands. *Biogeochemistry*. 2024;167(4):461-77. <https://doi.org/10.1007/s10533-023-01095-y>
- Rao DLN, Balachandar D. Nitrogen inputs from biological nitrogen fixation in Indian agriculture. In: Abrol YP, Adhya TK, Aneja VP, Raghuram N, Pathak H, Kulshrestha U, et al., editors. *The Indian Nitrogen Assessment*: Elsevier; 2017. 117-32. <https://doi.org/10.1016/B978-0-12-811836-8.00008-2>
- Li Y, Shang J, Zhang C, Zhang W, Niu L, Wang L, et al. The role of freshwater eutrophication in greenhouse gas emissions: A review. *Sci Total Environ*. 2021;768. <https://doi.org/10.1016/j.scitotenv.2020.144582>
- Murrell TS, Mikkelsen RL, Sulewski G, Norton R, Thompson ML. Improving potassium recommendations for agricultural crops. *Springer Nature*. 2021. <https://doi.org/10.1007/978-3-030-59197-7>
- Shakoor A, Shahbaz M, Farooq TH, Sahar NE, Shahzad SM, Altaf MM, et al. A global meta-analysis of greenhouse gases emission and crop yield under no-tillage as compared to conventional tillage. *Sci Total Environ*. 2021;750. <https://doi.org/10.1016/j.scitotenv.2020.142299>
- Gupta K, Kumar R, Baruah KK, Hazarika S, Karmakar S, Bordoloi N. Greenhouse gas emission from rice fields: a review from Indian context. *Environ Sci Pollut Res*. 2021;28(24):30551-72. <https://doi.org/10.1007/s11356-021-13935-1>
- Meng Q, Yue S, Hou P, Cui Z, Chen X. Improving yield and nitrogen use efficiency simultaneously for maize and wheat in China: A review. *Pedosphere*. 2016;26(2):137-47. [https://doi.org/10.1016/S1002-0160\(15\)60030-3](https://doi.org/10.1016/S1002-0160(15)60030-3)
- Reijnders L, Huijbregts MAJ. Palm oil and the emission of carbon-based greenhouse gases. *J Clean Prod*. 2008;16(4):477-82. <https://doi.org/10.1016/j.jclepro.2006.07.054>
- Adhisankaran K, Sakthivel S. Carbon sequestration. In: Priya S, Ninama Atulkumar R, Chiragkumar MB, Narinder P, Vishnu M, editors. *Emerging Trends in Agricultural Practices*: ND Global Publication House; 2024. p. 48-63.
- Mason ARG, Salomon MJ, Lowe AJ, Cavagnaro TR. Microbial solutions to soil carbon sequestration. *J Clean Prod*. 2023;417. <https://doi.org/10.1016/j.jclepro.2023.137993>
- Francaviglia R, Almagro M, Vicente-Vicente JL. Conservation agriculture and soil organic carbon: principles, processes, practices and policy options. *Soil Systems*. 2023;7(1):17. <https://doi.org/10.3390/soilsystems7010017>
- Piccoli I, Chiarini F, Carletti P, Furlan L, Lazzaro B, Nardi S, et al. Disentangling the effects of conservation agriculture practices on the vertical distribution of soil organic carbon. Evidence of poor carbon sequestration in North-Eastern Italy. *Agric Ecosyst Environ*. 2016;230:68-78. <https://doi.org/10.1016/j.agee.2016.05.035>
- Gonzalez-Sanchez EJ, Veron-Gonzalez O, Conway G, Moreno-Garcia M, Kassam A, Mkomwa S, et al. Meta-analysis on carbon sequestration through conservation agriculture in Africa. *Soil Tillage Res*. 2019;190:22-30. <https://doi.org/10.1016/j.still.2019.02.020>
- Tadiello T, Acutis M, Perego A, Schillaci C, Valkama E. Soil organic carbon under conservation agriculture in Mediterranean and humid subtropical climates: Global meta-analysis. *European J Soil Science*. 2023;74(1):e13338. <https://doi.org/10.1111/ejss.13338>
- Mehra P, Baker J, Sojka RE, Bolan N, Desbiolles J, Kirkham MB, et al. A review of tillage practices and their potential to impact the soil carbon dynamics. *Adv Agron*. 150: Elsevier. 2018;p. 185-230. <https://doi.org/10.1016/bs.agron.2018.03.002>
- Yang S, Wang Y, Wang Z, Yan X, Feng M, Xiao L, et al. Interactive effects of conservation tillage on the aggregate stability and soil organic carbon. *J Plant Nutr Soil Sci*. 2022;185(4):505-12. <https://doi.org/10.1002/jpln.202200044>
- Briedis C, De Moraes Sá JC, Lal R, De Oliveira Ferreira A, Franchini JC, Milori DMBP. Preservation of labile organic compounds is the pathway for carbon storage in a 23-year continuous no-till system on a Ferralsol in southern Brazil. *Geoderma Regional*. 2023;33:e00643. <https://doi.org/10.1016/j.geodrs.2023.e00643>

27. Liu X, Tan S, Song X, Wu X, Zhao G, Li S, et al. Response of soil organic carbon content to crop rotation and its controls: A global synthesis. *Agric Ecosyst Environ.* 2022;335:108017. <https://doi.org/10.1016/j.agee.2022.108017>
28. Guo L, Shi J, Lin W, Liang J, Lu Z, Tang X, et al. Soil bacteria mediate soil organic carbon sequestration under different tillage and straw management in rice-wheat cropping systems. *Agriculture.* 2022;12(10):1552. <https://doi.org/10.3390/agriculture12101552>
29. Zhang Q, Zhang Y, Wang X, Li H, Liu P, Wang X, et al. Change of tillage system affects the soil carbon pools characters, reduces carbon emissions and improves maize yield in the Loess Plateau. *Eur J Agron.* 2022;141. <https://doi.org/10.1016/j.eja.2022.126614>
30. Zhu K, Ran H, Wang F, Ye X, Niu L, Schulin R, et al. Conservation tillage facilitated soil carbon sequestration through diversified carbon conversions. *Agric Ecosyst Environ.* 2022;337:108080. <https://doi.org/10.1016/j.agee.2022.108080>
31. Wang X, Xu X, Qiu S, Zhao S, He P. Deep tillage enhanced soil organic carbon sequestration in China: A meta-analysis. *J Clean Prod.* 2023;399. <https://doi.org/10.1016/j.jclepro.2023.136686>
32. Zhao J, Liu Z, Lai H, Yang D, Li X. Optimizing residue and tillage management practices to improve soil carbon sequestration in a wheat-peanut rotation system. *J Environ Manage.* 2022;306:114468. <https://doi.org/10.1016/j.jenvman.2022.114468>
33. Sharma S, Thind HS, Yadvinder S, Sidhu HS, Jat ML, Parihar CM. Effects of crop residue retention on soil carbon pools after 6 years of rice-wheat cropping system. *Environ Earth Sci.* 2019;78(10):296. <https://doi.org/10.1007/s12665-019-8305-1>
34. Aravindh S, Chinnadurai C, Malathi P, Sanjivkumar V, Pandian PS, Thiyageshwari S, et al. Nutrient management and cropping pattern influence the carbon sequestering ability of semi-arid tropical soils. *Environmental Sustainability.* 2023;6(1):87-98. <https://doi.org/10.1007/s42398-023-00264-x>
35. Li S, Li Y, Li X, Tian X, Zhao A, Wang S, et al. Effect of straw management on carbon sequestration and grain production in a maize-wheat cropping system in Anthrosol of the Guanzhong Plain. *Soil Tillage Res.* 2016;157:43-51. <https://doi.org/10.1016/j.still.2015.11.002>
36. Haas E, Carozzi M, Massad RS, Butterbach-Bahl K, Scheer C. Long term impact of residue management on soil organic carbon stocks and nitrous oxide emissions from European croplands. *Sci Total Environ.* 2022;836:154932. <https://doi.org/10.1016/j.scitotenv.2022.154932>
37. Blanco-Canqui H, Lal R. Crop residue management and soil carbon dynamics. In: Lal R, Follett RF, editors. *SSSA Special Publications*. Madison, WI, USA: American Society of Agronomy and Soil Science Society of America; 2009. p. 291-309. <https://doi.org/10.2136/sssaspecpub57.2ed.c17>
38. Aditi K, Abbhishek K, Chander G, Singh A, Falk T, Mequanint MB, et al. Assessing residue and tillage management options for carbon sequestration in future climate change scenarios. *Curr Res Environ Sustain.* 2023;5. <https://doi.org/10.1016/j.crsust.2023.100210>
39. Karan SK, Woolf D, Azzi ES, Sundberg C, Wood SA. Potential for biochar carbon sequestration from crop residues: A global spatially explicit assessment. *GCB Bioenergy.* 2023;15(12):1424-36. <https://doi.org/10.1111/gcbb.13102>
40. Zhang X, Hou H, Yin J, Fang Y, Yu X, Wang H, et al. Crop rotation with plastic mulching increased soil organic carbon and water sustainability: A field trial on the Loess Plateau. *Soil Use Manag.* 2023;39(2):717-28. <https://doi.org/10.1111/sum.12873>
41. Giacometti C, Mazzon M, Cavani L, Triberti L, Baldoni G, Ciavatta C, et al. Rotation and fertilization effects on soil quality and yields in a long term field experiment. *Agronomy.* 2021;11(4):636. <https://doi.org/10.3390/agronomy11040636>
42. Zheng F, Liu X, Ding W, Song X, Li S, Wu X. Positive effects of crop rotation on soil aggregation and associated organic carbon are mainly controlled by climate and initial soil carbon content: A meta-analysis. *Agric Ecosyst Environ.* 2023;355:108600. <https://doi.org/10.1016/j.agee.2023.108600>
43. De Stefano A, Jacobson MG. Soil carbon sequestration in agroforestry systems: a meta-analysis. *Agroforest Syst.* 2017;92:285-99. <https://doi.org/10.1007/s10457-017-0147-9>
44. Ajit, Dhyani SK, Handa AK, Newaj R, Chavan SB, Alam B, et al. Estimating carbon sequestration potential of existing agroforestry systems in India. *Agroforest Syst.* 2017;91(6):1101-18. <https://doi.org/10.1007/s10457-016-9986-z>
45. Rahman MM, Alam MS, Islam MM, Kamal MZU, Rahman GKMM, Haque MM, et al. Potential of legume-based cropping systems for climate change adaptation and mitigation. *Advances in Legumes for Sustainable Intensification: Elsevier.* 2022;381-402. <https://doi.org/10.1016/B978-0-323-85797-0.00030-6>
46. Quintarelli V, Radicetti E, Allevato E, Stazi SR, Haider G, Abideen Z, et al. Cover crops for sustainable cropping systems: A review. *Agriculture.* 2022;12(12). <https://doi.org/10.3390/agriculture12122076>
47. McClelland SC, Paustian K, Schipanski ME. Management of cover crops in temperate climates influences soil organic carbon stocks: a meta-analysis. *Ecol Appl.* 2021;31(3). <https://doi.org/10.1002/eap.2278>
48. Lal R. Cover cropping and the "4 per thousand" proposal. *J Soil Water Conserv.* 2015;70(6):141A-A. <https://doi.org/10.2489/jswc.70.6.141A>
49. Sharma P, Singh A, Kahlon CS, Brar AS, Grover KK, Dia M, et al. The role of cover crops towards sustainable soil health and agriculture -A review paper. *AJPS.* 2018;09(09):1935-51. <https://doi.org/10.4236/ajps.2018.99140>
50. Zhu S, Sainju UM, Zhang S, Tan G, Wen M, Dou Y, et al. Cover cropping promotes soil carbon sequestration by enhancing microaggregate-protected and mineral-associated carbon. *Sci Total Environ.* 2024;908. <https://doi.org/10.1016/j.scitotenv.2023.168330>
51. Poeplau C, Zopf D, Greiner B, Geerts R, Korvaar H, Thumm U, et al. Why does mineral fertilization increase soil carbon stocks in temperate grasslands? *Agric Ecosyst Environ.* 2018;265:144-55. <https://doi.org/10.1016/j.agee.2018.06.003>
52. Gopinath KA, Rajanna GA, Venkatesh G, Jayalakshmi M, Kumari VV, Prabhakar M, et al. Influence of crops and different production systems on soil carbon fractions and carbon sequestration in rainfed areas of semiarid tropics in India. *Sustainability.* 2022;14(7):4207. <https://doi.org/10.3390/su14074207>
53. Chinnadurai C, Gopalaswamy G, Balachandrar D. Impact of long-term organic and inorganic nutrient managements on the biological properties and eubacterial community diversity of the Indian semi-arid Alfisol. *Arch Agron Soil Sci.* 2014;60(4):531-48. <https://doi.org/10.1080/03650340.2013.803072>
54. Bai X, Tang J, Wang W, Ma J, Shi J, Ren W. Organic amendment effects on cropland soil organic carbon and its implications: A global synthesis. *CATENA.* 2023;231. <https://doi.org/10.1016/j.catena.2023.107343>
55. Li X, Zhu W, Xu F, Du J, Tian X, Shi J, et al. Organic amendments affect soil organic carbon sequestration and fractions in fields with long-term contrasting nitrogen applications. *Agric Ecosyst Environ.* 2021;322. <https://doi.org/10.1016/j.agee.2021.107643>
56. Wang G, Luo Z. Organic amendments alter long-term turnover and stability of soil carbon: Perspectives from a data-model integration. *Agronomy.* 2021;11(11):2134. <https://doi.org/10.3390/agronomy11112134>
57. Hou P, Xue L, Wang J, Petropoulos E, Xue L, Yang L. Green manure amendment in paddies improves soil carbon sequestration but

- cannot substitute the critical role of N fertilizer in rice production. *Agronomy*. 2022;12(7):1548. <https://doi.org/10.3390/agronomy12071548>
58. Zhang C, Zhao Z, Li F, Zhang J. Effects of organic and inorganic fertilization on soil organic carbon and enzymatic activities. *Agronomy*. 2022;12(12). <https://doi.org/10.3390/agronomy12123125>
 59. Li S, Wei W, Liu S. Long-term organic amendments combined with nitrogen fertilization regulates soil organic carbon sequestration in calcareous soil. *Agronomy*. 2023;13(2):291. <https://doi.org/10.3390/agronomy13020291>
 60. Liang S, Sun N, Wang S, Colinet G, Longdoz B, Meersmans J, et al. Manure amendment acts as a recommended fertilization for improving carbon sequestration efficiency in soils of typical drylands of China. *Front Environ Sci*. 2023;11. <https://doi.org/10.3389/fenvs.2023.1173509>
 61. Luo R, Kuzyakov Y, Liu D, Fan J, Luo J, Lindsey S, et al. Nutrient addition reduces carbon sequestration in a Tibetan grassland soil: Disentangling microbial and physical controls. *Soil Biol Biochem*. 2020;144:107764. <https://doi.org/10.1016/j.soilbio.2020.107764>
 62. Chen Z, Geng S, Zhou X, Gui H, Zhang L, Huang Z, et al. Nitrogen addition decreases soil aggregation but enhances soil organic carbon stability in a temperate forest. *Geoderma*. 2022;426: <https://doi.org/10.1016/j.geoderma.2022.116112>
 63. Meng D, Cheng H, Shao Y, Luo M, Xu D, Liu Z, et al. Progress on the effect of nitrogen on transformation of soil organic carbon. *Processes*. 2022;10(11):2425. <https://doi.org/10.3390/pr10112425>
 64. Lu X, Vitousek PM, Mao Q, Gilliam FS, Luo Y, Turner BL, et al. Nitrogen deposition accelerates soil carbon sequestration in tropical forests. *Proc Natl Acad Sci USA*. 2021;118(16). <https://doi.org/10.1073/pnas.2020790118>
 65. Liang Z, Cao B, Jiao Y, Liu C, Li X, Meng X, et al. Effect of the combined addition of mineral nitrogen and crop residue on soil respiration, organic carbon sequestration and exogenous nitrogen in stable organic matter. *Appl Soil Ecol*. 2022;171. <https://doi.org/10.1016/j.apsoil.2021.104324>
 66. Jin S. Recommended nitrogen fertilization enhances soil carbon sequestration in China's monsoonal temperate zone. *PeerJ*. 2018;6:e5983. <https://doi.org/10.7717/peerj.5983>
 67. Li M, Hu H, He X, Jia J, Drosos M, Wang G, et al. Organic carbon sequestration in soil humic substances as affected by application of different nitrogen fertilizers in a vegetable-rotation cropping system. *J Agric Food Chem*. 2019;67(11):3106-13. <https://doi.org/10.1021/acs.jafc.8b07114>
 68. Farooqi ZUR, Sabir M, Zeeshan N, Naveed K, Hussain MM. Enhancing carbon sequestration using organic amendments and agricultural practices. In: Agarwal RK, editor. *Carbon Capture, Utilization and Sequestration: InTech*; 2018. p. 17-35. <https://doi.org/10.5772/intechopen.79336>
 69. Wani S, Chand S, Najjar G, Teli M. Organic farming: As a climate change adaptation and mitigation strategy. *Curr Agri Res*. 2013;1(1):45-50. <https://doi.org/10.12944/CARJ.1.1.06>
 70. Muller A, Aubert C. The potential of organic agriculture to mitigate the influence of agriculture on global warming—A review. In: Bellon S, Penvern S, editors. *Organic Farming, Prototype for Sustainable Agricultures*. Dordrecht: Springer Netherlands; 2014. p. 239-59. https://doi.org/10.1007/978-94-007-7927-3_13
 71. Patle GT, Badyopadhyay KK, Kumar M. An overview of organic agriculture: A potential strategy for climate change mitigation. *JANS*. 2014;6(2):872-79. <https://doi.org/10.31018/jans.v6i2.548>
 72. Scialabba NE-H, Müller-Lindenlauf M. Organic agriculture and climate change. *Renew Agric Food Syst*. 2010;25(2):158-69. <https://doi.org/10.1017/S1742170510000116>
 73. Gopinath KA, Visha Kumari V, Venkatesh G, Jayalakshmi M, Prabhamani PS, Ravindra Chary G. Organic agriculture: Potentials in managing abiotic stresses in crop production. In: Bal SK, Mukherjee J, Choudhury BU, Dhawan AK, editors. *Advances in Crop Environment Interaction*. Singapore: Springer Singapore; 2018. 229-43. https://doi.org/10.1007/978-981-13-1861-0_9
 74. Panwar P, Mahalingappa DG, Kaushal R, Bhardwaj DR, Chakravarty S, Shukla G, et al. Biomass production and carbon sequestration potential of different agroforestry systems in India: A critical review. *Forests*. 2022;13(8):1274. <https://doi.org/10.3390/f13081274>
 75. Bogale GA, Bekele SE. Sustainability of agroforestry practices and their resilience to climate change adaptation and mitigation in sub-Saharan Africa: A review. *Ekológia (Bratislava)*. 2023;42(2):179-92. <https://doi.org/10.2478/eko-2023-0021>
 76. Duguma LA, Minang PA, Watson C, Nath AJ, Muthee KW, Van Noordwijk M, et al. Agroforestry as a key intervention to achieve nationally determined contribution (NDC) targets. In: Dagar JC, Gupta SR, Sileshi GW, editors. *Agroforestry for Sustainable Intensification of Agriculture in Asia and Africa*. Singapore: Springer Nature Singapore; 2023. 641-64. https://doi.org/10.1007/978-981-19-4602-8_19
 77. Gupta SR, Dagar JC, Sileshi GW, Chaturvedi RK. Agroforestry for climate change resilience in degraded landscapes. In: Dagar JC, Gupta SR, Sileshi GW, editors. *Agroforestry for Sustainable Intensification of Agriculture in Asia and Africa*. Singapore: Springer Nature Singapore; 2023. 121-74. https://doi.org/10.1007/978-981-19-4602-8_5
 78. Kumar R, Kumar R, Karmakar S, Kumar A, Singh AK, Kumar A, et al. Impact of amide fertilizer on carbon sequestration under the agroforestry system in the Eastern Plateau region of India. *Sustainability*. 2023;15(12):9775. <https://doi.org/10.3390/su15129775>
 79. Carranca C, Pedra F, Madeira M. Enhancing carbon sequestration in Mediterranean agroforestry systems: A review. *Agriculture*. 2022;12(10):1598. <https://doi.org/10.3390/agriculture12101598>
 80. Swamy SL, Tewari VP. Mitigation and adaptation strategies to climate change through agroforestry practices in the tropics. In: Dagar JC, Tewari VP, editors. *Agroforestry*. Singapore: Springer Singapore; 2017. 725-38. https://doi.org/10.1007/978-981-10-7650-3_29
 81. Shah S. Agroforestry practices and its contribution to combat climate change in subtropical region of Pakistan. *PAB*. 2020;9(1). <https://doi.org/10.19045/bspab.2020.90042>
 82. Ghosh PK, Mahanta SK, Mandal D, Mandal B, Ramakrishnan S. Carbon management in tropical and sub-tropical terrestrial systems: Springer; 2020. <https://doi.org/10.1007/978-981-13-9628-1>
 83. Liang C, Balser TC. Warming and nitrogen deposition lessen microbial residue contribution to soil carbon pool. *Nat Commun*. 2012;3(1):1222. <https://doi.org/10.1038/ncomms2224>
 84. Liu X, Wu X, Liang G, Zheng F, Zhang M, Li S. A global meta-analysis of the impacts of no-tillage on soil aggregation and aggregate-associated organic carbon. *Land Degrad Dev*. 2021;32(18):5292-305. <https://doi.org/10.1002/ldr.4109>
 85. Sommer R, Ryan J, Masri S, Singh M, Diekmann J. Effect of shallow tillage, moldboard plowing, straw management and compost addition on soil organic matter and nitrogen in a dryland barley/wheat-vetch rotation. *Soil Tillage Res*. 2011;115-116:39-46. <https://doi.org/10.1016/j.still.2011.06.003>
 86. Dikgwatlhe SB, Chen Z-D, Lal R, Zhang H-L, Chen F. Changes in soil organic carbon and nitrogen as affected by tillage and residue management under wheat-maize cropping system in the North China Plain. *Soil Tillage Res*. 2014;144:110-18. <https://doi.org/10.1016/j.still.2014.07.014>
 87. Liu C, Lu M, Cui J, Li B, Fang C. Effects of straw carbon input on carbon dynamics in agricultural soils: a meta-analysis. *Glob Change Biol*. 2014;20(5):1366-81. <https://doi.org/10.1111/gcb.12517>

88. Song K, Yang J, Xue Y, Lv W, Zheng X, Pan J. Influence of tillage practices and straw incorporation on soil aggregates, organic carbon and crop yields in a rice-wheat rotation system. *Sci Rep.* 2016;6(1). <https://doi.org/10.1038/srep36602>
89. Dhaliwal SS, Naresh RK, Gupta RK, Malhotra SK, Kumar A, Kumar A, et al. Impact of conservation tillage and intensifying crop rotations in enhancing soil carbon, microbial cycling and aggregation in semiarid agro-eco systems: A review. *Prog Agri.* 2019;19(2):165. <https://doi.org/10.5958/0976-4615.2019.00051.6>
90. Yang X, Ren W, Sun B, Zhang S. Effects of contrasting soil management regimes on total and labile soil organic carbon fractions in a loess soil in China. *Geoderma.* 2012;177-178:49-56. <https://doi.org/10.1016/j.geoderma.2012.01.033>
91. Fan J, Ding W, Xiang J, Qin S, et al. Carbon sequestration in an intensively cultivated sandy loam soil in the North China Plain as affected by compost and inorganic fertilizer application. *Geoderma.* 2014;230-231:22-28. <https://doi.org/10.1016/j.geoderma.2014.03.027>
92. UNFCCC. Emissions Trading [Internet]. 2024 [updated 14 Feb 2024; cited 14 Feb 2024]. Available from: <https://unfccc.int/process/the-kyoto-protocol/mechanisms/emissions-trading>
93. Yu D, Liu L, Gao S, Yuan S, Shen Q, Chen H. Impact of carbon trading on agricultural green total factor productivity in China. *J Clean Prod.* 2022;367. <https://doi.org/10.1016/j.jclepro.2022.132789>
94. Hua J, Zhu D, Jia Y. Research on the policy effect and mechanism of carbon emission trading on the total factor productivity of agricultural enterprises. *IJERPH.* 2022;19(13):7581. <https://doi.org/10.3390/ijerph19137581>
95. Bryan E, Akpalu W, Yesuf M, Ringler C. Global carbon markets: Opportunities for sub-Saharan Africa in agriculture and forestry. *Clim Dev.* 2010;2(4):309-31. <https://doi.org/10.3763/cdev.2010.0057>
96. World Bank. State and trends of carbon pricing dashboard [Internet]. World Bank; 2024 [updated 2024 Aug 11; cited 2024 Aug 11]. Available from: <https://carbonpricingdashboard.worldbank.org/>
97. Stiglitz JE, Stern N, Duan M, Edenhofer O, Giraud G, Heal GM, et al. Report of the high-level commission on carbon prices. 2017. <https://doi.org/10.7916/d8-w2nc-4103>
98. Isermeyer F, Heidecke C, Osterburg B, Isermeyer F, et al. Integrating agriculture into carbon pricing. 2021. <https://doi.org/10.22004/AG.ECON.310017>
99. Ellis J. Agriculture produces just 1 % of carbon credits, data suggests [Internet]. *AgFunderNews*: 2021 [updated 14 Feb 2024; cited 14 Feb 2024]. Available from: <https://agfundernews.com/carbon-credits-just-one-percent-from-agriculture>