



RESEARCH ARTICLE

# Multivariate analysis and multi-trait index-based selection of maize (*Zea mays* L.) inbreds for agromorphological and yield components

N Kumari Vinodhana\*, S Sivakumar & M Himakara Datta

Department of Millets, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

\*Correspondence email - [soundhini@yahoo.co.in](mailto:soundhini@yahoo.co.in)

Received: 29 June 2024; Accepted: 26 October 2024; Available online: Version 1.0: 22 July 2025; Version 2.0: 29 July 2025

**Cite this article:** Kumari VN, Sivakumar S, Himakara DM. Multivariate analysis and multi-trait index-based selection of maize (*Zea mays* L.) inbreds for agromorphological and yield components. Plant Science Today. 2025; 12(3): 1-10. <https://doi.org/10.14719/pst.4206>

## Abstract

Germplasm is pivotal in breeding, significantly impacting targeted efforts through genetic diversity. Assessing genetic parameters aids researchers in precise germplasm selection for enhanced gains. Ideotype-based breeding refines this by tailoring plant types to specific goals. The Multi Trait Genotype Ideotype Distance Index (MGIDI) defines ideotype parameters and aids in selecting genotypes closely aligned with desired traits. A study conducted at Tamil Nadu Agricultural University during 2022-23 utilized 55 maize germplasm lines in a Randomized Block Design, evaluating nine biometrical traits. Analysis of Variance revealed significant genotype differences for all traits, with grain yield positively associated with plant height, cob length, cob diameter, kernel row number and kernels per row. The principal component analysis identified three PC with eigenvalue >1 explaining 74.44 % cumulative variance, associating the first component with yield and cob traits, the second with flowering traits and the third PC with shelling percentage and plant height. Two ideotypes based on maturity i.e. medium and late maturity were defined and MGIDI was calculated. Selection gains were positive for all the traits and the genotype B. NO 1265-6-2 (G8) was ranked as the closest to the medium maturing ideotype. Similarly, the genotype UMI 1003-2-3 (G10) was closest in resemblance to the late maturing ideotype and the selection gains were positive for all the traits except plant height. This study emphasizes the critical role of germplasm selection and ideotype-based breeding in enhancing genetic gains in maize breeding programs.

**Keywords:** association analysis; ideotype; MGIDI; selection gain; variability

## Introduction

Maize (*Zea mays* L.), often known as corn, is a key grain crop in the Poaceae family with enormous worldwide agricultural relevance. Its origin may be traced back to the Tehuacan Valley in Mexico, where it was first domesticated (1). Maize has a genome size ranging from 2.4 to 2.7 gigabase pairs (Gbp) and is notable for being one of the first plant genomes of gigabase size to be sequenced using novel approaches such as omics technology (2). This crop is critical for producing food, animal feed and low-cholesterol edible oil for human and cattle use. Its domestication required considerable morphological changes, allowing the plant to adapt from its tropical origins to survive in a variety of environmental situations (3).

In India, maize is cultivated around 10 million ha, yielding 34.3 million metric tonnes and contributing just 2 % to world production in the 2022-2023 timeframe (4). Accurately estimating and projecting grain output is critical for guaranteeing food security and developing effective food policy (5). Understanding the relationship between yield and its constituent parts is essential for increasing agricultural productivity through strategic breeding (6). To precisely gauge the level of genetic diversity within a population, Genotypic

Coefficients of Variation (GCV), Phenotypic Coefficients of Variation (PCV), Broad-sense Heritability ( $h^2$ ) and Genetic Advance (GA) must be meticulously measured. Efficient selection methods rely on a parent population with a high degree of diversity. Metrics such as PCV and GCV give information on the amount of variation in a population, whereas heritability represents the proportion of a trait that is passed down to future generations (7). Understanding heritability helps guide selection approaches, forecast improvements and evaluate the significance of genetic influences (8). The level of development observed in a certain characteristic under specific selection forces is measured by genetic advancement. Higher genetic progress combined with enhanced heritability gives ideal circumstances for selection operations.

Grain yield in maize is a complex quantitative trait, hence direct selection for this trait is not fruitful. The correlation analysis points to the interrelationships between the yield and other traits taken under study. The information on these interrelationships can be utilized to structure a selection strategy to improve the yield through the selection of yield-associated traits (9). The Principal Component Analysis (PCA) is a dimensionality reduction technique that is

useful for inferring or identifying the patterns amongst the germplasm and the traits responsible for the variation (10). It can be used for the genetic improvement of important traits contributing to the variability (11).

An ideotype can be defined as a plant type that has a combination of all the ideal traits (12). The complexity of the selection of a superior genotype with all the favourable traits can be attributed to the quantitative nature of the inheritance of those traits. Several selection indices have been proposed for selecting superior genotypes based on the defined selection criteria (13). The limitation of these indices is the effective conversion of the economic value of these traits into weightage for the selection of genotypes. This roadblock was overcome by through the development of a multi-trait selection index called the Multi Trait Genotype Ideotype Distance Index (MGIDI) which is based on factorial analysis (14). This index assigns weightage to the individual traits according to the breeding goal and aids in the selection of superior genotypes. Using a selection index to simultaneously select yield-attributing traits by providing appropriate weightage to all the components could efficiently improve grain yield. The MGIDI index has been used by multiple researchers for the selection of superior genotypes in barley, oats, sesame and maize (15-18).

The correlation studied between yield and yield components is a prerequisite to planning a meaningful breeding program. A significant association suggests that such characters could be improved simultaneously. Principal Component Analysis (PCA) is better utilized for studying the diversity among the genotypes in various crops. The number of variables is reduced through Principal Component (PC) analysis to linear functions known as canonical vectors, which account for the majority of the variation created by the traits under investigation. The current study aims to estimate the genetic variability present in the maize germplasm, mine the trait associations through correlation analysis and selection of superior genotypes by defining ideotypes with differential maturity i.e. late and medium maturity with high yield and employ the MGIDI index to select superior genotypes which are close to the defined ideotype.

## Materials and Methods

### Experimental material and layout

A total of 55 maize inbred lines were obtained from the maize unit, Department of Millets, Tamil Nadu Agricultural University, Coimbatore to carry out the research study (Table 1). The experiment was performed in a Randomised Block Design with three replications during Rabi 2022-23 in the experimental farm area of 0.04 ha of the Department of Millets. Standard agronomic procedures and plant protection practices were implemented throughout the crop season. To carry out the current investigation, the biometrical observation on days to 50 % tasseling, days to 50 % silking, plant height (cm), cob length (cm), cob diameter (cm), Number of kernel rows, number of kernels per row, shelling percentage (%) and grain yield (g) were recorded in selected five plants per genotype.

### Statistical analysis

For each of the traits studied, an analysis of variance was performed. Burton and De Vane's formula was used to calculate the coefficient of variation for these qualities (19). This variance was then categorized as high (more than 20 %), moderate (10 % - 20 %), or low (less than 10 %). The broad sense heritability and the projected genetic advance (GA) were calculated using the method suggested by Johnson et al. and categorized as low, moderate, or high (20). Correlation analysis was carried out according to the methods described by an earlier study (21). Principal Component Analysis, a dimension reduction approach developed by Massey; and Jolliffe was used to compress variables while keeping important information (22, 23). PCA was critical in decreasing data dimensionality and gaining insights from the dataset.

The MGIDI distance index as established by a previous research, was used to find genotypes that were closely related to the suggested ideotype (14). This ideotype is specially designed for early and late maturity periods, to produce a high yield. As a result, three ideotype reference models representing early, middle and late maturity were developed. These reference models were used to identify genotypes that closely matched the target ideotype's attributes. To begin, each characteristic ( $rX_{ij}$ ) was rescaled using the following equation (Eqn. 1).

$$rX_{ij} = \frac{\eta_{nj} - \varphi_{nj}}{\eta_{oj} - \varphi_{oj}} \times (\theta_{ij} - \eta_{oj}) + \eta_{nj} \quad (\text{Eqn. 1})$$

where  $\varphi_{oj}$  and  $\eta_{oj}$  are the original minimum and maximum values for the trait  $j$ , respectively;  $\varphi_{nj}$  and  $\eta_{nj}$  are the new minimum and maximum values for trait  $j$  after rescaling, respectively; and  $\theta_{ij}$  is the original value for  $j$ th trait of the  $i$ th

**Table 1.** List of 55 genotypes used under study

Code	Genotypes	Code	Genotypes
G1	B. No 9119-1-1	G29	B. No 9233-1
G2	UMI 1151-2	G30	UMI 1113
G3	UMI 653-2-3	G31	B. No 1125-7
G4	Hyd No 1075-4-1-1	G32	B. No 1258-7
G5	UMI 819-3	G33	UMI 96
G6	B. No 1265-6-2	G34	UMI 1051
G7	UMI 1101	G35	Hyd No. 2009-2-2-15
G8	B. No 1076-5-1	G36	B. No 1110-8
G9	B. No 71810	G37	B. No 1076-5-4-1
G10	UMI 1003-2-3	G38	Hyd No. 1082-2
G11	UMI 178	G39	UMI 142
G12	B. No 72183-9-2	G40	B. No 426-3
G13	B. No 1043-7	G41	B. No 71806
G14	B. No 1131-5	G42	Hyd No. 1075-4-2
G15	B. No 1421-5-1	G43	B. No 1076-5-4-3
G16	UMI 164	G44	UMI 346-1
G17	B. No 1118-3	G45	9119-1-2-1
G18	UMI 406	G46	UMI 504
G19	UMI 697-2	G47	B. No 1253-8
G20	UMI 1131-1	G48	UMI 920
G21	B. No 1075-2	G49	UMI 1098-4
G22	B. No 1917-2-1-1	G50	UMI 823
G23	UMI 1105	G51	UMI 1223
G24	B. No 1064-5	G52	UMI 1210
G25	B. No 1266-7	G53	UMI 1205
G26	UMI 1009-2-2	G54	UMI 1220
G27	B. No 1048-7	G55	UMI 1201
G28	B. No 1076-5-4-2		

genotype. The values for  $\varphi_{ij}$  and  $\eta_{ij}$  are chosen as follows: for the traits in which positive gains are desired,  $\varphi_{ni} = 0$  and  $\eta_{nj} = 100$  should be used, while for the traits in which negative gains are desired,  $\varphi_{nj} = 100$  and  $\eta_{ij} = 0$  should be used (14). Subsequently, a factor analysis (FA) was carried out to facilitate the reduction of data dimensionality and to explore the underlying relationship structure. This analysis adhered to the following model (Eqn. 2).

$$F = Z(A^T R^{-1})^T \quad (\text{Eqn. 2})$$

Where F is a  $g \times f$  matrix with the factorial score; Z is a  $g \times p$  matrix with the rescaled means; A is a  $p \times f$  matrix of canonical loading and R is a  $p \times p$  correlation matrix between the traits. Furthermore, g, f and p indicate the number of genotypes, factor retained and measured traits, respectively. In the third step, a  $[1 \times p]$  vector was considered as the ideotype matrix. The MGIDI index was calculated by calculating the Euclidean distance between the genotype scores and the ideal genotype values. The following equation (Eqn. 3) was used to achieve this calculation:

$$MGIDI_i = \sqrt{\sum_{j=1}^f (F_{ij} - F_j)^2} \quad (\text{Eqn. 3})$$

where MGIDI<sub>i</sub> is the index for the *i*th genotype; *F<sub>ij</sub>* is the score of the *i*th genotype in the *j*th factor (*i* = 1, 2, ..., *g*; *j* = 1, 2, ..., *f*), being *g* and *f* the number of genotypes and factors, respectively and *F<sub>j</sub>* is the *j*th score of the ideotype. The genotype with the lowest MGIDI is then closer to the ideotype and therefore should present desired values for all the analyzed traits.

GRAPES, an R-based programme created by (24), was used for the ANOVA and genetic variability parameters analysis. The "metan" package inside the R programme, (25), was used for Correlation analysis and MGIDI calculations. Furthermore, Principal Component Analysis (PCA) was performed with a mix of tools and packages. GRAPES was used in conjunction with the programmes ggplot2, corplot and facto extra. These packages, used together, aided in the implementation of the PCA (26-28).

## Results

### Per se performance of the genotypes

Analysis of variance showed the presence of significant differences between all the traits indicating the presence of ample genetic variation for further exploitation (Table 2). The per se performance (Table 3 & Fig. 1) for the observed traits in the experiment revealed the following trends. Plant height ranged between 108 to 182 cm with a mean of 142.46 cm, while days to 50 % tasseling ranged between 50 to 65 days categorizing the germplasm into medium and late duration types, while the days to 50 % silking ranged between 52 to 68 days. The cob length ranged between 10.0 to 18.0 cm with an average length of 14.7 cm, while the cob diameter ranged between 3.50 cm to 8.10 cm with an average diameter of 6.40 cm. Kernel rows per cob ranged between 10 to 18 rows with a mean of 14 rows per cob, while the kernel number per row ranged between 12 to 37 kernels per row with an average of 28 kernels per row. The shelling percentage ranged from 75.0 to 82.5 % with an average of 79 %. The grain yield ranged between 42.4 g to 77.5 g with a mean yield of 61.20 g.

**Table 2.** Analysis of variance for nine characters under Randomised Block Design

Source of variance	Df	Mean sum of squares								
		PH	DFT	DFS	CL	CD	KRN	KN	SP	GY
Treatment	54	1068.09**	26.14**	26.08**	15.14**	2.403**	12.52**	107.87**	9.96*	311.78**
Replication	2	5.07	2.96	2.76	0.733	0.158	8.19	3.87	10.775	4.08
Error	108	20.663	0.97	1.43	0.549	0.537	3.7	7.05	7.32	17.945

Df: Degrees of freedom; PH = Plant height (cm); DFT = Days to 50 % flowering; DFS = Days to 50 % silking; CL = Cob length (cm); CD = Cob diameter (cm); NKR = Number of kernel rows; NK = Number of kernels per row; SP = Shelling percentage (%); GY = Grain yield (g); \*\*: Level of Significance at 5%; \*: Level of Significance at 10%

**Table 3.** Best performing maize inbred lines for yield and yield attributing traits

S. No	Genotype	GY	PH	DFT	DFS	CL	CD	NKR	NKP	SP
1	UMI 1009-2-2	77.5	143.7	57	60	17.2	7.0	18	33	79.0
2	UMI 1131-1	75.3	182.0	55	57	17.0	7.9	18	32	78.6
3	B. No 1048-7	75.0	162.2	56	59	18.0	7.1	16	37	78.9
4	B. No 1076-5-1	74.7	134.8	51	54	17.7	7.0	16	35	79.8
5	UMI 96	74.5	168.7	54	57	17.4	6.6	16	37	82.0
6	B. No 1110-8	74.5	108.0	53	55	16.5	7.0	16	32	80.2
7	9119-1-2-1	74.3	126.1	57	59	17.6	6.5	18	32	80.7
8	UMI 1201	74.1	162.4	53	56	17.0	6.2	16	37	79.4
9	Hyd No. 2009-2-2-15	73.6	118.8	58	62	17.0	6.8	16	36	79.0
10	UMI1210	73.0	158.0	52	55	16.0	7.0	16	35	79.2
Minimum		42.4	108	50	52	10.1	3.45	10	12	75
Maximum		77.5	182	65	68	18	8.05	18	37	82.5
Mean		61.20	142.46	55	58	14.75	6.40	14	28	79.00
PCV (%)		17.59	13.5	5.54	5.36	15.69	16.88	18.15	22.9	3.64
GCV (%)		16.17	13.12	5.24	4.95	14.88	12.37	12.08	20.82	1.19
h <sup>2</sup> <sub>bs</sub> (%)		84.5	94.4	89.6	85.2	89.9	53.7	44.3	82.6	10.7
GAM (%)		30.6	26.3	10.2	9.4	29.1	18.7	16.6	39	0.8

PH = Plant height (cm); DFT = Days to 50 % flowering; DFS = Days to 50 % silking; CL = Cob length (cm); CD = Cob diameter (cm); NKR = Number of kernel rows; NK = Number of kernels per row; SP = Shelling percentage (%); GY = Grain yield (g)

Genetic variability

The components of genetic variability such as PCV, GCV and Broad-sense heritability are given in Table 3 and discussed below.

Components of genetic variance

PCV is higher than the GCV indicating the presence of environmental effect in the trait expression. Further, high values for both PCV and GCV were exhibited by Kernel number per row. Kernel row number, grain yield, cob diameter, cob length and plant height exhibited moderate values for both PCV and GCV, while days to 50 % tasseling, days to 50 % silking and shelling percentage showed low PCV and GCV values. The magnitude of difference between the PCV and GCV estimates can be used to determine the amount of variance that occurs due to the environment.

Broad sense heritability (h2bs) and Genetic Advance as percent of Mean (GAM)


Plant height, cob length, kernels per row and grain yield exhibited high heritability coupled with high genetic advance. Days to 50 % tasseling and days to 50 % silking exhibited high heritability with moderate GAM. Cob diameter and kernel row number exhibited moderate values for both heritability and genetic advancement. While shelling percentage exhibited low heritability and genetic advance.

Correlation

Correlation analysis for the traits studied revealed various significant trait associations (Fig. 2). Days to 50 % tasseling and days to 50 % silking showed a high positive significant correlation (0.97) between each other. Yield is a trait that is influenced by multiple dependent factors, hence an analysis of the trait association with the Grain yield reveals the traits



Fig. 1. Best performing maize inbred lines.



PH - Plant height (cm); DFT - Days to 50 % flowering; DFS - Days to 50 % silking; CL - Cob length (cm); CD - Cob diameter (cm); NKR - Number of kernel rows; NK - Number of kernels per row; SP - Shelling percentage (%); GY - Grain yield (g)

Fig. 2. Correlation analysis for all the nine traits under study.



to be improved to achieve higher yields. In this study, the grain yield was found to have a significant positive association with Plant height (0.28), cob length (0.72), cob diameter (0.60), number of kernel rows (0.53) and number of kernels per row (0.73). Hence selection for these traits can be utilized to improve grain yield.

### Principal component analysis

The Principal Component Analysis (PCA) conducted in this study effectively partitioned the total variability into nine principal components, where three components displayed eigenvalues surpassing one, signifying their substantial contribution in explaining the dataset's variance as evidenced in the scree plot (Fig. 3a) (29). Collectively, these three principal components accounted for a noteworthy 74.44 % of the overall variance (Table 4). Notably, the first PC emerged as the most influential, explaining 37.37 % of the total variance, followed by the second (22.89 %) and the third (14.17 %) PCs. The identified trait contributions delineated distinct patterns: the first PC was characterized by traits associated with grain yield and cob characteristics, except the number of kernel rows, while the second PC predominantly encapsulated flowering traits. The third PC prominently featured shelling percentage and plant height, elucidating their influence on this component. The trait number of kernel rows was the major contributor to the variance explained by PC6 (Fig. 3b).


PCA biplot exhibits the relation between the traits and between the genotypes (Fig. 4). The acute angle between two traits reveals the positive association between them, while an angle less than or more than 90 ° indicates a negative association. The flowering traits are in high positive correlation with each other, while the yield and cob traits such as cob length, cob diameter, kernel row number and kernels per row showed positive associations amongst them. Based on their scatter position on the biplot, the genotypes UMI 1003-2-3 (G10), B. NO 1421-5-1 (G15), B. NO 1917-2-1-1 (G22), G3, B. NO 1125-7 (G31), UMI 920 (G48) are placed the farthest in the biplot indicating the diverse nature of these genotypes.

### Multi-trait Genotype Ideotype Index

Initially, two ideotypes were defined with varying maturity i.e. medium and late, maturing with high yield. The MGIDI distances to select the genotype were calculated for each ideotype. A selection intensity of 15 % is opted for the selection criteria.

### Medium maturing ideotype

The selection gains (Table 4) were positive for all the traits with kernels per row having the highest gain at 18.60 % followed by grain yield at 18.30 % and cob length (14.80 %). Shelling percentage has the least gains at 0.07 %. Cumulative selection gain was recorded at 70.21 %. Eight genotypes were selected at 15 % SI, which are B. NO 1265-6-2 (G8), UMI 1210



**Fig. 3.** (a) Scree Plot for all principal components; (b) Contribution of different traits to the variance explained by each principal component.

**Table 4.** Factor analysis and Multi-Trait genotype ideotype distance index for two ideotypes

Trait	Factor	Eigen Value	PVE	CPVE	SG <sub>perc</sub>		Genotypes selected			
					Medium	Late	Medium	MGIDI	Late	MGIDI
CL	FA1	3.36	37.37	37.37	14.80	12.40	G8	1.10	G10	0.71
CD					5.98	5.78	G52	1.59	G35	2.44
KRN					7.40	5.99	G33	1.69	G26	2.50
NKP					18.60	18.80	G55	1.73	G42	2.61
GY					18.30	17.00	G49	1.98	G27	2.81
DFT	FA2	2.06	22.89	60.26	1.47	4.47	G36	2.12	G28	3.00
DFS					1.23	4.59	G24	2.12	G45	3.07
PH					2.36	-5.31	G53	2.17	G37	3.21
SP	FA3	1.28	14.17	74.44	0.07	0.03				
Cumulative selection gain					70.21	69.06				
Negative selection gain						-5.31				

PH - Plant height (cm); DFT - Days to 50 % flowering; DFS - Days to 50 % silking; CL - Cob length (cm); CD - Cob diameter (cm); NKR - Number of kernel rows; NK - Number of kernels per row; SP - Shelling percentage (%); GY - Grain yield (g); PVE - Percent variance explained; CPVE - Cumulative percent variance explained

(G52), UMI 96 (G33), UMI 1201 (G55), UMI 1098-4 (G49), B. No 1110-8 (G36), B. NO 1064-5 (G24) and UMI 1205 (G53). The genotype B. NO 1265-6-2 (G8) was the closest with the least MGIDI value of 1.10 followed by UMI 1210 (G52) (1.59) and UMI 96 (G33) (1.69) (Table 4 & Fig. 5).


### Late maturing ideotype

All the traits showed positive selection gains (Table 4) ranging from 18.80 % for Kernel number to 0.03 % for shelling percentage, except for plant height which showed a negative selection gain (-5.31). A cumulative selection gain was recorded at 69.06 % and a negative selection gain of -5.31 %.

Similarly, eight genotypes were selected based on SI of 15 % UMI 1003-2-3 (G10) HYD NO. 2009-2-2-15 (G35), UMI 1009-2-2 (G26), Hyd No. 1075-4-2 (G42), B. NO 1048-7 (G27), B. NO 1076-5-4-2 (G28), 9119-1-2-1 (G45), B. NO 1076-5-4-1 (G37) with UMI 1003-2-3 (G10) having the closest resemblance to the proposed ideotype based on its MGIDI values (0.71) (Table 4 & Fig. 5).


### Strength and weakness view of selected genotypes

The radar plot depicts the strengths and weaknesses of the selected genotypes for various maturity periods (Fig. 6). Smaller proportions explained by a factor that is placed



PH - Plant height (cm); DFT - Days to 50% flowering; DFS - Days to 50% silking; CL - Cob length (cm); CD - Cob diameter (cm); NKR - Number of kernel rows; NK - Number of kernels per row, SP - Shelling percentage (%); GY - Grain yield (g).

**Fig. 4.** PCA Biplot for both traits and genotypes.



A) MGIDI view for Late maturing ideotype; B) MGIDI view for medium maturing ideotype.

**Fig. 5.** Multi-trait Genotype Ideotype Distance Index for late and medium maturing ideotypes.

closer to the external edge indicate that the trait within that factor is more similar to the ideotype (18). A view on strength and weakness under medium maturity showed that the genotypes B. NO 71806 (G41) and G56 showed strengths related to factor 1 which holds yield and yield attributing traits, whereas B. NO 1064-5 (G24) and B. NO 1265-6-2 (G8) showed strengths for the factor 2 which holds flowering traits (DFT, DFS) and UMI 96 (G33), UMI 1098-4 (G49) and B. NO 1265-6-2 (G8) showed strengths in factor 3 which holds SP and PH.

Similarly, for late maturity, the genotypes 9119-1-2-1 (G45) and B. NO 1048-7 (G27) showed strength in FA1, while UMI 1003-2-3 (G10) showed strength in FA2 and B. NO 1076-5-4-1 (G37), B. NO 1076-5-4-2 (G28) and UMI 1009-2-2 (G26) showed strength in FA3.


## Discussion

A successful maize breeding program relies on the amount of genetic variability present in the available germplasm. The wider presence of genetic variability aids in the selection of desirable genotypes. The estimates of PCV and GCV explain the amount of variation present as expressed by either the genotype or the environment. In this study, high values for both PCV and GCV were exhibited by kernel number per row. Kernel row number, grain yield, cob diameter, cob length and plant height exhibited moderate values for both PCV and GCV. The magnitude of difference between the PCV and GCV estimates can be used to determine the amount of variance that occurs due to the environment. These results were supported by the findings of other researchers who reported moderate phenotypic and genotypic coefficient of variation for the number of kernels per row, number of kernel rows per cob, cob length, cob girth and grain yield per plant (30–34). Thus, selection for these traits could bring about an improvement in the yield attributes of maize.

Estimates of genetic variability coupled with heritability and genetic advance estimates aid us in selection and improvement programs to obtain desirable genetic gains (35). The values of heritability and genetic advance as mean, together can be used to determine the mode of gene action that is underlying the expression of the particular trait. A high heritability and GAM value indicate additive gene action, while a high heritability and low GAM percent indicate non-additive gene action. Low heritability and high GAM values indicated additive gene action but the heritability value could be due to environmental influence. Low heritability and GAM values show non additive gene action and also the prevalence of environmental influence in the manifestation of the character under study. In this study, traits like plant height, cob length, kernels per row and yield are influenced by additive gene action, so selection for improvement of these traits can be followed to achieve desired gains. Shelling percentage is based on its heritability and GAM values; it can be deduced that the underlying gene action controlling its expression is of a non-additive type. Many previous studies reported similar results where they reported high heritability and genetic advance for the number of kernels per row, no of kernel rows/cob, cob length, cob girth and grain yield per plant (31, 36-38).

Traits that contribute to yield are termed as yield attributing traits and these are generally in positive association with the yield. Positive, significant and high associations can be observed between cob length, cob diameter, number of kernel rows and number of kernels per row with yield (39-42). Hence, selection for improvement of these traits can be used to achieve yield improvement.

The study utilized Principal Component Analysis (PCA) to decompose total variability into nine components, where three components, accounting for 74.44 % of variance, exhibited eigenvalues > 1. Trait contributions highlighted distinct patterns:



A) Strength and weakness view of genotypes selected for late maturing ideotype; B) Strength and weakness view of genotypes selected for medium maturing. Ideotype.

**Fig. 6.** Strength and weakness view of genotypes selected for late and medium maturing ideotype.

PC1 correlated with grain yield and cob traits; PC2 predominantly linked to flowering traits, while PC3 emphasized shelling percentage and plant height while the variation captured by PC6 was explained by the number of kernel rows. Similar work has been done where they reported principal components with an Eigenvalue of more than 1 accounted for 70.56 % of the overall variations (43-45). PCAs displayed that the total genetic diversity was distributed throughout the agromorphological traits and the identification of a high level of genetic diversity during the current study could be implied for maize germplasm characterization, conservation and further improvement in maize breeding.

Varimax rotation reinforced the 'number of kernel rows' association with PC1. High commonality values (98 % to 53 %) indicated substantial collective variance explanation for traits. The placement of the genotypes far from the origin revealed their diversity compared to other genotypes. The genotypes UMI 1003-2-3 (G10), B. NO 1421-5-1 (G15), B. NO 1917-2-1-1 (G22), G3, B. NO 1125-7 (G31) and UMI 920 (G48) were placed in both extremes of the biplot, hence explaining their diverse nature. The genotypes can be employed for their utilization in breeding programs.

The identification of medium and late-maturing genotypes is crucial for producing hybrids preferred in specific Indian regions, emphasizing the importance of selecting genotypes with targeted traits in breeding programs. The MGIDI index offers a clear and effective selection method with practical applications for long-term genetic improvement. Assessing strengths and weaknesses using this approach provides a valuable tool to pinpoint areas for trait enhancement, distinguishing it from other indices. For instance, genotype UMI 1003-2-3 (G10) excels in flowering traits aligning with the proposed late maturing ideotype, while B. NO 1265-6-2 (G8), despite underperforming in flowering traits, suggests potential for improving yield traits while maintaining medium maturity. This approach was utilized in earlier studies in which they used different crops to select genotypes that closely resemble the ideotypes suited for their breeding objective (16, 46-48).

## Conclusion

The presence of genetic variability is a requisite for any successful breeding program. Successful exploitation of the available genetic variability depends on the efficiency of selection based on the breeding objectives. Genotypes UMI 1009-2-2, UMI 1131-1, B. No 1048-7, B. No 1076-5-1, UMI 96, B. No 1110-8, 9119-1-2-1, Hyd No. 2009-2-2-15 and UMI 1210 showed higher yield amongst other genotypes. Assessment of the association of traits provides an understanding, of trait interrelationships which can aid in further improvement. Every breeding objective has an ideotype around which the objective is focused. Selection for single traits may increase the gains of that particular trait, but in process, there is a chance of overlooking complex interactions between the studied traits. MGIDI is a comprehensive approach that enables the simultaneous selection of genotypes based on multiple traits. Cumulative positive selection gains obtained for medium maturing ideotype was 70.21 %, while it was

69.06 % for late maturing ideotype with grain yield and number of kernels per row contributing the highest selection gains. Negative selection gains were observed in plant height for late maturing ideotype. At a selection intensity of 15 %, eight genotypes were selected which lie closer to the ideotype. Hence, the utilization of a multi trait-based selection index can enable the effective selection of genotypes without any loss of variability.

## Acknowledgements

We gratefully thank our Maize unit, Department of Millets, Centre for Plant Breeding and Genetics, TNAU, Coimbatore for carrying out the field experiment.

## Authors' contributions

NK carried out the research work and drafted the manuscript, SS was involved in the selection of promising genotypes and MH carried out the statistical analysis.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interest to declare.

**Ethical issues:** None

## References

- Shikha K, Shahi JP, Vinayan MT, Zaidi PH, Singh AK, Sinha B. Genome-wide association mapping in maize: status and prospects. 3 Biotech. 2021;11(5):244. <https://doi.org/10.1007/s.13205-021-02799-4>
- Rabinowicz PD, Bennetzen JL. The maize genome as a model for efficient sequence analysis of large plant genomes. Curr Opin Plant Biol. 2006;9(2):149–56. <https://doi.org/10.1016/j.pbi.2006.01.015>
- Gálvez Ranilla LFM. The Application of metabolomics for the study of cereal corn (*Zea mays* L.). Metabolites. 2020;10(8):300. <https://doi.org/10.3390/metabo10080300>.
- United States. U.S. National Agricultural Statistics Service NASS. United States; [Web Archive] 2023 Retrieved from the Library of Congress. <https://www.loc.gov/item/lcwaN0019059>.
- Ren Y, Li Q, Du X, Zhang Y, Wang H, Shi G, Wei M. Analysis of corn yield prediction potential at various growth phases using a process-based model and deep learning. Plants. 2023;12(3):446. <https://doi.org/10.3390/plants12030446>.
- Datta MH, Yadu S, Nandedkar K, Rawte S, Saxena RR. Exploration of potential donors for machine-amenable traits in desi chickpea (*Cicer arietinum* L.): towards enhancing agricultural automation and efficiency. Electron J Plant Breed. 2023;14(3):1246–54. <https://doi.org/10.37992/2023.1403.130>
- Adhikari BN, Shrestha J, Dhakal B, Joshi BP, Bhatta NR. Agronomic performance and genotypic diversity for morphological traits among early maize genotypes. Int J Appl Biol. 2018;2(2):33–43. <https://doi.org/10.20956/ijab.v2i2.5633>
- Hadi BH, Hassan WA. Evaluating the performance of introduced varieties of maize (*Zea mays* L.) and estimating some genetic parameters. Int J Agric Stat Sci. 2021;17(1):85–91. <https://connectjournals.com/03899.2021.17.85>
- Pavlov J, Delic N, Markovic K, Crevar M, Camdzija, Stevanovic M. Path analysis for morphological traits in maize (*Zea mays* L.).



- Genetika. 2015;47(1):295–301. <https://doi.org/10.2298/GENSRI501295P>
10. Guei RG, Sanni KA, Fawole AFJ. Genetic diversity of rice (*Oryza sativa* L.). Agron Afr. 2005;5:17–28.
  11. Das S, Das SS, Chakraborty I, Roy N, Nath MK, Sarma D. Principal component analysis in plant breeding. Biomol Rep-Int eNewsletter. 2017;BR/09/17/03.
  12. Rocha JRASC, Machado JC, Carneiro CS. Multi-trait index based on factor analysis and ideotype-design: proposal and application on elephant grass breeding for bioenergy. GCB Bioenergy. 2018;10:52–60. <https://doi.org/10.1111/gcbb.12443>
  13. Céron-Rojas JJ, Crossa J. Linear selection indices in modern plant breeding. Springer Nature. 2018; p. 256. <https://doi.org/10.1007/978-3-319-91223-3>
  14. Olivoto T, Nardino M. MGIDI: toward an effective multivariate selection in biological experiments. Bioinformatics. 2020. <https://doi.org/10.1093/bioinformatics/btaa981>
  15. Pour-Aboughadareh A, Sanjani S, Nikkiah-Chamanabad H, Mehrvar MR, Asadi A, Amini A. Identification of salt-tolerant barley genotypes using multiple-traits index and yield performance at the early growth and maturity stages. Bull Natl Res Cent. 2021;45(1):1–16. <https://doi.org/10.1186/s42269-021-00576-0>
  16. Klein LA, Marchioro VS, Toebe M, Olivoto T, Meira D, Meier C, et al. Selection of superior black oat lines using the MGIDI index. Crop Breed Appl Biotechnol. 2023;23. <https://doi.org/10.1590/1984-70332023v23n3a25>
  17. Ahsan AFMS, Alam Z, Ahmed F, Akter S, Khan MAH. Selection of waterlogging tolerant sesame genotypes (*Sesamum indicum* L.) from a dataset using the MGIDI index. Data Brief. 2024;110176. <https://doi.org/10.1016/j.dib.2024.110176>
  18. Singamsetti A, Zaidi PH, Seetharam K, Vinayan MT, Olivoto T, Mahato A, Shikha K. Genetic gains in tropical maize hybrids across moisture regimes with multi-trait-based index selection. Front Plant Sci. 2023;14:1147424. <https://doi.org/10.3389/fpls.2023.1147424>
  19. Burton GW, Devane E. Quantitative inheritance in grass pea. Proc 6th Int Grassland Cong. 1953;1:277–83.
  20. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variations in Soybean. Agron J. 1955;47:314–18. <https://doi.org/10.2134/agronj1955>
  21. Miller PA, Williams JE, Robinson HF, Comstock RE. Estimates of variance and co-variance in upland cotton and their implications in selection. Agron J. 1958;50:126–31. <https://doi.org/10.2134/agronj1958.00021962005000030004x>
  22. Massey WF. Principal components regression in exploratory statistical research. J Am Stat Assoc. 1965;60:234–46. <https://doi.org/10.1080/01621459.1965.10480787>
  23. Jolliffe IT. Principal component analysis. Springer Series in Statistics. 1986.. <https://doi.org/10.1007/978-1-4757-1904-8>.
  24. Gopinath PP, Parsad R, Joseph B, Adarsh VS. GRAPES: General Rshiny based analysis platform empowered by Statistics. Version 1.0.0. 2020. <https://doi.org/10.5281/zenodo.4923220>
  25. Olivoto T, Lúcio ADC. Metan: An R package for multi-environment trial analysis. Methods Ecol Evol. 2020;11(6):783–89. <https://doi.org/10.1111/2041-210X.13384>.
  26. Wickham H. ggplot2: Elegant graphics for data analysis. Springer-Verlag New York; 2016 ISBN 978-3-319-24277-4. <https://ggplot2.tidyverse.org>
  27. Wei T, Simko V. R package 'corrplot': Visualization of a correlation matrix. Version 0.92;2021. <https://github.com/taiyun/corrplot>.
  28. Kassambara A, Mundt F. Factoextra: Extract and visualize the results of multivariate data analyses. R Package Version 1.0.7. 2020. <https://CRAN.R-project.org/package=factoextra>.
  29. Venujayakanth B, Dudhat AS, Swaminathan B, Anurag ML. Assessing crop genetic diversity using principal component analysis: A review. Trends Biosci. 2017;10(2):523–28.
  30. Netaji SVSRK, Satyanarayana E, Suneetha V. Heterosis studies for yield and yield component characters in maize (*Zea mays* L.). Andhra Agric J. 2000;47:39–42.
  31. Nagabhushan NM, Mallikarjuna CH, Shashibhaskar MS, Prahalada GD. Genetic variability and correlation studies for yield and related characters in single cross hybrids of maize (*Zea mays* L.). Curr Bio. 2011;5:157–63.
  32. Kapoor R, Batra C. Genetic variability and association studies in maize (*Zea mays* L.) for green fodder yield and quality traits. Electron J Plant Breed. 2015;6(1):233–40. <https://www.ejplantbreeding.org/index.php/EJPB/article/view/654>
  33. Kandel BP, Sharma BK, Sharma S, Shrestha J. Genetic variability, heritability and genetic advance estimates in maize (*Zea mays* L.) genotypes in Nepal. Agricultura. 2018;107(3-4):29–35.
  34. Prakash R, Ravikesavan R, Vinodhana NK, Senthil A. Genetic variability, character association and path analysis for yield and yield component traits in maize (*Zea mays* L.). Electron J Plant Breed. 2019;10(2):518–24. <https://doi.org/10.5958/0975-928X.2019.00065.6>
  35. Prasanna BM. Diversity in global maize germplasm: Characterization and utilization. J Biosci. 2012;37:843–55. <https://doi.org/10.1007/s12038-012-9227-1>.
  36. Yadav PK, Tripathi MK, Tiwari S, Chauhan S, Tripathi N, Sikarwar RS, et al. Genetic components and variability assessment for grain yield and its accrediting traits in maize (*Zea mays* L.). Intern J Environ and Climate Change. 2023;13(9):772–84. <https://doi.org/10.9734/ijec/2023/v13i92298>.
  37. Sofi PA, Rather AG, Dar Z. Association of heterotic expression for grain yield and its component traits in maize (*Zea mays* L.). Int J Agric Res. 2007;2(5):500–03. <https://doi.org/10.3923/ijar.2007.500.503>
  38. Panwar LL, Mahawar RK, Narolia RS. Genetic variability and interrelationships among grain yield and yield components in maize. Ann Plant Soil Res. 2013;15(1):15–18.
  39. Rahman M, Hoque A, Hossain MA, Al Bari MA. Variability and trait association analyses in maize (*Zea mays* L.) genotypes. Agriculturist. 2017;15(2):101–14. <https://doi.org/10.3329/agric.v15i2.35473>.
  40. Rafiq CM, Rafique M, Hussain A, Altaf M. Studies on heritability, correlation and path analysis in maize (*Zea mays* L.). J Agric Res. 2010;48(1):35–38.
  41. Munawar M, Shahbaz M, Hammad G, Yasir M. Correlation and path analysis of grain yield components in exotic maize (*Zea mays* L.) hybrids. Int J Sci Basic Appl Res. 2013;12(1):22–27.
  42. Lakshmi MS, Jagadev PN, Das S, Lenka D, Swain D, Tripathy SK. Genetic variability and association analysis of maize hybrids under excessive soil moisture condition. Int J Curr Microbiol App Sci. 2018; 7(9):2935–41. <https://doi.org/10.20546/ijcmas.2018.709.365>
  43. Rai R, Khanal P, Chaudhary P, Dhital R. Genetic variability, heritability and genetic advance for growth, yield and yield-related traits in maize genotypes. J Agric Appl Biol. 2021;2(2):96–104. <https://doi.org/10.11594/jaab.02.02.04>
  44. Okporie EO. Characterization of maize (*Zea mays* L.) germplasm with principal component analysis. Agro-Sci. 2008;7(1):66–71. <https://doi.org/10.4314/as.v7i1.1587>
  45. Iqbal J, Shinwari ZK, Rabbani MA. Maize (*Zea mays* L.) germplasm agro-morphological characterization based on descriptive, cluster and principal component analysis. Pak J Bot. 2015;47:255–64.
  46. Bhusal TN, Lal GM, Marker S, Synrem GJ. Discrimination of maize (*Zea mays* L.) inbreds for morpho-physiological and yield traits by D2 statistics and principal component analysis (PCA). Asian J Bio Sci. 2016;11(1):77–84. <https://doi.org/10.15740/HAS/AJBS/11.1/77-84>

47. Shirzad A, Asghari A, Zali H, Sofalian O, Chamanabad MH. Application of the multi-trait genotype-ideotype distance index in the selection of top barley genotypes in the warm and dry region of Darab. *J Crop Breed*. 2022;14(44):65–76. <https://doi.org/10.52547/jcb.14.44.65>
48. Palaniyappan S, Ganesan KN, Manivannan N, Ravichandran V, Senthil N. Multi-trait genotype-ideotype distance index-A tool for the identification of elite parental inbreds for developing heterotic hybrids of fodder maize (*Zea mays* L.). *Electron J Plant Breed*. 2023; 14(3):841–49. <https://doi.org/10.37992/2023.1403.098>

#### Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonpublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonpublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc  
See [https://horizonpublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

**Publisher information:** Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.