



RESEARCH ARTICLE

# Impact of drip fertigation on nutrient use efficiency, dynamics, water productivity and bulb yield in *Aggregatum* onion

S K Natarajan<sup>1</sup>, T Sampathkumar<sup>2\*</sup>, J Bhuvaneswari<sup>3</sup>, M Manikandan<sup>4</sup> & S Thenmozhi<sup>5</sup>

<sup>1</sup>Department of Agronomy, Tapioca and Castor Research Station, Yethapur 636 119, Salem, Tamil Nadu, India

<sup>2</sup>Department of Agronomy, Agricultural College and Research Institute, Madurai 625 104, Tamil Nadu, India

<sup>3</sup>Department of Agronomy, VOC Agricultural College and Research Institute, Killikulam 628 252, Vallanad, Tamil Nadu, India

<sup>4</sup>Department of Soil and Water Conservation, Agricultural Research Station, Kovilpatti 628 501, Tuticorin, Tamil Nadu, India

<sup>5</sup>Department of Soil Science and Agricultural Chemistry, ICAR-KVK, Woodhouse farm, Dodabetta, Ooty 643 002, The Nilgiris Tamil Nadu, India

\*Email: [sampath.t@tnau.ac.in](mailto:sampath.t@tnau.ac.in)



## ARTICLE HISTORY

Received: 22 November 2024

Accepted: 20 December 2024

Available online

Version 1.0 : 29 December 2024

Version 2.0 : 01 January 2025



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Natarajan SK, Sampathkumar T, Bhuvaneswari J, Manikandan M, Thenmozhi S. Impact of drip fertigation on nutrient use efficiency, dynamics, water productivity and bulb yield in *Aggregatum* onion. Plant Science Today. 2025; 12(1): 1-7. <https://doi.org/10.14719/pst.5937>

## Abstract

A field experiment was conducted on aggregatum onion (*Allium cepa* L. var. *aggregatum* Don.) under a drip irrigation system to evaluate the water productivity, nutrient use efficiency, nutrient dynamics and yield of onion in order to optimize the best fertigation level. The experiment was designed in a Randomized Block Design (RBD), with treatments replicated 3 times. The treatment details are as follows: T<sub>1</sub>-100% nitrogen (N) through fertigation with 100% phosphorus (P) and potassium (K) applied as basal; T<sub>2</sub>- 100% N and K through venturi as fertigation and basal application of 100% P; T<sub>3</sub>- 100% N, P and K through fertigation; T<sub>4</sub>- 75% N, P and K through fertigation with basal application of 25% N, P and K; T<sub>5</sub>- 75% N through fertigation with 75% P and K applied as basal; T<sub>6</sub>-75% N and K through venturi as fertigation with basal application of 75% P; T<sub>7</sub>- 75% N, P and K through fertigation; T<sub>8</sub>- 75% of N, P and K through fertigation and 25% of 75 % N, P and K applied as basal; T<sub>9</sub>- 50 % N 100 % P and K applied as basal and 50% N applied on 30 DAT; T<sub>10</sub>- absolute control (no fertilizer). The results indicated that the bulb yield of aggregatum onion was significantly higher in 100% N, P and K through fertigation (23.24 to 23.47 t ha<sup>-1</sup>). Among the different fertigation treatments, the 100% N, P and K through fertigation recorded higher water use efficiency (75.08 to 78.44 kg/hamm), water productivity (7.841 to 7.508 kg m<sup>-3</sup>) and nutrient use efficiency, which was comparable to the 75% N, P and K through fertigation.

## Keywords

aggregatum onion; drip fertigation; nutrient dynamics; nutrient use efficiency; water use efficiency

## Introduction

Small aggregatum onion (*Allium cepa* L. var. *aggregatum* Don.) is a popular vegetable in south India and produced commercially in Tamil Nadu. Tamil Nadu is one of the major onion producing states with production of 3.01 lakh tonnes in an area of 0.28 lakh ha. Small onion occupies nearly 75% of the total onion produced in India with a productivity of 12 t ha<sup>-1</sup> (1). However, productivity remains low in India, mainly due to lack of scientific management of irrigation water and nutrients which determined the yield of the crop. Its production is rapidly increasing under irrigated conditions. The area under onion cultivation in the country is increasing every year, mainly

due to its high profitability. However, the average yield is low due to insufficient information on proper fertilizer and irrigation schedules and lack of appropriate technology. This can be improved through the use of micro-irrigation systems, which increase water and nutrient use efficiency compared to conventional irrigation systems.

In India, the average productivity of aggregatum onion is about 16.41 t ha<sup>-1</sup>, which is lower than world's average of 20.08 t ha<sup>-1</sup> which needs crop production strategies to increase the productivity of the crop (1). One of the reasons for this low productivity is the surface irrigation method predominantly adopted by farmers, which results in low productivity. Onion, a shallow-rooted crop, requires frequent irrigation, but surface irrigation methods result in low water use efficiency. By adopting a drip irrigation system, high water use efficiency and better productivity can be achieved by maintaining moisture near the root zone at field capacity. Drip irrigation resulted in water savings of up to 39%, with 520 mm of water applied compared to 840 mm for surface irrigation (2). The field water use efficiency of drip irrigation was 2.5 times higher when compared to conventional methods. Given the limited research on onion cultivation using drip fertigation, this study aims to optimize drip fertigation techniques to improve both water and nutrient use efficiency, ultimately boosting onion yields.

## Materials and Methods

An on-farm research experiment was conducted on aggregatum onion (*Allium cepa* L. var. *aggregatum* Don.) under a drip irrigation system to study the growth, yield and economics of onion cultivation under different fertigation treatments and to optimize the best fertigation level. The study was carried out at the Agriculture Research Station, Bhavanisagar, during 2017-2018 and 2018-2019 growing seasons. The experimental field's soil was sandy clay loam in texture. Soil samples were collected at a depth of 15cm prior to the experiment and analyzed for various physico-chemical characteristics. The results indicated that the soil was low in available nitrogen (N) (250 kg ha<sup>-1</sup>), high in available phosphorus (P) (24 kg ha<sup>-1</sup>) and medium in available potassium (K) (352 kg ha<sup>-1</sup>). The on-farm research experiment was laid out in a Randomized Block Design (RBD) with 3 replications and a plot size of 22.5 m × 0.9 m.

The treatment details are as follows

T <sub>1</sub>	100% N through venturi with 100% basal application of P and K
T <sub>2</sub>	100% N and K through venturi with 100% basal application of P
T <sub>3</sub>	100% N, P and K through fertigation
T <sub>4</sub>	75% N, P and K through fertigation and basal application of 25% N, P and K
T <sub>5</sub>	75% N through fertigation with 75% P and K applied as basal
T <sub>6</sub>	75% N and K through venturi as fertigation and basal application of 75% P
T <sub>7</sub>	75% N, P and K through fertigation
T <sub>8</sub>	75% of 75% N, P and K through fertigation and 25% of 75% N, P and K applied as basal
T <sub>9</sub>	50% N 100% P and K applied as basal and 50% N applied on 30 DAT
T <sub>10</sub>	Absolute control (No Fertilizer)

Irrigation water was pumped using a 7.5 HP motor and water was conveyed to the main line, made up of PVC pipes with 63 mm OD (outer diameter), after filtering through sand and disc filters. A venturi system was installed in the mainline for fertigation. Sub-mains, constructed with PVC pipes of 40 mm OD, were connected to the main pipes. Lateral lines made of 12 mm LDPE pipes were spaced at 1.2 m intervals, with inline emitters positioned 50 cm apart, discharging at a rate of 4 Litre per hour (L/hr). Individual control valves were installed for each treatment to facilitate different fertigation applications. A single lateral line was used for each bed, which contained four rows of onion crops. The wetting diameter of a single dripper was 0.5 m. The sub-mains and laterals pipes were sealed with end caps. After installation, a test run was conducted to measure the mean dripper discharge and irrigation efficiency. Based on the observations, irrigation time was determined. During the cropping period, the average irrigation efficiency was recorded as approximately 90%.

## Drip irrigation

The first irrigation was applied before transplanting and subsequent irrigations were scheduled on alternate days using the daily pan evaporation values. Irrigation was applied at 100% Crop Evapotranspiration (ET<sub>c</sub>). The water requirement (WR) was calculated as follows

$$WR \text{ (mm/irrigation)} = \frac{PE \times K_p \times K_c}{IE} \times 100$$

$$WR \text{ (L/unit area/irrigation)} = \frac{PE \times K_p \times K_c \times A}{IE} \times 100$$

where,

PE = Pan evaporation (mm)

K<sub>p</sub> = Pan co-efficient (0.70)

K<sub>c</sub> = Crop co-efficient (3)

Initial stage (1-20 DAT) 0.70

Mid-season stage (21-60 DAT) 11.05

Late season stage (61-80 DAT) 0.80

A = Area (m<sup>2</sup>)

IE = Irrigation efficiency (90%)

The operation time of the drip irrigation system to deliver the required quantity of irrigation water was calculated using the formula,

Time of application (min) =

$$\frac{\text{Volume of required water (L / plot)} \times 60}{\text{Emitter discharge (L hr}^{-1}\text{)} \times \text{No. of emitters/ plot}}$$

Fertigation was scheduled based on the soil test crop response (STCR) based fertilizer prescription equation developed for onion in the Irugur soil series. The STCR method is a technique that helps to determine the amount of fertilizers to specific location or similar soil in a particular agro-eco region (4).

### Fertilizer Prescription Equations for onion under STCR

STCR equation for requirement of major nutrients viz., NPK  
 $FN = 0.9 T - 0.37 SN$

$FP2O_5 = 0.58 T - 1.43 SP$

$FK_2O = 0.67 T - 0.44 SK$

where,

FN,  $FP_2O_5$  and  $FK_2O$  are N,  $P_2O_5$  and  $K_2O$  fertilizers (in  $kg\ ha^{-1}$ ) respectively

T is the yield target in  $q\ ha^{-1}$

SN, SP and SK are available soil N, P and K (in  $kg\ ha^{-1}$ ) respectively.

The STCR approach to fertilizer application is a scientific method used to optimize crop productivity while maintaining soil health. This approach increased the crop productivity, efficient use of fertilizer and cost effective. STCR NPK approach was used for calculation of fertilizer dose by substituting initial soil test values viz., N (245), P (24) and K (350). The calculated fertilizer dose was 120\*\*: 99: 60\*\* N,  $P_2O_5$ ,  $K_2O$ ,  $kg\ ha^{-1}$  respectively adopted for application to crop. (\*\* maximum dose 200% of RDF). The targeted yield of onion was fixed as 230  $q\ ha^{-1}$ .

The fertilizers used for drip fertigation included urea (46% N), poly feed (19:19:19), mono ammonium phosphate (12:61:0) and sulphate of potash (50% K). For basal application, conventional fertilizers such as urea (46% N), single super phosphate (16% P) and muriate of potash (60% K) were used. The required quantities of fertilizers were dissolved in water and applied from the 2<sup>nd</sup> to the 11<sup>th</sup> week after transplanting, as per the fertigation schedule (refer to Table 1).

### Nutrient use efficiency (NUE)

NUE was calculated using the following formula and expressed as yield (kg) per kg of nutrients (N, P and K) applied per ha applied.

$$NUE = \frac{\text{Yield (kg ha}^{-1}\text{)}}{\text{Nutrient applied (kg/ha)}}$$

**Table 1.** Fertigation schedule as per growth stage

Sl. No.	Crop stage	Duration (in days)	Fertigation frequency days	% requirement		
				N	P	K
1	Vegetative stage	1-30	7	30	50	20
2	Bulb formation stage	31-60	7	40	30	50
3	Bulb development stage	61-90	7	30	20	30
<b>Total</b>		<b>90</b>		<b>100</b>	<b>100</b>	<b>100</b>

### Soil nutrient dynamics under drip irrigation

Soil samples were collected from 2 horizontal points relative to emitters (0 cm and 25 cm on both sides) and 2 vertical points (0-15 cm and 15-30 cm) below the soil surface on both sides of each lateral emitter. Soil samples were collected at 60 DAT using a screw auger for three treatments:

- **T3:** 100% N, P and K through fertigation
- **T7:** 75% N, P and K through fertigation
- **T8:** 75% NPK via fertigation and 25% as basal application

The samples air dried, ground, sieved (2 mm) and stored in polythene bags for the estimation of nutrients viz., available N, P and K ( $kg\ ha^{-1}$ ). The available N, P and K were mapped in both horizontal and vertical dimensions by using SURFER-version 7.0 software (developed by Golden Software of USA) which show the three-dimensional view of soil nutrient distributions from the emitter. Surfer software was a contouring package includes 3D surface mapping program which is compatible with Microsoft Windows Operating System.

## Results and Discussion

### Nitrogen use efficiency

The application of different drip fertigation treatments significantly influenced NUE across both years (Table 2). The treatment with 75% N, P and K through fertigation ( $T_7$ ) resulted in the highest NUE (252.7 and 254.0  $kg\ ha^{-1}$ ). Moderately high NUE was noticed in treatment  $T_8$  (75% of 75% N, P and K through venturi fertigation and 25% of 75% N, P and K as basal application) (226.2 and 227.3  $kg\ ha^{-1}$ ) and  $T_6$  (75% N and K through venturi as fertigation with 75% P applied as basal) (213.3 and 214.4  $kg\ ha^{-1}$ ). Lower NUE values were recorded in treatments  $T_9$  (50% N 100% P and K applied as basal and 50% N applied on 30 DAT 50 % N 100% P and K applied as basal and 50% N applied on 30 DAT) (147.9 and 148.6  $kg\ kg^{-1}$ ),  $T_1$  (100% N through fertigation with basal application of 100% P and K; 150.9 and 151.7  $kg\ kg^{-1}$ ) and  $T_2$  (100% N and K through fertigation with basal application of 100% P) (166.8 and 167.6  $kg\ kg^{-1}$ ), indicating the importance of balanced nutrient application for optimal nitrogen use and Lower NUE values were recorded in treatment of absolute control.

### Phosphorus use efficiency

Phosphorus use efficiency (PUE) was also significantly influenced by the fertigation treatments in both the experiments (Table 2). The higher PUE was found in both the years with 75% N, P and K through fertigation ( $T_7$ ) (307.3 and 308.8  $kg\ kg^{-1}$ ). PUE was moderately increased and comparable with drip fertigation of 75% of 75% N, P and K through fertigation and 25% of basal application of 75% N, P and K applied ( $T_8$ ) (275.2 and 276.6  $kg\ kg^{-1}$ ), 75% N and K through venturi as fertigation with basal application of 75% P ( $T_6$ ) (259.4 and 260.7  $kg\ kg^{-1}$ ) and 100% N, P and K through fertigation ( $T_3$ ) (258.3 and 259.6  $kg\ kg^{-1}$ ). Distinctly lower and comparable values of PUE was recorded in the treatment in which basal application of 50% N, 100% P and K and 50% N applied on 30 DAT ( $T_9$ ) (179.2 and 180.1  $kg\ kg^{-1}$ ) and 100% N through fertigation with 100% P and K applied as basal ( $T_1$ ) (182.9 and 183.8  $kg\ kg^{-1}$ ), barring absolute control.

### Potassium use efficiency

Drip fertigation has shown significant influence on the potassium use efficiency (KUE) (Table 2) during both the years of study. The highest KUE was noticed with 75% N, P and K through fertigation ( $T_7$ ) (505.4 and 507.9  $kg\ kg^{-1}$ ) which

**Table 2.** Nutrient use efficiency (kg kg<sup>-1</sup>) of onion as influenced by drip fertigation treatments

Treatment	First experiment			Second experiment		
	N	P	K	N	P	K
T <sub>1</sub> - 100% N through fertigation with 100% P and K applied as basal	150.9	182.9	301.8	151.7	183.8	303.3
T <sub>2</sub> - 100% N and K through fertigation with 100% P applied as basal	166.8	222.4	333.7	167.6	223.5	335.4
T <sub>3</sub> - 100% N, P and K through fertigation	193.7	258.3	387.5	194.7	259.6	389.4
T <sub>4</sub> - 75% N, P and K through fertigation and 25% N, P and K applied as basal	171.9	229.2	343.8	172.8	230.3	345.5
T <sub>5</sub> - 75% N through fertigation with 75% P and K applied as basal	191.2	232.5	382.4	192.2	233.7	384.3
T <sub>6</sub> - 75% N and K through fertigation with 75% P applied as basal	213.3	259.4	426.6	214.4	260.7	428.7
T <sub>7</sub> - 75% N, P and K through fertigation	252.7	307.3	505.4	254.0	308.8	507.9
T <sub>8</sub> - 75% of 75% N, P and K through fertigation and 25% of 75% N, P and K applied as basal	226.2	275.2	452.5	227.3	276.6	454.8
T <sub>9</sub> - 50% N and 100% P, K applied as basal and 50% N applied on 30 DAT	147.9	179.2	295.8	148.6	180.1	297.3
<b>SEd</b>	<b>10.7</b>	<b>13.9</b>	<b>27.4</b>	<b>10.8</b>	<b>14.0</b>	<b>27.5</b>
<b>CD (P=0.05)</b>	<b>22.6</b>	<b>29.5</b>	<b>58.1</b>	<b>22.7</b>	<b>29.6</b>	<b>58.4</b>

was at par with 75% of 75% N, P and K through fertigation and 25% of 75% N, P and K applied as basal (T<sub>8</sub>) (452.5 and 454. kg kg<sup>-1</sup>), 75% N and K through fertigation with 75% P applied as basal (T<sub>6</sub>) (426.6 and 428.7 kg kg<sup>-1</sup>), registered moderate potassium use efficiency and was at par with 75% of 75% N, P and K through fertigation with 25% of 75% N, P and K applied as basal (T<sub>8</sub>). Substantially lower and comparable KUE was recorded in 50% N 100% P and K applied as basal and 50% N applied on 30 DAT (T<sub>9</sub>) (295.8 and 297.3 kg kg<sup>-1</sup>), 100% N through fertigation with 100% P and K applied as basal (T<sub>1</sub>) (301.8 and 303.3 kg kg<sup>-1</sup>), 100% N and K through fertigation with 100% P applied as basal (T<sub>2</sub>) (333.7 and 335.4 kg kg<sup>-1</sup>) and 75% N, P and K through fertigation with 25% N, P and K applied as basal (T<sub>4</sub>) (343.8 and 345.5 kg kg<sup>-1</sup>). Higher nutrient use efficiencies in treatments like T<sub>3</sub> (100% N, P and K through fertigation) were attributed to enhanced nutrient uptake due to minimized leaching losses and efficient root zone utilization. This corroborate the findings of another work (5). Under drip irrigation, nutrient use efficiency was higher and fertilizer saving was around 30-50% (5, 6).

#### Soil nutrient dynamics under drip fertigation

The nutrients applied at each crop stage should be readily available in the soil to ensure their absorption without any hindrance. Leaching of nutrients, volatilization of fertilizers and fixation of nutrients in the soil are some of the factors that affect the availability of nutrients in the soil. The mobility of nutrients in the soil depends on the source of fertilizers, the levels of applied fertilizers and the ionic forms of the nutrient. Soil moisture also significantly influences the availability of N, P and K in the soil. The mobility of the applied nutrients can be assessed from soil samples taken at various distances from the emitter, both horizontally (0 to 25 cm) and vertically (0-15 and 15-30 cm) at 60 DAT for mapping. The mobility of nutrients in the soil depends on the source of fertilizer, application dose, forms of nutrient ions, soil moisture and other reactive ions present in soil solution.

#### Nitrogen dynamics

Irrespective of soil depth analysed for nutrients, more available nitrogen was accumulated in periphery regions of wetting zone. Among the fertigation treatments, peak

available N was recorded in 100% N, P and K through fertigation (T<sub>3</sub>) (253.8 kg ha<sup>-1</sup>) at a distance of 25 cm away from emitter and a depth 15-30 cm soil layer. The same trend in terms of magnitude similarity was observed in other fertigation treatments also. Being mobile nutrient in soil, higher amount of nitrogen was accumulated in the periphery region of wetted zone. The mobility of available nitrogen as influenced by drip fertigation for 2 years study is depicted in Table 3 and Fig. 1. In the present investigation, more nitrogen was accumulated in the periphery region of wetted zone of soil layer 15-30 cm below the emitter. This was due to movement of nitrogen in soil solution which led to more accumulation of available nitrogen in lower layer than surface layer of soil. Under drip irrigation, high soil moisture beneath the dripper increases the availability of nitrogen in the soil (6). Similar finding was noticed as higher NO<sub>3</sub>-N concentration was found mostly in 15-30 cm soil layer (6). Nitrogen mobility in the soil occurred due to the concentration gradient, causing nutrients to move from higher concentration to lower one. The available soil moisture in the root zone plays a vital role in the mobility of nutrients (7). The study observed that higher soil moisture below the emitter corresponded to an increase in available nitrogen. A direct relationship was found between nitrogen availability and soil moisture. The nitrate ion as being highly mobile in nature, it has a tendency to move away from the emitter to the periphery of the waterfront (8, 9).

#### Phosphorus dynamics

The mobility of phosphorus as average of two experiments as influenced by drip fertigation is depicted in Table 3 and Fig. 1. The higher available P in soil was confined to top layer 0-15 cm just below the emitter and decreased thereafter under all fertigation levels. The available P was higher in drip fertigation with 100% N, P and K through fertigation (T<sub>3</sub>) (22.5 kg ha<sup>-1</sup>) than other fertigation treatments. Generally, phosphorus was immobile in soil, soil applied Pas water soluble form resulted in increased mobility of P to some distance. Among the different nutrients required by onion crop, phosphorus is one of the highly demanded nutrients for higher yield (10). The higher quantity of available P was



**Table 3.** Nutrient dynamics under drip fertigation in onion (pooled data)

Nutrient (kg/ha)	Soil depth (cm)	75% of 75% RDF			75% RDF			100% RDF		
		25 cm	Emitter	25 cm	25 cm	Emitter	25 cm	25 cm	Emitter	25 cm
<b>Nitrogen</b>	<b>0-15</b>	224.5	211.5	221.5	236.5	230.8	232.5	248.6	245.5	249.5
	<b>15-30</b>	230.8	220.5	228.6	248.7	245.5	249.5	253.8	250.4	254.2
<b>Phosphorus</b>	<b>0-15</b>	17.8	19.4	18.1	19.6	21.8	19.1	21.5	22.5	20.9
	<b>15-30</b>	15.8	17.5	15.2	17.4	18.4	17.8	18.4	21.1	18.2
<b>Potassium</b>	<b>0-15</b>	400.2	412.2	398.5	420.5	431.5	419.5	432.5	438.7	431.5
	<b>15-30</b>	394.7	401.5	392.5	415.2	421.5	412.5	427.5	431.5	424.6

RDF- recommended dose of fertilizer; \*Data not statistically analysed

observed right below the emitter and thereafter decreased both vertically and horizontally in the present study. These results are in agreement with another findings (11). It was also found from the present investigation that higher availability of phosphorus was confined to 0-15 cm of soil layer. The available P decreased with increase in horizontal distance away from emitter. Similar finding was also reported earlier (7). The restricted mobility of phosphorus in soil might be attributed to its strong reactivity, as reported earlier (12). Subsequently, frequent fertilizer applications in smaller splits within the root zone increased phosphorus availability under drip fertigation (13). Further, study by (14) supported the hypothesis that continuous application of P through drip irrigation system increased P availability when compared other application methods.


#### Potassium dynamics

The mobility of potassium influenced by drip fertigation, averaged over two experiments, is depicted in Table 3 and Fig. 1. The potassium availability was higher in the top layers. The availability of potassium varied with soil layers and distance from the emitter point. Higher potassium availability was recorded just below the emitter (0-15 cm depth). Potassium availability was more under 100% N, P and K through fertigation ( $T_3$ ) (438.7 kg ha<sup>-1</sup>) than 75% RDF at 0-15 cm depth of soil layer in the emitter point. Available potassium was decreased with increasing depth from the emitter with irrespective of the distances. Potassium is less mobile in nature than nitrate but distributed uniformly in the wetted soil volume through interactions soil particles (15). In the present study, potassium mobility varied under and across the emitter under drip fertigation, as the highest available potassium was found in soil depth of 0-15 cm. This result was in line with findings of another study (16). Due to slow downward movement of potassium might be partially attributed to net upward flux of water in soil. This corroborates the findings of another work (17). Similar finding was also reported earlier (18, 19), who observed that potassium distribution in the soil profile is characterized by decreasing potassium content with increasing soil depth, with higher potassium content noticed at the 0-15 cm soil profile. This was mainly due to slow movement of surface applied potassium in soil. Additionally, (20) also reported that the available potassium was higher in the surface layer due to entrance of potassium ions on soil exchange complex resulted in slow movement to deep soil layers.

#### Water use efficiency and Water Productivity in onion

The water uses efficiency (WUE) and water productivity of onion were highly influenced by drip fertigation during both years due to the application of fertilizers at different levels as fertigation (Table 4). WUE was higher in 100% N, P and K

applied through venturi as fertigation ( $T_3$ ) (78.44 and 75.08 kg/ha mm) and was followed by 75% N, P and K through fertigation ( $T_7$ ) (76.72 and 73.48 kg /ha mm). Conspicuously lower water use efficiency was registered in 75% N through fertigation with 75% P and K applied as basal ( $T_5$ ) (58.05 and 55.57 kg /ha mm). Application of 100% N, P and K through fertigation ( $T_3$ ) registered substantially higher water productivity (7.841 and 7.508 kg m<sup>-3</sup>) which was followed by 75% N, P and K through fertigation ( $T_7$ ) (7.670 and 7.348 kg / m<sup>3</sup>). Barring absolute control, 75% N through venturi as fertigation with basal application of 75% P and K ( $T_5$ ) (5.803 and 5.557 kg m<sup>-3</sup>) registered perceptively lower and comparable water productivity. Consumptive use of irrigation water calculated by adding the amount of irrigation water required to meet the demand of evapotranspiration and effective rainfall during the crop growing season. The data pertaining to water use indicated that a significantly higher WUE and water productivity were registered in 100%

**Fig. 1.** Nutrient dynamics under drip fertigation.

**Table 4.** Influence of drip fertigation on water use, water use efficiency (WUE) and water productivity (WP) of onion

Treatment	First experiment			Second experiment		
	*Total water use (mm)	WUE (kg/ha mm)	WP (kg/m <sup>3</sup> )	*Total water use (mm)	WUE (kg/ha mm)	WP kg m <sup>-3</sup> )
T <sub>1</sub> - 100% N through fertigation with 100% P and K applied as basal	296.4	61.08	6.106	312.6	58.48	5.848
T <sub>2</sub> - 100% N and K through fertigation with 100% P applied as basal	296.4	67.54	6.752	312.6	64.65	6.465
T <sub>3</sub> - 100% N, P and K through fertigation	296.4	78.44	7.841	312.6	75.08	7.508
T <sub>4</sub> - 75% N, P and K through fertigation and 25% N, P and K applied as basal	296.4	69.59	6.957	312.6	66.63	6.663
T <sub>5</sub> - 75% N through fertigation with 75% P and K applied as basal	296.4	58.05	5.803	312.6	55.57	5.557
T <sub>6</sub> - 75% N and K through fertigation with 75% P applied as basal	296.4	64.77	6.475	312.6	62.00	6.200
T <sub>7</sub> - 75% N, P and K through fertigation	296.4	76.72	7.670	312.6	73.48	7.348
T <sub>8</sub> - 75% of 75% N, P and K through fertigation and 25% of 75% N, P and K applied as basal	296.4	68.70	6.868	312.6	65.77	6.577
T <sub>9</sub> - 50% N and 100% P, K applied as basal and 50% N applied on 30 DAT	296.4	59.87	5.985	312.6	57.33	5.733
T <sub>10</sub> - Absolute control	296.4	46.96	4.695	312.6	44.95	4.495

Total water use includes effective rainfall also; \*data not statistically analysed

N, P and K through fertigation (T<sub>3</sub>) and 75% N, P and K through fertigation (T<sub>7</sub>). This might due higher yield (Fig. 1) and frequent application of irrigation water through drip irrigation system, which helped in maintaining the soil moisture of crop root zone near field capacity and reduced the movement of water beyond root zone. Similar findings were also reported, indicating higher water use efficiency (WUE) for daily and alternate-day drip irrigation (21, 22).

#### Effect of drip fertigation on bulb yield of onion

Application of nutrients through drip fertigation significantly influenced the bulb yield and the data are presented for 2 years in Table 5. The highest bulb yield was noticed in 100% N, P and K through fertigation (T<sub>3</sub>) (23.24 and 23.47 t ha<sup>-1</sup>) and it was comparable with 75% N, P and K through fertigation (T<sub>7</sub>) (22.74 and 22.97 t ha<sup>-1</sup>). Moderate yields were recorded in 75% N, P and K through fertigation and 25% N, P and K applied as basal (T<sub>4</sub>) (20.62 and 20.83 t ha<sup>-1</sup>) and was comparable with 75% N, P and K through fertigation (T<sub>7</sub>). The increase in yield might be due to higher number of bulbs, bulb weight due to increase in the availability of all the major nutrients (NPK) in the soil solution which in turn increased the uptake of nutrients and effective translocation of assimilates from source to sink led to higher crop yield. This was in line with the findings of a previous study which stated that application of 100% WSF as fertigation increased the fruit yield significantly over drip irrigation with soil application of

nutrients (23). Similarly, another study also reported higher onion yield in both daily and alternate day fertigation (21). Application of 75% N through venturi as fertigation and basal application of 75% P and K (T<sub>5</sub>) (17.20 and 17.37 t ha<sup>-1</sup>), 50% N, 100% P and K applied as basal and 50% N applied on 30 DAT (T<sub>9</sub>) (17.74 and 17.92 t ha<sup>-1</sup>), 100% N through fertigation with basal application of 100% P and K (T<sub>1</sub>) (18.10 and 18.28 t ha<sup>-1</sup>) and 75% N and K through fertigation with basal application of 75% P (T<sub>6</sub>) (19.19 and 19.38 t ha<sup>-1</sup>) registered significantly lower and comparable yield. The substantially lower yield observed in 75% N through fertigation with 75% P and K applied as basal (T<sub>5</sub>) might be attributed to lower nutrient absorption efficiency of granular fertilizer. When applied through broadcasting these fertilizers are prone to leaching and volatilization losses, leading to reduced availability of plant uptake. This is consistent with the findings of a previous study (24).

#### Conclusion

The conclusion drawn out from the above investigation are as follows: Drip fertigation with 100% N, P and K through venturi recorded substantially higher nutrient use efficiency, which was comparable with 75% N, P and K through fertigation. Irrespective of depth significantly higher amount of available N accumulated near the periphery region of the wetting zone. For P and K, higher concentrations were observed directly

**Table 5.** Yield attributes and yield of onion as influenced by drip fertigation treatments

Treatment	First experiment	Second experiment
	Bulb yield (t ha <sup>-1</sup> )	Bulb yield (t ha <sup>-1</sup> )
T <sub>1</sub> - 100% N through fertigation with 100% P and K applied as basal	18.10	18.28
T <sub>2</sub> - 100% N and K through fertigation with 100% P applied as basal	20.01	20.21
T <sub>3</sub> - 100% N, P and K through fertigation	23.24	23.47
T <sub>4</sub> - 75% N, P and K through fertigation and 25% N, P and K applied as basal	20.62	20.83
T <sub>5</sub> - 75% N through fertigation with 75% P and K applied as basal	17.20	17.37
T <sub>6</sub> - 75% N and K through fertigation with 75% P applied as basal	19.19	19.38
T <sub>7</sub> - 75% N, P and K through fertigation	22.74	22.97
T <sub>8</sub> - 75% of 75% N, P and K through fertigation and 25% of 75% N, P and K applied as basal	20.36	20.56
T <sub>9</sub> - 50% N and 100% P, K applied as basal and 50% N applied on 30 DAT	17.74	17.92
T <sub>10</sub> - Absolute control	13.91	14.05
<b>SEd</b>	<b>1.05</b>	<b>1.06</b>
<b>CD (P=0.05)</b>	<b>2.21</b>	<b>2.23</b>

below the emitter. Application of 100% N, P and K through venturi as fertigation and 75% N, P and K through fertigation recorded higher water use efficiency (78.44 kg/ha mm) and water productivity (7.841 kg m<sup>-3</sup>). Onion bulb yield was higher in 100% N, P and K through fertigation (23.24 t ha<sup>-1</sup>). In aggregatum onion cultivation, application of 75% N, P and K through fertigation using water soluble fertilizers in weekly intervals was identified as the optimal fertigation treatment to achieve higher nutrient use efficiency, nutrient dynamics, water use efficiency, water productivity and bulb yield. These results offer a highly efficient, eco-friendly alternative to traditional irrigation and fertilizer application in onion cultivation.

## Acknowledgements

Authors acknowledge ICAR-AICRP for providing financial support for this work.

## Authors' contributions

SKN, JB conceived idea, planned the work and carried out the experiments and statistical analysis. TS, MM and ST drafted the manuscript and corrected. All authors read and approved the final manuscript.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** None

## References

1. FAO. FAO statistical database; 2009. <http://www.fao.org/faostat/agriculture.html>
2. Dingre S, Pawar D, Kadam K. Productivity, water use and quality of onion (*Allium cepa*) seed production under different irrigation scheduling through drip. Ind J Agron. 2012;57(2):186–90. <https://doi.org/10.59797/ija.v57i2.4620>
3. FAO. Irrigation and drainage paper 56. Crop Evapotranspiration - Guidelines for Computing Crop Water Requirements; 2014 <https://www.fao.org/4/x0490e/x0490e00.htm>
4. Sellamuthu KM, Santhi R, Maragatham S, Dey P. Validation of soil test target based balanced fertilizer prescription model for rainfed maize on inceptisol. Madras Agri J. 2016;103(4-6):114–19. <https://doi.org/10.29321/MAJ.10.001002>
5. Kumar SP. Performance evaluation of drip fertigation system with water soluble fertilizers on water, fertilizer use and yield in hybrid capsicum. M. Sc. (Ag.) Thesis, Tamil Nadu Agric. Univ. Coimbatore; 2000
6. Rekha KB, Mahavishnan K. Drip fertigation in vegetable crops with emphasis on lady's finger (*Abelmoschus esculentus* (L.) Moench)- A review. Agri Rev. 2008;29(4):298–305. <https://www.cabidigitallibrary.org/doi/pdf/10.5555/20113215346>
7. Bangar A, Chaudhari B. Nutrient mobility in soil, uptake, quality and yield of Suru sugarcane as influenced by drip-fertigation in medium vertisols. J Ind Society of Soil Sci. 2004;52(2):164–71.
8. Badr AE, Abuarab ME. Soil moisture distribution patterns under surface and subsurface drip irrigation systems in sandy soil using neutron scattering technique. Irrigation Sci. 2013;31(3):317–32. <https://doi.org/10.1007/s00271-011-0306-0>
9. Li J, Zhang J, Rao M. Wetting patterns and nitrogen distributions as affected by fertigation strategies from a surface point source. Agri Water Manage. 2004;67(2):89–104. <https://doi.org/10.1016/j.agwat.2004.02.002>
10. Ryan I. Efficient use of phosphate fertilizers for sustainable crop production in WANA. IMPHOS: Phosphate Newsletter. 2002;2–5.
11. Rauschkolb R, Rolston D, Miller R, Carlton A, Burau R. Phosphorus fertilization with drip irrigation. Soil Sci Society of America J. 1976;40(1):68–72. <https://doi.org/10.2136/sssaj1976.03615995004000010021x>
12. Singh HNK, Narda, Chawla JK. Efficacy of phosphorus through trickle fertigation of potato (*Solanum tuberosum*). Ind J Agri Sci. 2004;74(8):476–78. <https://www.cabidigitallibrary.org/doi/full/10.5555/20053020981>
13. Pawar D, Dingre S, Durgude A. Enhancing nutrient use and sugarcane (*Saccharum officinarum*) productivity with reduced cost through drip fertigation in western Maharashtra. Ind J Agri Sci. 2014;84(7):844–49. <https://doi.org/10.56093/ijas.v84i7.41997>
14. Grant C, Flaten D, Tomasiewicz D, Sheppard S. The importance of early season phosphorus nutrition. Canadian J Plant Sci. 2001;81(2):211–24. <http://doi/pdf/10.4141/P00-093>
15. Kafkafi U, Bar-Yosef B, Rosenberg R, Sposito G. Phosphorus adsorption by kaolinite and montmorillonite: Organic anion competition. Soil Sci Society of America J. 1988;52(6):1585–89. <https://doi.org/10.2136/sssaj1988.03615995005200060012x>
16. Singh R, Namdeo K. Effect of fertility levels and herbicides on growth, yield and nutrient uptake of direct seeded rice (*Oryza sativa*). Ind J Agron. 2004;49(1):34–36. <https://doi.org/10.59797/ija.v49i1.5150>
17. Zeng Q, Brown PH. Soil potassium mobility and uptake by corn under differential soil moisture regimes. Plant and Soil. 2000;221(2):121–34. <https://doi.org/10.1023/A:1004738414847>
18. Mmolawa K, Or D. Root zone solute dynamics under drip irrigation: A review. Plant and soil. 2000;222(1-2):163–90. <https://doi.org/10.1023/A:1004756832038>
19. Veeranna H, Khalak A, Sujith G. Effect of fertigation and irrigation methods on yield, water and fertilizer use efficiencies in chilli (*Capsicum annuum* L.). South Indian Horticulture. 2001;49:101–04.
20. Suganya S, Anitha A, Appavu K. Moisture and nutrient distribution system under drip fertigation systems. In: 3<sup>rd</sup> International Ground Water Conference on “Water, Environment and Agriculture- Present Problems and Future Challenges 2007; 7th to 10th February 2007.
21. Ankush, Singh V, Sharma SK. Response of tomato (*Solanum lycopersicum* L.) to fertigation by irrigation scheduling in drip irrigation system. J Appl and Nat Sci. 2017;9(2):1170–75. <https://doi.org/10.31018/jans.v9i2.1342>
22. Rajput TBS, Patel N. Water and nitrate movement in drip-irrigated onion under fertigation and irrigation treatments. Agri Water Manage. 2006;79(3):293–311. <https://doi.org/10.1016/j.agwat.2005.03.009>
23. Hebbar S, Ramachandrapa B, Nanjappa H, Prabhakar M. Studies on NPK drip fertigation in field grown tomato (*Lycopersicon esculentum* Mill.). European J Agron. 2004;21(1):117–27. [https://doi.org/10.1016/S1161-0301\(03\)00091-1](https://doi.org/10.1016/S1161-0301(03)00091-1)
24. Kumar A, Singh A. Improving nutrient and water use efficiency through fertigation. J Water Manage. 2002;10:42–48.