



RESEARCH ARTICLE

# Exploring the potential of seaweed extract in paddy seed presoaking: A pathway to improve crop performance

R Elamparathi<sup>1\*</sup>, K Sujatha<sup>1\*</sup>, V Alex Albert<sup>1</sup>, T Sivakumar<sup>2</sup>, A Gurusamy<sup>2</sup> & ML Mini<sup>3</sup>

<sup>1</sup>Department of Seed Science and Technology, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India

<sup>2</sup>Department of Agronomy, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India

<sup>3</sup>Department of Biotechnology, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India

\*Email: [sujathakseed@tnau.ac.in](mailto:sujathakseed@tnau.ac.in); [elamparathi27798@gmail.com](mailto:elamparathi27798@gmail.com)



## ARTICLE HISTORY

Received: 19 October 2024

Accepted: 17 November 2024

Available online

Version 1.0 : 31 December 2024

Version 2.0 : 27 August 2025



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Elamparathi R, Sujatha K, Albert VA, Sivakumar T, Gurusamy A, Mini ML. Exploring the potential of seaweed extract in paddy seed presoaking: A pathway to improve crop performance. Plant Science Today.2024;11 (sp4):01-11. <https://doi.org/10.14719/pst.5981>

## Abstract

Paddy (*Oryza sativa* L.) is a globally important staple crop, and achieving high yield is closely linked to effective seed treatments. In this study, seeds of improved kavuni CO 57 were treated with seaweed extracts (SE) from *Sargassum myricocystum* (brown algae) and *Kappaphycus alvarezii* (red algae) at various concentrations to assess their impact on seed performance. The treated seeds were evaluated for physiological and biochemical improvements. Notably, seeds soaked in a 0.5% methanol extract of *S. myricocystum* (T<sub>6</sub>) showed significant improvements compared to the control group, including a higher germination rate (94%), increased root length (19.51 cm), enhanced shoot length (9.29 cm), higher dry matter production (0.155 g/seedling), and a marked increase in seedling vigour index (2707). Biochemical analysis revealed significant enhancements in enzyme activities, with α-amylase (2.41 mg maltose min<sup>-1</sup>), catalase (3.15 μmol H<sub>2</sub>O<sub>2</sub> (hydrogen peroxide) reduced min<sup>-1</sup> g<sup>-1</sup>), and peroxidase (0.332 moles tetra guaiacol min<sup>-1</sup> g<sup>-1</sup>) all exhibiting higher levels in treated seeds. Additionally, Gas Chromatography-Mass Spectrometry (GC-MS) analysis identified key secondary metabolites in the treated seeds, with hexadecanoic acid (21.14%) and octadecanoic acid (10.86%) as dominant compounds. These compounds, known for their antimicrobial, antiviral, antibacterial, and antifungal properties, suggest enhanced resilience in the treated plants. Overall, the findings highlight the potential of SE as a sustainable alternative to conventional seed treatments, offering a promising approach for enhancing crop growth and yield in organic and sustainable agricultural systems.

## Keywords

improved kavuni CO 57; *Kappaphycus alvarezii*; paddy; presoaking; *Sargassum myricocystum*; seaweed

## Introduction

Paddy (*Oryza sativa* L.) is one of the world's most important staple crops and remains a key source of food for more than half of the global population (1). Organic produce, particularly organic rice, has experienced unprecedented demand in recent years due to increasing consumer awareness (2). Organic rice production emphasizes the use of natural inputs and ecological farming approaches, which contribute to improved soil health and biodiversity (3). In this context, organic seed production for paddy holds significant

potential, enabling farmers to operate sustainably while improving the quality of their output.

Quality seeds are essential for maximizing agricultural production and yield, particularly in organic farming systems, which often limit or prohibit the use of conventional inputs. Organic seed production requires careful selection of high-quality seeds and practices that enhance genetic diversity, pest resistance, and local adaptation. These factors are critical in developing resilient crops capable of withstanding both biotic and abiotic stresses, which is particularly important for yields in organic systems that depend on biodiversity and natural regulatory mechanisms (4, 5).

Presoaking technology is an advanced agronomic technique used to improve seed germination and crop establishment. It involves submerging seeds in water or nutrient solutions prior to sowing, which induces physiological changes that promote earlier and more predictable germination (6). Presoaking offers numerous benefits for many crops, including rice, especially in terms of enhancing seed viability and increasing resilience to environmental stresses (7).

The advantage of presoaking seeds is that it improves the seed's water absorption capacity (8). Water absorption activates the necessary metabolic processes that must occur before germination, such as the mobilization of stored carbohydrates and proteins (9). Experiments have demonstrated that this process reduces the germination time, leading to faster seedling establishment. This is particularly crucial in paddy cultivation, where timely germination allows plants to take advantage of favorable weather conditions, enabling them to grow to their full potential and thereby increasing yields (10).

SE derived from marine algae (brown, red, and green) contain bioactive substances, including hormones (auxins, cytokinins, abscisic acid, and gibberellins), vitamins, and minerals, all of which can positively influence plant growth and physiological responses (11). Treating seeds with SE can promote seed germination vigour and enhance resistance to environmental stress (12, 13). It has been reported that presowing treatments with SE improve germination percentages, root development, and overall plant robustness, which is especially beneficial for organic paddy cultivation.

Several studies have shown that SE regulate germinating seeds in various physiological ways, such as enhancing water uptake, hormonal signaling, and nutrient absorption (14). These extracts promote early root development, which aids in anchorage and nutrient uptake once the plant is established. The auxins and cytokinins in SE are critical for stimulating cell division and elongation, which ultimately strengthens root and shoot growth (15). Pre-soaking seeds in SE also accelerates water uptake, reducing germination times. This approach enables agro-scientists to promote more balanced and uniform plant distribution, which is essential for maximizing yields. By shortening the germination lag phase, it minimizes competition among seedlings, contributing to healthier crop establishment.

The improved kavuni CO 57 variety of paddy represents an innovative advancement in rice cultivation, specifically developed to address the challenges posed by modern agricultural demands. This variety is distinguished by its high yield potential, nutritional benefits, and adaptability to a range of climatic conditions. The kavuni CO 57 variety is also well-suited for organic farming practices, aligning with the growing global trend toward sustainable and environmentally friendly agriculture (16). Its resilience to various biotic and abiotic stresses enables farmers to reduce their reliance on synthetic agrochemicals, fostering a more sustainable food production system. This adaptability not only benefits the ecosystem but also aids organic farmers in maintaining soil health and biodiversity.

This study examines the effects of presoaking paddy seeds of the improved kavuni variety with SE derived from two species, using water and methanol as solvents at different concentrations. By applying SE treatments prior to sowing, the study seeks to explore their potential in enhancing the physiological and biochemical performance of rice seedlings. Physiological parameters such as germination percentage and seedling vigour were assessed, alongside key biochemical markers.

The selection of *S. myricocystum* (brown algae) and *K. alvarezii* (red algae) was based on their known bioactive properties and previous research indicating their potential in promoting plant growth. *S. myricocystum* is recognized for its rich composition of growth-promoting compounds such as alginates, fucoidans, and phytohormones, which are beneficial for enhancing seed vigour and resilience. *K. alvarezii*, on the other hand, is valued for its high carrageenan content and various micronutrients, which contribute to improved germination rates, root development, and stress tolerance in plants.

Furthermore, metabolite profiling was performed using GC-MS to gain deeper insight into the metabolic changes induced by SE treatments. GC-MS provides a comprehensive analysis of small molecules, enabling the identification of potential metabolites that may contribute to enhanced seed performance (17).

In this context, the present study aims to standardize presoaking techniques to improve seed germination and seedling vigour in the improved kavuni CO 57 variety. This study addresses a specific research gap by investigating the effectiveness of presoaking paddy seeds, conducting both physiological and biochemical analyses, and identifying metabolic compounds in presoaked seeds compared to non-soaked seeds. Paddy serves as an ideal model for this research due to its agricultural significance and the need to improve its germination rates and overall crop performance. The findings from this research will contribute to sustainable farming practices and provide valuable insights into enhancing paddy cultivation through presoaking.

## Materials and Methods

### Seed materials and seaweed

A total of 2 kg of improved kavuni CO 57 paddy seeds were

collected from the Department of Rice, Tamil Nadu Agricultural University, Coimbatore. The laboratory experiment was conducted at the Department of Seed Science and Technology, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai, during the 2023-24 period. The seaweeds *S. myricocystum* (brown algae) and *K. alvarezii* (red algae) were collected from the Mandapam coast in Ramanathapuram, Tamil Nadu.

### Preparation of seaweed extract

The seaweed was initially washed with seawater to remove macroscopic epiphytes and sand particles, followed by a rinse with fresh water to eliminate any adhering salt. It was then shade-dried for one day, sun-dried for one week, and finally oven-dried (Model: Binder FD 115, Company: Binder GmbH, Germany) at 50°C for 24 hours. The dried seaweed was ground using a Willey mill, and the pulverized powder was sieved through a 0.25 mm mesh to obtain a fine powder.

For the preparation of SE, 10 g of the seaweed powder was homogenized with 100 mL of a solvent mixture (water and methanol, 1:1) in a conical flask. The flask was kept at ambient temperature, and the homogenized material was placed in a water bath at 60°C for 1 hour. Afterward, the mixture was centrifuged (Eppendorf 5804, Company: Eppendorf AG, Germany) at 3000 rpm for 20 minutes. The supernatant was then collected as the SE (18).

### Experimental design

In this experiment, the seeds were subjected to pre-soaking in SE at different concentrations for 28 hours. The treatment details are as follows: T<sub>0</sub>-control (without soaking) it's an absolute control, T<sub>1</sub>- seeds soaked with water, T<sub>2</sub>- seeds soaked with 5% *S. myricocystum* water extract, T<sub>3</sub>- seeds soaked with 7.5% *S. myricocystum* water extract, T<sub>4</sub>- seeds soaked with 10% *S. myricocystum* water extract, T<sub>5</sub>- seeds soaked with 0.25% *S. myricocystum* methanol extract, T<sub>6</sub>- seeds soaked with 0.5% *S. myricocystum* methanol extract, T<sub>7</sub>- seeds soaked with 0.75% *S. myricocystum* methanol extract, T<sub>8</sub>- seeds soaked with 5% *K. alvarezii* water extract, T<sub>9</sub>- seeds soaked with 7.5% *K. alvarezii* water extract, T<sub>10</sub>- seeds soaked with 10% *K. alvarezii* water extract, T<sub>11</sub>- seeds soaked with 0.25% *K. alvarezii* methanol extract, T<sub>12</sub>- seeds soaked with 0.5% *K. alvarezii* methanol extract, T<sub>13</sub>- seeds soaked with 0.75% *K. alvarezii* water extract. SE at higher concentrations were found to affect seed quality parameters.

### Assessment of seed physiological quality parameters

A standard germination test was conducted using the roll towel method, following International Seed Testing Association (ISTA) guidelines (19). Four sets of 100 seeds were used in this experiment. The seeds were placed on moistened germination paper and incubated in a germinator set at 25 ± 2°C with a relative humidity of 95 ± 2%. After 14 days, the germination % was assessed and expressed as a percentage (20).

$$\text{Germination (\%)} = \frac{\text{Number of normal seedlings}}{\text{Total number of seeds sown}} \times 100$$

.....(Eqn. 1)

The average root and shoot lengths were measured in cm to evaluate seedling growth.

To determine the dry weight, ten healthy seedlings, selected for measuring root and shoot lengths, were placed in paper covers and air-dried in the shade for 24 hours. They were then dried in a hot air oven at 85 ± 1°C for an additional 24 hours. The average weight was recorded in grams per ten seedlings. Additionally, the seedling vigour index was calculated, and the mean values were presented as whole numbers (21).

$$\text{Vigour index I} = \text{Germination (\%)} \times \text{Total seedling length (cm)}$$

.....(Eqn. 2)

$$\text{Vigour index II} = \text{germination (\%)} \times \text{Dry matter production (g per 10 seedlings)}$$

.....(Eqn. 3)

### Biochemical analysis during seed germination

Biochemical assays such as α -amylase, catalase and peroxidase were carried out (22-24).

#### α -amylase (mg maltose min<sup>-1</sup>)

The fresh and aged seeds were pre-germinated using the top-of-paper method and allowed to undergo radicle emergence. In this method, 500 mg of pre-germinated seeds were homogenized in 1.8 mL of ice-cold 0.02 M sodium phosphate buffer (pH 6.0). The homogenate was then centrifuged at 20,000 rpm for 20 minutes to collect the extract. Subsequently, 1 mL of a 0.0067% starch solution was added to the extract and incubated for 10 minutes at 25°C. The reaction was terminated by adding 1 mL of iodine-HCl solution containing 60 mg KI (potassium iodide) and 6 mg I<sub>2</sub> (iodine) dissolved in 100 mL of 0.05 N HCl (hydrochloric acid). The color change was observed at 620 nm, and the enzyme activity was calculated and expressed as milligrams of maltose per minute (22).

#### Catalase (μmol of reduced H<sub>2</sub>O<sub>2</sub> g<sup>-1</sup> min<sup>-1</sup>)

The enzyme extract was prepared by finely grinding 0.5 g of pre-germinated seed sample with 5 mL of ice-cold 50 mM phosphate buffer (pH 7.0). The homogenized sample was then centrifuged at 15,000 rpm for 20 minutes at 4°C. The collected supernatant was used as the enzyme extract. To 0.3 mL of enzyme extract, 1.5 mL of 50 mM potassium phosphate buffer and 1.2 mL of 12.5 mM hydrogen peroxide were added. The mixture was then incubated for 10-15 minutes. Subsequently, the optical density (OD) values at 240 nm were measured every 15 seconds for 1 minute using a Ultraviolet-Visible spectrophotometer. Based on the reduction of H<sub>2</sub>O<sub>2</sub>, catalase activity was calculated and compared with known concentrations of hydrogen peroxide using a standard curve. The enzyme ac-

tivity was calculated as the amount of  $\text{H}_2\text{O}_2$  reduced (initial reading – final reading = quantity of  $\text{H}_2\text{O}_2$  reduced) and expressed as  $\mu\text{mol}$  of reduced  $\text{H}_2\text{O}_2$  per gram per minute (23).

#### Peroxidase ( $\text{m mol tetra guaiacol min}^{-1}\text{g}^{-1}$ )

A 0.5 g sample of pre-germinated seeds was homogenized with 1.5 mL of 60 mM phosphate buffer using a pestle and mortar. The sample was then centrifuged at 10,000 rpm for 10 minutes at 4°C. To 0.1 mL of enzyme extract, 0.5 mL of 1% hydrogen peroxide and 0.5 mL of 96 mM guaiacol were added. The mixture was then diluted with 0.4 mL of water to make a final volume of 3 mL and incubated at 25°C for 10 minutes. The optical density (OD) value was recorded at 470 nm using a UV-Vis spectrophotometer, based on the coefficient of oxidized tetra-guaiacol. Peroxidase activity was calculated from the absorbance and expressed as mmol of tetra-guaiacol per minute per gram (24).

$$\text{Peroxidase activity} = \frac{\text{Difference in OD value} / 10 \text{ min} \times 1000 / 500 \times 60}{\dots\dots\dots} \text{(Eqn. 4)}$$

#### Identification of metabolic compounds through GC-MS analysis

Paddy seeds were presoaked with 0.5% *S. myricocystum* methanol extract ( $T_6$ ) (the best treatment) and water-soaked seeds ( $T_2$ ) were used for GC-MS analysis. The extraction protocol for polar compounds was modified and carried out as follows: 50 mg of seed material was ground using a grinder. For primary metabolite profiling, solute extraction was performed with 400  $\mu\text{L}$  of methanol ( $-20^\circ\text{C}$ ), containing 200 nmol of cis-inositol as an internal standard. The mixture was processed in a thermomixer (Grant Instruments) at 70°C for 10 minutes at 950 rpm. Subsequently, 200  $\mu\text{L}$  of chloroform was added, and the solution was shaken for another 5 minutes at 70°C and 950 rpm. Finally, 400  $\mu\text{L}$  of ultra-pure water was added, followed by vortexing for 20 seconds and centrifugation for 10 minutes at 7400 $\times$ g. A 50  $\mu\text{L}$  aliquot of the methanol supernatant was dried in a speed vacuum for subsequent GC-MS analysis (25).

The samples were subjected to GC-MS analysis using an Agilent 7890A Gas Chromatograph (GC) and an Agilent 5975C Mass Spectrometer (MS) (Agilent Technologies, USA), widely used for analyzing complex chemical mixtures and detecting volatile and semi-volatile compounds with high sensitivity. It is commonly applied in environmental testing, food safety, and forensic analysis.

A 1  $\mu\text{L}$  aliquot of the reaction mixture was injected directly into the GC-MS system. The operating conditions were as follows: the initial temperature was set to 80°C for 1 minute, then raised to 250°C at a rate of 8°C per minute, followed by a further increase to 300°C at 12°C per minute, where it was held for 5 minutes. The total run time for the GC was 30 minutes, and the injector temperature was maintained at 240°C.

#### Statistical design

The observed data were recorded and subjected to ANOVA

(analysis of variance) at a 5% level of significance. The percentage values were transformed to arc-sine values before analysis. The critical difference (CD) was calculated at both 1% and 5% probability levels and tested for statistical significance. Graphs were generated using Microsoft Excel (2019). The data were further analyzed using principal component analysis (PCA) in R software.

## Results

#### Efficacy of seaweed extract on seedling growth parameters

The results demonstrated that seeds pre-soaked with SE exhibited significant improvements ( $p < 0.05$ ) in seedling attributes, such as germination %, root and shoot lengths (cm), and vigour indices I and II, compared to untreated seeds. The most pronounced increase in germination (94%) was observed in seeds pre-soaked in 0.5% *S. myricocystum* methanol extract ( $T_6$ ) for 28 hours, followed by  $T_5$  (92%) compared to the control (Table 1) (Fig. 1). Furthermore, seeds pre-soaked with 0.5% *S. myricocystum* methanol extract ( $T_6$ ) showed substantial improvements in root length (19.51 cm), shoot length (9.29 cm), dry matter production (0.155 g per 10 seedlings), and seedling vigour index (2707). Statistically significant differences in germination and seedling vigour were observed between pre-soaked and untreated seeds. Overall, the pre-soaking treatments increased germination rates and seedling vigour relative to the control. However, higher concentrations of soaking agents tended to negatively affect germination and seed quality parameters.

#### Effects of presoaking on biochemical changes during germination


The activities of  $\alpha$ -amylase ( $\text{mg maltose min}^{-1}$ ), catalase ( $\mu\text{mol H}_2\text{O}_2$  reduced  $\text{min}^{-1}\text{g}^{-1}$ ), and peroxidase (mmoles of tetra-guaiacol  $\text{min}^{-1}\text{g}^{-1}$ ) in paddy seeds were significantly influenced by the pre-soaking treatments with SE, in comparison to untreated seeds. Seeds soaked in 0.5% *S. myricocystum* methanol extract ( $T_6$ ) for 28 hours exhibited the highest  $\alpha$ -amylase activity at 2.41 mg maltose  $\text{min}^{-1}$ , followed by  $T_5$  (0.25% *S. myricocystum* methanol extract for 28 hours) at 2.38 mg maltose  $\text{min}^{-1}$ , while non-soaked seeds showed only 2.15 mg maltose  $\text{min}^{-1}$  (Fig. 2A). This increase in  $\alpha$ -amylase activity was associated with higher germination rates. Catalase activity also increased due to the nutrient content in the extracts, with  $T_6$  recording 3.15  $\mu\text{mol H}_2\text{O}_2$  reduced  $\text{min}^{-1}\text{g}^{-1}$ , followed by  $T_5$  at 3.12  $\mu\text{mol H}_2\text{O}_2$  reduced  $\text{min}^{-1}\text{g}^{-1}$ , and the lowest value in non-soaked seeds at 2.88  $\mu\text{mol H}_2\text{O}_2$  reduced  $\text{min}^{-1}\text{g}^{-1}$  (Fig. 2B). Peroxidase levels were similarly higher in  $T_6$  (0.332 mmoles of tetra-guaiacol  $\text{min}^{-1}\text{g}^{-1}$ ), with  $T_5$  close behind at 0.328 mmoles of tetra-guaiacol  $\text{min}^{-1}\text{g}^{-1}$ , compared to 0.308 mmoles of tetra-guaiacol  $\text{min}^{-1}\text{g}^{-1}$  in untreated seeds (Fig. 2C). A strong positive correlation was found between germination percentage and  $\alpha$ -amylase activity (Fig. 2D).



**Table 1.** Effect of seed pre-soaking on physiological parameters of improved kavuni CO 57

Treat-ments	Germination (%)	Root length (cm)	Shoot length (cm)	Dry matter production (g 10 seedlings <sup>-1</sup> )	Vigour index I	Vigour index II
T <sub>0</sub>	74	16.74	7.54	0.130	1797	9.6
T <sub>1</sub>	74	16.83	7.67	0.131	1813	9.7
T <sub>2</sub>	84	17.66	8.21	0.140	2173	11.8
T <sub>3</sub>	88	18.44	8.54	0.144	2374	12.7
T <sub>4</sub>	76	17.03	7.84	0.134	1890	10.2
T <sub>5</sub>	92	19.12	9.16	0.151	2602	13.9
T <sub>6</sub>	94	19.51	9.29	0.155	2707	14.6
T <sub>7</sub>	82	17.49	8.05	0.138	2094	11.3
T <sub>8</sub>	84	17.54	8.16	0.138	2159	11.6
T <sub>9</sub>	86	18.35	8.34	0.141	2295	12.1
T <sub>10</sub>	80	17.19	7.95	0.136	2011	10.9
T <sub>11</sub>	90	18.48	8.70	0.144	2446	13.0
T <sub>12</sub>	90	18.87	8.92	0.148	2501	13.3
T <sub>13</sub>	80	17.38	8.02	0.136	2032	10.9
<b>SE</b>	1.792	0.424	0.157	0.003	36.637	0.185
<b>CD (0.05)</b>	3.276	0.868	0.322	0.007	75.032	0.378

T<sub>0</sub>-Control (Without soaking) T<sub>1</sub>- Seeds soaked with water T<sub>2</sub>- Seeds soaked with 5% *S. myricocystum* water extract T<sub>3</sub>- Seeds soaked with 7.5% *S. myricocystum* water extract T<sub>4</sub>- Seeds soaked with 10% *S. myricocystum* water extract T<sub>5</sub>- Seeds soaked with 0.25% *S. myricocystum* methanol extract T<sub>6</sub>- Seeds soaked with 0.5% *S. myricocystum* methanol extract T<sub>7</sub>- Seeds soaked with 0.75% *S. myricocystum* methanol extract T<sub>8</sub>- Seeds soaked with 5% *K. alvarezii* water extract T<sub>9</sub>- Seeds soaked with 7.5% *K. alvarezii* water extract T<sub>10</sub>- Seeds soaked with 10% *K. alvarezii* water extract T<sub>11</sub>- Seeds soaked with 0.25% *K. alvarezii* methanol extract T<sub>12</sub>- Seeds soaked with 0.5% *K. alvarezii* methanol extract T<sub>13</sub>- Seeds soaked with 0.75% *K. alvarezii* water extract.


**Fig. 1.** Effect of seed pre-soaking on germination and seedling vigour of improved kavuni CO 57.

### GC-MS analyses of metabolite compounds in presoaked seeds

GC-MS analysis of metabolite compounds from seeds pre-soaked in 0.5% *S. myricocystum* methanol extract (T<sub>6</sub>) re-


vealed distinct differences in the composition of metabolite blends released by the SE compared to those released by the water-soaked control.

Sixty compounds identified from the water-soaked



**Fig. 2.** The effect of seed pre-soaking on (A)  $\alpha$ -amylase (B) catalase (C) peroxidase (D) correlation analysis between germination and  $\alpha$ -amylase activity.


seeds included 2-cyclopenten-1-one, 2-hydroxy; oxirane; oxalic acid, monoamide; n-propyl tetradecyl ester; 2-propenoic acid, 3-(4-methoxyphenyl); hexadecanoic acid, methyl ester; n-hexadecanoic acid; and benzene. These identified compounds exhibit antimicrobial, antiviral, antibacterial, and antifungal activities. The primary chemical constituents were hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester, with a peak area of 21.14% and a retention time of 16.507 minutes, and octadecanoic acid, methyl ester, with a peak area of 10.86% and a reten-




**Fig. 3.** GC-MS chromatogram of metabolite profiling for water-soaked seeds of the improved kavuni CO 57.

tion time of 14.285 minutes (Fig. 3).

In contrast, 60 chemical constituents were identified from paddy seeds presoaked with the SE, including decane, 2-pyrrolidine methanol, rhodopin, acetamide, hexanedioic acid, 1,2-benzene dicarboxylic acid, bis (2-methylpropyl) and mono (2-ethylhexyl) ester, campesterol, stigmasterol, and gamma-sitosterol. The identified compounds possess antimicrobial, antiviral, antibacterial, antifungal, antioxidant, and insecticidal activities. The primary chemical constituents were 2-propenoic acid, 3-(4-methoxyphenyl)-, ethyl ester, with a peak area of 13.51% and a retention time of 11.875 minutes, and 2-propenoic acid, 3-(4-methoxyphenyl)-, ethyl ester, with a peak area of



**Fig. 4.** GC-MS chromatogram of metabolite profiling for 0.5% *S. myricocystum* methanol extract-soaked seeds of the improved kavuni CO 57.




**Fig. 5.** Heatmap visualization - the colour gradient from yellow to blue represents the range of metabolite expression levels, with yellow indicating lower levels and blue indicating higher levels. Treatments are T<sub>2</sub> - control (Water-soaked seeds); T<sub>6</sub> - seed pre-soaked with 0.5% *S. myricocystum* methanol extract.

12.13% and a retention time of 12.986 minutes (Fig. 4). Additionally, heatmap analysis revealed significant differences in metabolite expression patterns between the two groups (Fig. 5).

### Principal component analysis

Principal Component Analysis (PCA) was used to examine the relationships among the measured seedling growth parameters (Fig. 6). This statistical method is particularly useful for simplifying datasets with many correlated variables by reducing them to a smaller number of principal



**Fig. 6.** The principal component analysis shows treatments' impact on variables such as seed quality and biochemical parameters of improved kavuni CO 57.

components. In this analysis, the first principal component (PC1) accounts for 98.2% of the total variance, while the second principal component (PC2) captures 0.9%. Together, PC1 and PC2 explain 99.1% of the variance in the data.

Germination,  $\alpha$ -amylase, catalase, and peroxidase showed strong positive contributions to both PC1 and PC2, indicating that these factors played a major role in the variability captured by these components. Additionally, vigour indices I and II, root length, shoot length, and dry matter production exhibited weaker associations with PC1 but still contributed to the overall variation.

The points labeled T<sub>0</sub>, T<sub>1</sub>–T<sub>13</sub> represent different treatment groups, positioned according to their similarity in variable response. Treatment T<sub>2</sub>, located at the top near the germination vector, suggests a high germination rate and a positive association with catalase activity.

Treatments T<sub>6</sub> and T<sub>5</sub>, positioned further down and to the right, exhibit a distinctive response pattern with a stronger influence from variables like root length and shoot length.

Treatment T<sub>1</sub>, positioned in the bottom left quadrant, is negatively associated with most variables shown in the PCA plot, indicating it may have lower overall performance for the measured parameters.

This PCA visualization helps identify which treatments (T<sub>6</sub>, T<sub>5</sub>) may be optimal for specific growth parameters. Treatments that align closely with growth-promoting variables, such as seedling vigour index or germination rate, may be favourable for enhancing overall seedling development. Conversely, treatments with distinct separations, like T<sub>1</sub>, could indicate unique or less effective responses, warranting further investigation or exclusion from recommended practices.

### Discussion

Seaweeds, or macroalgae, are multicellular marine organisms that play a crucial role in coastal marine ecosystems. They are classified into three primary groups based on their pigmentation: Phaeophyta (brown), Rhodophyta (red), and Chlorophyta (green). Extracts obtained from these seaweeds are rich in a variety of bioactive compounds (26), including polysaccharides, pigments, phenolic compounds, proteins and bioactive peptides, phytohormones, as well as both micro- and macronutrients (26–29). Numerous studies have highlighted the potential benefits of using SE as biostimulants under both normal and stressed environmental conditions (26–30).

SE are abundant in phytohormones, sterols such as fucosterol, carbohydrates, polysaccharides, sugars, polyphenols (including flavonoids), macro- and micronutrients, vitamins, lipids, amino acids, and proteins, including enzymes (26–30).

Soluble alginates and protein hydrolysates derived from seaweeds have been shown to enhance the aggregation of soil particles, thereby improving nutrient availability, aeration, and water retention in the soil (26–31). Beyond the direct benefits to plants, SE positively influence the soil microbiome. Studies indicate that microbes can absorb free amino acids more effectively than plants; in some crops, only 6% to 25% of the flagged amino acids were taken up by the roots, with the remainder absorbed by soil microorganisms (32).

Various phenolic compounds have been identified in seaweeds, with brown seaweeds primarily containing phlorotannins, while red and green species are richer in bromophenols, flavonoids, and phenolic acids (33). Phlorotannins, in particular, contain a higher number of phenolic rings compared to other phenolic compounds,

which is associated with enhanced antioxidant activity (34).

Micronutrients are also present in seaweed products, whether in their fresh, dried, or extracted forms (35). Another group of bioactive molecules found in SE is plant hormones. The hormonal composition of seaweeds is similar to that of terrestrial plants. Although the mechanisms of action for these hormones in seaweeds are not fully understood, various phytohormones, including bioactive forms of auxins, cytokinins (CK), abscisic acid (ABA), and gibberellins, have been identified (36). Additionally, ethylene, brassinosteroids, salicylic acid, jasmonates (JA), and strigolactones have also been detected in SE (28).

The compounds present in SE may function as signaling molecules that regulate key pathways at both the transcriptional and post-translational levels (via microRNAs), leading to the differential expression of essential genes in crops. This regulation can enhance plant growth by affecting genes related to cell metabolism, including those involved in lipid, amino acid, and nucleotide metabolism, glycolysis, and transport, as well as cell and cell wall development (37).

SE have shown significant positive effects on crop growth, yield, and quality in various studies. Foliar application of SE has enhanced growth parameters and yield components in different field crops (38). In bean plants, lower concentrations of *Fucus spiralis* and *Ulva rigida* extracts improved shoot and root length, chlorophyll content, and protein levels (39). Similarly, wheat plants irrigated with *Ascophyllum nodosum* extract exhibited increased height, dry mass, and spike number. SE contain multiple growth regulators, as well as macro- and micronutrients essential for plant development (40). These biostimulants can serve as an eco-friendly alternative to inorganic fertilizers, promoting early seed germination, improving crop performance, and enhancing resistance to biotic and abiotic stresses (38–40). However, high concentrations of SE may have negative effects on plant growth (39).

In the present study, presoaked seeds ( $T_6$ ) showed improvements in average quality parameters compared to the control. These findings align with earlier studies, which indicated that enhancing shoot and root length in bean plants led to significant improvements in seed germination, seedling growth, and biochemical parameters (39).

Seeds treated with seaweed nanopowder showed an increase in germination % in pigeon pea (41). In soybean, the application of SE at a 15% concentration resulted in taller plants compared to the control group (42). Similarly, foliar spraying of 0.4% SE led to increased plant height in green gram (43).

A combination of kappaphycus sap at 10% with the recommended dose of fertilizer (RDF) significantly increased the leaf area index in maize compared to the control group (44). In rice, soil application of SE gel at 12.5 kg/ha, combined with a foliar spray of 0.5% seaweed extract at the tillering and panicle initiation stages, resulted in a higher leaf area index (45).

In green gram, applying 15% kappaphycus sap along with the recommended fertilizer dose led to higher dry matter production (46), while seeds treated with seaweed nanopowder produced greater dry matter in pigeon pea (41). The use of biostimulants, such as Algex derived from *Ascophyllum nodosum*, significantly enhanced dry matter production in red clover compared to untreated plants (47).

SE have demonstrated positive effects on crop growth and biochemical parameters when applied at optimal concentrations. Studies on various crops, including wheat, beans, and legumes, have shown enhanced growth metrics such as shoot and root length, as well as increased dry weight (48). Biochemical parameters, such as chlorophyll content, protein levels, and enzyme activities, were also improved. Notably, catalase activity increased with higher SE concentrations in wheat (48), while peroxidase activity was enhanced at moderate concentrations. Alpha-amylase activity increased in cowpea treated with *Ulva lactuca* extract. The optimal concentration for most beneficial effects was generally found to be between 3–10% of SE, with higher concentrations sometimes showing inhibitory effects (49). These studies suggest that SE can serve as effective biofertilizers, promoting crop growth and enhancing biochemical parameters.

In rice, soil application of SE gel at 12.5 kg/ha, combined with foliar spraying of 0.5% extract at critical growth stages (tillering and panicle initiation), resulted in a significantly higher 1000-grain weight (45). Furthermore, the same treatment led to a substantial improvement in the harvest index, reaching 44%. Additionally, the application of SE gel at 25 kg/ha increased the uptake of nutrients such as nitrogen, phosphorus, potassium, iron, zinc, copper, and manganese in rice compared to the control group (45).

Based on GC-MS analysis results, 2-propenoic acid, 3-(4-methoxyphenyl)-, ethyl ester plays a significant role in seed germination and plant growth by acting as a growth promoter and disease inhibitor. Research indicates that derivatives of 4-methoxyphenyl compounds enhance germination energy and biometric parameters in various seeds, suggesting their effectiveness in improving seed viability and growth parameters (50). Additionally, compounds such as 2-(4-methoxyphenyl) propionic acid can stimulate stem growth while inhibiting root development, optimizing the edible yield of sprouts. Furthermore, these compounds exhibit potential in preventing plant diseases by inhibiting phytopathogen virulence without adversely affecting plant growth. Overall, integrating such compounds into agricultural practices could lead to improved crop yields and healthier plants (51).

Octadecanoic acid, methyl ester, is a significant compound found in various plant sources, including seaweeds. Research indicates that seaweed species like *Cladophora rupestris* contain novel octadecadienoic fatty acids, which can be analyzed through oxidative ozonolysis techniques to identify their structures (52). Furthermore, octadecanoic acid, methyl ester has demonstrated antiviral properties, particularly when combined with ribavirin



against the measles virus, showing enhanced efficacy at lower concentrations (53). Additionally, the photo-oxidation of lipids in dried seaweed can influence the volatile compounds produced, which may include derivatives of octadecanoic acid. Overall, the incorporation of octadecanoic acid, methyl ester in SE presoaked seeds presents potential for both nutritional and therapeutic applications.

## Conclusion

The study demonstrated that pre-soaking seeds with SE significantly enhanced seed growth parameters in improved kavuni CO 57. Seeds soaked in a 0.5% methanol extract of *S. myricocystum* for 28 hours exhibited higher germination rates, longer root and shoot lengths, and a greater seedling vigour index compared to untreated seeds. Biochemical analysis revealed elevated levels of  $\alpha$ -amylase, catalase, peroxidase, and other key enzymes in the treated seeds. These results suggest that seaweed extract serves as an effective and eco-friendly alternative to chemical fertilizers, promoting sustainable agriculture while improving crop quality and yield. Further research should explore the long-term impacts of seaweed pre-soaking on overall crop yield and investigate the method's applicability across different crops. Such studies would help validate these findings and provide valuable insights for broader agricultural applications.

## Acknowledgements

Authors wish to thank Department of Seed Science and Technology, AC & RI, (TNAU), Madurai, for providing financial support during all research activities and manuscript preparation.

## Authors' contributions

RE carried out the experimental work and drafted the original draft of the writing. KS participated in the conceptualization of the study. VAA and TS supervised the work, drafted and reviewed the manuscript. AG participated in the sequence alignment and editing. MLM participated in the visualization. All authors read and approved the final manuscript.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** None

## References

- Muthayya S, Sugimoto JD, Montgomery S, Maberly GF. An overview of global rice production, supply, trade, and consumption. *Ann N Y Acad Sci.* 2014;1324(1):7-14. <https://doi.org/10.1111/nyas.12540>
- Lernoud J, Willer H. Organic Agriculture Worldwide: Key results from the FiBL survey on organic agriculture worldwide 2019. Research Institute of Organic Agriculture (FiBL), part. 2019;1.
- Lori M, Symnaczik S, Mäder P, De Deyn G, Gättinger A. Organic farming enhances soil microbial abundance and activity—A meta-analysis and meta-regression. *PloS one.* 2017;12(7):e0180442. <https://doi.org/10.1371/journal.pone.0180442>
- Van Bueren EL, Jones SS, Tamm L, Murphy KM, Myers JR, Leifert C, Messmer MM. The need to breed crop varieties suitable for organic farming, using wheat, tomato and broccoli as examples: A review. *NJAS-Wagen J Life Sci.* 2011;58(3-4):193-205. <https://doi.org/10.1016/j.njas.2011.08.001>
- Crowder DW, Jabbour R. Relationships between biodiversity and biological control in agroecosystems: Current status and future challenges. *Bio Control.* 2014;75:8-17. <https://doi.org/10.1016/j.biocontrol.2013.10.010>
- Paparella S, Araújo SS, Rossi G, Wijayasinghe MA, Carbonera D, Balestrazzi A. Seed priming: State of the art and new perspectives. *Plant Cell Rep.* 2015;34:1281-93. <https://doi.org/10.1007/s00299-015-1784-y>
- Hussain S, Khan F, Hussain HA, Nie L. Physiological and biochemical mechanisms of seed priming-induced chilling tolerance in rice cultivars. *Front Plant Sci.* 2016;7:116. <https://doi.org/10.3389/fpls.2016.00116>
- Li C, Nonogaki H, Barrero. Seed Dormancy, Germination and Pre-Harvest Sprouting. *Frontiers Media SA-2019.* <https://doi.org/10.3389/978-2-88963-898-2>
- Rajjou L, Duval M, Gallardo K, Catusse J, Bally J, Job C, Job D. Seed germination and vigor. *Annu Rev Plant Biol.* 2012;63(1):507-33. <https://doi.org/10.1146/annurev-arplant-042811-105550>
- Basra SM, Farooq M, Tabassam R, Ahmad N. Physiological and biochemical aspects of pre-sowing seed treatments in fine rice (*Oryza sativa* L.). *SST.* 2005;33(3):623-8. <https://doi.org/10.15258/sst.2005.33.3.16>
- Battacharyya D, Babgohari MZ, Rathor P, Prithiviraj B. Seaweed extracts as biostimulants in horticulture. *Sci Hortic.* 2015;196:39-48. <https://doi.org/10.1016/j.scienta.2015.09.012>
- de Carvalho ME, de Camargo PR, Gallo LA, Junior MV. Seaweed extract provides development and production of wheat. *Agrarian.* 2014;7(23):166-70. <http://dx.doi.org/10.5281/zenodo.51607>
- Kumar G, Sahoo D. Effect of seaweed liquid extract on growth and yield of *Triticum aestivum* var. Pusa Gold. *J Appl Phycol.* 2011;23:251-55. <https://doi.org/10.1007/s10811-011-9660-9>
- Kocira A, Świeca M, Kocira S, Złotek U, Jakubczyk A. Enhancement of yield, nutritional and nutraceutical properties of two common bean cultivars following the application of seaweed extract (*Ecklonia maxima*). *Saudi J Biol Sci.* 2018;25(3):563-71. <https://doi.org/10.1016/j.sjbs.2016.01.039>
- Ertani A, Francioso O, Tinti A, Schiavon M, Pizzeghello D, Nardi S. Evaluation of seaweed extracts from *Laminaria* and *Ascophyllum nodosum* spp. as biostimulants in *Zea mays* L. using a combination of chemical, biochemical and morphological approaches. *Front Plant Sci.* 2018;9:428. <https://doi.org/10.3389/fpls.2018.00428>
- Udhaya A, Radhamani S, Ravichandhran V, Janaki P, Manonmani S. Evaluation of integrated nutrient management practices on growth, yield parameters and yield of improved traditional rice in the western zone of Tamil Nadu. *J Appl Nat Sci.* 2024;16(3):1233-9. <https://doi.org/10.31018/jans.v16i3.5784>
- Tanna B, Yadav S, Patel MK, Mishra A. Metabolite profiling, biological and molecular analyses validate the nutraceutical potential of green seaweed *Acrosiphonia orientalis* for human health. *Nutrients.* 2024;16(8):1222. <https://doi.org/10.3390/nu16081222>
- Godlewska K, Michalak I, Tuhy Ł, Chojnacka K. Plant growth biostimulants based on different methods of seaweed extraction with water. *Biomed Res Int.* 2016;2016(1):5973760. <https://doi.org/10.1155/2016/597376>
- International Seed Testing Association (ISTA). Germination percentage. 2019.

20. Abdul-Baki AA, Anderson JD. Vigor determination in soybean seed by multiple criteria 1. *Crop Sci.* 1973;13(6):630-33. <https://doi.org/10.2135/cropsci1973.0011183X001300060013x>
21. ISTA 2013. International rules for seed testing. *Seed Sci. & Technol.* 2013;27:25-30.
22. Paul AK, Mukherji S, Sircar SM. Metabolic changes in rice seeds during storage. 1970.
23. Luck H. *Methods in Enzymatic Analysis*. II Edition, Bergmeyer Publication. 1974:885-90.
24. Malik CP, Singh M. *Plant enzymology and histo-enzymology*. Kalyani Publishers. New Delhi; 1980.
25. Quéro A, Molinié R, Mathiron D, Thiombiano B, Fontaine JX, Brancourt D et al. Metabolite profiling of developing *Camelina sativa* seeds. *Metabolomics*. 2016;12:1-4. <https://doi.org/10.1007/s11306-016-1135-1>
26. Khan W, Rayirath UP, Subramanian S, Jithesh MN, Rayorath P, Hodges DM, et al. Seaweed extracts as biostimulants of plant growth and development. *J Plant Growth Regul.* 2009;28:386-99. <https://doi.org/10.1007/s00344-009-9103-x>
27. Craigie JS. Seaweed extract stimuli in plant science and agriculture. *J Appl Phycol.* 2011;23:371-93. <https://doi.org/10.1007/s10811-010-9560-4>
28. Stirk WA, Van Staden J. Plant growth regulators in seaweeds: Occurrence, regulation and functions. In *Adv Bot Res.* 2014; 71:125-59.
29. El Boukhari ME, Barakate M, Bouhia Y, Lyamlouli K. Trends in seaweed extract based biostimulants: Manufacturing process and beneficial effect on soil-plant systems. *Plants.* 2020;9(3):359. <https://doi.org/10.3390/plants9030359>
30. Shukla PS, Borza T, Critchley AT, Prithiviraj B. Carrageenans from red seaweeds as promoters of growth and elicitors of defense response in plants. *Front Mar Sci.* 2016;3:81. <https://doi.org/10.3389/fmars.2016.00081>
31. Colla G, Hoagland L, Ruzzi M, Cardarelli M, Bonini P, Canaguier R, Rouphael Y. Biostimulant action of protein hydrolysates: Unraveling their effects on plant physiology and microbiome. *Front Plant Sci.* 2017;8:2202. <https://doi.org/10.3389/fpls.2017.02202>
32. Moe LA. Amino acids in the rhizosphere: From plants to microbes. *Am J Bot.* 2013;100(9):1692-705. <https://doi.org/10.3732/ajb.1300033>
33. Cotas J, Leandro A, Monteiro P, Pacheco D, Figueirinha A, Gonçalves AM, et al. Seaweed phenolics: From extraction to applications. *Marine drugs.* 2020;18(8):384. <https://doi.org/10.3390/md18080384>
34. Sathya R, Kanaga N, Sankar P, Jeeva S. Antioxidant properties of phlorotannins from brown seaweed *Cystoseira trinodis* (Forsskål) C. Agardh. *Arab J Chem.* 2017;10:S2608-14. <https://doi.org/10.1016/j.arabjc.2017.01.002>
35. Sharma HS, Fleming C, Selby C, Rao JR, Martin T. Plant biostimulants: A review on the processing of macroalgae and use of extracts for crop management to reduce abiotic and biotic stresses. *J Appl Phycol.* 2014;26:465-90. <https://doi.org/10.1007/s10811-013-0101-9>
36. Lu Y, Xu J. Phytohormones in microalgae: A new opportunity for microalgal biotechnology. *Trends Plant Sci.* 2015;20(5):273-82. <https://doi.org/10.1016/j.tplants.2015.01.006>
37. Goni O, Fort A, Quille P, McKeown PC, Spillane C, O'Connell S. Comparative transcriptome analysis of two *Ascophyllum nodosum* extract biostimulants: Same seaweed but different. *J Agric Food Chem.* 2016;64(14):2980-89. <https://doi.org/10.1021/acs.jafc.6b00621>
38. Rani A, Saini KC, Fartyal M, Jaitak V, Bast F. A concise review on the bioactive potential of the genus *Gracilaria* (Rhodophyta). *The Nucleus.* 2024;68:161-77. <http://dx.doi.org/10.1007/s13237-024-00471-9>
39. Latique S, Chernane H, Mansori M, El Kaoua M. Seaweed liquid fertilizer effect on physiological and biochemical parameters of bean plant (*Phaseolus vulgaris* variety Paulista) under hydroponic system. *Eur Sci J.* 2013.
40. Begum M, Bordoloi BC, Singha DD, Ojha NJ. Role of seaweed extract on growth, yield and quality of some agricultural crops: A review. *Agri Rev.* 2018;39(4):321-26. <https://doi.org/10.18805/ag.R-1838>
41. Ambika S, Sujatha K. Organic seed treatment with seaweed nano powders weed nano powders on physiological quality and enzyme activities in aged seeds of pigeon pea. *The Bioscan.* 2016;11(Supplement 1):353-58.
42. Rathore SS, Chaudhary DR, Boricha GN, Ghosh A, Bhatt BP, Zodape ST, Patolia JS. Effect of seaweed extract on the growth, yield and nutrient uptake of soybean (*Glycine max*) under rain-fed conditions. *S Afr J Bot.* 2009;75(2):351-55. <https://doi.org/10.1016/j.sajb.2008.10.009>
43. Kavipriya R, Dhanalakshmi PK, Jayashree S, Thangaraju N. Seaweed extract as a biostimulant for legume crop, green gram. *J Ecobiotechnol.* 2011;3(8).
44. Garai S, Brahmachari K, Sarkar S, Kundu R, Pal M, Pramanick B. Crop growth and productivity of rainy maize-garden pea coping sequence as influenced by Kappaphycus and Gracilaria Saps at alluvial soil of West Bengal, India. *Curr J Appl Sci Technol.* 2019;36(2):1-11. <http://dx.doi.org/10.9734/cjast/2019/v36i230227>
45. Deepana P, Bama KS, Santhy P, Devi TS. Effect of seaweed extract on rice (*Oryza sativa* var. ADT53) productivity and soil fertility in Cauvery delta zone of Tamil Nadu, India. *J Appl & Nat Sci.* 2021;13(3):1111-20. <https://doi.org/10.31018/jans.v13i3.2906>
46. Pramanick B, Brahmachari K, Ghosh A. Effect of seaweed saps on growth and yield improvement of green gram. *Afr J Agric Res.* 2013;8(13):1180-86. <https://doi.org/10.5897/AJAR12.1894>
47. Godlewska A, Ciepiela GA. Yield performance and content of selected organic compounds in *Trifolium pratense* treated with various biostimulants against the background of nitrogen fertilisation. *Legume Research-An International Journal.* 2020;43(6):850-55. <http://dx.doi.org/10.18805/LR-522>
48. Ghafarizadeh A, Gilani A. Synergistic effect of seaweed extract and urea fertilizer on growth and biochemical parameters of *Triticum aestivum* at vegetative stage. *Eur Online J Nat Soc.* 2017;6(4):545.
49. Bharath B, Nirmalraj S, Mahendrakumar M, Perinbam K. Biofertilizing efficiency of *Sargassum polycystum* extract on growth and biochemical composition of *Vigna radiata* and *Vigna mungo*. *Asian Pac J Reprod.* 2018;7(1):27-32. <http://dx.doi.org/10.4103/2305-0500.220982>
50. Mikhedkina EI, Melnik II, Tsygankov AV, Kozhich DT, Vasyleiko MV. Synthesis of ethyl 3, 5-dimethyl-4-(aryliminomethyl)-1h-pyrrole-2-carboxylates and their reactions with thioglycolic acid. *CNCH-2018.* 2018:113.
51. Epifano F, Genovese S, Menghini L, Curini M. Chemistry and pharmacology of oxyprenylated secondary plant metabolites. *Phytochemistry.* 2007;68(7):939-53. <https://doi.org/10.1016/j.phytochem.2007.01.019>
52. Ratnayake WN, Ackman RG. Fatty alcohols in capelin, herring and mackerel oils and muscle lipids: II. A comparison of fatty acids from wax esters with those of triglycerides. *Lipids.* 1979;14(9):804-10. <https://doi.org/10.1007/bf02533519>
53. Linton RE, Jerah SL, Ahmad IB. The effect of combination of octadecanoic acid, methyl ester and ribavirin against measles virus. *IJSTR.* 2013;2(10):181-84.