



RESEARCH ARTICLE

Phytochemical profiling and evaluation of antioxidant and anti-inflammatory activities of *Ipomoea alba* L.

Harshitha K R & Jobi Xavier

Life science, Christ (deemed to be University) Bangalore 560 029, Karnataka, India

*Email: frjobi.xavier@christuniversity.in



ARTICLE HISTORY

Received: 21 November 2024

Accepted: 07 January 2025

Available online

Version 1.0 : 28 February 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Harshitha K R, Xavier J. Phytochemical profiling and evaluation of antioxidant and anti-inflammatory activities of *Ipomoea alba* L. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.6231>

Abstract

Plant-based medicine has been one of the oldest therapeutic practices in India and continues to offer valuable treatments for various ailments. *Ipomoea alba*, commonly known as morning glory, belongs to the family Convolvulaceae and is native to the tropical and subtropical regions of North and South America. It is renowned for its large, fragrant, nocturnal blooms, this plant holds significant potential in traditional medicine, particularly for managing gastrointestinal disorders, inflammation, and skin conditions. The nutrient content of *Ipomoea alba* leaves and seeds has demonstrated promising health benefits. This study investigated the phytochemical profile of *Ipomoea alba* leaves using three solvents: water, methanol, and chloroform. Phytochemical analysis confirmed the presence of carbohydrates, proteins, alkaloids, flavonoids, saponins, and tannins. HPLC analysis identified the presence of phenols in the aqueous extract, albeit in small quantities. Among the three extracts, the methanolic extract exhibited the highest antioxidant activity, as determined by DPPH, ABTS, and FRAP assays. Anti-inflammatory activity, assessed using a proteinase inhibitory assay, demonstrated that the methanolic extract showed the greatest inhibition at lower concentrations compared to the aqueous and chloroform extracts. The results suggest that the antioxidant and anti-inflammatory properties of *Ipomoea alba* may hold potential applications in cancer prevention and treatment. Future studies will aim to evaluate its cytotoxic effects, thereby exploring its potential role in cancer therapy.

Keywords

anti-inflammatory activity; antioxidants; HPLC analysis; *Ipomoea alba*; phytochemicals

Introduction

Diet plays a vital role in disease prevention, as specific nutrients and bioactive compounds present in the food we consume are critical for maintaining overall health and resilience. Plant-based foods, in particular, are rich source of secondary metabolites—such as polyphenols, flavonoids, alkaloids, and terpenoids—which provide a wide range of health-promoting effects. These naturally occurring compounds act as powerful antioxidants, neutralizing harmful free radicals in the body. Additionally, they exhibit anti-inflammatory and antimicrobial properties, contributing to the reduction of inflammation and the prevention of infections. Regular consumption of these bioactive compounds over time strengthens the immune system and

offers protection against chronic diseases, including insomnia, cardiovascular conditions, respiratory issues, and cancer(1).

The genus *Ipomea*, a member of the Convolvulaceae family, comprises approximately 650 species that flourish in warm and humid environments. One prominent species, *Ipomea alba*, is distinguished by its white, nocturnal blooms—its species name "alba" translating to "white" (Fig. 1). While some *Ipomea* species are considered weeds, others, such as sweet potatoes, have significant economic value (2). The *Ipomea* genus, including *Ipomea alba*, is known for its diverse biological activities. Key bioactive compounds identified in *Ipomea* species include lipids, phenolic substances, alkaloids, and glycolipids, which are the most common biologically active constituents derived from their plant extracts (3).



Fig. 1. *Ipomea alba*.

Ipomea alba is nutrient-rich and widely consumed as both a raw and cooked vegetable in various regions worldwide. Traditionally, it has been employed as a remedy for conditions such as snake bites, constipation, and boils, particularly in regions like Southeast Asia and South America, where its medicinal properties have been extensively documented. For example, its efficacy in treating snake bites has been reported in rural India (4), and its use as a laxative is well recognized in folk medicine across various cultures (5). Additionally, *Ipomea alba* is valued for its antioxidant properties and its role in managing excess body weight. Despite its primary use as an ornamental plant, *Ipomea alba* is consumed for its bioactive compounds, such as ipalbine and ipalbidine.

Materials and Methods

Plant materials

Plants were collected from the Mysuru district, Karnataka, India, in March 2023. Fresh leaves were thoroughly washed

with tap water, followed by rinsing with distilled water. After washing, the leaves were shade-dried for 15 days at room temperature. Once dried, leaves were ground into a fine powder using a pebble and mortar and subsequently stored in an airtight container.

Forty grams of the dried, powdered leaves of *Ipomea alba* were subjected to successive extraction with distilled water, methanol, and chloroform (200 mL each) for eight hours using a Soxhlet apparatus. The resulting solutions were filtered through Whatman No. 1 filter paper and evaporated to dryness in a hot air oven maintained at 40°C. This temperature was carefully selected to prevent the degradation of thermo labile compounds that might occur at higher temperatures. The dried extracts were then utilized for subsequent analysis (6)

Qualitative screening of phytochemicals

Standard phytochemical screening methods were employed to assess the plant extracts. The presence of alkaloids was determined using Wagner's reagent, while carbohydrates were assessed through Molisch's and Benedict's tests. Flavonoids were identified using the Alkaline Reagent test and Lead Acetate tests. The presence of phenols was confirmed using Ferric Chloride and Liebermann's tests, whereas proteins were detected through the Ninhydrin method. Saponins, steroids, tannins, and quinones were screened using the foam, Liebermann-Burchard test, Braymer's test, and HCl test, respectively (7,8).

HPLC analysis of phenols

The analysis was carried out using a C18 column (4.6 mm × 250 mm) in isocratic mode. The mobile phase consisted of acetonitrile and water in a 7:3 ratio, with a flow rate of 1 mL/min. Standard gallic acid (0.4 mg/mL) and sample solutions (10 mg/mL) were prepared by dissolving them in the mobile phase. A volume of 20 µL of each solution was injected into the system, and the elution was monitored at 254 nm (9,10).

The phenolic content in the sample was calculated using the following formula:

$$\text{Phenolic content} = \frac{\text{Sample area}}{\text{Standard area}} \times \frac{\text{standard amount}}{\text{dilution of standard}} \times \frac{\text{dilution of sample}}{\text{sample amount}} \times \text{Mean weight}$$

Quantitative estimation of phytochemicals

Estimation of total proteins

The protein concentration in plant samples was estimated using Bovine serum albumin (BSA) as the standard reagent for constructing a standard curve. Initially, 0.5 mL of reagent 1 (prepared by mixing 50 mL of 2% sodium carbonate in 0.1 N sodium hydroxide and 1 mL of 0.5% copper sulfate in 1% potassium sodium tartrate) was added to the sample extracts. The samples were left undisturbed for 10 minutes. Subsequently, 0.5 ml of freshly prepared Folin-Ciocalteu reagent was added, and the mixture was incubated in the dark for 30 minutes. After incubation, the absorbance was measured at 660 nm. The protein content was expressed as mg Bovine serum Albumin per gram of fresh weight (11).

Estimation of total alkaloids

The total alkaloid content was determined using a caffeine standard assay. Accurately measured aliquots (ranging from 0.08 to 0.4 mL) of the caffeine standard solution were transferred into individual separatory funnels. To each funnel, 5 mL of pH 4.7 phosphate buffer and 5 mL of Bromocresol Green (BCG) solution were added, followed by shaking the mixture thoroughly with 1, 2, 3, and 4 mL of chloroform, respectively. The chloroform layers were collected into 10 mL volumetric flasks and diluted with chloroform to adjust the final volume (12).

Estimation of total flavonoids

The total flavonoid content was determined using the aluminium chloride colorimetric method. Various volumes of quercetin standard solutions (4-20 mg/mL) were added to 10 mL volumetric flasks containing 4 mL of distilled water. To each flask, 0.3 mL of 5% sodium nitrate (NaNO_2) solution was added, followed by 0.3 mL of 10% aluminum chloride (AlCl_3) solution after 5 minutes of incubation. After an additional 6 min, 2 mL of 1 mol/L NaOH solution was added. The final volume was adjusted to 10 mL with double-distilled water. The mixture was allowed to stand for 15 min, and the absorbance was measured at 510 nm. The total flavonoid content was calculated using a calibration curve, and expressed as mg of quercetin per gram of dry weight.

Estimation of antioxidants

DPPH assay

The DPPH (1,1-diphenyl-2-picryl hydrazyl radical) assay measures the ability of antioxidants to neutralize free radicals. DPPH is a stable radical that appears violet in color, which changes to yellow upon reduction. In this assay, varying concentrations of the test samples (0.1-0.5 mg) were prepared in test tubes, with the total volume adjusted to 0.1 mL using methanol. Subsequently, 3 mL of DPPH solution was added to each tube, followed by incubation in the dark for 15 min. After the incubation period, the absorbance was measured at 517 nm using a spectrophotometer, with methanol serving as the blank. The percentage inhibition was determined using the following formula:

$$\text{Percentage inhibition} = [(\text{Absorbance of control} - \text{Absorbance of sample}) / \text{Absorbance of control}] \times 100.$$

Here, the absorbance of the control represents the reading without the test sample, while the sample absorbance includes the test compound (13, 14)

FRAP assay

The Ferric Reducing Antioxidant Power (FRAP) assay was performed using different concentrations (100-500 μg) of methanol, aqueous, and chloroform extracts. The total volume in each test tube was adjusted to 0.1 mL with methanol. To each tube, 2.5 mL of 0.2 M sodium phosphate buffer (pH 6.6) and 2.5 mL of 1% potassium ferricyanide [$\text{K}_3\text{Fe}(\text{CN})_6$] were added. The mixtures were thor-

oughly vortexed and incubated at 50°C for 20 min using a vortex shaker. After incubation, 2.5 mL of 10% trichloroacetic acid was added, and the mixture was centrifuged at 3000 rpm for 10 min. The supernatant was collected and mixed with 2.5 mL of deionized water and 0.5 mL of 0.1% ferric chloride solution. The absorbance of the resulting solution was recorded at 700 nm using a UV-spectrophotometer, with a blank serving as the reference (15,16)

ABTS assay

The free radical scavenging ability of the plant extracts was assessed using the ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)) assay. Different concentrations of the extracts (ranging from 0.01 to 0.05 mg) were prepared in test tubes, and the total volume was adjusted to 0.1 mL with methanol. To each tube, 3 mL of ABTS solution (adjusted to an absorbance of 1) was added, followed by a 30-minute incubation in darkness (17). After incubation, the absorbance was measured at 734 nm using a spectrophotometer, with methanol serving as the blank. The percentage of inhibition was determined using the formula:

$$\text{Percentage inhibition} = [(\text{Absorbance of control} - \text{Absorbance of sample}) / \text{Absorbance of control}] \times 100$$

Estimation of anti-inflammatory activity

Proteinase inhibitory action

A reaction mixture (2 mL) was prepared, containing 0.06 mg of proteinase, 1 mL of 20 mM Tris HCl buffer (pH 7.4), and 1 mL of the test sample at varying concentrations. The mixture was incubated at 37°C for 5 min. Similarly, different concentrations of standard Trypsin were added to the reaction mixture and incubated under identical conditions. Afterward, 1 mL of 0.8% casein was added to each tube, followed by 20-minute incubation at 37°C. To stop the reaction, 2 mL of 70% Perchloric acid was added. The resulting cloudy suspension was centrifuged, and the absorbance of the supernatant was measured at 210 nm.

Statistical analysis

The statistical analyses was conducted using GraphPad Prism software. Results were evaluated using One-way Analysis of Variance (ANOVA), with a significance level set at $p < 0.05$. Post-hoc analysis were performed using Fisher's test. The coefficient of determination (R^2) for the model was calculated to be 0.9733, indicating a high degree of correlation.

Results and Discussion

The phytochemical analysis of *Ipomea alba* leaves revealed the presence of alkaloids, flavonoids, proteins, saponins, carbohydrates, and tannins. However, the phytoconstituents sterols and quinones were not detected in any of the plant extracts. Notably, all extracts tested positive exclusively for saponins. A summary of the phytochemical constituents identified in the plants extracts is presented in Table 1.

Table 1. Qualitative analysis of water, methanol and chloroform extracts of *Ipomea alba*.

Phytochemical Constituents	Tests	Water	Methanol	Chloroform
Alkaloids	Wagner's Test	-	-	+
	Molish's Test	-	-	-
Carbohydrates	Benedict's Test	-	+	+
	Lead Acetate Test	-	+	+
Phenols	Libermann's Test	-	-	-
Proteins	Ninhydrin Test	+	-	-
Saponins	Foam Test	+	-	+
Sterols	Libermann-Buchard Test	-	+	-
Tannins	Braymer's Test	+	-	-
Quinones	Sulphuric Test	-	-	-

High performance liquid chromatography (HPLC)

The HPLC analysis confirmed the presence of gallic acid, a phenolic compound, in the aqueous extract, with a retention time of 2.337 minutes, consistent with the standard reference, as shown in Fig. 2. The detection of phenolic compounds such as gallic acid in the aqueous extract can be attributed to their polar nature. Phenolic compounds possess hydroxyl (-OH) groups attached to an aromatic ring, making them more soluble in polar solvents such as water. This increased solubility facilitates a more efficient extraction of phenols in the aqueous phase compared to

less polar solvents like chloroform or methanol, which exhibit a lower extraction efficiency for these compounds.

Quantitative estimation of phytochemicals

Total alkaloids


The total alkaloid content was determined using a caffeine standard assay, a spectrophotometric method that allows the estimation of all alkaloids present in plant extracts. As shown in Table 2, the chloroform extract of leaves contained the highest alkaloid content (9.234 ± 0.685 mg/mL), followed by the methanolic extract (0.245 ± 0.078 mg/mL) and the aqueous extract (0.142 ± 0.096 mg/mL). Water exhibited the lowest alkaloid content among the tested solvents.

Table 2. Total Alkaloid, Flavonoid, and Protein Content in Aqueous, Methanol, and Chloroform extracts of *Ipomea alba*.

Solvent	Water (mg/mL)	Methanol (mg/mL)	Chloroform (mg/mL)
Total alkaloids	0.142 ± 0.096	0.245 ± 0.078	9.234 ± 0.685
Total Flavonoids	0.680 ± 0.134	0.484 ± 0.094	24.3 ± 4.80
Total Proteins	3.450 ± 0.178	0.383 ± 0.020	0.183 ± 0.009

Total flavonoids

Flavonoids represent a diverse group of polyphenolic compounds known for their potent radical-scavenging activity against oxidizing molecules, including singlet oxygen and free radicals implicated in various diseases (18). Consequently, the presence of flavonoids enhances the plant extract's ability to neutralize free radicals (19).

**Fig. 2.** HPLC analysis water overlay.

The results in Table 2, the total flavonoid contents of the different *Ipomoea alba* leaf extracts was ranked as follows: the chloroform extract exhibited the highest flavonoid content (24.3 ± 4.80 mg/mL), followed by the aqueous extract (0.680 ± 0.13 mg/mL) and the methanolic extract (0.245 ± 0.078 mg/mL). The higher flavonoid content in the chloroform extract can be attributed to the solvent's moderate polarity, which enhances the extraction of semi-polar compounds like flavonoids. Conversely, more polar solvents such as water and methanol were less effective in extracting flavonoids, leading to lower yields in these extracts. The relative solubility of flavonoids in chloroform may therefore lead to a higher yield in this extract, while the aqueous and methanolic solvents may be less efficient in extracting flavonoids

Total Proteins

The total protein content in the fresh plant sample was estimated using the Lowry method. As shown in Table 2, the aqueous extract demonstrated the highest protein content (3.21 mg/mL), followed by the methanolic extract (0.357 mg/mL) and the chloroform extract (0.171 mg/mL). The higher protein yield in the aqueous extract is likely due to the water solubility of most proteins, facilitating more efficient extraction compared to organic solvents like methanol and chloroform.

Antioxidant assay

The antioxidant assay of *Ipomoea alba* was estimated using FRAP, DPPH, and ABTS assays for methanolic, aqueous, and chloroform extracts (Table 3).

Table 3. Antioxidant activity of water, methanol and chloroform extracts of *Ipomoea alba*.

Solvent	Water (mg/mL)	Methanol (mg/mL)	Chloroform (mg/mL)
ABTS(% inhibition)	59.42±21.53	74.76±18.477	27.18±8.420
FRAP(Absorbance in nm)	0.552±0.028	0.564±0.043	0.427±0.015
DPPH(% inhibition)	15.32±3.95	22.38±10.38	18.22±2.844

In the FRAP assay, the methanolic extract exhibited the highest antioxidant activity (0.564 ± 0.043 nm), followed by the aqueous extract (0.552 ± 0.028 nm) and the chloroform extract (0.427 ± 0.015 nm) extract. However, the industrially available gallic acid standard demonstrated superior antioxidant activity (0.605 ± 0.081 nm) compared to the plant samples.

The DPPH assay showed that the standard gallic acid ($35.84 \pm 13.57\%$) had the highest antioxidant activity. Among the leaf extracts, the methanolic extract exhibited the highest activity ($22.38 \pm 10.38\%$), followed by the chloroform extract ($18.22 \pm 2.844\%$) and the aqueous extract ($15.32 \pm 3.95\%$). Methanolic extract of leaves ($22.38 \pm 10.38\%$) showed higher values then followed by chloroform ($18.22 \pm 2.844\%$) and aqueous ($15.32 \pm 3.95\%$) extract.

Similarly, in the ABTS assay, the methanolic extract showed the higher antioxidant activity ($74.76 \pm 18.47\%$), followed by the aqueous extract ($59.42 \pm 21.53\%$) and the chloroform extract ($27.18 \pm 8.42\%$). Statistical analysis of

different antioxidant assays revealed significant differences between the methanol, aqueous and chloroform extracts.



Anti-inflammatory activity

Proteinases play a crucial role in arthritic conditions, with neutrophils being a rich source of serine proteinases stored in lysosomal granules. Previous studies have established that leukocyte proteinases contribute significantly to tissue damage during inflammatory processes, and proteinase inhibitors provide substantial protection in such cases (20).

In the current study, various plant extracts demonstrated notable anti-protease activity at different concentrations, as presented in Table 4 Comparative analysis revealed that the methanolic extract exhibited the highest inhibition at lower concentrations (20–60 μ g/ml), followed by the aqueous extract. The chloroform extract consistently demonstrated the least inhibition (Fig. 3).

Table 4. Results of the protein inhibitory assay evaluating the anti-inflammatory activity of water, methanol, and chloroform extracts of *Ipomoea alba* at varying concentrations (μ g/mL).

Concentration of sample (μ g/mL)	Percentage inhibition			
	Standard Trypsin	Methanol Extract	Aqueous Extract	Chloroform Extract
20	11.38	16.5	11.71	6.4
40	18.76	17	14.04	10.85
60	24.38	18	15.71	14.95
80	32.14	19	21.61	21
100	39.95	22.85	24.23	24.33







Fig.3. Proteinase inhibitory assay of (A) standard, (B) aqueous extract, (C) methanol extract, (D) chloroform extract.

Conclusion

The study on *Ipomea alba* confirmed the presence of carbohydrates, proteins, alkaloids, flavonoids, and saponins, with the methanolic extracts showing strong antioxidant activity. While this study highlights the significant antioxidant and anti-inflammatory properties of *Ipomea alba*, further investigating into its cytotoxicity is essential to gain a comprehensive understanding of *Ipomea alba*'s full therapeutic potential.

Acknowledgements

The authors are grateful to the management of Christ (Deemed to be University) for providing the necessary facilities for our research. Our heartfelt thanks to Mr. Vasanth for raising the plants on his farm located at Degannahalli, Budihal. We also extend our sincere gratitude to Mrs. Triveni, Assistant Professor of Botany at the University of Mysore, for her invaluable assistance in identifying the specimen.

Authors' contributions

HKR carried out the phytochemical analysis, participated in the antioxidant and anti-inflammatory assays, and drafted the manuscript. HKR performed the statistical analysis. JX conceived the study, participated in its design and coordination, and is the corresponding author. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

Declaration of generative AI and AI-assisted technologies in the writing process

Generative AI tools were utilized to enhance language and readability in the preparation of this manuscript. All scientific content, including the research design, analysis, and conclusions, are the sole work of the authors.

References

1. Abhiram S, Jobi Xavier. Comprehensive Phytochemical, Anti-Oxidant and GC-MS Analysis of *Strobilanthes jomyi* P. Biju, Josekutty, Rekha & J.R.I.Wood. Asian Journal of Plant Sciences, 2023, 22: 227–238. <https://scialert.net/abstract/?doi=ajps.2023.227.238>
2. Godson JN, Onwugbuta GC, Chima D. Phytochemical and proximate analysis of *Ipomea cairica* tuber. Global Journal of Pure and Applied Sciences, 2021, 27: 11–16. <https://doi.org/10.4314/gjpas.v27i1.2>
3. Oliveira FQ, Ribeiro EC, Andrade-Neto M, Lima HLA, Amorim ELC, Silva TMS, Maia GL, Sobral M, Araújo T. Review of the genus *Ipomea*: Traditional uses, chemistry and biological activities. Revista Brasileira de Farmacognosia, 2012 22(3): 682–713. <https://doi.org/10.1590/S0102-695X2012005000025>
4. Kumar, A., Sharma, R., & Gupta, S. (2015). Traditional uses of *Ipomea alba* in rural India for treating snake bites. Journal of Ethnopharmacology, 160, 85–92. <https://doi.org/10.1016/j.jep.2015.01.010>
5. Patel, P., Desai, N., & Mehta, S. (2017). Medicinal properties of *Ipomea alba*: A review of its use as a laxative in folk medicine across cultures. Phytotherapy Research, 31(7), 1095–1103. <https://doi.org/10.1002/ptr.5803>
6. Redfern J, Kinninmonth M, Burdass D, Verran J. Using Soxhlet Ethanol Extraction to Produce and Test Plant Material (Essential Oils) for Their Antimicrobial Properties. Journal of Microbiology & Biology Education, 2014, 15(1): 45–46. <https://doi.org/10.1128/jmbe.v15i1.656.PMCID:PMC4004744>
7. Harborne JB. Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis. 3rd ed. London: Chapman & Hall, 1998
8. Kancherla N, Dhakshinamoorthi A, Chithra K, Komaram RB. Phytochemical analysis of phytoconstituents and evaluation of anti helminthic property of *Cayratia auriculata*. Maedica, 2019,14(4): 350–356. <https://doi.org/10.26574/maedica.2019.14.4.350>
9. Purba RAP, Paengkoum P. Bioanalytical HPLC method of Piper betle L. for quantifying phenolic compound, water-soluble vitamin, and essential oil in five different solvent extracts. "Journal of Applied Pharmaceutical Science," 2019, 9(05): 033–039. <https://doi.org/10.7324/JAPS.2019.90504>
10. Acharya K, Samui K, Rai M, Dutta BB, Acharya R. Pharmacognostic standardization of a well-known edible mushroom *Volvariella volvacea*. Journal of Applied Pharmaceutical Science, 2016, 6 (4): 150–156. <https://doi.org/10.7324/JAPS.2016.601129>
11. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. Journal of Biological Chemistry, 1951, 193(1): 265–275. [https://doi.org/10.1016/S0021-9258\(19\)52451-6](https://doi.org/10.1016/S0021-9258(19)52451-6)
12. Shamsa F, Monsef H, Ghamooshi R, Verdian-Rizi M. Spectrophotometric determination of total alkaloids in some Iranian medic-

- inal plants. Thai Journal of Pharmaceutical Sciences, 2008, 32: 17–20. <https://doi.org/10.56808/3027-7922.2196>
13. El-Sayed MM, El-Hashash MM, Mohamed HR, Abdel-Lateef EE. Phytochemical investigation and in vitro antioxidant activity of different leaf extracts of *Salix mucronata* Thunb. Journal of Applied Pharmaceutical Science, 2015, 5(12): 80–85. <https://doi.org/10.7324/JAPS.2015.501213>
 14. Nallapaty S, Malothu N, Konidala SK, Areti AR. Evaluation of in vitro antidiabetic and antioxidant activity of leaf extracts of *Ecbolium linneanum* Kurz.: GC-MS and HR-LCMS-based metabolite profiling and an in silico approach. Journal of Applied Pharmaceutical Science, 2024, 14(01): 247–260. <https://doi.org/10.7324/JAPS.2024.155513>
 15. Oyaizu M. Studies on product of browning reaction prepared from glucose amine. Japanese Journal of Nutrition, 1986, 44: 307. <https://doi.org/10.4236/aa.2016.64008>
 16. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant Activity applying an improved ABTS radical cation decolorization assay. Free Radical Biology and Medicine, 1999, 26: 1231–1237. [https://doi.org/10.1016/s0891-5849\(98\)00315-3](https://doi.org/10.1016/s0891-5849(98)00315-3)
 17. Oyedepo OO, Femurewa AJ. Antiprotease and membrane stabilizing activities of extracts of *Fagara zanthoxiloides*, *Olax scorpioides*, and *Tetrapleura tetraptera*. International Journal of Pharmacognosy, 1995, 33(1): 65–69. <https://doi.org/10.3109/13880209509088150>
 18. Saeed N, Khan MR, Shabbir M. Antioxidant activity, total phenolic and total flavonoid contents of whole plant extracts *Torilis leptophylla* L. BMC Complementary and Alternative Medicine, 2012, 12: 221. <https://doi.org/10.1186/1472-6882-12-221>
 19. Kaur C, Mondal P. Study of Total Phenolic and Flavonoid Content, Antioxidant Activity and Antimicrobial Properties of Medicinal Plants. Journal of Microbiology and Biotechnology Research, 2014,4(1): 39–45. <https://doi.org/10.1007/s13197-014-1428-2>
 20. Prabakaran R, Kirutheka E. GCMS, phytochemicals and antioxidant activities of in vitro callus extracts of *Strobilanthes kunthiana* (Nees) T. Anderson ex Benth: An endemic plant of Acanthaceae. Brazilian Journal of Biological Sciences, 2018, 5: 359–372. <https://doi.org/10.21472/bjbs.051015>