



RESEARCH ARTICLE

Enhancing plant growth, yield and nutritional quality of brinjal through Arka Microbial Consortium (AMC) applications

Utpal Das¹, S Ramesh Kumar¹, Madhusmita Dishri¹, Manglambika B¹, Gomathi A¹, Shiva Sai Prasad² & Satya Prakash Barik^{3*}

¹Department of Horticulture and Food Science, VIT School of Agricultural Innovations and Advanced Learning, Vellore Institute of Technology, Vellore 632 014, Tamil Nadu, India

²Department of Agriculture, Koneru Lakshmaiah Education Foundation, Green Fields, Vaddeswaram 522 302, Andhra Pradesh, India

³Department of Agriculture and Allied Sciences, CV Raman Global University, Bhubaneswar 752 054, Odisha, India

*Correspondence email - satyahort@gmail.com

Received: 09 January 2025; Accepted: 25 April 2025; Available online: Version 1.0: 08 August 2025

Cite this article: Utpal D, Ramesh KS, Madhusmita D, Manglambika B, Gomathi A, Shiva SP, Satya PB. Enhancing plant growth, yield and nutritional quality of brinjal through Arka Microbial Consortium (AMC) applications. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.7139>

Abstract

Brinjal (*Solanum melongena* L.) is the "poor man's vegetable" and prized for its affordability, nutritional value and medicinal uses. The Elavambadi brinjal (VRM-1) is a Geographical Indication (GI)-tagged variety from Tamil Nadu, India. It is known for its tender texture, fewer seeds and health benefits, particularly for managing diabetes and liver ailments. This study investigated the impact of the Arka Microbial Consortium (AMC), a carrier-based bio-fertilizer containing nitrogen-fixing, phosphorus, zinc solubilizing and plant growth-promoting microbes, on the growth, yield and quality of Elavambadi brinjal. The treatment T₅, which combined soil drenching, rhizosphere soil and foliar application of AMC, resulted in the highest performance across multiple parameters. It recorded the maximum plant height, branching, stem diameter, fruit yield (8704.67 g/plant), ascorbic acid content (107.81 mg/100 g) and phenol content (90.49 mg Gallic Acid Equivalent (GAE)). Additionally, T₅ demonstrated superior cooking quality and reduced weight loss during frying (65.96 %) and baking (11.97 %). These results highlight AMC's ability to enhance nutrient uptake, plant growth and the synthesis of bioactive compounds. The findings emphasize that AMC is a potentially effective bio-fertilizer for improving brinjal production, enhancing nutritional quality and promoting sustainable agricultural practices.

Keywords: empirical model; global models; hydrological model; sensitivity; uncertainty

Introduction

Brinjal (*Solanum melongena* L.) (2n=24) is widely known as the poor man's vegetable (1) due to its cheap source of vitamins and minerals. It is used as a principal component in the pharmaceutical industry for the treatment of diseases like high cholesterol, liver and skin diseases and throat problems (2, 3). Brinjal is the third most widely cultivated crop in the Solanaceae family after tomato and potato. It is grown in an area of about-1.87 million hectares with a total production of 56.62 million tons in 2020-21, with a productivity of 30.27 tonnes/ha (4). Elavambadi brinjal (VRM-1) or Mullu Katharikai is a Geographical Indication (GI) tagged cultivar of brinjal predominantly grown in the Vellore and Tiruvannamalai districts of Tamil Nadu, India, known for its tender texture, fewer seeds and slightly sweet taste (5). The Elavambadi brinjal is characterized by a typical purple hue, spines on the stem, calyx and leaves (6). In the Vellore district, the cultivation of Elavambadi brinjal spans approximately 1397 ha with an annual production of around 41100 tonnes. The unripe fruits of this variety are commonly used as cooked vegetables and are reputed for their health benefits. Notably, Elavambadi brinjal is

considered beneficial for diabetic patients. Additionally, when the fruits are fried in sesame oil, they are believed to alleviate toothaches and serve as a remedy for individuals suffering from liver ailments (7).

Soil microorganisms enhance food production by influencing nutrient accumulation, organic matter decomposition, carbon dynamics and plant resilience to abiotic and biotic stresses. Microbial Consortia (MC), which consist of multiple microbial species or strains coexisting symbiotically, often exhibit superior performance compared to single-species inoculants (8). In brinjal cultivation, biofilm-based bio-fertilizers and plant growth-promoting rhizobacteria within MC have demonstrated potential benefits. These microbial consortia contribute to nutrient solubilisation from the soil's native reserves, enhance root growth by producing beneficial compounds such as siderophores, phytohormones and mitigate the effects of growth-inhibitory hormones, ultimately improving plant health and productivity. The current experiment aimed to maximize AMC efficiency in increasing the growth and yield of Elavambadi brinjal by testing different inoculation methods (9).

Materials and Methods

The experiment was carried out in the Instructional Farm Area (Latitude: 12.967886°, Longitude: 79.162115°), Department of Horticulture and Food Science, Vellore Institute of Technology, Vellore to study the effect of Arka Microbial Consortium (AMC) in seedling growth, yield and quality parameters of Elavambadi brinjal (VRM-1). The nursery-raising method was adopted to raise healthy, disease-free seedlings of brinjal. The media consisted of cocopeat and vermicompost (7:3 ratio). The seedlings were ready for transplanting at 35 days after sowing. The main field was cultivated to a fine tilth and well-decomposed farmyard manure was incorporated @ 20 tonnes /ha. Brinjal seedlings were transplanted at a spacing of 60 cm (between plants) x 75 cm (between rows) on the tops of the ridges. Brinjal cultivation was carried out following the recommended fertilization protocol. Farmyard manure (FYM) and chemical fertilizers were applied at 125 kg N, 50 kg P₂O₅ and 70 kg K₂O per hectare. The full dose of phosphorus, along with 20 % of both nitrogen and potassium, was incorporated as a basal application. The remaining 80 % of nitrogen and potassium was administered in two equal split doses at 20 and 40 days after transplanting, following weeding and hoeing. The treatment details include T₁: Seedling dipping with AMC, T₂: Soil drenching with AMC, T₃: rhizosphere soil application of AMC, T₄: Foliar application of AMC, T₅: Soil drenching + Rhizosphere soil + foliar application of AMC, T₆: Seedling dipping + Soil drenching + Rhizosphere soil + foliar application of AMC and T₇: Control (no application of AMC). For seedling dipping, AMC was mixed with water @ 20 g L⁻¹ and the roots of the seedlings were dipped in AMC for 15 min before transplanting. For soil drenching, AMC was mixed with water @ 20 g L⁻¹ and applied in the soil after 10 days of transplanting. For rhizosphere application, 20 g of AMC was applied to each plant equally near the root zone. For foliar application, AMC solution was prepared @ 10g L⁻¹ of water and sprayed 15 days after flowering. The experiment was carried out in a Randomized Block Design (RBD) with seven treatments replicated thrice. Ten healthy plants from each treatment and replication were selected randomly and tagged for recording the observations.

The plant height was measured from the ground level to the top of the inflorescence at the time of maturity and their mean value was taken for analysis of this character at 30, 60 and 90 days after transplanting (DAT). The total number of branches in each of the ten tagged plants was counted and their mean value was taken as the number of branches per plant at 30, 60 and 90 DAT. Stem diameter was measured with the help of a vernier calliper at 30, 60 and 90 days after transplanting. Days to first flowering were recorded by calculating the number of days taken from the date of transplanting till plants in the plot start flowering. The number of fruits from each plant in the population was counted and the mean was taken. For fruit yield per plant, fruits harvested from the tagged plants were weighed and the mean was taken and expressed in grams. The lengths of twenty fruits from the tagged population were measured in cm from the attachment end (stalk end) to the tip (stylar end) and the mean values were calculated as fruit length. After recording the fruit length, the same fruits were also used for measuring the fruit girth. Fruit girth was measured at the widest diameter of the fruit. Average

fruit weight was calculated by dividing the total fruit yield by a total number of fruits harvested.

The ascorbic acid content was determined using a volumetric titration method. A 5 mL aliquot of the standard solution was placed in a 100 mL conical flask, followed by the addition of 10 mL of 4 % oxalic acid. The solution was titrated against a dye solution until a stable pink endpoint was achieved (V₁), with the volume of dye consumed corresponding to the ascorbic acid content. For sample analysis, 1 g of plant material was weighed, homogenized in 4 % oxalic acid and filtered using Whatman No.1 filter paper. The final volume was adjusted to 100 mL. The ascorbic acid concentration was calculated and expressed in milligrams per 100 g of fruit pulp.

Ascorbic acid content (mg/100 g) =

$$\frac{0.5 \text{ mg} \times V_2 \times 100 \text{ mL} \times 100}{V_1 \text{ mL} \times 5 \text{ mL} \times \text{weight of sample}}$$

The total phenolic content (TPC) of the brinjal extracts was determined calorimetrically following the method described with some modifications (10). A five-gram sample was pulverized and extracted in 80 % acetone the volume was made up to 50 mL. The extract was then filtered using Whatman's No. 1 filter paper and centrifuged for 15 min at 4000 rpm. 0.15 mL aliquot of the supernatant was mixed with 0.15 mL of 0.25 N Folin-Ciocalteu reagent, 2.4 mL of distilled water in a glass test tube properly mixed and incubated for 3 min. 0.3 mL of 1N sodium carbonate was added and the mixture was incubated at room temperature for 2 hr under dark conditions. Absorbance was measured using a UV spectrophotometer at 760 nm with distilled water as the blank. A gallic acid standard was prepared and the total phenolic content of the brinjal fruit extract was expressed as mg GAE g⁻¹ fresh weight of brinjal (11, 12).

Results and Discussion

Plant growth parameters

The growth performance of brinjal was significantly influenced by the different treatments of the AMC (Table 1). Among all treatments, T₅ (Soil drenching + Rhizosphere soil + Foliar application of AMC) exhibited the highest plant growth, with the maximum plant height observed at 30 days after transplanting (DAT) (27.70 cm), 60 DAT (51.67 cm) and 90 DAT (71.28 cm). This was followed by T₆ (Seedling dipping + Soil drenching + Rhizosphere soil + Foliar application of AMC), with plant heights of 25.91 cm, 42.08 cm and 60.34 cm at 30, 60 and 90 DAT, respectively. The least growth was observed in T₁ (Seedling dipping with AMC), with a plant height of 13.99 cm at 30 DAT and 52.85 cm at 90 DAT.

The number of branches per plant was highest in T₅ at 30 DAT (5.33), 60 DAT (7.67) and 90 DAT (10.43), followed by T₇ (Control) with values of 4.52 (30 DAT), 7.71 (60 DAT) and 9.57 (90 DAT). T₄ (Foliar Application of AMC) showed the least number of branches, particularly at 30 DAT (3.43). T₁ displayed fewer branches at 60 DAT (5.52) and 90 DAT (5.05). This indicates that multiple applications of AMC through various methods (soil drenching, rhizosphere and foliar treatments) positively impacted plant branching.

Table 1. Effect of AMC on morphological traits of brinjal

	Plant height (30 DAT) (cm)	Plant height (60 DAT) (cm)	Plant height (90 DAT) (cm)	Number of branches (30 DAT)	Number of branches (60 DAT)	Number of branches (90 DAT)	Stem diameter (30 DAT) (cm)	Stem diameter (60 DAT) (cm)	Stem diameter (90 DAT) (cm)
T ₁	13.99 ^d	33.75 ^d	52.85 ^d	3.66 ^{cd}	5.05 ^d	5.52 ^d	1.30 ^d	2.60 ^d	3.66 ^c
T ₂	18.63 ^c	31.64 ^e	57.69 ^{bc}	4.38 ^{bc}	6.15 ^c	9.57 ^{ab}	1.40 ^{cd}	2.39 ^d	3.30 ^c
T ₃	15.32 ^d	24.83 ^f	54.48 ^{cd}	3.44 ^d	6.29 ^c	7.22 ^c	1.74 ^{bcd}	3.38 ^{bc}	2.79 ^d
T ₄	18.76 ^c	40.19 ^c	58.03 ^{bc}	3.43 ^d	6.76 ^b	5.95 ^d	1.91 ^{abc}	3.11 ^c	3.23 ^{cd}
T ₅	27.70 ^a	51.67 ^a	71.28 ^a	5.33 ^a	7.67 ^a	10.43 ^a	2.45 ^a	4.53 ^a	5.54 ^a
T ₆	25.91 ^{ab}	42.08 ^b	60.34 ^b	4.50 ^b	6.15 ^c	9.38 ^b	1.25 ^d	3.47 ^{bc}	4.88 ^b
T ₇	24.18 ^b	41.97 ^{bc}	57.06 ^{bc}	4.52 ^b	7.71 ^a	7.33 ^c	2.03 ^{ab}	3.64 ^{bc}	4.70 ^b
S. Em.	0.64	0.59	1.19	0.23	0.09	0.30	0.19	0.17 ^b	0.15
S. Ed.	0.91	0.83	1.69	0.33	0.13	0.43	0.27	0.23	0.22
C.D. 0.05	1.98*	1.82**	3.68**	0.72*	0.28**	0.93**	0.60**	0.51**	0.47**
CV (%)	5.41	2.69	3.52	9.72	2.40	6.62	19.61	8.77	6.59

*- statistically significant differences ($p \leq 0.05$); **- statistically significant differences ($p \leq 0.01$) (Duncan's multiple range test)

The stem diameter, a key indicator of plant robustness, was also highest in T₅, with values of 2.45 cm (30 DAT), 4.53 cm (60 DAT) and 5.54 cm (90 DAT). This was followed by T₇ (Control), which exhibited a diameter of 2.03 cm (30 DAT), 3.64 cm (60 DAT) and T₆ at 90 DAT (4.88 cm). T₆ showed the lowest stem diameter at 30 DAT (1.25 cm), while T₂ (Soil drenching with AMC) showed reduced stem growth at 60 DAT (2.39 cm) and T₃ showed the lowest at 90 DAT (2.79 cm). The increased stem diameter in T₅ may be attributed to the synergistic effect of AMC treatments that promote nutrient absorption and stimulate cellular growth in plants.

Flowering and fruit characteristics

The timing of flowering varied significantly across treatments. T₇ (Control) exhibited the earliest flowering at 50 DAT, followed by T₃ (54.76 DAT), while the delayed flowering was observed in T₅ (68.10 DAT) (Table 2). The delayed flowering period may be perceived as a disadvantage, but it is indicative of a robust physiological response to multiple AMC applications, potentially contributing to improved fruit development later in the growth cycle.

Arka Microbial Consortium significantly enhanced the yield of numerous vegetable crops such as radish, lettuce and basil (13-15). In terms of fruit characteristics, T₅ demonstrated the maximum fruit length (7.09 cm), fruit diameter (4.30 cm), fruit weight (75.90 g) and fruit yield (8704.67 g/plant). These values were significantly higher than other treatments, indicating that the combined application of AMC via soil drenching, rhizosphere soil application and foliar spraying strongly impacted fruit quality and yield. T₆ showed a moderate yield (4388.60 g/plant), while T₁ (Seedling dipping with AMC) had the lowest fruit weight (41.43 g) and fruit yield (1202.75 g/plant). This suggests that the AMC application through multiple routes is essential for achieving higher fruit production.

Table 2. Effect of AMC on flowering and fruiting attributes of brinjal

	Days to first flowering	Fruit length (cm)	Fruit diameter (cm)	Fruit weight (cm)	Number of fruits per plant	Fruit yield per plant (g)
T ₁	55.76 ^{de}	5.18 ^c	3.70 ^b	47.14 ^{cd}	24.00 ^d	1202.75 ^e
T ₂	59.71 ^{bc}	5.26 ^c	2.80 ^d	50.29 ^{bc}	34.00 ^c	2235.26 ^d
T ₃	54.76 ^c	6.41 ^b	3.18 ^c	41.43 ^e	48.33 ^b	3401.02 ^c
T ₄	57.90 ^{cd}	6.19 ^b	3.80 ^b	52.48 ^b	28.33 ^d	2484.31 ^d
T ₅	68.10 ^a	7.09 ^a	4.30 ^a	75.90 ^a	76.67 ^a	8704.65 ^a
T ₆	61.76 ^b	6.20 ^b	3.23 ^c	53.81 ^b	38.00 ^c	4388.60 ^b
T ₇	50.19 ^f	5.37 ^c	2.61 ^d	46.43 ^d	36.33 ^c	2541.05 ^d
S. Em.	0.84	0.17	0.12	1.22	1.74	142.05
S. Ed.	1.19	0.23	0.16	1.72	2.47	200.89
C.D. 0.05	2.58**	0.51**	0.36**	3.75**	5.37**	437.71**
CV (%)	2.49	4.84	5.99	4.02	7.40	6.90

*- statistically significant differences ($p \leq 0.05$); **- statistically significant differences ($p \leq 0.01$) (Duncan's multiple range test)

Nutrient content and cooking quality

The treatments exhibited significant variations in ascorbic acid content, with T₅ attaining the highest value of 107.81 mg/100 g, indicating superior antioxidant properties (Fig. 1). This result suggests that the Arka Microbial Consortium (AMC) enhances nutrient uptake, as evidenced by the increased ascorbic acid content in treatments such as T₅ and T₄. In contrast, T₆ had the lowest ascorbic acid content (74.19 mg/100 g). This demonstrates the potential of AMC in enhancing the nutritional quality of brinjal by increasing its ascorbic acid content, which is important for human health due to its antioxidant properties. Research demonstrated that the application of bio-fertilizers on brinjal resulted in an increase in ascorbic acid content from 75 mg/100 g to 105 mg/100 g, closely aligning with the highest recorded value of 107.81 mg/100 g in T₅ in the present study. This demonstrates the comparable or superior efficacy of AMC in enhancing nutritional quality (16).

The phenol content of the brinjal was significantly influenced by AMC (Fig. 2). T₅ exhibited the highest phenol content at 90.49 mg GAE, followed by T₄ at 69.57 mg GAE, indicating a substantial enhancement in the synthesis of bioactive compounds. This elevated phenol level underscores the functional improvement of the brinjal attributed to the microbial consortium. Conversely, treatments such as T₇ recorded the lowest phenol content at 43.31 mg GAE, demonstrating the varying effects of AMC on different treatments. This enhancement in phenolic content is crucial for the nutritional and medicinal value of brinjal. The AMC treatment (particularly T₅) positions brinjal as a more phenolic-dense and functionally improved vegetable compared to black beans, especially after cooking. While black beans are still nutritionally valuable, the enhanced phenolic levels and likely stability of AMC-treated brinjal under appropriate cooking conditions (17).




Fig. 1. Effect of AMC on ascorbic acid content of brinjal.




Fig. 2. Effect of AMC on the phenol content of brinjal.

The frying process revealed substantial differences in weight loss among the treatments (Fig. 3). T₅ exhibited the lowest weight loss of 65.96 %, indicating superior retention of structural integrity and moisture, whereas T₇ demonstrated the highest weight loss of 81.73 %. These variations may be attributed to differences in the cell wall composition influenced by the microbial activity of AMC. The reduced weight loss in treatments such as T₅ resulted in improved cooking quality, which is essential for consumer satisfaction and nutritional value. Their research indicates a weight loss by frying resulted in a lower loss of phenolic compounds compared to grilling, indicating that frying may help retain more of the brinjal's nutritional value despite some weight loss (18), compared to research findings of 65.96 % in T₅, suggesting that AMC might offer enhanced retention of structural integrity and moisture during cooking.

Baking-induced weight loss exhibited significant variation, with T₅ demonstrating a minimum weight loss of 11.97 %, while T₇ recorded a maximum weight loss of 40.13 % (Fig. 3). The minimal weight loss observed in treatments such as T₅ and T₂ can be attributed to the enhanced water retention capacity and improved cellular structure, potentially facilitated by the application of AMC. This observation underscores the role of AMC in preserving the structural integrity of the brinjal during cooking. The present study recorded a baking-induced weight loss of 15 %, whereas the study about baking resulted in a significant loss of phenolic compounds, with a reported degradation of 61.4 % after the baking process, indicates the loss of weight, which was higher compared to the present study (18).




Fig. 3. Effect of AMC on cooking quality of brinjal.

Conclusion

The application of Arka Microbial Consortium (AMC) through different treatment methods significantly influenced the growth, development and yield of plants, with T₅ (Soil drenching + Rhizosphere soil + Foliar application of AMC) providing the best results. The study demonstrated that AMC enhanced plant height, branching, stem diameter and fruit yield, while improving the nutritional composition, particularly ascorbic acid and phenol content. Furthermore, AMC treatments contributed to better post-harvest quality by reducing weight loss during frying and baking. These findings suggest that AMC can be an effective tool for improving the growth, productivity and quality of crops, offering potential benefits for sustainable agriculture.

Acknowledgements

The author acknowledges the financial assistance from the Vellore Institute of Technology, Vellore, India, under the SEED Grant research project (F. No.: SG20220046).

Authors' contributions

UD and SPB conceptualized the research work. RKS and MD conducted the field experiment. MB, SSP and GA conducted the nutritional quality assessment. UD and SPB prepared the first draft. All the authors reviewed and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: The authors do not have any conflict of interest to declare.

Ethical issues: None

References

- Som MG, Maity JK. Brinjal. In: Bose TK, Kabir J, Maity TK, Parthasarathy VA, editors. Vegetable Crops. Som MG and Prokash N Publishers, Kolkata. 2002;3(1):265–344
- Mak G. Health benefits of aubergine; 2013 [Accessed: 22 Oct 2015]. Retrieved from: <https://www.healthbenefitstimes.com/>
- Oladosu Y, Rafii MY, Arolu F, Chukwu SC, Salisu MA, Olaniyan BA, Muftaudeen TK. Genetic diversity and utilization of cultivated eggplant germplasm in varietal improvement. Plants. 2021;10(8):1714. <https://doi.org/10.3390/plants10081714>
- FAO FAOSTAT Database Collections. FAO: Rome, Italy; 2020. <https://www.fao.org/faostat/en/#home>
- Arun S, Malaisamy A. An economic analysis of Vellore GI spiny brinjal in organic and inorganic vegetable farming. Asian J Agri Ext Eco Soc. 2024;42(6):262–69. <https://doi.org/10.9734/ajaees/2024/v42i62488>
- Nanthakumar S. Effect of different growth regulators and their response to physiological and biochemical attributes in brinjal (*Solanum melongena* L.). Pharma Innov. 2021;10(11):1328–32. <https://doi.org/10.9734/jsrr/2024/v30i62056>
- Bharathi KI, Shanthi A, Manikandan M. Assessment of genetic diversity of brinjal (*Solanum melongena* L.) in the coastal region of Karaikal, India. J Sci Res Reports. 2024;30(6):407–11. <https://doi.org/10.9734/jsrr/2024/v30i62056>
- Santoyo G, Guzmán-Guzmán P, Parra-Cota FI, Santos-Villalobos SDL, Orozco-Mosqueda M DC, Glick BR. Plant growth stimulation by microbial consortia. Agron. 2021;11(2):219. <https://doi.org/10.3390/agronomy11020219>
- Sadasivam S, Manickam A. Biochemical methods for agricultural science. Willey Eastern Limited, New Delhi; 1992.
- Kriengsak T, Unaroj B, Kevin C, Luis C, David HB. Comparison of ABTS+, DPPH, FRAP and ORAC assays for estimating antioxidant activity from guava fruit extracts. Food Comp Anal. 2006;19(6–7):669–75. <https://doi.org/10.1016/j.jfca.2006.01.003>
- Moehninsi NDA, Brown CR. Phytonutrient content of *Solanum sisymbriifolium* Lam. berries. J Food Comp Anal. 2015;44:73–79. <https://doi.org/10.1016/j.jfca.2015.07.004>
- Ponnam N, Kumari M, Ganesan S, Barik S, Pattanaik M, Adamala AK, Acharya GC. Breeding leafy amaranth (*Amaranthus* spp.) for white rust resistance. South Africa J Bot. 2021;163:794–804. <https://doi.org/10.1016/j.sajb.2023.10.020>
- Teja GS, Nayak MH, Prasanth P, Mamatha A, Praneeth S. Influence of inorganic nutrients and biofertilizers on growth and yield of tropical radish (*Raphanus sativus* L.) Cv. Pusa Chetki. Pharma Innov J. 2021;10:1810–12.
- Aswathi NC, Prasad VM, Sreekumar G. Effect of different concentrations of Arka microbial consortium on the growth, yield and quality of lettuce (*Lactuca sativa*) cv. Grand Rapid under the poly house in Prayagraj agro-climatic condition. Int J Curr Microbiol Appl Sci. 2020;9(12):1135–43. <https://doi.org/10.20546/ijcmas.2020.912.137>
- Baraa AM, Kalaivanan D. Soil microbial community dynamics as influenced by integrated nutrient management practices in sweet basil (*Ocimum basilicum* L.) cultivation. J Horti Sci. 2021;16(1):103–13. <https://doi.org/10.24154/JHS.2021.v16i01.011>
- Singh RP, Kasera S, Singh D. Effect of bio-fertilizers on growth, yield and quality of brinjal (*Solanum melongena* L.) cv. Kashi Uttam. Chem Sci Rev Letters. 2020;9(35):786–91.
- Melini F, Lisciani S, Camilli E, Marconi S, Melini V. Effect of cooking on phenolic compound content and *in vitro* bio accessibility in sustainable foods: A case study on black beans. Sustainability. 2023;16(1):279. <https://doi.org/10.3390/su16010279>
- Martini S, Conte A, Cattivelli A, Tagliazucchi D. Domestic cooking methods affect the stability and bio accessibility of dark purple eggplant (*Solanum melongena*) phenolic compounds. Food Chem. 2021;341:128298. <https://doi.org/10.1016/j.foodchem.2020.128298>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.