



RESEARCH ARTICLE

The soil nutrient status of Shirpur taluka of Dhule district, Maharashtra

Nitin Misal*, Indra Raj Singh & Rajap Shiva Kumar

Department of Soil Science and Agricultural Chemistry, SVKM's College of Agriculture, Shirpur 425 405, affiliated to Mahatma Phule Krushi Vidyapeeth, Rahuri, Maharashtra, India

*Email: nmmisal70@gmail.com



ARTICLE HISTORY

Received: 29 January 2025 Accepted: 16 March 2025 Available online Version 1.0: 15 April 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an openaccess article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (https://creativecommons.org/licenses/by/4.0/)

CITE THIS ARTICLE

Nitin M, Indra RS, Rajap SK. The soil nutrient status of Shirpur taluka of Dhule district, Maharashtra. Plant Science Today (Early Access). https://doi.org/10.14719/pst.7504

Abstract

Soil and water are the most important natural resources and all the basic needs of human beings and plants depend on these natural resources. Soil testing is an essential method for evaluating the soil's nutrient-supplying capacity. In the present study, 37 soil samples were collected from different locations of Tajpuri and Hisale villages of Shirpur Taluka, Dhule district. The chemical parameters such as soil pH, EC, organic carbon, available nitrogen, phosphorus and potassium from soil were determined by using standard procedures. The result shows that the pH of all the soil samples from both the villages was reported in the neutral to alkaline range with mean values of both the villages Hisale and Tajpuri were 7.47 and EC were in normal range with mean values of 0.27 dSm⁻¹ and 0.25 dSm⁻¹ and fit for crop production. The organic carbon content in both villages found in the medium range with the mean of Hisale and Tajpuri villages were 0.59 % and 0.60 %. In Hisale village the available nitrogen and a greater number of samples were in low range, available phosphorus and potassium in medium range. Concern to Tajpuri village most of the samples of available nitrogen, phosphorus and potassium were in medium range.

Keywords

nutrient; nitrogen; organic carbon; phosphorus and potassium

Introduction

Healthy soil functions as a dynamic living system that provides numerous ecosystem services, including supporting water quality, enhancing plant productivity, regulating nutrient recycling and decomposition and mitigating greenhouse gases by removing them from the atmosphere (1). India plants about 142 million hectare of which about 67 million hectare are problematic and nutrient deficient. Soil analysis provides nutrient availability index for judicious crop planning. Soil-test based nutrient management maximizes farm profitability and reduces environmental pollution. Major nutrients in soil like nitrogen, phosphorus and potassium have a pilot role in crop growth and production. Understanding the nutrient status of farms will enable farmers and land managers to optimize crop yield through efficient fertilizer management. The nutrient content of soil is influenced by various factors such as parent material, climate and land management practices. For instance, nutrient availability depends upon the weathering of parent material, decomposition of organic matter and microbial activity. Assessing and monitoring soil nutrient status is crucial for sustainable agricultural practices. It helps identify nutrient deficiencies, prevent over-fertilization and mitigate environmental impacts, thereby promoting balanced nutrient management for long-term soil health and productivity. This study aims to analyze the nutrient status of soil in specific regions to provide a framework for sustainable agricultural recommendations.

NITIN ET AL 2

Materials and Methods

The Shirpur tehsil study area spans 1,106 square kilometres and is located between 19 °2' to 22 °3' North latitude and 74 °10' to 75 °11' East longitude. Situated in the Khandesh region of Maharashtra, Shirpur is the northernmost tehsil of Dhule district, bordering Madhya Pradesh to the north. The tehsil lies within the Tapi basin, with the Tapi River flowing westward through its southern part. The river is joined by tributaries from the north, including the Aner, Arunawati, Kordiand Nandi rivers, depositing alluvial soil along its banks. The climate of Shirpur tehsil is predominantly dry, except during the southwest monsoon season. It falls within the assured rainfall zone, receiving 700 to 900 mm of rainfall annually, with 75 % of the total precipitation occurring during the monsoon, which supports Kharif crop cultivation.

The surface soil samples (0-22.5 cm depth) were collected from 37 locations in two villages (Fig. 1). 19 soil samples from Hisale village and 18 samples from Tajpuri village of Shirpur taluka were collected based on soil variation and the exact sample location was recorded using a GPS. Soil samples were analyzed by using standard procedures. Soil pH and Electrical conductivity were determined in 1:2.5 suspension using a standard pH meter and EC meter by potentiometry and conductometry (2) respectively. Soil organic carbon was analysed by using the wet oxidation method (3). Soil available N was determined by the modified alkaline permanganate method (4), available P by 0.5M NaHCO₃ method (5) and available K by flame photometer (1N NH₄OAc pH 7) method (1).

Results

Status of soil pH and EC

The soil pH range of selected samples from Hisale village ranges from 6.50 to 8.45 with a mean value 7.47 and Tajpuri village ranges from 6.40 to 8.50 with a mean value 7.47 (Table 1). Most of the samples from both the villages were found in between neutral to slightly alkaline range. The pH value is affected by various factors such as parent material, climatic conditions, type of fertilizer used and management practices during crop production. The EC of various soil samples of both the villages were ranged from 0.10 to 0.75 dS m⁻¹ with averages 0.27 dS m⁻¹ in Hisale and 0.25 dS m⁻¹ in Tajpuri village (Table 1).




Fig. 1. Map of Hisale and Tajpuri villages.

Status of organic carbon and available NPK

The organic carbon ranges from 0.27 % to 0.88 % in both the villages with mean of Hisale and Tajpuri villages were 0.59 and 0.60 respectively (Table 2). The organic carbon most of the samples were found in medium range in both the villages (Fig. 2 and 3). The available nitrogen of Shirpur taluka was varied from 125 to 439 kg ha⁻¹ with averages of 245 and 303 kg ha⁻¹ in Hisale and Tajpuri villages, respectively. In Hisale village the available nitrogen status as 68 % of samples are in low and 32 % in medium category and in Tajpuri village the available nitrogen status are 39 % samples are in low, 61 % medium, 0 % are in high category (Fig. 1 and 2). The range of available phosphorus 26.88 to 69.44 kg ha⁻¹ with average 47.91 kg ha⁻¹ and 33.60 to 67.20 kg ha⁻¹ with average 51.02 kg ha⁻¹ in the villages of Hisale and Tajpuri respectively (Table 2). Out of 19 samples, 5 % in low, 79 % medium, 16 % high samples were found in Hisale village and 0 % in low, 67 % medium and 33 % high samples were found in Tajpuri village. Concerning potassium, most of the samples were found in the medium range (32 % low, 68 % medium, 0 % high in Hisale village and 6 % low, 83 % medium, 11 % high in Tajpuri village).

Table 1. The status of pH and EC in the soils of Hisale and Tajpuri village of Shirpur taluka

Village name	Parameter	Range	Mean	
Hisale	pH (1:2.5)	6.50 - 8.45	7.97	_
riisate	EC (ds m ⁻¹)	0.10 - 0.75	0.27	
Tainuri	рН	6.40 - 8.50	7.86	
Tajpuri	EC (ds m ⁻¹)	0.11 - 0.49	0.25	

 $\textbf{Table 2.} \ \ \text{The status organic carbon, available Nitrogen, available } \ P_2O_5 \ \ \text{and available } \ K_2O \ \text{in the soils of Hisale and Tajpuri village of Shirpur taluka}$

Parameter	Range	Mean
Organic carbon (%)	0.27-0.88	0.60
Available Nitrogen (kg ha ⁻¹)	125-376	245
Available P ₂ O ₅ (kg ha ⁻¹)	26.88-69.44	47.91
Available K ₂ O (kg ha ⁻¹)	96.32-239.68	157.98
Organic carbon (%)	0.42-0.78	0.60
Available Nitrogen (kg ha ⁻¹)	125-439	303
Available P ₂ O ₅ (kg ha ⁻¹)	33.60-67.20	51.02
Available K ₂ O (kg ha ⁻¹)	123.20-338.24	230.10
	Organic carbon (%) Available Nitrogen (kg ha ⁻¹) Available P ₂ O ₅ (kg ha ⁻¹) Available K ₂ O (kg ha ⁻¹) Organic carbon (%) Available Nitrogen (kg ha ⁻¹) Available P ₂ O ₅ (kg ha ⁻¹)	Organic carbon (%) 0.27-0.88 Available Nitrogen (kg ha $^{-1}$) 125-376 Available P ₂ O ₅ (kg ha $^{-1}$) 26.88-69.44 Available K ₂ O (kg ha $^{-1}$) 96.32-239.68 Organic carbon (%) 0.42-0.78 Available Nitrogen (kg ha $^{-1}$) 125-439 Available P ₂ O ₅ (kg ha $^{-1}$) 33.60-67.20






Fig. 2. Organic carbon and NPK status of Hisale village of Shirpur taluka.




Fig. 3. Organic carbon and NPK status of Hisale village of Shirpur taluka.

Discussion

The pH of both the villages was found neutral to alkaline range. Generally, soils were alkaline in nature might be due to medium black soil with well-irrigated conditions and pH less than 8 because of soils were light in texture. Similar result was reported in soils of Parola tehsil of Jalgaon District (6) and (7) in soils of Sindkheda tehsil of Dhule district. Concern to EC all 100 % samples were in non-saline range and fit for crop production. This may be due to the basaltic type of parent material with a high content of basic cations. Similar observations were found in previous works (8), from Andhra Pradesh (9), Gujarat (10) and in Maharashtra (11) black soil regions. The organic carbon in most of the samples was low to medium range. This might be due to the rapid decomposition of organic matter as attributed to high temperatures during the summer season of Shirpur tehsil. Similar findings were reported in early findings (12, 13). In available nitrogen, most of the samples from both villages fall under the low to medium category. This might be due to the semi-arid climate of Shirpur taluka with high temperatures which favour faster decomposition of organic matter and decline of organic matter, resulting in low to medium status of available nitrogen. Similar results were found in soils of Guntur District of Andhra Pradesh (7, 14, 15) . The higher number of samples of available phosphorus from both the villages were found in the medium category. This might be due to the soil of both the villages being alkaline in nature and the high pH of the soil the calcium precipitate with phosphorus as Ca-phosphate and reduce phosphorus availability in soil. (9, 16, 17) were found similar results. The medium to high level of available potassium content in the soil can be attributed to the dissolution and diffusion of potassium from the internal crystal lattice of silicate clay minerals, particularly in soils with a high clay content, notably montmorillonite clay minerals (18). The similar result was also explained in early works of available potassium (8, 19, 20, 21).

Conclusion

The pH is neutral to slightly alkaline and EC are safe for all the selected samples, so no need to add any amendments. The overall status of organic carbon, available nitrogen, phosphorus and potassium in Hisale and Tajpuri villages of Dhule district were low to medium category. Thus, the need to increase the recommended dose of fertilizer by 25 %, the site where the soil samples were reported to have a low level of nutrient status. Along with chemical fertilizer, adequate organic fertilizers such as FYM, compost and vermicompost need to be added for improvement of nutrient status as well as organic matter status of soils for sustainable crop production.

Acknowledgements

The authors are grateful to the SVKM's College of Agriculture, Shirpur for providing research area and lab facilities.

Authors' contributions

The authors wish to express their sincere gratitude to IS, for his invaluable guidance and constructive feedback throughout the research process and RSK, NBM for helping for preparation of this manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

References

- Ahat M, Alananbeh MK, Othman YA, Leskovar DI. Soil health and sustainable agriculture. Sustainability. 2020;12(12):4859. https:// doi.org/10.3390/su12124859
- Jackson ML. Soil chemical analysis. New Delhi: Prentice Hall of India. 1973.
- Nelson DW, Sommer LE. Total carbon and organic matter. In: Page AL, editor. Methods of soil analysis, Part II. Agronomy Monograph No. 9. Madison, WI: American Society of Agronomy. 1982.
- 4. Subbiah BV, Asija GL. A rapid procedure for the estimation of available nitrogen in soils. Curr Sci. 1956;25:259-60.
- Watanabe FS, Olsen SR. Test of ascorbic acid methods for phosphorus in water and sodium bicarbonate extract of soil. Proc Soil Sci Soc Am. 1965;21:677-78. https://doi.org/10.2136/ sssaj1965.03615995002900060025x
- Kadu PP. A survey of fertility status of soils of Parola Tahsil of Jalgaon District. Paper presented at: State Level Seminar on Soil Health Enhancement for Food and Environmental Security, Parbhani, India. 2007:12–3.
- 7. Parhad SL, Kondvilkar NB, Khupse SM, Sale RB, Patil TD. Management of soil quality through assessment of macro and secondary nutrient status of Sindkheda Tehsil of Dhule District (M.S.). Int J Chem Stud. 2018;6(3):3098–103.
- 8. Waikar SL, Malewar GU, More SD. Elemental composition of humic and fulvic acid in soils of Marathwada region of Maharashtra. J Maharashtra Agric Univ. 2004;29(2):127–29.
- Desavathu RN, Nadipena AR, Peddada JR. Assessment of soil fertility status in Paderu Mandal, Visakhapatnam District of

NITIN ET AL 4

Andhra Pradesh through geospatial techniques. Egypt J Remote Sens Space Sci.

- Sharma RP, Singh RS, Singh SK, Arora S. Land resource inventory (LRI) for development of sustainable agricultural land use plans using geospatial techniques: A case study of Pata-Meghpar village, Jamnagar District, Gujarat. J Soil Water Conserv. 2018;17 (1):15–24. https://doi.org/10.5958/2455-7145.2018.00003.6
- Naitam RK, Kharche VK, Kadu PR, Mohrana PC, Sharma RP. Field-scale spatial variability of physical properties of black soils of Purna Valley, India, using geostatistical approach. J Soil Water Conserv. 2018;17(4):325–34. https://doi.org/10.5958/2455-7145.2018.00050.4
- Sharma PK, Sood A, Setia RK, Tur NS, Mehara D, Singh H. Mapping of micronutrients in soils of Amritsar District (Punjab): A GIS approach. J Indian Soc Soil Sci. 2008;56(1):34–41.
- Chandrakar K, Srivastava LK, Devdas D. Estimation of available micronutrients on the basis of correlation between physicochemical properties of pH, EC, OCand available Fe, Mn, Cuand Zn in Inceptisol of Akaltara Block of Janjgir District of Chhattisgarh. Asian J Soil Sci. 2013;8(2):416–18.
- 14. Shinde LV, Phalke GB. Chemical composition of soil from Godavari Basin of Beed (M.S.), India. Biosci Discov. 2013;5(1):15–8.
- 15. Ratnakumari S, Subbaramaniam P, Reddy AN. Survey on fertility

- status of cotton-growing soils of Guntur District in Andhra Pradesh. Indian J Agric Chem. 2006;39(1):17–24.
- Meena HB, Sharma RP, Rawat US. Status of macro and micronutrients in some soils of Tonk District of Rajasthan. J Indian Soc Soil Sci. 2006;54:508–12.
- Bhattacharyya T, Pal DK, Easter M. Modeled soil organic carbon stocks and changes in the Indo-Gangetic Plains of India from 1980 to 2030. Agric Ecosyst Environ. 2007;122:84–94. https:// doi.org/10.1016/j.agee.2007.01.010
- Durgude AG. Morphology, characterization, classificationand mapping of salt-affected soils of central campus, Research Farms, Mahatma Phule Krishi Vidyapeeth, Rahuri [PhD thesis]. Rahuri, India: MPKV.1999.
- Jangir A, Sharma RP, Tiwari G, Vasu D, Chattaraj S, Dash B, et al. Status of available major and micronutrients in soils of Kelapur Block, Yavatmal District, Maharashtra. J Soil Water Conserv. 2019;18 (3):241–45. https://doi.org/10.5958/2455-7145.2019.00034.1
- 20. Jibhakate SB, Bhende SN, Kharche VK, Selvalakshmi V. Physicochemical status of soils of Katol Tehsil in Nagpur District. J Soils Crops. 2009;19(1):122–28.
- 21. Patil SS, Patil VC, Al-Gaadi KA. Spatial variability in fertility status of soil surface soils. World Appl Sci J. 2011;14(7):1020–24.