



REVIEW ARTICLE

Carbon vaults on farmlands: Unveiling the carbon sequestration potential of trees for climate resilience

Yoga Soundraraja V G¹, S Radhakrishnan^{2*}, A Balasubramanian¹, K T Parthiban³, M Sivaprakash^{1*}, B Sivakumar¹, C N Hariprasath¹, Mathivanan M⁴, S Navaneetha Krishnan¹ & Ashick Rajah R³

¹Department of Silviculture & Natural Resource Management, Forest College and Research Institute, Tamil Nadu Agricultural University, Mettupalayam 641 301, Tamil Nadu, India

²Department of Natural Resource Management, Horticulture College and Research Institute, Tamil Nadu Agricultural University, Periyakulam 625 604, Tamil Nadu, India

³Department of Agroforestry, Forest College and Research Institute, Tamil Nadu Agricultural University, Mettupalayam 641 301, Tamil Nadu, India

⁴Department of Agronomy, Anbil Dharmalingam Agricultural College and Research Institute, Tamil Nadu Agricultural University, Tiruchirappalli 620 027, Tamil Nadu, India

*Email: radhakrishnan.s@tnau.ac.in, sivaprakash.m@tnau.ac.in



ARTICLE HISTORY

Received: 11 February 2025

Accepted: 23 March 2025

Available online

Version 1.0 : 06 May 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Yoga SVG, Radhakrishnan S, Balasubramanian A, Parthiban KT, Sivaprakash M, Sivakumar B, Hariprasath CN, Mathivanan M, Navaneetha SK, Ashick RR. Carbon vaults on farmlands: Unveiling the carbon sequestration potential of trees for climate resilience. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.7694>

Abstract

Farmlands are emerging as a crucial landscape for sequestering carbon, while trees acting as a natural carbon vault that capture and store atmospheric carbon dioxide in biomass and soil. Understanding the potential of trees in sequestering carbon is essential for assessing their role in climate resilience and sustainable management practices. This review evaluates the carbon storage capacity of various tree species, examining factors such as species allocation, carbon accumulation and sequestration rates. By exploring the existing research, it highlights how various agroforestry systems contribute to long term carbon storage, with sequestration rates ranging from 0.29 to 15.21 Mg C ha⁻¹ year⁻¹ in aboveground biomass and 30 to 300 Mg C ha⁻¹ year⁻¹ in soil. Short-rotation species have demonstrated rapid carbon uptake, while long-rotation species contribute to sustained sequestration over the decades. This review also highlights the challenges in precisely quantifying carbon stocks, emphasizing the need for advanced allometric models, remote sensing and standardized methodologies. Additionally, the potential for monetizing farmland carbon stocks through carbon credits and offset trading is explored, emphasizing the economic viability of tree based carbon sequestration. While policies support afforestation and tree farming implementation gaps need to be filled. In conclusion, trees in farmlands hold immense promise as carbon sink, reinforcing the need for research driven strategies, policy support and financial incentives to enhance their role in mitigating climate change and strengthening climate resilience.

Keywords

agroforestry; biomass; carbon; carbon sequestration; farmland; greenhouse gas

Introduction

There has been an increase in greenhouse gas concentrations in the atmosphere and annual anthropogenic Green House Gas (GHG) emissions which is a significant driver of global climate change, as highlighted in the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (1). Anthropogenic GHG emissions, primarily from fossil fuel combustion, have reached record levels, intensified the greenhouse effect and contributed to global warming (2). Recent data indicate that global per capita GHG emissions rose by 0.4 % in 2022, culminating in an 8.3 % increase from 1990 levels.

Simultaneously, emissions intensity per unit of Gross Domestic Product (GDP) reached a 52-year low in 2022, reflecting a modest decoupling of emissions from economic growth. However, these achievements are insufficient to meet global climate targets (3).

Atmospheric gases like carbon dioxide (CO_2), methane (CH_4) and nitrous oxide (N_2O) are trapped in the atmosphere of the Earth and because of this, the atmosphere amplifies the greenhouse effect, heating the Earth (4). CO_2 is the most important of the total GHG emissions and it comprises 81 % of this total (5). The highest level of atmospheric CO_2 in these recent years is 421 ppm in 2023, a record high, due to human activities such as fossil fuel combustion, deforestation and land use change of the natural carbon cycle (6). The Intergovernmental Panel on Climate Change (IPCC) reports that reaching global carbon neutrality by 2050 with carbon negativity by 2100 requires capturing 6 billion metric tons of CO_2 each year by 2050 and increasing this to 14 billion metric tons annually by 2100 to keep temperature rise below 1.5 °C (1). Carbon sequestration decreases the warming potential of greenhouse gases by capturing and storing atmospheric CO_2 (7). Trees cultivated on agricultural lands within agroforestry systems receive significant interest due to their dual capability of storing carbon in biomass and soil while delivering other ecosystem benefits (8). The IPCC Special Report on Climate (9) demonstrates how agroforestry can mitigate 0.08 to 5.7 Gt CO_2 -eq annually across the world. Planting trees within agricultural systems helps capture carbon while advancing sustainable farming techniques and creating additional revenue streams for farmers alongside conserving biological diversity (10). Integrating trees into farmland enhances crop productivity and soil health by elevating soil fertility through nutrient cycling and microclimate regulation. This integration reduces the impact on food production unlike large-scale afforestation projects and provides environmental and rural livelihood benefits (11). Carbon sequestration capabilities in these systems differ according to tree species and management practices alongside climate conditions and socio-economic circumstances which require comprehensive understanding to maximize their impact on climate change mitigation (12). This review intends to provide a thorough examination of how trees on farmlands can capture carbon. This study will assess how different agroforestry systems perform while investigating implementation challenges and limitations and scrutinizing policy frameworks and incentives that support their adoption. Through synthesizing current knowledge this review aims to deliver practical insights for policymakers as well as researchers and practitioners who address climate change with nature-based strategies.

The carbon sequestration process in trees

Carbon sequestration is the phenomenon of extraction of atmospheric CO_2 and storing them in other carbon pools such as oceanic, pedologic, biotic and geological strata for longer period (Fig. 1). The main aim is to reduce the accumulation of excessive CO_2 in the atmosphere (13).

The primary process of carbon sequestration in trees involves the conversion of atmospheric carbon dioxide into




Fig.1. Carbon sequestration process.

plant biomass through the process of photosynthesis (14), which is then followed by the conversion of biomass into persistent Soil Organic Carbon (SOC) facilitated by the formation of organo-mineral complexes. After the decomposition of plant biomass in soil, a portion of it contributes to the formation of Soil Inorganic Carbon (SIC) in the form of bicarbonates and carbonates (15).

Carbon allocation and partitioning

Once fixed, carbon is distributed among tree components through physiological mechanisms influenced by genetic and environmental factors (16).

Source-sink dynamics

All photosynthetic tissues are sources of carbon and all developing tissues and storage organs are sinks (17). The sink strength of developing meristems is an important factor in governing the movement of resources from sources towards sinks (18). Plant hormones such as auxins and cytokinins also regulate the allocation of carbon in response to dynamic environmental and developmental cues (19).

Biomass allocation patterns

Carbon allocation patterns differ considerably between tree species and are affected by factors such as age, nutrient availability and environmental stressors (20). Trade-offs occur between the distribution of biomass among leaves, stems and roots and these vary among plant species. The more limiting resource controls how much biomass a plant will allocate to the organ that is most critical in obtaining that resource (21). On average, plants allocate 46 % of their Biomass to aboveground parts of the plant (leaves, branches and stems), 21 % to coarse roots and 33 % to their fine roots (22).

Carbon pools in tree-based systems

Knowledge of how carbon is distributed in tree-based systems is necessary for optimizing tree-based system's role on carbon sequestration. These systems contribute to three primary carbon pools viz., aboveground biomass, belowground biomass and SOC.

1. Aboveground Biomass : Often, aboveground biomass accounts for the largest share of carbon storage in trees (23). Carbon content varies among different tree components and species but is approximately 45-50 % of the dry biomass. The scale at which trees store carbon is determined by wood density, with denser woods storing

more carbon per unit volume (24). Since modern remote sensing technologies, particularly LiDAR (Light Detection and Ranging), provide new opportunities for mapping aboveground biomass and carbon storage on large landscapes at high spatial resolution, there is interest and considerable need for developing algorithms that improve scaling these technologies (25).

2. Belowground biomass : Although extremely difficult to measure carbon sequestration accurately, root systems play a critical role in it. Tree biomass is about 20 to 47 % belowground (26) and is a function to a large degree of the age, species and environmental conditions (27). Under the conditions of nutrient poor soils or younger trees the proportion of biomass in root tends to increase to maximise nutrient uptake (26). Belowground biomass also helps to stable soils, retain water and improve the cycling of nutrients in agroforestry systems (28). These benefits contribute to the resilience and productivity of agricultural systems, making belowground biomass a critical focus in tree-based farming.

3. Soil Organic Carbon (SOC): The large pool of organic carbon in terrestrial ecosystems is stored mostly in SOC, which is subject to vegetation inputs and soil processes and positively affects soil fertility for sustaining sustainable agricultural practices (29). Therefore SOC in trees is derived from root exudates, rooted turnover and aboveground litter deposition, which together determine the composition and stability of the soil organic matter (30). SOC dynamics are strongly influenced by roots, whereby the mineralization of soil carbon and the formation of organo-mineral complexes are promoted (31).

Carbon sequestration in agroforestry systems

Integrating trees on farms can improve agricultural productivity, increase carbon sequestration, prevent environmental degradation and support healthy soil and ecosystems (32). Through a meta-analysis Stefano and Jacobson (33), concluded that a shift in simplified land use system such as agricultural systems to a complex land use systems like agroforestry systems had increased the SOC stocks.

The potential for carbon sequestration within agroforestry systems varies according to the type of agroforestry system being implemented, the ecological region and the characteristics of the soil with levels ranging from 0.29 to 15.21 Mg ha⁻¹ year⁻¹ above-ground and 30-300 Mg C ha⁻¹ year⁻¹ in the soil up to 1 m depth (34). Trees can be incorporated into agricultural landscapes through various agroforestry systems, including alley cropping, silvipasture, windbreaks and shelterbelts and home gardens, each offering unique carbon sequestration potentials.

1. Alley cropping : Alley cropping system incorporates parallel rows of trees or shrubs alongside agricultural crops, has demonstrated considerable promise for carbon sequestration. This approach improves carbon retention in both above ground and below ground environments by integrating woody perennials into cultivated areas (35). Incorporation of agricultural crops and tree crops caused SOC stocks to rebound significantly (36). Research has shown that

these systems can sequester 2-4 times more carbon than conventional agricultural practices (37). SOC levels also rise in alley cropping systems. The recurrent addition of organic matter from tree litter and crop residues, along with minimized soil disturbance, fosters SOC accumulation. In temperate regions alley cropping systems could increase SOC stocks by 0.24 to 0.72 Mg C ha⁻¹ yr⁻¹ compared to treeless croplands (38). The carbon sequestration potential of alley cropping varies depending on factors such as tree species, crop type, climate and management practices. Alley cropping systems implemented in Costa Rica exhibit a superior soil organic carbon reservoir and an elevated annual rate of SOC accumulation relative to southern Canada, primarily attributed to the increased contribution from tree prunings (35).

2. Silvipasture system : Silvipasture system combines trees and grazed pastures, while enhancing carbon sequestration, as well as the production of livestock (39). Studies conducted in various countries, including the USA, Brazil, Mexico and India, it is shown that the system is highly effective at sequestering carbon through biomass accumulation and deep root systems (40). A study conducted in Southern Mexico shows that the total carbon stock was 8 % higher in silvipasture with *Leucaena* trees (276.3 Mg C ha⁻¹) compared to open pasture (255.3 Mg C ha⁻¹) (41). Moreover, SOC stocks in grasslands converted to silvipastoral systems have been shown to increase significantly (42) by as much as 36.3 % to 60 % in relation to tree-only systems and as much as 27.1 % to 70.8 % in comparison to pasture systems (43). Total carbon stock sequestered in two silvipasture systems namely *Chrysopogon fulvus* and *Hardwickia binata* based system were around 55.18 t/ha in India during a period of five years (44). In addition silvipasture has also been shown to have broader indirect benefits (differential impact only) including improved soil composition, reduced erosion and biodiversity, all of which indirectly facilitate greater net carbon sequestration and climate change mitigation (39). Therefore, it underlines the superiority of silvipastoral systems as a better alternative for atmospheric carbon sequestration compared to systems based on trees as well as pastures.

3. Windbreaks and shelterbelts : Linear plantings of trees and shrubs, windbreaks and shelterbelts, have long been used to moderate wind velocity, change exchange of moisture and heat and retard pollutant diffusion (45). In addition, these practices not only increase agricultural productivity but also contribute to quality of the environment (46). Agroforestry systems in China, particularly shelterbelts, have been shown to effectively increase SOC stocks in both topsoil and subsoil, with soil type playing a crucial role (47). In Hungary doubling the extent of windbreak plantations established across a combined expanse of 14256 hectares would cumulatively sequester around of 913 kilotons of carbon (48).

4. Home garden systems : Home gardens are advanced agroforestry systems characterized by multi-layered canopy structures that maximize space utilization and carbon capture potential (49). Typically found in tropical regions, these gardens consist of tall trees, medium-sized fruit trees, shrubs, herbs and ground covers, fostering significant

agrobiodiversity (50). The home gardens of Kerala on an average tend to possess 16 to 36 Mg ha⁻¹ of aboveground carbon (51). Smaller gardens tend to have higher SOC concentrations (46.85 Mg ha⁻¹), while larger gardens show greater vegetative biomass carbon (60.38 Mg ha⁻¹) (52).

5. Industrial agroforestry : Industrial agroforestry systems focus on cultivating fast-growing tree species such as *Populus deltoides*, *Eucalyptus tereticornis*, *Melia dubia* and *Casuarina equisetifolia*. In these systems, capturing carbon and supplying industrial wood are dual advantages (53). Industrial agroforestry has become popular in India since the enforcement of Forest Conservation Act of 1980 and National Forest Policy of 1988, when wood-based industries were required to obtain their own raw materials (54). These systems combine structured arrangements of woody perennials to improve the above and below ground pools of carbon storage. Moreover, there are species including *Leucaena leucocephala* and *Dalbergia sissoo* that further enhance soil fertility through nitrogen fixation and enhance long term carbon storage (55).

Carbon sequestration potential of different tree species

1. Short rotation crops : Fast-growing species are trees that grow quickly and accumulate biomass at a fast rate and can achieve a mean annual increment of at least 10 m³ per hectare under favourable conditions, provided proper techniques are used for site preparation, planting and management (56). Short rotation forestry represents a silvicultural approach that employs fast growing species planted at higher densities, accompanied by more intensive management practices compared to conventional forestry, with harvesting cycles ranging from 2 to 25 years (57). Block and boundary plantations are known for their incredibly high carbon sequestration potential. In poplar-based agroforestry systems, block plantations have been shown to sequester 5.45 t ha⁻¹ yr⁻¹ and boundary plantations sequester 1.84 t ha⁻¹ yr⁻¹ and thus represent a means of reducing fossil fuel-based carbon emissions (58). In Coromandel coasts of India, the plantations of *Casuarina equisetifolia* (12 years old) are known to store 2.792 tonnes of total carbon (59). Meanwhile in Gujarat the *Casuarina equisetifolia* plantations sequestered 1609.91 kg of CO₂ per tree by a total of 767.97 t ha⁻¹ in six years which can be witnessed in Fig. 2 (60). In semi-arid region of Northwestern India, *Eucalyptus tereticornis* plantations stored between 114.1 and 118.8 Mg C ha⁻¹ (61). This effectiveness in capturing the atmospheric carbon while supplying sustainable wood and bioenergy products, makes it attractive to farmers.

Leucaena based farm forestry systems in Andhra Pradesh stored 62 t ha⁻¹ and *Eucalyptus* stored 34 t ha⁻¹ for four years with the highest rate of carbon accumulation in the third year (62). Particularly, fast growing species can be used as alternatives to fossil fuels and are very valuable for short term carbon storage (63).




Fig. 2. Carbon sequestration potential of fast-growing species.

2. Long rotation species : Long-lived species grow more slowly than fast growing species, however they store more carbon in larger amounts through longer periods of time (64) (Table 1). These trees play a critical role in both biomass and soil carbon storage, especially when they are grown for extended rotations (69). Studies shows that increasing the rotation period of long-lived species increases carbon storage. Extension of the rotation period from 120 to 150 years resulted in a 12 % increase in the average carbon stock of the forest ecosystem, encompassing both tree biomass and soil carbon (70). Long-duration rotational systems, including agroforestry, homegardens and boundary plantings, possess the capacity to sequester substantial amounts of carbon within both plant biomass and durable wood products (71).

Traditional agroforestry systems in Timor-Leste store an average of 156 Mg C ha⁻¹, up to 210 Mg C ha⁻¹ in forest gardens, comparable to old growth tropical forest (72). As a potential long-term carbon sinks, long lived species of trees are indispensable in the climate change mitigation strategies, as they can store carbon for decades and even centuries (73).

Quantification of carbon sequestration

The measurement of carbon storage in trees helps us understand their impact on climate change mitigation. Different methods and models are utilized for quantifying the carbon sequestration task with its distinct benefits and drawbacks. The simplest approach requires multiplying tree biomass with the average carbon content in dry biomass which generally falls between 47 % and 50 % (74). This basic method delivers rapid assessments even though it fails to provide accurate results when applied to extensive projects (75).

Scientists use allometric equations as a precise approach because these equations link tree growth parameters to carbon storage amounts (76). Tree height and Diameter at Breast Height (DBH) data from a two-parameter equation can be reduced to one-parameter data just utilizing only DBH. This enables the assessment of carbon storage levels without felling the tree by measuring its basal area alone (77). Allometric models serve as essential tools for site-specific and

Table 1. Carbon sequestration potential of long rotation trees in farmlands

Tree Species	Biomass carbon storage (t ha ⁻¹)	Rotation age (Years)	Methodology	Reference
Teak (<i>Tectona grandis</i>)	330.00	20-30	Non-destructive method	(65)
Mahogany (<i>Swietenia macrophylla</i>)	150.17	15-20	Non-destructive method	(66)
Rosewood (<i>Dalbergia latifolia</i>)	42.95	40-50	Destructive Sampling and regression modelling	(67)
Sandalwood (<i>Santalum album</i>)	21.05	30-40	Non-destructive method	(68)

species-specific assessments which makes them broadly useful in forestry research (78).

Remote sensing technologies including LiDAR and multi-spectral aerial imagery represent advanced methods to assess tree coverage, structural attributes and species classification. Experts utilize these methods to evaluate carbon storage and sequestration potential across large geographical areas (79). Studies conducted across urban areas in the U.S. showed that urban forests maintain 7.69 kg C m^{-2} of carbon storage while capturing 0.28 kg C m^{-2} every year (80).

However, different models often yield varying estimates. A study conducted in Bolzano, Italy, showed that the Urban Forest Effects Model produced conservative carbon storage numbers while CUFR Tree Carbon Calculator (CTCC) generated more positive sequestration estimates (81). The US Forest Service's Center for Urban Forest Research's CTCC model generated increased sequestration estimates than satellite-based methods for the North Little Rock, Arkansas region (82). The variation between different methodologies demonstrates why researchers should choose methods that fit their study scale and available resources while meeting precision needs (83). Future carbon accounting will see revolutionary changes through emerging technologies such as machine learning, artificial intelligence and blockchain along with traditional techniques (84, 85).

Monetization of carbon stocks

The monetization of carbon stocks in farmland trees presents significant opportunities for financial incentives, especially for farmers engaging in sustainable practices. This process involves mechanisms such as carbon credit trading and participation in carbon markets, where the quantified sequestered carbon in agroforestry systems can be converted into tradable carbon credits (Fig. 3). These credits, representing sequestered carbon in both trees and soil, are increasingly sought in voluntary and compliance markets (86).

Carbon offset trading

Agroforestry provides opportunities to the farmers in yielding economic benefits through carbon offset trading mechanism. The programs like Chicago Climate Exchange (CCX) allowed industries to offset their emissions by balancing them through investing in carbon reduction activities. This activity not only opened marketplace for sustainable practices but also supports climate change mitigation (87). Additionally, the

REDD+ (Reducing Emissions from Deforestation and Forest Degradation) program, managed by United Nations provides financial assistance to promote agroforestry practices that mitigate greenhouse gas emissions. These programs are well suited for tropical regions where agroforestry is a vital force for sustaining development (88).

Carbon credits trading

Carbon credits are a tradable permit within a market-based framework and the holder has the right to emit one metric tonne of carbon dioxide equivalent (89). So that organizations that emit less than their cap can sell the unused credits. This creates a monetary value for sustainable practices, including agroforestry (90). This gives a monetary price upon sustainable practices like agroforestry. Carbon credits are generated and traded by means of global mechanisms such as the Clean Development Mechanism (CDM) and the Verified Carbon Standard (VCS) (91). These projects are rewarded in the market through credits available from converting these reductions of greenhouse gas emissions, including agroforestry initiatives, which actively reduce carbon. There are 27 cap-and-trade programs in the world, but only a few include forest-based carbon offsets. This indicates an untapped potential for agroforestry systems to expand their role in climate mitigation through carbon credit markets (92).

Policy implications and incentives

Tree-based systems and agroforestry for carbon sequestration have gained global governmental acceptance. The Kyoto Protocol under Article 3.3 (93) supports afforestation and reforestation initiatives as well as wood product utilization to reduce climate change effects (94). Since 1986 Costa Rica has become a leader in reforestation while its payment for environmental services program established in 1996 offers financial rewards for planting native trees to demonstrate the effectiveness of government-backed incentives in carbon sequestration projects (95). The Conservation Reserve Program (CRP) in the United States functions as an essential tool for advancing both conservation initiatives and carbon sequestration practices. The CRP program works to get farmers to convert their cropland into grasslands or forestlands to capture carbon and lower greenhouse gas emissions (96). In the southern high plains region, the program produced yearly soil organic carbon gains of $69.82 \text{ kg C ha}^{-1}$ in the top 10 cm and $132.87 \text{ kg C ha}^{-1}$ in the top 30 cm (97).

The National Forest Policy in India sets the target of growing forest coverage from 23 % to 33 % across all land areas. Silviculture along with agri-silviculture and agri-horticulture practices within agroforestry systems function as key elements to reach this target especially for both rain fed and irrigated lands. The ICAR works to promote agroforestry systems so the nation can better sequester carbon (98).

Several countries have incorporated agroforestry into their climate strategies, but additional targeted approaches are necessary for others. Specific provisions for agroforestry initiatives could be included in the US Farm Bill (99). Unlocking more opportunities for agroforestry to aid in climate change mitigation can be achieved through the simplification of forestry carbon project regulations combined with the examination of voluntary carbon markets and REDD+ programs. Agroforestry stands out as a powerful solution for




Fig. 3. Carbon trading.

Table 2. Carbon sequestration potential of agroforestry tree species

Sl. No	Potential agroforestry tree species	Carbon sequestration rate (kg tree ⁻¹ yr ⁻¹)
1	<i>Neolamarckia cadamba</i>	14.84
2	<i>Melia dubia</i>	6.85
3	<i>Toona ciliata</i>	2.64
4	<i>Tectona grandis</i>	4.48
5	<i>Lagerstroemia lanceolata</i>	6.11
6	<i>Swietenia macrophylla</i>	5.22
7	<i>Dalbergia latifolia</i>	3.54
8	<i>Psidium guajava</i>	1.08
9	<i>Syzygium cumini</i>	1.61
10	<i>Moringa oleifera</i>	1.65

carbon capture (100) (Table 2). To fully tap into the potential benefits of agroforestry and create lasting impacts on global climate goals we need improved policy support along with simplified procedures and financial incentives.

Conclusion

Trees planted in farmlands serve as essential elements for carbon sequestration while helping reduce climate impact and supporting practices for sustainable land management. The ability of tree species to capture carbon varies since short-rotation trees allow quick carbon capture while long-lived trees maintain carbon storage for extended periods. These trees enhance soil fertility along with biodiversity and improve farmers' livelihoods aside from their ability to store carbon. Accurate measurements of carbon stocks remain problematic because of the uneven distribution of biomass and changing soil dynamics. The accuracy of carbon stock measurement is improving because of the advancements in the field of remote sensing and machine learning which leads to increased economic benefits through carbon credits and encourage farmer participation. The implementation of robust policies combined with research funding and the integration of agroforestry practices into climate strategies will improve its effectiveness in global carbon reduction initiatives.

Acknowledgements

I heartily thank my chairman, Dr. S. Radhakrishnan and advisory committee members for guiding me throughout the process, and I am grateful to the Forest College and Research Institute, TNAU, Mettupalayam, for providing me all the facilities.

Authors' contributions

VGYS and SR involved in collecting the articles relevant to the topic and framing the overall outline. AB, KTP and MS, BS, CNH participated in enhancing manuscript scientifically and grammatically. SNK, RAR, MM, helped in creating pictorial representation. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of

interests to declare.

Ethical issues: None

References

- Lee H, Calvin K, Dasgupta D, Krinner G, Mukherji A, Thorne P, et al. IPCC. Climate change 2023: Synthesis report, Summary for policymakers. Contribution of working Groups I, II and III to the sixth assessment report of the Intergovernmental panel on climate change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland. 2023.
- Le Quéré C, Peters GP, Friedlingstein P, Andrew RM, Canadell JG, Davis SJ, et al. Fossil CO₂ emissions in the post-COVID-19 era. *Nature Climate Change*. 2021;11(3):197-99. <https://doi.org/10.1016/j.jclepro.2024.140875>
- Crippa M, Guizzardi D, Solazzo E, Muntean M, Schaaf E, Monforti-Ferrario F, et al. GHG emissions of all world countries. Publications Office of the European Union. 2021.
- Kweku DW, Bismark O, Maxwell A, Desmond KA, Danso KB, Oti-Mensah EA, et al. Greenhouse effect: Greenhouse gases and their impact on global warming. *Journal of Scientific Research and Reports*. 2018;17(6):1-9. <https://doi.org/10.9734/JSRR/2017/39630>
- Amarpuri L, Yadav N, Kumar G, Agrawal S. Prediction of CO₂ emissions using deep learning hybrid approach: A case study in Indian context. In 2019 twelfth international conference on contemporary computing (IC3). IEEE. 2019:1-6.
- Adak S, Mandal N, Mukhopadhyay A, Maity PP, Sen S. Current state and prediction of future global climate change and variability in terms of CO₂ levels and temperature. In *Enhancing resilience of dryland agriculture under changing climate: Interdisciplinary and convergence approaches*, Singapore. Springer Nature Singapore. 2023:15-43.
- Lal R. Carbon sequestration. *Philosophical transactions of the royal society. Biological Sciences*. 2008;363(1492):815-30. <https://doi.org/10.1098/rstb.2007.2185>
- Jose S, Bardhan S. Agroforestry for biomass production and carbon sequestration: An overview. *Agroforestry systems*. 2012;86:105-11.
- Shukla PR, Skeg J, Buendia EC, Masson-Delmotte V, Pörtner HO, Roberts DC, et al. Climate change and land: An IPCC special report on climate change, desertification, land degradation, sustainable land management, food security and greenhouse gas fluxes in terrestrial ecosystems.
- Kumar A, Malik MS, Shabnam S, Kumar R, Karmakar S, Das SS, et al. Carbon sequestration and credit potential of gamhar (*Gmelina arborea* Roxb.) based agroforestry system for zero carbon emission of India. *Scientific Reports*. 2024;14(1):4828.
- Ramachandran Nair PK, Mohan Kumar B, Nair VD. Agroforestry as a strategy for carbon sequestration. *Journal of Plant nutrition and Soil Science*. 2009;172(1):10-23.
- Lorenz K, Lal R. Soil organic carbon sequestration in agroforestry systems. A review. *Agronomy for Sustainable Development*. 2014;34:443-54.
- Lal R, Negassa W, Lorenz K. Carbon sequestration in soil. *Current Opinion in Environmental Sustainability*. 2015;15:79-86. <https://doi.org/10.1016/j.cosust.2015.09.002>
- Anderson JM, Spencer T. Carbon, nutrient and water balances of tropical rain forest ecosystems subject to disturbance: management implications and research proposals.
- Lal R, Smith P, Jungkunst HF, Mitsch WJ, Lehmann J, Nair PR, et al. The carbon sequestration potential of terrestrial ecosystems. *Journal of Soil and Water Conservation*. 2018;73(6):145A-52A.
- Zhou Q, Shi H, He R, Liu H, Zhu W, Yu D, et al. Prioritized carbon

- allocation to storage of different functional types of species at the upper range limits is driven by different environmental drivers. *Science of the Total Environment*. 2021;773:145581. <https://doi.org/10.1016/j.scitotenv.2021.145581>
17. Guillemot J, Martin-StPaul NK, Dufrêne E, François C, Soudani K, Ourcival JM, et al. The dynamic of the annual carbon allocation to wood in European tree species is consistent with a combined source-sink limitation of growth: Implications for modelling. *Biogeosciences*. 2015;12(9):2773-90.
 18. Kaitaniemi P, Honkanen T. Simulating source-sink control of carbon and nutrient translocation in a modular plant. *Ecological Modelling*. 1996;88(1-3):227-40. [https://doi.org/10.1016/0304-3800\(95\)00109-3](https://doi.org/10.1016/0304-3800(95)00109-3)
 19. Lemoine R, Camera SL, Atanassova R, Dédaldéchamp F, Allario T, Pourtau N, et al. Source-to-sink transport of sugar and regulation by environmental factors. *Frontiers in Plant Science*. 2013;4:272. <https://doi.org/10.3389/fpls.2013.00272>
 20. Poorter H, Nagel O. The role of biomass allocation in the growth response of plants to different levels of light, CO₂, nutrients and water: a quantitative review. *Functional Plant Biology*. 2000;27(12):1191.
 21. McCarthy MC, Enquist BJ. Consistency between an allometric approach and optimal partitioning theory in global patterns of plant biomass allocation. *Functional Ecology*. 2007;713-20. <https://doi.org/10.1111/j.1365-2435.2007.01276.x>
 22. Poorter H, Niklas KJ, Reich PB, Oleksyn J, Poot P, Mommer L. Biomass allocation to leaves, stems and roots: Meta-analyses of interspecific variation and environmental control. *New Phytologist*. 2012;193(1):30-50. <https://doi.org/10.1111/j.1469-8137.2011.03952.x>
 23. Thomas SC, Martin AR. Carbon content of tree tissues: a synthesis. *Forests*. 2012;3(2):332-52. <https://doi.org/10.3390/f3020332>
 24. Bredin YK, Peres CA, Haugaasen T. Forest type affects the capacity of Amazonian tree species to store carbon as woody biomass. *Forest Ecology and Management*. 2020;473:118297. <https://doi.org/10.1016/j.foreco.2020.118297>
 25. Lu D, Chen Q, Wang G, Liu L, Li G, Moran E. A survey of remote sensing-based aboveground biomass estimation methods in forest ecosystems. *International Journal of Digital Earth*. 2016;9(1):63-105. <https://doi.org/10.1080/17538947.2014.990526>
 26. Ma H, Mo L, Crowther TW, Maynard DS, van den Hoogen J, Stocker BD, et al. The global distribution and environmental drivers of aboveground versus belowground plant biomass. *Nature Ecology & Evolution*. 2021;5(8):1110-22. <https://www.nature.com/articles/s41559-021-01485-1>
 27. Peichl M, Arain MA. Allometry and partitioning of above-and belowground tree biomass in an age-sequence of white pine forests. *Forest Ecology and Management*. 2007;253(1-3):68-80. <https://doi.org/10.1016/j.foreco.2007.07.003>
 28. Das DK, Chaturvedi OP. Root biomass and distribution of five agroforestry tree species. *Agroforestry systems*. 2008;74:223-30.
 29. Suárez JC, Segura M andrade HJ. Agroforestry systems affect soil organic carbon stocks and fractions in deforested landscapes of Amazonia. *Agroforestry Systems*. 2024;1-3.
 30. Leppälampi-Kujansuu J, Aro L, Salemaa M, Hansson K, Kleja DB, Helmisaari HS. Fine root longevity and carbon input into soil from below-and aboveground litter in climatically contrasting forests. *Forest Ecology and Management*. 2014;326:79-90. <https://doi.org/10.1016/j.foreco.2014.03.039>
 31. Wang Q, Zhong M. Composition and mineralization of soil organic carbon pools in four single-tree species forest soils. *Journal of Forestry Research*. 2016;27:1277-85.
 32. Castle SE, Miller DC, Merten N, Ordóñez PJ, Baylis K. Evidence for the impacts of agroforestry on ecosystem services and human well-being in high-income countries: A systematic map. *Environmental Evidence*. 2022;11(1):10.
 33. De Stefano A, Jacobson MG. Soil carbon sequestration in agroforestry systems: A meta-analysis. *Agroforestry Systems*. 2018;92:285-99. <https://doi.org/10.1007/s10457-017-0147-9>
 34. Nair PR, Nair VD, Kumar BM, Showalter JM. Carbon sequestration in agroforestry systems. *Advances in Agronomy*. 2010;108:237-307. [https://doi.org/10.1016/S0065-2113\(10\)08005-3](https://doi.org/10.1016/S0065-2113(10)08005-3)
 35. Oelbermann M, Voroney RP, Thevathasan NV, Gordon AM, Kass DC, Schlönvoigt AM. Soil carbon dynamics and residue stabilization in a Costa Rican and southern Canadian alley cropping system. *Agroforestry Systems*. 2006;68:27-36. <https://doi.org/10.1007/s10457-005-5963-7>
 36. Hombegowda HC, Van Straaten O, Köhler M, Hölscher D. On the rebound: soil organic carbon stocks can bounce back to near forest levels when agroforests replace agriculture in southern India. *Soil Discuss*. 2015;2:871-902.
 37. Udawatta RP, Jose S. Agroforestry strategies to sequester carbon in temperate North America. *Agroforestry Systems*. 2012;86:225-42.
 38. Cardinael R, Chevallier T, Cambou A, Béral C, Barthès BG, Dupraz C, et al. Increased soil organic carbon stocks under agroforestry: A survey of six different sites in France. *Agriculture, Ecosystems & Environment*. 2017;236:243-55. <https://doi.org/10.1016/j.agee.2016.12.011>
 39. Jose S, Walter D, Mohan Kumar B. Ecological considerations in sustainable silvopasture design and management. *Agroforestry Systems*. 2019;93(1):317-31. <https://doi.org/10.1007/s10457-016-0065-2>
 40. Torres B, Herrera-Feijoo R, Torres Y, García A. Global evolution of research on silvopastoral systems through bibliometric analysis: Insights from Ecuador. *Agronomy*. 2023;13(2):479.
 41. López-Hernández JC, Aryal DR, Villanueva-López G, Pinto-Ruiz R, Reyes-Sosa MB, Hernández-López A, et al. Carbon storage and sequestration rates in *Leucaena leucocephala*-based silvopasture in Southern Mexico. *Agroforestry Systems*. 2024;98(5):1105-21.
 42. Feliciano D, Ledo A, Hillier J, Nayak DR. Which agroforestry options give the greatest soil and above ground carbon benefits in different world regions? *Agriculture, Ecosystems & Environment*. 2018;254:117-29. <https://doi.org/10.1016/j.agee.2017.11.032>
 43. Mangalassery S, Dayal D, Meena SL, Ram B. Carbon sequestration in agroforestry and pasture systems in arid northwestern India. *Current Science*. 2014:1290-93.
 44. Ram SN, Manjanagouda S, Ghosh A, Shukla A. Effect of pruning and grasses on forage productivity, carbon storage, soil health and economics of Anjan tree (*Hardwickia binata*) based silvopasture systems. *Indian Journal of Agronomy*. 2024;69(3). <https://doi.org/10.59797/ija.v69i3.5526>
 45. Sheetal KR, Renjith PS, Subbulakshmi V, Patel A, Soni ML, Kumar R. Shelterbelt and Windbreak research in Arid India: A review of research advances and future directions. *Indian Journal of Agricultural Research*. 2024;58. <https://arccjournals.com/journal/indian-journal-of-agricultural-research/A-6321>
 46. Aili A, Hailiang X, Waheed A, Bakayisire F, Yingying X. Synergistic windbreak efficiency of desert vegetation and oasis shelter forests. *Plos One*. 2024;19(10):e0312876. <https://doi.org/10.1371/journal.pone.0312876>
 47. Hübner R, Kühnel A, Lu J, Dettmann H, Wang W, Wiesmeier M. Soil carbon sequestration by agroforestry systems in China: A meta-analysis. *Agriculture, Ecosystems & Environment*. 2021;315:107437. <https://doi.org/10.1016/j.agee.2021.107437>
 48. Király É, Bidló A, Keserű Z, Borovics A. Climate benefit assessment of doubling the extent of windbreak plantations in Hungary. *Earth*. 2024;5(4):654-69.
 49. Direba L. Agroforestry homegarden's impact on food security and

- income generation. *Global Research in Environment and Sustainability*. 2024;2(4):1-19.
50. Paranavithana TM, Wasalamuni WA, Ratnayake RR. Harnessing home gardens for sustainable agroforestry: A promising approach to reducing greenhouse gas emission. In *Decarbonization strategies and drivers to achieve carbon neutrality for sustainability*. Elsevier. 2024;481-99. <https://doi.org/10.1016/B978-0-443-13607-8.00022-5>
 51. Kumar BM. Species richness and aboveground carbon stocks in the homegardens of central Kerala, India. *Agriculture, Ecosystems & Environment*. 2011;140(3-4):430-40. <https://doi.org/10.1016/j.agee.2011.01.006>
 52. Subba MO, Pala NA, Shukla GO, Chakravarty SU. Study of the variability of home gardens influencing carbon stock under sub-humid tropical zone of West Bengal, India. *Indian Forester*. 2018;144(1):60-6.
 53. Parthiban KT, Fernandaz CC, Sudhagar RJ, Sekar I, Kanna SU, Rajendran P, et al. Industrial agroforestry-A sustainable value chain innovation through a consortium approach. *Sustainability*. 2021;13(13):7126.
 54. Parthiban KT, Umarani R, Kanna SU, Sekar I, Rajendran P, Durairasu P. Industrial agroforestry perspectives and Prospectives. Scientific Publishers; 2014.
 55. Chavan SB, Dhillon RS, Ajit, Rizvi RH, Sirohi C, Handa AK, et al. Estimating biomass production and carbon sequestration of poplar-based agroforestry systems in India. *Environment, Development and Sustainability*. 2022;1-29. <https://link.springer.com/article/10.1007%2Fs10668-021-01996-8>
 56. Badalamenti E, Sferlazza S, Veca DS, Maetzke F, Sala G, La Mantia T. The evolution in time of the concept of fast-growing tree species: is it possible to use a definition applicable to all environmental conditions?. *Annals of Silvicultural Research*. 2020;45(1):53-61.
 57. Hansen EA. Energy plantations in north central United States: Status of research and development plantations. *Energy Sources*. 1991;13(1):105-10. <https://doi.org/10.1080/00908319108908973>
 58. Joshi R, Sharma B, Singh H, Dhakal N, Ayer S, Maraseni T. Poplar Plantation as an Agroforestry Approach: Economic Benefits and Its Role in Carbon Sequestration in North India. *Journal of Resources and Ecology*. 2024;15(4):880-88. <https://doi.org/10.5814/j.issn.1674-764x.2024.04.009>
 59. Anbarashan M, Padmavathy A, Alexandar R, Dhatchanamoorhty N. Survival, growth, aboveground biomass and carbon sequestration of mono and mixed native tree species plantations on the Coromandel Coast of India. *Geology, Ecology and Landscapes*. 2020;4(2):111-20. <https://doi.org/10.1080/24749508.2019.1600910>
 60. Chaudhary K, Jat JR, Joshi T, Chauhan AH. Growth, biomass production and CO₂ sequestration of some important multipurpose trees under rainfed conditions. <https://doi.org/10.33545/26174693.2024.v8.i6Sd.1289>
 61. Kumar P, Mishra AK, Chaudhari SK, Sharma DK, Rai AK, Singh K, et al. Carbon sequestration and soil carbon build-up under Eucalyptus plantation in semi-arid regions of North-West India. *Journal of Sustainable Forestry*. 2021;40(4):319-31. <https://doi.org/10.1080/10549811.2020.1749856>
 62. Prasad JV, Srinivas K, Rao CS, Ramesh C, Venkatravamma K, Venkateswarlu B. Biomass productivity and carbon stocks of farm forestry and agroforestry systems of leucaena and eucalyptus in Andhra Pradesh, India. *Current Science*. 2012;536-40.
 63. Farrelly DJ, Everard CD, Fagan CC, McDonnell KP. Carbon sequestration and the role of biological carbon mitigation: A review. *Renewable and Sustainable Energy Reviews*. 2013;21:712-27. <https://doi.org/10.1016/j.rser.2012.12.038>
 64. Shimamoto CY, Botosso PC, Marques MC. How much carbon is sequestered during the restoration of tropical forests? Estimates from tree species in the Brazilian Atlantic forest. *Forest Ecology and Management*. 2014;329:1-9. <https://doi.org/10.1016/j.foreco.2014.06.002>
 65. Reddy MC, Priya RM, Madiwalar SL. Carbon sequestration potential of teak plantations of different agro-climatic zones and age-gradations of Southern India. *Current World Environment*. 2014;9(3):785-8.
 66. Dinesha S, Panda MR, Pradhan D, Rakesh S, Dey AN, Bhat JA, et al. Ecosystem carbon budgeting under Swietenia macrophylla King plantation in sub humid foothills of Eastern Himalayans of India. *Environment, Development and Sustainability*. 2024;26(2):4661-77. <https://doi.org/10.1007/s10668-022-02902-6>
 67. Chopra N, Tewari LM, Tewari A, Wani ZA, Asgher M, Pant S, et al. Estimation of biomass and carbon sequestration potential of *Dalbergia latifolia* Roxb. and *Melia composita* Willd. Plantations in the Tarai Region (India). *Forests*. 2023;14(3):646. <https://doi.org/10.3390/f14030646>
 68. Venkatesh L, Maheshnaik BL, Rathod R. Studies on Sandalwood Tree (L.) Based Santalum album Agroforestry System in Haveri District of Karnataka, India. *Indian Journal of Ecology*. 2023;50(5):1602-07. <https://doi.org/10.55362/IJE/2023/4102>
 69. Brien RJ, Caldwell L, Duchesne L, Voelker S, Barichivich J, Baliva M, et al. Forest carbon sink neutralized by pervasive growth-lifespan trade-offs. *Nature Communications*. 2020;11(1):4241. <https://doi.org/10.1038/s41467-020-17966-z>
 70. Kaul M, Mohren GM, Dadhwal VK. Carbon storage and sequestration potential of selected tree species in India. *Mitigation and Adaptation Strategies for Global Change*. 2010;15:489-510. <https://doi.org/10.1007/s11027-010-9230-5>
 71. Albrecht A, Kandji ST. Carbon sequestration in tropical agroforestry systems. *Agriculture, Ecosystems & Environment*. 2003;99(1-3):15-27. [https://doi.org/10.1016/S0167-8809\(03\)00138-5](https://doi.org/10.1016/S0167-8809(03)00138-5)
 72. Piponiot C, Cogné M, Freycon V, Thoumazeau A, Gusmão M, Peltier R. Traditional agroforestry systems in Timor-Leste can store large amounts of carbon in both soil and biomass. <http://dx.doi.org/10.21203/rs.3.rs-4460614/v1>
 73. Pugh TA, Jones CD, Huntingford C, Burton C, Arneth A, Brovkin V, Ciais P, Lomas M, Robertson E, Piao SL, Sitch Set al. A large committed long-term sink of carbon due to vegetation dynamics. *Earth's Future*. 2018;6(10):1413-32. <https://doi.org/10.1029/2018EF000935>
 74. Khan MN, Islam MR, Rahman A, Azad MS, Mollick AS, Kamruzzaman M, et al. Allometric relationships of stand level carbon stocks to basal area, tree height and wood density of nine tree species in Bangladesh. *Global Ecology and Conservation*. 2020;22:e01025. <https://doi.org/10.1016/j.gecco.2020.e01025>
 75. Muthmainnah M, Rusyidi S, Abdullah A, Ridha A. Estimation of biomass potential, carbon stocks and carbon sequestration of *Trigona* sp honey bees feed. *IOP Conference Series: Earth and Environmental Science*. 2021;886(1):012072. <https://doi.org/10.1088/1755-1315/886/1/012072>
 76. N  var J. Allometric equations for tree species and carbon stocks for forests of northwestern Mexico. *Forest Ecology and Management*. 2009;257(2):427-34. <https://doi.org/10.1016/j.foreco.2008.09.028>
 77. Shupe HA, Jensen K, Ludewig K. Adapting a *Quercus robur* allometric equation to quantify carbon sequestration rates on the Middle Elbe floodplain. *MethodsX*. 2022;9:101800. <https://doi.org/10.1016/j.mex.2022.101800>
 78. Paul KI, Roxburgh SH, England JR, Ritson P, Hobbs T, Brooksbank K, et al. Development and testing of allometric equations for estimating above-ground biomass of mixed-species environmental plantings. *Forest Ecology and Management*. 2013;310:483-94. <https://doi.org/10.1016/j.foreco.2013.08.054>
 79. Klein LJ, Zhou W, Albrecht CM. Quantification of carbon sequestration in urban forests. *arXiv preprint arXiv:2106.00182*.

2021. <https://doi.org/10.48550/arXiv.2106.00182>
80. Nowak DJ, Greenfield EJ, Hoehn RE, Lapoint E. Carbon storage and sequestration by trees in urban and community areas of the United States. *Environmental Pollution*. 2013;178:229-36. <https://doi.org/10.1016/j.envpol.2013.03.019>
 81. Russo A, Escobedo FJ, Timilsina N, Schmitt AO, Varela S, Zerbe S. Assessing urban tree carbon storage and sequestration in Bolzano, Italy. *International Journal of Biodiversity Science, Ecosystem Services & Management*. 2014;10(1):54-70. <https://doi.org/10.1080/21513732.2013.873822>
 82. Sandefur HN, Matlock MD, Johnston RZ. Comparison of methods for quantifying carbon sequestration in the urban forest of North little rock, Arkansas. *Biological Engineering Transactions*. 2014;6(4):221-31.
 83. Kuyah S, Mbow C, Sileshi GW, van Noordwijk M, Tully KL, Rosenstock TS. Quantifying tree biomass carbon stocks and fluxes in agricultural landscapes. *Methods for measuring greenhouse gas balances and evaluating mitigation options in smallholder agriculture*. 2016:119-34.
 84. Temesgen H, Affleck D, Poudel K, Gray A, Sessions J. A review of the challenges and opportunities in estimating above ground forest biomass using tree-level models. *Scandinavian Journal of Forest Research*. 2015;30(4):326-35.
 85. Wutzler T, Köstner B, Bernhofer C. Spatially explicit assessment of carbon stocks of a managed forest area in eastern Germany. *European Journal of Forest Research*. 2007;126:371-83. <https://doi.org/10.1007/s10342-006-0155-1>
 86. Somarriba E, Cerda R, Orozco L, Cifuentes M, Dávila H, Espin T, et al. Carbon stocks and cocoa yields in agroforestry systems of Central America. *Agriculture, Ecosystems & Environment*. 2013;173:46-57. <https://doi.org/10.1016/j.agee.2013.04.013>
 87. Holderieath J, Valdivia C, Godsey L, Barbieri C. The potential for carbon offset trading to provide added incentive to adopt silvopasture and alley cropping in Missouri. *Agroforestry Systems*. 2012;86:345-53.
 88. Atangana A, Khasa D, Chang S, Degrande A, Atangana A, Khasa D, et al. Carbon sequestration in agroforestry systems. *Tropical agroforestry*. 2014:217-25. https://doi.org/10.1007/978-94-007-7723-1_10
 89. Somasundaram D, Elango A, Karthikeyan S. Estimation of carbon credits of fishing boat diesel engine running on diesel-ethanol-bio-diesel blends with nano alumina doped ceria-zirconia. *Materials Today: Proceedings*. 2020;33:2923-28. <https://doi.org/10.1016/j.matpr.2020.02.887>
 90. Arya S, Garg AK, Mudgal RK. An empirical study of carbon credits trading practices among Indian organizations. *Abhigyan*. 2019;37(3):11-22. https://doi.org/10.56401/Abhigyan_37.3.2019.11-22
 91. Von Avenarius A, Devaraja TS, Kiesel R. An empirical comparison of carbon credit projects under the clean development mechanism and verified carbon standard. *Climate*. 2018;6(2):49.
 92. Kim C, Daniels T. California's success in the socio-ecological practice of a forest carbon offset credit option to mitigate greenhouse gas emissions. *Socio-Ecological Practice Research*. 2019;1:125-38. <https://link.springer.com/article/10.1007/s42532-019-00017-3>
 93. Protocol K. Kyoto protocol", UNFCCC Website [Internet]. 1997.
 94. Oluwadare AO. *Climate change policies and issues: A tool for increased wood production in developing countries*. Springer Berlin Heidelberg. 2011.
 95. Redondo-Brenes A. Growth, carbon sequestration and management of native tree plantations in humid regions of Costa Rica. *New Forests*. 2007;34:253-68.
 96. Obembe OS, Hendricks NP. Marginal cost of carbon sequestration through forest afforestation of agricultural land in the southeastern United States. *Agricultural Economics*. 2022;53(S1):59-73. <https://doi.org/10.1111/agec.12719>
 97. Li C, Fultz LM, Moore-Kucera J, Acosta-Martínez V, Horita J, Strauss R, et al. Soil carbon sequestration potential in semi-arid grasslands in the Conservation Reserve Program. *Geoderma*. 2017;294:80-90. <https://doi.org/10.1016/j.geoderma.2017.01.032>
 98. Dhyani SK, Ram A, Dev I. Potential of agroforestry systems in carbon sequestration in India. 2016:1103-12. <https://doi.org/10.56093/ijas.v86i9.61348>
 99. Schoeneberger MM. Agroforestry: working trees for sequestering carbon on agricultural lands. *Agroforestry Systems*. 2009;75:27-37. <https://doi.org/10.1007/s10457-008-9123-8>
 100. Rachana PM, Sekar I, Parthiban KT, Raj SV, Akshay FM, Singh D, et al. Eco-investments: Quantification of carbon sequestration potential and economic valuation of multifunctional agroforestry system. *Horizon*. 2025;12(1):1-0. <https://doi.org/10.14719/pst.4820>