



RESEARCH ARTICLE

Influence of tillage and herbicide strategies on weed dynamics and wheat (*Triticum aestivum* L.) performance in saline soils of Northern India

Mohammad Vaheed^{1*}, Chandra Bhushan¹, Sunil Kumar Verma¹, SB Singh², Sudhir Kumar Rajpoot¹, Sidra Qidwai² & Sameer Shrivastava¹

¹Department of Agronomy, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi 221 005, Uttar Pradesh, India

²Department of Agronomy, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya 224 229, Uttar Pradesh, India

*Correspondence email - vaheedias@bhu.ac.in

Received: 29 April 2025; Accepted: 30 June 2025; Available online: Version 1.0: 14 August 2025

Cite this article: Vaheed M, Bhushan C, Verma SK, Singh SB, Rajpoot SK, Qidwai S, Shrivastava S. Influence of tillage and herbicide strategies on weed dynamics and wheat (*Triticum aestivum* L.) performance in saline soils of Northern India. Plant Science Today (Early Access).

<https://doi.org/10.14719/pst.9195>

Abstract

A field experiment was conducted during the *rabi* season of 2019-20 at the Agronomy Research Farm, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya, to assess the effects of tillage and herbicide combinations on weed dynamics, crop growth and wheat yield under saline soil conditions. The study involved 15 treatment combinations arranged in a split-plot design (SPD) with three replications. The main plots included three tillage operations: T₀-zero tillage (ZT), T₁-minimum tillage (MT) and T₂-conventional tillage (CT). The sub-plots consisted of five weed management treatments: pendimethalin 1.5 kg active ingredient/ha (pre-emergence), pendimethalin 1.0 kg active ingredient/ha (pre-emergence) followed by sulfosulfuron (SFS) at 0.025 kg active ingredient/ha (post-emergence), SFS + metsulfuron-methyl (MSM) at 0.032 kg active ingredient/ha (post-emergence), a weedy check and a weed-free control. The results revealed that ZT significantly reduced weed density and dry weight while enhancing weed control efficiency (WCE) compared to MT and CT. Among the herbicide treatments, the sequential application of pendimethalin (1.0 kg active ingredient/ha) followed by SFS (0.025 kg active ingredient/ha) was the most effective in suppressing weed growth and improving WCE and enhancing crop growth and yield. This treatment was statistically on par with the SFS + MSM application. The combination of ZT with pendimethalin followed by SFS (T₀W₂) recorded the lowest weed infestation and the highest values for plant height, dry matter accumulation, tillers per m row length and grain yield. This was closely followed by the ZT combined with SFS + MSM treatment (T₀W₃), highlighting the synergistic benefits of conservation tillage and sequential herbicide application.

Keywords: herbicide; saline soil; tillage; weed control efficiency; wheat; yield

Introduction

After rice, wheat (*Triticum aestivum* L.) is India's second-largest cereal crop and a major contributor to the nation's food security. It constitutes 33.53 % of India's total food grain production. Wheat production in India during 2022-23 was 110.55 MT with an area of 31.40 Mha with a productivity of 6.40 t/ha. Over the decades, India's wheat production has risen substantially, rising from 6.40 MT in 1949-50 to 110.55 MT in 2022-23. Madhya Pradesh, Punjab and Uttar Pradesh are the three main wheat-producing states; Uttar Pradesh alone accounts for 31.77 % of the nation's wheat production (1).

Despite this impressive progress, wheat production in India faces several challenges, particularly traditional tillage practices and heavy weed infestation (2, 3). These issues are exacerbated by the delayed harvesting of preceding rice crops, which often delays wheat sowing, especially under conventional tillage systems that require intensive land ploughing (4). Late sowing has been shown to significantly reduce yield, by

approximately 38 to 78 kg/ha/day when sown after 14th November (5).

Weeds also play a critical role in limiting wheat productivity. Globally, weed competition causes more crop losses than the combined effects of insect pests and disease. In India, severe competition from diverse weed flora can reduce wheat yields by 30-80 % (6). Traditional weed control methods, such as manual weeding, are labour-intensive, time-consuming and expensive. Consequently, herbicide-based weed management has gained popularity because of its lower labour costs and effective weed suppression (7).

CT, particularly ZT and reduced tillage (RT), has emerged as a sustainable solution to overcome these challenges. ZT allows timely sowing of wheat by reducing the need for extensive land preparation. Additionally, it affects weed dynamics and improves yields by conserving water; energy and labour (8, 9), reduces soil degradation, improve nutrient cycle and promote environmental sustainability (10).

However, despite these advantages, the adoption of CT has been hindered by high weed infestation, particularly during the initial years. Small-seeded weed species tend to thrive under reduced tillage systems, necessitating the use of diverse herbicidal strategies (11). The effective use of herbicides during the initial stages of conservation tillage is therefore crucial to minimize the weed-seed bank and ensure long-term sustainability. Therefore, an effort was made to assess the effect of herbicides and tillage on dynamics of weeds and wheat yield in saline soil.

Materials and Methods

A field study was conducted during the *rabi* season of 2019-20 at the Agronomy Research Farm of Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya, Uttar Pradesh, India. The geographical coordinates were located at 26°32'55"N latitude, 81°50'14"E longitude and at an altitude of 113 m above sea level (Fig. 1). During the experimental period, the maximum recorded temperatures ranged between 14.0 °C and 36.6 °C, while the minimum temperatures ranged from 5.3 °C to 18.7 °C. The total rainfall during the study period was 184.6 mm. The surface-soil layer was silty-loam, with a pH of 8.08 and low organic carbon content (0.40 %).




Fig. 1. Location of the experimental site. The figure illustrates the geographical context of the study area, beginning with a map of India, followed by zoomed-in views of Uttar Pradesh and Ayodhya district and concluding with a photograph of the experimental field where the research was conducted.

The study comprised fifteen treatment combinations arranged in an SPD with three replications. The main plots included three tillage treatments: T₀: ZT-no ploughing, T₁: MT-two ploughing and T₂: CT-four ploughing with broadcast seeding and in sub-plots were herbicides i.e. W₁: pendimethalin at 1.5 kg active ingredient/ha as a pre-emergence (PE), W₂: pendimethalin at 1.0 kg active ingredient/ha (PE) followed by SFS 0.025 kg active ingredient/ha as post-emergence (PoE), W₃: SFS + MSM at 0.032 kg active ingredient/ha (PoE), W₄: weedy-check and W₅: weed-free (up to 60 days).

In ZT (T₀) plots, glyphosate was sprayed at 1.0 kg active ingredient/ha 10 days prior to sowing to control the persistent weeds. In MT and CT plots, seedbeds were prepared through two and four ploughings respectively. Utilizing a ZT-seed drill to open slits the wheat variety NW-5054 was seeded at the seed rate of 100 kg/ha utilizing various crop establishment techniques with a 20 cm row spacing.

To meet the crops' nutritional needs, the prescribed fertilizer dosage of 120, 60, 40 kg/ha of nitrogen (N), phosphorous (P₂O₅) and potash (K₂O) was applied. Half the dose of N and the full dose of P₂O₅ and K₂O were applied at sowing, while the remaining N was applied in two equal top dressings; once after first irrigation and second at the tillering stage respectively. All other agronomic practices were carried out as per crop requirements to ensure successful crop establishment and growth.

Observations

The relative composition of each group of weeds was determined from weedy plots (weedy checks) by calculating the proportion of the single group of weeds in total number of all groups of weeds multiplied by 100. Weed density (WD) was determined at 30, 60, 90 days after sowing (DAS) and at harvest, using a quadrant of 50 × 50 cm size. Four random samples were collected from each treatment plot. All weeds within the sampled area were collected individually and categorized into 3 major groups: grasses, broad-leaf weeds (BLWs) and sedges. The total density of weeds was then determined.

After collection, individual weed species were washed twice, first with clean water and then with distilled water and air dried for 2-3 hr. These air-dried samples were kept in large brown paper bags and further dried in a hot-air oven at 70 °C till the constant weight was obtained. An electronic balance was used to record the dry weight of individual groups of weeds. The dry weight of the individual group of weeds was summed up to get total weed dry weight (WDW).

The mature crop was harvested from the net-plot area (4 × 3.2 m), excluding a 0.5 m border on all side, in the second week of April 2020. Harvested crops were sun-dried and weighed to get biological yield. After threshing, all grains were cleaned and weighed to determine grain yield and reported at 14 % moisture content. The grain yield was subtracted from biological yield to get straw yield.

The relative composition of weed was further computed based on the proportion of each major weed group to the total weed population. Weed control efficiency (%) at 30, 60 and 90 DAS of crops was determined by using the expression (12).

$$WCE (\%) = \frac{DMC - DMT}{DMC} \times 100$$

Where, WCE is weed control efficiency; DMC is dry weight of weeds (gm⁻²) for control plot (weedy check) and DMT is dry weight of weeds (gm⁻²) for the treated plot.

Statistical analysis

All recorded data were average replication-wise and subjected to statistical analysis using analysis of variance (ANOVA) (13) and mean comparisons were based on the least significant difference (LSD) at a 5 % probability level ($p < 0.05$). Data pertaining to WD and WDW were first transformed using the square-root-transformation $\sqrt{(x + 0.5)}$ and then analyzed. 'F-test' was used to establish the significance of treatment effects.

Results and Discussion

Relative composition of weeds

The wheat field was affected by *Phalaris minor*, *Avena ludoviciana* *Chenopodium album*, *Anagallis arvensis* and *Cyperus rotundus*. Among the grassy weeds, *P. minor* was the dominant weed followed by *A. ludoviciana*. Of the two dominant broad-leaf weeds, *C. album* showed 21.78 % highest density over *A. arvensis* 16.22 % (Fig. 2). In total, narrow-leaf weeds comprised the largest proportion (39 %) of the overall weeds followed by broad-leaf weeds (38 %) and sedges (23 %) (Fig. 3).

Weed density

Data were analyzed using analysis of variance (ANOVA) and treatment means were compared using the least significant difference (LSD) test at a 5 % significance level. Error bars indicate standard errors based on the error degrees of freedom from ANOVA. Treatment differences were considered significant at $p \leq 0.05$. The density of all weed groups was lowest at 30 DAS, but as the crop progressed towards maturity, the weed density gradually increased, reaching its peak at 60 DAS. All the tillage operations significantly affected WD with ZT resulting in the lowest WD across all groups, followed by MT, while CT exhibited the highest density. The superior performance of ZT in reducing weed infestation and enhancing wheat yield aligns with previous studies (2, 14). Reduced soil disturbance under ZT minimises the upward movement of weed seeds from deeper soil layers, thereby lowering the chances of germination. Conversely, intensive ploughing in CT facilitates weed emergence by exposing buried seeds to favourable environmental conditions (15-17). This mechanistic explanation underscores the sustainability of conservation tillage in saline soil conditions.

Among herbicidal treatment, W₁-W₃ and W₅ had noticeable negative effects on weeds and were shown to have reduced the densities of all weed groups compared to the untreated weedy-check (Fig. 4-6). Application of pendimethalin at 1.5 kg active ingredient/ha followed by SFS at 0.025 kg active ingredient/ha (W₂) significantly reduced WD and it was statistically at par with SFS + MSM at 0.032 kg active ingredient/ha (W₃). The density of grasses, BLWs and sedges was significantly impacted by the interaction between tillage and herbicidal treatment. The lowest density of all weed groups was recorded under ZT along with the application of




Fig. 2. Relative composition of weed species at 60 days after sowing.




Fig. 3. Relative composition of weed groups at 60 days after sowing.




Fig. 4. Interaction effect of tillage and herbicide treatments on weed density (No./m²) at 30 days after sowing. The figure represents five dominant weed species: *Phalaris minor*, *Avena ludoviciana*, *Chenopodium album*, *Anagallis arvensis* and *Cyperus rotundus*.




Fig. 5. Interaction effect of tillage and herbicides on density of weeds (No./m²) at 60 days after sowing. The figure represents five dominant weed species: *Phalaris minor*, *Avena ludoviciana*, *Chenopodium album*, *Anagallis arvensis* and *Cyperus rotundus*.




Fig. 6. Interaction effect of tillage and herbicides on density (No./m²) of weeds at 90 days after sowing.

The figure represents five dominant weed species: *Phalaris minor*, *Avena ludoviciana*, *Chenopodium album*, *Anagallis arvensis* and *Cyperus rotundus*.

pendimethalin at 1.0 kg active ingredient/ha followed by SFS at 0.025 kg active ingredient/ha (T_0W_2), outperforming all other tillage and herbicide treatment combinations (6, 18, 19).

Total weed density, weed dry weight and weed control efficiency

There is a strong correlation observed between WD and WDW, with both being highest in the weedy-check plots. Among the tillage treatment, ZT had the lowest overall WD, WDW and maximum WCE, followed by MT and CT respectively. The enhanced soil structure (i.e. tilling) and the upward movement of weed seeds to the surface layer in conventional tillage likely resulted in a higher weed population, leading to increased weed biomass and reduced WCE in the CT system (2, 8, 20).

Among herbicidal treatments, W_2 was the most effective in reducing both total weed density and dry weight and recorded the highest WCE across treatments (Fig. 7-9). It was due to the lowest weed population and dry weight observed under this treatment. These outcomes are consistent with the other findings (17).

Tillage and herbicides had a significant interaction effect in the T_0W_2 followed by T_0W_3 treatment combinations. However, total WD at 60 DAS was significantly lower (38.10 to 72.16 %); in T_0 (ZT) plots treated with (W_2) pendimethalin at 1.0 kg active ingredient/ha followed by SFS at 0.025 kg active ingredient/ha compared to the T_2 plots $\times$ (W_1 , W_2 , W_3) combinations and being at par with T_1 plots (MT). When compared with the weedy check (W_4), the reduction in weedy density under the T_0W_2 combination ranged from 73.71 to 90.39 %.

Furthermore, under W_1 and W_2 treatments, WDW in T_0 plots was 16.87 % and 37.80 % lower than T_1 plots respectively (21). Even under weedy-check condition, T_0 plots recorded 10.13 % lower WDW than T_2 plots. WDW from T_2 plots treated with W_1 was significantly greater than most other tillage operations (T) $\times$ herbicides combinations except for T_0W_3 . Apart from weedy-check and weed-free plots; the increase in WDW $T_2 \times W_1$ combinations ranged from 7.60 % to 58.46 % (22, 23).




Fig. 7. Interaction of tillage and herbicides on total weed density (No./m²).




Fig. 8. Interaction of tillage and herbicides on total weed dry weight (g/m²).




Fig. 9. Interaction effect of tillage and herbicides on WCE (%).

Crop growth

Tillage operations and herbicides treatments had a significant effect on the growth characteristics of wheat. Among tillage operation, treatment ZT showed the highest leaf area index (LAI), as shown in Fig. 10, along with superior plant height, number of tillers and dry matter accumulation (DMA), as summarized in Table 1. This was followed by MT and CT respectively. The enhanced crop growth and yield under ZT can be attributed to improved soil structure, better moisture conservation and more efficient nutrient cycling (10, 24). Moreover, reduced weed interference in ZT systems allows wheat plants to allocate greater resources to growth and grain formation, corroborating findings reported from similar agro-ecological conditions (25, 26).

Among the herbicidal treatment, pendimethalin at 1.0 kg active ingredient/ha followed by SFS at 0.025 kg active ingredient/ha recorded the highest values for LAI, plant height, tillers and DMA and was at par with SFS + MSM at 0.032 kg active ingredient/ha. These improvements in crop growth can

be attributed to reduced crop-weed competition for light, water and nutrients, thereby enhancing physiological and morphological growth parameters (27, 28).

Yield

On average, ZT increased grain yield by 13.59 % over MT and 20.32 % over CT, respectively (2, 4, 14). Herbicidal treatments also had a significant impact on grain yield. The sequential application of pendimethalin at 1.0 kg active ingredient/ha followed by SFS at 0.025 kg active ingredient/ha resulted in a substantial increase in grain yield compared to both the weedy-check and the sole application of pendimethalin at 1.5 kg active ingredient/ha and was at par with SFS + MSM at 0.032 kg active ingredient/ha. Evidently, the highest grain yield was produced in weed-free plots, which were noticeable higher than yield under any herbicidal treatment (Table 2). Nevertheless, the initial phase of ZT adoption is often challenged by persistent weed pressure, necessitating the use of effective herbicide regimes during the early stages (11, 25).




Fig. 10. Effect of tillage and herbicides on leaf area index.

Table 1. Effects of tillage and herbicides on plant height, tillers, dry matter accumulation at harvest and grain yield

Treatments	Plant height (cm)	Tillers (No./m ²)	DMA (g/m ²)	Grain yield (kg/ha)
Tillage operations (T)				
T ₀	81.83	295.46	862.37	3834
T ₁	78.82	280.28	803.58	3376
T ₂	67.73	274.28	738.03	3187
SEm±	1.04	1.80	8.03	44
CD (p=0.05)	4.08	7.05	31.51	172
Herbicides (Ws)				
W ₁	72.66	271.83	752.54	3162
W ₂	79.51	292.64	859.59	3687
W ₃	78.37	286.19	821.18	3488
W ₄	65.11	254.14	700.76	2988
W ₅	84.98	311.89	872.55	4003
SEm±	0.99	2.63	9.28	38
CD (p=0.05)	2.88	7.69	27.09	110
Interaction (T×Ws)				
SEm±	1.71	4.56	16.07	65
CD (p=0.05)	4.99	13.32	46.92	190

T₀: Zero-tillage (No tillage), T₁: Minimum tillage (2 ploughing), T₂: Conventional tillage (4 ploughing) with broadcast seeding; W₁: Pendimethalin at 1.5 kg active ingredient/ha (PE), W₂: Pendimethalin at 1.0 kg active ingredient/ha (PE) followed by sulfosulfuron at 0.025 kg active ingredient/ha (PoE), W₃: Sulfosulfuron + metsulfuron at 0.032 kg active ingredient/ha (PoE), W₄: Weedy-check, W₅: Weed-free.

Table 2. Interaction tillage operations and herbicides on total weed density and dry weight at 60 DAS and grain yield of wheat

T/Ws	Total weed density (No./m ²)			Total weed dry weight (g/m ²)			Grain yield (kg/ha)		
	T ₀	T ₁	T ₂	T ₀	T ₁	T ₂	T ₀	T ₁	T ₂
W ₁	6.48 (41.50)*	6.87 (46.80)	7.20 (51.36)	5.09 (25.49)	5.60 (30.97)	5.83 (33.51)	3617	3045	2824
W ₂	3.84 (14.30)	4.19 (17.10)	4.86 (23.10)	3.80 (13.92)	4.46 (19.43)	4.90 (23.51)	3921	3700	3440
W ₃	4.25 (17.60)	4.71 (21.70)	5.45 (29.40)	4.03 (15.77)	4.63 (20.97)	4.96 (24.10)	3807	3358	3300
W ₄	12.51 (156.00)	13.35 (178.00)	13.54 (183.00)	7.07 (49.58)	7.22 (51.69)	7.46 (55.17)	3378	2820	2765
W ₅	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	0.71 (0.00)	4449	3954	3605
CD (p=0.05)	0.50 (10.30)			0.27 (3.60)			190		

*Data under parenthesis are the original $\sqrt{(x+0.5)}$ value which were transformed by for analysis.

Tillage and herbicides have the significant interaction effect on wheat grain yield. T₀W₁ produced the highest wheat grain yield, which was 12.27 % more than all MT and herbicides combinations, including the weed-free plots. Additionally, the yield under this combination of treatments was the same as the yield from T₁W₅. Furthermore, as per the findings of Table 2, the yields under both the weed-free and weedy-check plots were higher in MT by 8.83 % and 1.95 % respectively than under CT (29, 30).

Conclusion

The findings of this study demonstrate that tillage practices and herbicide applications significantly influence weed dynamics and wheat growth. ZT increased grain yield by 12.29 % and 18.63 % compared to MT and CT respectively. Among the herbicide strategies tested, the sequential application of pendimethalin at 1.0 kg active ingredient/ha followed by sulfosulfuron at 0.025 kg active ingredient/ha (W₂) was the most effective, leading in the lowest weed infestation, highest weed control efficiency (WCE) and maximum grain yield.

Acknowledgements

Authors wish to thank Acharya Narendra Deva University of Agriculture and Technology, College of Agriculture for providing the necessary infrastructure and resources.

Authors' contributions

MV carried out the research work in the field and laboratory and drafted the manuscript. MV, SKV, CB and SKR developed the methodology and contributed to visualization. MV, SKV and SKR performed software analysis. MV, SKV, CB, SKR and SBS provided validation. MV, SKV, CB and SS conducted formal analysis. MV, SKV, CB, SKR and SQ carried out the investigation. MV, SKV, CB and SQ provided resources. SKV, MV and SBS curated the data. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None

References

- Directorate of Economics and Statistics. Ministry of Agriculture and Farmers Welfare, Government of India; 2024.
- Singh RP, Verma SK, Kumar S. Weed management for enhancing yield and economics of wheat in the Eastern India. *Indian J Agric Sci.* 2020;84(6):780–83.
- Minhas WA, Mumtaz N, Ur-Rehman H, Farooq S, Farooq M, Ali HM, et al. Weed infestation and productivity of wheat crop sown in various cropping systems under conventional and conservation tillage. *Front Plant Sci.* 2023;14:1176738. <https://doi.org/10.3389/fpls.2023.1176738>
- Thind HS, Sharma S, Sidhu HS, Singh V. Tillage, crop establishment and residue retention methods for optimising productivity and profitability under rice-wheat system. *Crop Pasture Sci.* 2023;74(9):828–42. <https://doi.org/10.1071/CP21595>
- Tomar SPS, Tomar SS, Srivastava SC. Yield and yield component response of wheat genotypes to different sowing dates in Gird region of Madhya Pradesh. *Int J Farm Sci.* 2014;4(2):1–6.
- Lakra K, Husain K, Pyare R, Verma SK, Meena RS, Singh PK, et al. Productivity and profitability of irrigated bread wheat (*Triticum aestivum* L.) are influenced by Irrigation scheduling and weed management approaches. *Gesunde Pflanzen.* 2022;75(4):801–14. <https://doi.org/10.1007/s10343-022-00748-6>
- Lakra K, Pyare R, Singh PK, Verma SK, Singh RK, Upadhyay PK, et al. Effect of irrigation schedule and herbicides application on growth and productivity of wheat in semi-arid environment. *Indian J Agron.* 2020;67(2):129–36. <https://doi.org/10.59797/ija.v67i2.102>
- Yadav DK, Verma SK, Singh RK, Bhushan C, Rajpoot SK, Upadhyay A, et al. Effect of crop establishment and weed management practices on soil properties in wheat. *Pharma Innov J.* 2023;12(6):4607–609.
- Alhammad BA, Roy DK, Ranjan S, Padhan SR, Sow S, Nath D, et al. Conservation tillage and weed management influencing weed dynamics, crop performance, soil properties and profitability in a rice-wheat-green gram system in the eastern Indo-Gangetic plain. *Agronomy.* 2023;13(7):1953. <https://doi.org/10.3390/agronomy13071953>
- Yadav DK, Verma SK, Pratap V, Yadav SP, Jaysawal PK. Available nutrients in soil are influenced by planting techniques and weed management options in wheat. *Int J Chem Stud.* 2020;8(4):2718–21. <https://doi.org/10.22271/chemi.2020.v8.i4af.10057>
- Yang M, Zhao Y, Yang H, Shen Y, Zhang X. Suppression of weeds and weed seeds in the soil by stubbles and no-tillage in an arid maize-winter wheat-common vetch rotation on the Loess Plateau of China. *J Arid Land.* 2018;10:809–20. <https://doi.org/10.1007/s40333-018-0063-5>
- Hasanuzzaman M, Islam MO, Bapari MS. Efficacy of different herbicides over manual weeding in controlling weeds in transplanted rice. *Aust J Crop Sci.* 2008;2(1):18–24.
- Gomez KA, Gomez AA. Statistical procedures for agricultural research. New York: John Wiley & Sons; 1984.
- Yadav DK, Verma SK, Bhushan C, Mahajan NC, Kumar P. Influence of crop establishment methods and weed management practices on growth and yield attributes of wheat. *Int J Plant Soil Sci.* 2023;35(18):1452–58. <https://doi.org/10.9734/ijpss/2023/v35i183412>
- Shyam R, Rohitashav S, Singh VK, Guru SK. Effect of wheat establishment methods and weed management practices on weed dynamics and productivity of wheat grown in succession to rice. *Indian J Weed Sci.* 2009;41(1):70–72.
- Khalique A, Shakeel M, Matloob A, Hussain S, Tanveer A, Murtaza G. Influence of tillage and weed control practices on growth and yield of wheat. *Philipp J Crop Sci.* 2013;38(3):54–62.
- Singh RP, Verma SK, Prasad SK, Singh H, Singh SB. Effect of tillage and weed management practices on grassy weeds in wheat. *Int J Sci Environ Technol.* 2017;6(1):404–12.
- Ghazikhanlou Sani Y, Yousefi AR, Jamshidi K, Shekari F, Gonzalez-Andujar JL, Korres NE. Weed response to ALS-inhibitor herbicide (Sulfosulfuron + metsulfuron methyl) under increased temperature and carbon dioxide. *Agronomy.* 2023;13(8):2084. <https://doi.org/10.3390/agronomy13082084>
- Verma SK, Bhushan C, Rajpoot SK, Chaudhary R, Meena R, Kashyap SK, et al. Herbicide and irrigation management options in conventionally-tilled wheat: Deciphering water and energy budgeting and grain and monetary output in north-Indian plains. *Sci Rep.* 2025;15(1):1612. <https://doi.org/10.1038/s41598-024-81578-6>
- Ahmad N, Virk AL, Hafeez MB, Ercisli S, Golokhvast KS, Qi Y, et al. Effects of different tillage and residue management systems on soil organic carbon stock and grain yield of rice-wheat double cropping system. *Ecol Indic.* 2024;158:111452.
- Sumitra G, Rajkumara S, Lamani KD. Sequential herbicidal application on weed community and yield of wheat. *Indian J Weed Sci.* 2024;56(3):243–50.
- Brar AS, Walia US. Weed dynamics and wheat (*Triticum aestivum* L.) productivity as influenced by planting techniques and weed control practices. *Indian J Weed Sci.* 2009;41(3-4):161–66.
- Hussain Z, Hashim S, Jabran K. Integrated effect of tillage and herbicides on wheat crop. *Gesunde Pflanzen.* 2021;73(2):239–47. <https://doi.org/10.1007/s10343-021-00549-3>
- Wang Y, Song J, Li W, Yan T, Wang D, Xue J, et al. Wheat yield, biomass and radiation interception and utilization under conservation tillage: Greater response to drip fertigation compared to intensive tillage. *Agronomy.* 2024;14(12):2849. <https://doi.org/10.3390/agronomy14122849>
- Verma SK, Singh RP, Kumar S. Effects of irrigation and herbicides on the growth, yield and yield attributes of wheat. *Bangladesh J Bot.* 2017;46(3):839–45.
- Choudhary PC, Sharma AR. Herbicides in zero-till wheat grown after greengram. *Indian J Weed Sci.* 2023;55(1):95–98. <https://doi.org/10.5958/0974-8164.2023.00016.3>
- Kushwaha AS, Gupta D, Singh L. Chemical weed management in wheat (*Triticum aestivum*) using ready-mix and tank-mix herbicides. *Indian J Agron.* 2018;63(4):462–66.

28. Rana SS, Badiyala D, Brari P. Evaluation of herbicide combinations for controlling complex weed flora in wheat. *Indian J Weed Sci.* 2017;49(4):335–40. <https://doi.org/10.5958/0974-8164.2017.00088.0>
29. Deshmukh JP, Kakade SU, Goud VV. Tillage and weed management effect on wheat in inceptisols grown under soybean-wheat cropping sequence. *Indian J Weed Sci.* 2023;55(4):448–52.
30. Buczek J. The effect of tillage systems and weed control methods on grain yield and gluten protein compositional and content-related changes in hybrid bread wheat. *Agriculture.* 2024;14(9):1558. <https://doi.org/10.3390/agriculture14091558>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.