



RESEARCH ARTICLE

Comparative study of proximate composition, antioxidant and FT-IR profiles of *Flourensia cernua* DC., *Jatropha dioica* Sessé and *Lippia graveolens* Kunth from Mexican semi-arid regions

Cecilia Castro-López¹, Polet Sifuentes-Gómez^{1,2}, Guillermo C G Martínez-Ávila³, Romeo Rojas^{3*} & Julio C Tafolla-Arellano^{1*}

¹Laboratorio de Biotecnología y Biología Molecular, Departamento de Ciencias Básicas. Universidad Autónoma Agraria Antonio Narro, Saltillo 25315, Coahuila, México

²Coordinación de Alimentos de Origen Vegetal. Centro de Investigación en Alimentación y Desarrollo A.C., Hermosillo 83304, Sonora, México

³Laboratorio de Química y Bioquímica, Facultad de Agronomía. Universidad Autónoma de Nuevo León, General Escobedo 66050, Nuevo León, México

*Correspondence email - romeo.rojasmln@uanl.edu.mx; jtafare@uaaan.edu.mx

Received: 24 May 2025; Accepted: 25 July 2025; Available online: Version 1.0: 26 November 2025; Version 2.0: 11 December 2025

Cite this article: Cecilia CL, Polet SG, Guillermo CGMA, Romeo R, Julio CTA. Comparative study of proximate composition, antioxidant and FT-IR profiles of *Flourensia cernua* DC., *Jatropha dioica* Sessé and *Lippia graveolens* Kunth from Mexican semi-arid regions. Plant Science Today. 2025; 12(4): 1-9. <https://doi.org/10.14719/pst.9665>

Abstract

Flourensia cernua DC ('Hojasén'), *Jatropha dioica* Sessé ('Sangre de Drago') and *Lippia graveolens* Kunth ('Orégano') are three of the main medicinal plants native to the semi-arid regions of northern Mexico. Adaptation to these environments influences their compositional profile and biological activities; however such aspects remain underexplored. Hence this study aimed to compare the proximate composition, antioxidant activity and FT-IR fingerprints of these species collected from different semi-arid regions. The results revealed significant differences ($p < 0.05$) among collection sites. Overall, samples from San Jerónimo (SJ) and Estanque de León (EL) locations showed higher dry matter, crude protein and nitrogen-free extract contents. Conversely, samples from La Tortuga (T) location exhibited higher moisture, ash, crude fat and crude fibre contents. Regarding antioxidant activity, samples from SJ and EL recorded lower IC₅₀ values in both DPPH[•] and ABTS^{•+} assays along with higher FRAP values. Finally, all locations exhibited comparable FT-IR spectra, confirming the presence of several functional groups such as C=O, C-H, C-O, C-N and N-H. Although variability was site-dependent, we highlight the nutritional and functional potential of *F. cernua*, *J. dioica* and *L. graveolens* for future applications.

Keywords: native plants; composition; phenolics; bioactivity; FT-IR

Introduction

Arid and semi-arid regions worldwide comprise unique ecosystems characterized by adverse environmental conditions such as drought, salinity, high temperature and intense solar radiation which often hinder plant growth (1). However certain plant species have successfully adapted to these environments. Evidence has shown that plants evolved intricate genetic, biochemical and physiological mechanisms to cope with stress. These include the synthesis of specialized metabolites, enhanced water retention, improved photosynthetic efficiency and altered leaf structure (2). Notably, such adaptations not only allow plants to survive but also contribute to their nutritional quality and functional properties (1).

In this context, the semi-arid regions of northern Mexico harbor several native plants that have traditionally been used for both food and medicinal purposes (3). Among the most representative species, *Flourensia cernua* DC, *Jatropha dioica* Sessé, *Lippia graveolens* Kunth are widely distributed in the area, easily accessible and collected by rural communities as an economic activity. Indeed,

owing to their wide ethnobotanical use, these species have attracted research interest for their complex chemical composition, promising biological activities and high added value (4). For instance, *F. cernua* (Asteraceae; "Hojasén") is a shrub used as an infusion to alleviate various respiratory and gastrointestinal disorders (e.g., asthma, bronchitis, diarrhea and indigestion) (5). Previous studies on crude extracts and fractions revealed the presence of phenolic acids, flavonoids, sesquiterpenes, fatty acids and benzofurans which are associated with antioxidant, anti-inflammatory, anti-obesity and antidiabetic properties (6, 7).

On the other hand, *J. dioica* (Euphorbiaceae; "Sangre de Drago") is a shrub widely recognized for its use as a macerate for the treatment of gastrointestinal and dental disorders, anemia, diabetes, rheumatism, ophthalmia, alopecia and wounds among others (8). Research into the composition of crude extracts has led to the identification of lignans, coumarins, alkaloids, flavonoids, terpenes, cyclic peptides, phytosterols, steroids, phenolic acids and esters. These compounds have been correlated with antioxidant,

antimicrobial, antifungal, antiviral, antitumor, anti-inflammatory, immunomodulatory, anti-obesity and nephroprotective effects (8-10).

Finally, *L. graveolens* (Verbenaceae; “Oregano”) is not only a valued food condiment but also a shrub utilized in the form of decoctions or infusions to treat respiratory, gastrointestinal and urinary disorders as well as menstrual and musculoskeletal conditions, allergies, bacterial infections, migraine and diabetes (11). Recent studies have demonstrated that essential oils, crude extracts and fractions contain a range of compounds including flavonoids, terpenes, quinones, steroids, saponins, tannins, coumarins, alkaloids and lactones. The antimicrobial, antifungal, antiviral, antimutagenic, antioxidant, anticancer, anti-inflammatory and photochemopreventive activities of these compounds have also been reported (12, 13).

Although current knowledge supports the use of these species, studies on their compositional profiles and biological activities under different environmental contexts remain limited. Indeed, increasing research has focused on exploring how geographic and environmental variations influence the nutritional and functional properties of plants from arid and semi-arid regions (14-16). Therefore, understanding this variability is essential not only for conservation efforts but also for the exploitation of *F. cernua*, *J. dioica* and *L. graveolens* within the food, nutraceutical and pharmaceutical fields. On this basis, the study aimed to compare the proximate composition, antioxidant activity and FT-IR fingerprints of *F. cernua*, *J. dioica* and *L. graveolens* collected from different semi-arid regions of Mexico.

Materials and Methods

Plant materials

Fresh leaves of *F. cernua*, *J. dioica* and *L. graveolens* were collected from three different locations in two semi-arid regions of Mexico (Fig. 1): San Jerónimo (SJ) in Zacatecas (latitude: 22°39'11.95", longitude:

102°29'24.32", elevation: 2275 m a.s.l.), Estanque de León (EL) in Coahuila (latitude: 25°42'35.55", longitude: 103°18'02.27", elevation: 1115 m a.s.l.) and La Tortuga (T) in Coahuila (latitude: 25°51'39.77", longitude: 101°16'30.53", elevation: 984 m a.s.l.). Plant species were authenticated by the herbarium of the Universidad Autónoma Agraria Antonio Narro (<https://plants.jstor.org/partner/ANSM>). All samples were washed with distilled water, dried at 50 °C for 24 h (Oven HS35-AIA; Novatech, Jalisco, Mexico), ground into powder (Electric grinder HC-500; Homend, Istanbul, Turkey) and sieved (0.6 mm particle size). The resulting powders were collected and stored in air-tight bags at room temperature until use.

Proximate composition analysis

Proximate composition of leaf samples was determined according to Association of Official Analytical Chemists (AOAC) standard methods (17). Briefly, dry matter and ash contents were assessed using the drying (AOAC 930.04) and incineration (AOAC 930.05) methods, respectively. Crude protein content was determined by the Kjeldahl method ($N \times 6.25$) (AOAC 984.13) while crude fat content was measured using the Soxhlet extraction method (AOAC 920.39). Crude fiber was analyzed by the acid-alkali digestion method (AOAC 962.09). Nitrogen-free extract was calculated by difference.

Phenolic extraction

Phenolic extraction was performed as reported by Ascacio-Valdés *et al.* (18) with slight modifications. Firstly, 20 g of leaf sample was mixed with 100 mL of distilled water (solid: liquid ratio of 1:5 w/v) and incubated at 60 °C for 30 min. The crude extract was then filtered through filter paper (Whatman N° 41) to remove solids. The resulting filtrate was partially purified by column chromatography using Amberlite™ XAD-16N as the stationary phase. Distilled water was used to remove water-soluble compounds followed by anhydrous ethanol to recover phenolic compounds. The solvent in the phenolic fraction was evaporated at 50 °C for 48 h. Finally, the dried extract was collected and stored in the dark at room temperature until use.




Fig. 1. Collected plant material: a) *F. cernua*, b) *J. dioica* and c) *L. graveolens*.

Antioxidant activity analysis

DPPH[•] radical scavenging

DPPH[•] (2,2-diphenyl-1-picrylhydrazyl) radical scavenging activity was assessed according to Brand-Williams *et al.* (19) with slight modifications. Briefly, 50 µL of phenolic extract (at concentrations of 250, 500, 750 and 1000 µg mL⁻¹) was mixed with 2950 µL of a DPPH[•] methanolic solution (60 µM) and incubated at room temperature for 30 min. The absorbance of the reaction mixture was measured at 517 nm (Spectrophotometer 6320D; Jenway, Essex, UK). The radical scavenging capacity was calculated using Eqn. 1 and results were expressed as IC₅₀ values (µg mL⁻¹), indicating the concentration of phenolic extract required to inhibit 50 % of DPPH[•] radical.

$$\text{Inhibition (\%)} = (A_0 - A_1) / A_0 \times 100 \quad (\text{Eqn. 1})$$

where A₀ and A₁ are the absorbances of the control and sample, respectively.

ABTS^{•+} radical scavenging

ABTS^{•+} (2,2'-Azino-bis [3-ethylbenzothiazoline-6-sulfonic acid]) radical scavenging activity was evaluated as suggested by van den Berg *et al.* (20) with slight modifications. Firstly, an ABTS aqueous solution (7 mM) was mixed with potassium persulfate (2.45 mM) (liquid: liquid ratio of 2:1 v/v) and incubated at room temperature for 12 h to generate the ABTS^{•+} radical. The resulting solution was then diluted with anhydrous ethanol until an absorbance of 0.700 ± 0.002 at 734 nm was obtained. Subsequently, 50 µL of phenolic extract (at concentrations of 250, 500, 750 and 1000 µg mL⁻¹) was mixed with 950 µL of the ABTS^{•+} solution and after 1 min of reaction, absorbance was measured at 734 nm. The radical scavenging capacity was calculated using Eqn. 2 and results were expressed as IC₅₀ values (µg mL⁻¹) indicating the concentration of phenolic extract required to inhibit 50 % of ABTS^{•+} radical.

$$\text{Inhibition (\%)} = (A_0 - A_1) / A_0 \times 100 \quad (\text{Eqn. 2})$$

where A₀ and A₁ are the absorbances of the control and sample, respectively.

Ferric reducing antioxidant power

Ferric reducing antioxidant power (FRAP) was determined according to Çelik *et al.* (21) with slight modifications. Briefly, 50 µL of phenolic extract (at concentrations of 250 and 500 µg mL⁻¹) was mixed with 120 µL of phosphate-buffered saline solution (1M, pH 7). Subsequently, 220 µL of potassium ferrocyanide (1 % w/v) was added and the mixture was homogenized and incubated at 50 °C for 20 min. After incubation, 120 µL of trichloroacetic acid (10 % w/v), 450 µL of distilled water and 100 µL of ferric chloride (0.1 % w/v) were added sequentially. The absorbance of the reaction mixture was measured at 734 nm. A standard curve was prepared using gallic

acid at concentrations ranging from 0 to 150 µg mL⁻¹. Results were expressed as µg of gallic acid equivalents per milliliter (µg GAE mL⁻¹) of phenolic extract.

Fourier transform infrared (FT-IR) spectroscopy analysis

Fourier transform infrared (FT-IR) spectroscopy was conducted to identify the functional groups present in the phenolic extract. Briefly, the sample was placed on the plate reader (Spectrometer Cary 630 FTIR; Agilent Technologies, Santa Clara, CA, USA) and the spectrum was recorded over the range of 4000 to 600 cm⁻¹ by co-adding 32 scans at a resolution of 2 cm⁻¹. Data analysis was performed using MicroLab Expert software (version 1.1.0.1).

Statistical analysis

Experimental data are presented as the mean ± standard deviation (SD) of three independent replicates. One-way analysis of variance (ANOVA) followed by Tukey-Kramer's post hoc test was used to determine statistically significant differences among mean values (*p* < 0.05). Statistical analysis was performed using InfoStat software (version 2020e; freely available at <http://www.infostat.com.ar>).

Results and Discussion

Proximate composition

Variations in the proximate composition of *F. cernua*, *J. dioica* and *L. graveolens* leaves are presented in Table 1. As noticed, dry matter and moisture contents ranged from 38.81 to 98.50 % and 1.50 to 61.19 %, respectively. Except for *J. dioica*, significant differences (*p* < 0.05) were found in *F. cernua* and *L. graveolens* collected from different locations. The highest dry matter content was recorded in *F. cernua* from SJ (98.50 %) followed by *L. graveolens* from EL (88.39 %). Inversely proportional, these samples also exhibited the lowest moisture content with values of 1.50 and 11.61 %, respectively. According to Ntshambiwa *et al.* (22), variations in dry matter may be attributed to precipitation and soil moisture conditions at the growing site. Plants exposed to water deficit have been shown to adapt by modifying root water absorption and leaf transpiration which leads to increased dry matter accumulation (23).

Regarding ash content, values ranged from 7.35 to 14.53 %. Significant differences (*p* < 0.05) were detected among samples from different locations. Notably, *F. cernua*, *L. graveolens* and *J. dioica* collected from T exhibited the highest and lowest ash content with values of 14.53, 10.16 and 7.35 %, respectively. Yang *et al.* (24) reported that ash content may vary according to climatic conditions (e.g., radiation, temperature and precipitation) as plants modulate their metabolic activity in response to climate

Table 1. Proximate composition parameters of *F. cernua*, *J. dioica* and *L. graveolens* leaves collected from different locations.

Plant species	Location	Dry matter (%)	Moisture (%)	Ash (%)	Crude protein (%)	Crude fat (%)	Crude fiber (%)	Nitrogen-free extract (%)
<i>F. cernua</i>	SJ	98.50 ± 0.20 ^a	1.50 ± 0.20 ^b	13.00 ± 0.17 ^b	1.98 ± 0.04 ^b	6.40 ± 2.61 ^a	11.23 ± 0.62 ^b	67.38 ± 2.88 ^a
	EL	97.10 ± 0.36 ^b	2.90 ± 0.36 ^a	14.13 ± 0.11 ^a	2.11 ± 0.04 ^{ab}	8.60 ± 0.47 ^a	11.00 ± 0.35 ^b	64.08 ± 0.55 ^b
	T	96.23 ± 0.30 ^b	3.77 ± 0.30 ^a	14.53 ± 0.15 ^a	2.12 ± 0.02 ^a	8.66 ± 0.72 ^a	13.33 ± 0.66 ^a	51.41 ± 16.78 ^b
<i>J. dioica</i>	SJ	38.81 ± 2.04 ^a	61.19 ± 2.04 ^a	9.26 ± 0.07 ^a	12.70 ± 0.22 ^a	1.99 ± 0.17 ^b	19.20 ± 2.27 ^b	56.85 ± 1.92 ^a
	EL	40.12 ± 1.68 ^a	59.88 ± 1.68 ^a	8.87 ± 0.30 ^a	8.04 ± 0.08 ^b	2.04 ± 0.16 ^b	28.25 ± 0.30 ^a	52.90 ± 0.56 ^b
	T	39.38 ± 1.38 ^a	60.62 ± 1.38 ^a	7.35 ± 0.04 ^b	9.81 ± 0.65 ^b	3.64 ± 0.28 ^a	26.08 ± 0.22 ^a	53.10 ± 0.90 ^b
<i>L. graveolens</i>	SJ	74.81 ± 0.63 ^c	25.19 ± 0.63 ^a	9.64 ± 0.12 ^b	9.26 ± 0.29 ^c	3.83 ± 0.42 ^a	14.37 ± 0.67 ^a	62.91 ± 0.34 ^a
	EL	88.39 ± 0.63 ^a	11.61 ± 0.63 ^c	8.90 ± 0.26 ^c	13.12 ± 0.28 ^a	3.60 ± 0.31 ^a	11.99 ± 1.82 ^a	62.47 ± 1.53 ^a
	T	85.41 ± 0.80 ^b	14.59 ± 0.80 ^b	10.16 ± 0.03 ^a	10.27 ± 0.22 ^b	3.49 ± 0.38 ^a	11.64 ± 0.86 ^a	64.44 ± 0.92 ^a

Different superscript letters within the same column indicate significant differences (*p* < 0.05) among locations of plant species. Parameters are expressed as dry basis. SJ, San Jerónimo; EL, Estanque de León; T, La Tortuga.

induced stress. Similarly, soil properties such as texture, pH and organic matter content play a crucial role in regulating the adsorption, solubility and translocation of mineral elements within the plant system (24).

Crude protein content ranged from 1.98 to 13.12 %. Significant differences ($p < 0.05$) were observed among samples from different locations. The highest crude protein content was recorded in *L. graveolens* from EL (13.12 %) followed by *J. dioica* from SJ (12.70 %); while the lowest content was determined in *F. cernua* from SJ (1.98 %). Plant density and nitrogen availability have been shown to influence crude protein as suggested by Leghari *et al.* (25). These factors can affect the metabolic pathways involved in protein biosynthesis and accumulation ultimately shaping the quality and properties of plant-derived protein (26).

About crude fat content, values ranged from 1.99 to 8.66 %. Except for *J. dioica*, no significant differences ($p > 0.05$) were found in *F. cernua* and *L. graveolens* collected from different locations. Despite the non-significance, *F. cernua* from T exhibited the highest crude fat content (8.66 %). *L. graveolens* from SJ and *J. dioica* from T showed comparable crude fat values (3.83 and 3.64 %, respectively) while *J. dioica* from SJ recorded the lowest (1.99 %). It is well-known that the maturity stage determines the accumulation and redistribution of plant macronutrients especially crude fat. As plants mature, they tend to store more lipids which are essential for membrane biogenesis and cell signaling and serve as energy and carbon sources (27).

Crude fiber content ranged from 11.00 to 28.25%. Except for *L. graveolens*, significant differences ($p < 0.05$) were observed in *F. cernua* and *J. dioica* collected from different locations. The highest crude fiber content was recorded in *J. dioica* from EL (28.25%) followed by *L. graveolens* from SJ (14.37 %) and *F. cernua* from T (13.33%). The lowest value was found in *F. cernua* from EL (11.00%). According to Busuttill-Griffin *et al.* (28), high temperatures and water deficit at the growing site may lead to variations in crude fiber. These conditions accelerate plant maturation during which physiological and structural changes cause the leaves to become woody and fibrous. This response is associated with the regulation of metabolic pathways involved in cellulose and hemicellulose biosynthesis (29).

Finally, nitrogen-free extract (NFE), also referred to as soluble carbohydrate content, ranged from 51.41 to 67.38 %, exhibiting the same significant trend as for crude fibre. *F. cernua* from SJ presented the highest NFE content (67.38 %) while *L. graveolens* from all locations showed comparable values (ca. 62-64 %). The lowest NFE was recorded in *F. cernua* from T (51.41 %).

Plants are known to sense solar radiation and adjust their carbohydrate content accordingly. Changes in radiation intensity and quality may influence the regulation of photosynthetic pigments, stomatal development and carbohydrate metabolism pathways, ultimately affecting the accumulation of soluble sugars in plant tissues (30). Overall, the proximate composition observed in this study is comparable to or even higher than that reported by Estell *et al.* (31), Afolabi *et al.* (32) and Djengue *et al.* (33) for other *Flourensia*, *Jatropha* and *Lippia* species, respectively.

Antioxidant activity

Table 2 summarizes the variations in antioxidant activity of *F. cernua*, *J. dioica* and *L. graveolens* phenolic extracts. As noticed, for DPPH[•] radical scavenging activity, IC₅₀ values ranged from 37.20 to 668.25 µg mL⁻¹. A lower IC₅₀ value indicates higher antioxidant activity of the extract and vice versa; thus significant differences ($p < 0.05$) were found among samples from different locations. The lowest IC₅₀ value was recorded in *L. graveolens* from SJ (37.20 µg mL⁻¹) followed by *J. dioica* from EL (155.02 µg mL⁻¹) and *F. cernua* from T (288.81 µg mL⁻¹). The highest value was determined in *J. dioica* from T (668.25 µg mL⁻¹).

Regarding ABTS^{•+} radical scavenging activity, IC₅₀ values ranged from 14.59 to 139.50 µg mL⁻¹ exhibiting the same significant trend as for DPPH[•] activity. Hence, the lowest IC₅₀ values were in the ranking order of *L. graveolens* from SJ (14.59 µg mL⁻¹) < *J. dioica* from EL (24.57 µg mL⁻¹) < *F. cernua* from T (112.48 µg mL⁻¹). The highest value was found in *F. cernua* from SJ (139.50 µg mL⁻¹). According to Itam *et al.* (34), the antioxidant activity of plant extracts can be classified based on IC₅₀ values as follows: very strong (IC₅₀ < 50 µg mL⁻¹), strong (IC₅₀ = 50-100 µg mL⁻¹), moderate (IC₅₀ = 100-250 µg mL⁻¹), weak (IC₅₀ = 250-500 µg mL⁻¹) and inactive (IC₅₀ > 500 µg mL⁻¹). Under this classification and regardless of the collection site, *L. graveolens* exerted very strong antioxidant activity while *F. cernua* and *J. dioica* displayed moderate to inactive activity.

Finally, FRAP values ranged from 18.07 to 44.69 µg GAE mL⁻¹. A higher FRAP value indicates higher antioxidant activity. Although all extracts showed a concentration-dependent increase, significant differences ($p < 0.05$) were observed among samples from different locations. *L. graveolens* from SJ exhibited the highest FRAP, with values of 31.36 and 43.76 µg GAE mL⁻¹ per 250 and 500 µg mL⁻¹ extract, respectively. Meanwhile, *F. cernua* and *J. dioica* showed similar FRAP values at both extract concentrations (ca. 18-20 and 20-24 µg GAE mL⁻¹, respectively) with only slight variations between species and within locations (Table 2). These results were consistent with those of DPPH[•] and ABTS^{•+} assays.

Table 2. Antioxidant activity of *F. cernua*, *J. dioica* and *L. graveolens* phenolic extracts from different locations.

Plant species	Location	DPPH [•] (IC ₅₀ , µg mL ⁻¹)	ABTS ^{•+} (IC ₅₀ , µg mL ⁻¹)	FRAP	
				(µg GAE mL ⁻¹ , per 250 µg mL ⁻¹ extract)	(µg GAE mL ⁻¹ , per 500 µg mL ⁻¹ extract)
<i>F. cernua</i>	SJ	325.76 ± 1.04 ^b	139.50 ± 0.45 ^a	20.52 ± 0.38 ^a	24.64 ± 0.25 ^a
	EL	349.16 ± 0.66 ^a	137.69 ± 0.56 ^b	19.54 ± 0.08 ^a	20.97 ± 0.51 ^a
	T	288.81 ± 0.85 ^c	112.48 ± 0.42 ^b	18.07 ± 0.69 ^b	20.48 ± 0.25 ^b
<i>J. dioica</i>	SJ	244.07 ± 1.98 ^b	37.87 ± 1.27 ^b	19.99 ± 0.34 ^a	21.12 ± 0.39 ^b
	EL	155.02 ± 2.26 ^c	24.57 ± 1.01 ^c	20.52 ± 1.26 ^a	24.55 ± 0.22 ^a
	T	668.25 ± 1.67 ^a	108.78 ± 0.63 ^a	19.50 ± 0.25 ^a	21.27 ± 0.29 ^b
<i>L. graveolens</i>	SJ	37.20 ± 0.85 ^b	14.59 ± 1.42 ^c	31.36 ± 0.61 ^a	43.76 ± 1.06 ^a
	EL	44.51 ± 0.58 ^b	22.19 ± 0.52 ^a	25.82 ± 0.25 ^b	33.56 ± 0.22 ^b
	T	51.93 ± 0.42 ^a	21.13 ± 0.46 ^b	30.08 ± 2.19 ^a	44.69 ± 1.18 ^a

Different superscript letters within the same column indicate significant differences ($p < 0.05$) among locations of plant species. SJ, San Jerónimo; EL, Estanque de León; T, La Tortuga.

Although a plant's phenolic profile is genotype- and environment-influenced, the antioxidant activity of its constituents depends on their specific reaction mechanisms. In the DPPH[•] and ABTS^{•+} assays, phenolics can stabilize free radicals through combined hydrogen atom (H[•]) donation (HAT) and single electron (e⁻) transfer (SET) mechanisms. Meanwhile, in the FRAP assay which measures the reduction of ferric ions (Fe³⁺) to ferrous ions (Fe²⁺), they rely exclusively on the SET mechanism (35). However, their reaction efficiency is determined by the chemical structure. On this basis, the superior activity of *L. graveolens* compared to that of *F. cernua* and *J. dioica* may be attributed to unique structural features.

For instance, Vuolo *et al.* (36) reported that the number and position of hydroxyl (O-H) groups (e.g., presence of two O-H groups in the aromatic ring) correlate with an improved effect. Similarly, the presence of a C2=C3 double bond, coupled with a 4-carbonyl group, improves molecular planarity and electron delocalization, facilitating electron transfer reactions. Even though glycosides have reduced activity compared to free aglycones, glycosylation, particularly in the C-glycoside form may retain greater activity than O-glycoside (36, 37). Conversely, aromatic rings susceptible to O-methyl substitution show decreased effect since methylation leads to steric hindrance that interferes with radical interaction sites (36). Altogether, these features may support the variations observed in this study. Interestingly, previous studies have observed comparable antioxidant activity in other *Flourensia* (7), *Jatropha* (38) and *Lippia* (39) species growing under adverse environmental conditions.

FT-IR fingerprinting

Fig. 2 to 4 depicts the variations in functional groups present in *F. cernua*, *J. dioica* and *L. graveolens* phenolic extracts, respectively. Overall, all locations exhibited comparable FT-IR spectral patterns with only slight differences in peak intensities. The typical absorption peaks for *F. cernua*, *J. dioica* and *L. graveolens* considered as their molecular fingerprint, were clearly in the range of 650-1700 cm⁻¹, 670-1580 cm⁻¹ and 680-1600 cm⁻¹, respectively. As observed, there were between eight and eleven major peaks, depending on the sample.

Notably, several functional groups corresponding to qualitatively different compounds were identified. Regardless of the sample, peak wavenumbers (Fig. 2 to 4) in the range of 600-950 cm⁻¹ were assigned to out-of-plane C-H bending vibration of aromatic compounds, alkenes and alkynes; 1020-1275 cm⁻¹ were attributed to C-O symmetric/asymmetric stretching vibration (alcohols, esters, ethers and carboxylic acids) and C-N stretching vibration (aliphatic amines); 1300-1450 cm⁻¹ were assigned to C-H symmetric/asymmetric bending vibration (alkanes), C-N stretching vibration (aromatic amines), C-O stretching vibration (amide) and N-O symmetric stretching vibration (nitro compounds); 1500-1650 cm⁻¹ were attributed to C=C stretching vibration (aromatic rings in phenolic compounds) and N-H bending vibration (amines); 1700-1760 cm⁻¹ were assigned to N-H bending vibration (amino acids) and C=O stretching vibration (aldehydes, ketones, esters and fatty acids); 2000-2150 cm⁻¹ were attributed to C≡C stretching vibration of alkynes; 2800-2930 cm⁻¹ were assigned to C-H symmetric/asymmetric stretching vibration of lipids (CH₂ and CH₃), methoxy derivatives, aldehydes and alkanes and 3000-3350 cm⁻¹ were attributed to O-H stretching vibration (water, hydrogen-bonded alcohols, phenols, carbohydrates and peroxide) and N-H stretching vibration (amines) (40).

It is noteworthy that the FT-IR fingerprints obtained in this study agree with those previously informed for other *Flourensia*, *Jatropha* and *Lippia* species. In the case of the *F. cernua* extracts, peak wavenumbers at ~3338 (O-H and N-H stretch), ~2929 (C-H stretch), ~1702 (C=O stretch), ~1652 (C=C stretch) and ~1017 (C-N stretch) cm⁻¹ matched with those reported by Aranda-Ledesma *et al.* (6) and are characteristic of compounds commonly found in *Flourensia* crude extracts. On the other hand, the *J. dioica* extracts exhibited peak wave numbers at ~3250 (O-H stretch), ~2924 (C-H stretch), ~1355 (C-H bend), ~1260 (C-O stretch) and ~1040 (C-N stretch) cm⁻¹, aligning with those described by Umar *et al.* (41). These peaks are used as authentication markers of *Jatropha* spp. Finally in the *L. graveolens* extracts, peak wavenumbers at ~3269 (O-H stretch), ~2925 (C-H stretch), ~1596 (C=O stretch), ~1025 (C-O stretch) and ~813 (C-H bend) cm⁻¹ were consistent with those observed by de Sá-Filho *et al.* (42), indicating the presence of major constituents found in *Lippia* essential oils and crude extracts.




Fig. 2. FT-IR fingerprints of *F. cernua* phenolic extracts from (SJ) San Jerónimo, (EL) Estanque de León and (T) La Tortuga locations.




Fig. 3. FT-IR fingerprints of *J. dioica* phenolic extracts from (SJ) San Jerónimo, (EL) Estanque de León and (T) La Tortuga locations.




Fig. 4. FT-IR fingerprints of *L. graveolens* phenolic extracts from (SJ) San Jerónimo, (EL) Estanque de León and (T) La Tortuga locations.

As mentioned above, plant chemical composition is genetically determined; however, it may be modulated in response to biotic and abiotic stressors at the growing site (43). To counteract stress, plants differentially activate or repress genes that can redirect the metabolic pathways involved in secondary metabolite biosynthesis and accumulation, thereby affecting the type, content and function of compounds produced (44). For instance, solar radiation and temperature stimulate the shikimic acid pathway, leading to the accumulation of different aromatic compounds which play key roles in cell signalling, hormonal regulation and oxidative stress alleviation (44).

According to the FT-IR analysis of phenolic extracts, Sukmawaty *et al.* (43) stated that the peak wavenumber indicates qualitatively the type of compound while the absorption intensity reflects its concentration. Therefore, the FT-IR spectral patterns observed in Fig. 2 to 4 suggest that the types of metabolites present in *F. cernua*, *J. dioica* and *L. graveolens* do not vary among locations, except at the concentration level. However, this statement should be further investigated. Consistent with our findings, studies have reported that *Moringa oleifera* (43), *Laurus nobilis* (45) and *Isotoma longiflora* (46) also maintained similar metabolite profiles despite differing growing sites.

Conclusion

Based on the findings, *F. cernua*, *J. dioica* and *L. graveolens* could be considered promising sources of macronutrients and bioactive compounds. However significant variations in their compositional profiles and antioxidant activity were demonstrated among collection sites. The variability is mainly attributed to plant physiology, as well as to the soil and environmental conditions prevailing in each location. Overall, this study could help identify suitable cultivation sites for species with enhanced nutritional and functional properties offering prospects for natural product development in the food, pharmaceutical and cosmetic industries (e.g., as additives, preservatives, new drugs and supplements). Despite this, further research should focus on determining the vitamin, mineral and fatty acid composition of *F. cernua*, *J. dioica* and *L. graveolens* leaves. Furthermore, additional studies are needed to evaluate other potential *in vitro* biological activities of phenolic extracts beyond that reported here. Finally, future *in vivo* and clinical studies are required to validate the safety, efficacy and bioavailability of extracts for human use.

Acknowledgements

Cecilia Castro-López would like to thank the Secretariat of Science, Humanities, Technology and Innovation (SECIHTI) of Mexico. The authors sincerely thank Mariana S. Lavreda-Lozano and Ever T. Astorga-Arzola (Laboratorio de Biotecnología y Biología Molecular- Universidad Autónoma Agraria Antonio Narro) for their valuable technical support in performing the experimental procedures. This study was supported by the Universidad Autónoma de Nuevo León through the Science, Technology and Innovation Support Program-ProACTI 2025, research project 129-CAAFE-2025. The authors thank the Directorate of Research of UAAAN for the facilities.

Authors' contributions

CCL, data collection and analysis, figure preparation and drafting the original manuscript. PSG, sample collection and experimental analysis. GCGMA and RR, methodology supervision and manuscript review. JCTA, conceptualization, methodology supervision, resources and manuscript review. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the author(s) used Grammarly to check the English grammar. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

References

- Joshi T, Deepa PR, Joshi M, Sharma PK. Matters of the desert: A perspective on achieving food and nutrition security through plants of the (semi) arid regions. *J Agric Food Res.* 2023;14:100725. <https://doi.org/10.1016/j.jafr.2023.100725>
- Mohanta TK, Mohanta YK, Kaushik P, Kumar J. Physiology, genomics and evolutionary aspects of desert plants. *J Adv Res.* 2024;58:63-78. <https://doi.org/10.1016/j.jare.2023.04.019>
- Márquez-Rangel I, Cruz M, Neira-Vielma AA, Ramírez-Barrón SN, Aguilar-Zarate P, Belmares R. Plants from arid zones of Mexico: Bioactive compounds and potential use for food production. *Resources.* 2025;14(1):13. <https://doi.org/10.3390/resources14010013>
- Dávila-Rangel IE, Charles-Rodríguez AV, López-Romero JC, Flores-López ML. Plants from arid and semi-arid zones of Mexico used to treat respiratory diseases: A review. *Plants.* 2024;13(6):792. <https://doi.org/10.3390/plants13060792>
- Linares-Braham A, Palomo-Ligas L, Nery-Flores SD. Bioactive compounds and pharmacological potential of hojasen (*Flourensia cernua*): A mini review. *Plant Sci Today.* 2023;10(sp2):304-12. <https://doi.org/10.14719/pst.2546>
- Aranda-Ledesma NE, González-Hernández MD, Rojas R, Paz-González AD, Rivera G, Luna-Sosa B, et al. Essential oil and polyphenolic compounds of *Flourensia cernua* leaves: Chemical profiling and functional properties. *Agronomy.* 2022;12(10):2274. <https://doi.org/10.3390/agronomy12102274>
- Jasso-de Rodríguez D, Torres-Moreno H, López-Romero JC, Vidal-Gutiérrez M, Villarreal-Quintanilla JA, Carrillo-Lomeli DA, et al. Antioxidant, anti-inflammatory and antiproliferative activities of *Flourensia* spp. *Biocatal Agric Biotechnol.* 2023;47:102552. <https://doi.org/10.1016/j.bcab.2022.102552>
- Majeed I, Rizwan K, Saber FR, Munir S, Soria-Lopez A, Otero P. Ethnotraditional uses and potential industrial and nutritional applications of secondary metabolites of genus *Jatropha* L. (Euphorbiaceae): A review. *J Agric Food Res.* 2025;21:101861. <https://doi.org/10.1016/j.jafr.2025.101861>
- Bastos-Cavalcante N, da Conceição-Santos AD, da Silva-Almeida JRG. The genus *Jatropha* (Euphorbiaceae): A review on secondary chemical metabolites and biological aspects. *Chem Biol Interact.* 2020;318:108976. <https://doi.org/10.1016/j.cbi.2020.108976>
- Rodríguez-Rodríguez DR, Mendoza-Hernández OH, Cordero-Pérez P, Rivas-Galindo VM, Moreno-Peña DP, Tijerina-Márquez R, et al. Nephroprotective and antioxidant effects of *Jatropha dioica* extract against ischemia-reperfusion injury in wistar rats. *Int J Mol Sci.* 2025;26(5):1838. <https://doi.org/10.3390/ijms26051838>
- Bautista-Hernández I, Aguilar CN, Martínez-Ávila GCG, Torres-León C, Iliina A, Flores-Gallegos AC, et al. Mexican oregano (*Lippia graveolens* Kunth) as source of bioactive compounds: A review. *Molecules.* 2021;26(17):5156. <https://doi.org/10.3390/molecules26175156>
- García-Bores AM, Espinosa-González AM, Reyna-Campos A, Cruz-Toscano S, Benítez-Flores JC, Hernández-Delgado CT, et al. *Lippia graveolens* photochemopreventive effect against UVB radiation-induced skin carcinogenesis. *J Photochem Photobiol B.* 2017;167:72-81. <https://doi.org/10.1016/j.jphotobiol.2016.12.014>
- Amador S, Nieto-Camacho A, Ramírez-Apan MT, Martínez M, Maldonado E. Cytotoxic, anti-inflammatory and α-glucosidase inhibitory effects of flavonoids from *Lippia graveolens* (Mexican oregano). *Med Chem Res.* 2020;29:1497-506. <https://doi.org/10.1007/s00044-020-02569-6>
- Yahia Y, Bagues M, Zaghdoud C, Al-Amri SM, Nagaz K, Guerfel M. Phenolic profile, antioxidant capacity and antimicrobial activity of *Calligonum arich* L., desert endemic plant in Tunisia. *S Afr J Bot.* 2019;124:414-19. <https://doi.org/10.1016/j.sajb.2019.06.005>
- Mohamed-Ouali D, Gaceb-Terrak R. Impact of environmental aridity on the phytochemical composition of phenolic extracts and

- essential oil from *Vitex agnus-castus* L. leaves acclimated in the Algerian Sahara. *Analele Univ din Oradea Fasc Biol.* 2023;2:105-16.
16. Jaradat N, Adwan L, K'aibni S, Zaid AN, Shtaya MJY, Shraim N, et al. Variability of chemical compositions and antimicrobial and antioxidant activities of *Ruta chalepensis* leaf essential oils from three Palestinian regions. *Biomed Res Int.* 2017;2017:2672689. <https://doi.org/10.1155/2017/2672689>
 17. Association of Official Analytical Collaboration International-AOAC International. *Official Methods of Analysis*. 17th ed. Washington (DC): AOAC International; 2000. Available from: <https://www.aoac.org>
 18. Ascacio-Valdés J, Aguilera-Carbó A, Martínez-Hernández J, Rodríguez-Herrera R, Aguilar C. *Euphorbia antisiphilitica* residues as a new source of ellagic acid. *Chem Pap.* 2010;64(4):528-32. <https://doi.org/10.2478/s11696-010-0034-6>
 19. Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. *LWT Food Sci Technol.* 1995;28(1):25-30. [https://doi.org/10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5)
 20. van den Berg R, Haenen GRMM, van den Berg H, Bast A. Applicability of an improved Trolox equivalent antioxidant capacity (TEAC) assay for evaluation of antioxidant capacity measurements of mixtures. *Food Chem.* 1999;66:511-7. [https://doi.org/10.1016/S0308-8146\(99\)00089-8](https://doi.org/10.1016/S0308-8146(99)00089-8)
 21. Çelik SE, Özyürek M, Güçlü K, Apak R. Solvent effects on the antioxidant capacity of lipophilic and hydrophilic antioxidants measured by CUPRAC, ABTS/persulphate and FRAP methods. *Talanta.* 2010;81(4-5):1300-9. <https://doi.org/10.1016/j.talanta.2010.02.025>
 22. Ntshambiwa KT, Seifu E, Mokhawa G. Nutritional composition, bioactive components and antioxidant activity of *Moringa stenopetala* and *Moringa oleifera* leaves grown in Gaborone, Botswana. *Food Prod Process Nutr.* 2023;5:7. <https://doi.org/10.1186/s43014-022-00124-x>
 23. Bhattacharya A. Dry matter production, partitioning and seed yield under soil water deficit: A review. In: Bhattacharya A, editor. *Soil water deficit and physiological issues in plants*. Singapore: Springer; 2021. p. 585-702.
 24. Yang Q, Yue K, Wu F, Hedenec P, Ni X, Wang D, et al. Global patterns and drivers of initial plant litter ash concentration. *Sci Total Environ.* 2022;830:154702. <https://doi.org/10.1016/j.scitotenv.2022.154702>
 25. Leghari SJ, Wahocho NA, Laghari GM, Hafeez-Laghari A, Mustafa-Bhabhan G, Hussain-Talpur K, et al. Role of nitrogen for plant growth and development: A review. *Adv Environ Biol.* 2016;10(9):209-19.
 26. Wan C, Gao L, Wang J, Lei X, Tao J, Feng B, et al. Effects of nitrogen fertilizer on protein synthesis, accumulation and physicochemical properties in common buckwheat. *Crop J.* 2023;11(3):941-50. <https://doi.org/10.1016/j.cj.2023.01.002>
 27. Murphy DJ. Plant storage lipids. *eLS.* 2016;1:1-7. <https://doi.org/10.1002/9780470015902.a0001918.pub3>
 28. Busuttil-Griffin F, Shoemaker C, Attard E, Azzopardi LM. Crude fibre determination of *Malva sylvestris* L. and evaluation of its faecal bulking and laxative properties in rats. *Int J Biol.* 2015;7(4):1-8. <https://doi.org/10.5539/ijb.v7n4p1>
 29. Le Gall H, Philippe F, Domon JM, Gillet F, Pelloux J, Rayon C. Cell wall metabolism in response to abiotic stress. *Plants.* 2015;4(1):112-66. <https://doi.org/10.3390/plants4010112>
 30. Wu W, Chen L, Liang R, Huang S, Li X, Huang B, et al. The role of light in regulating plant growth, development and sugar metabolism: A review. *Front Plant Sci.* 2025;15:1507628. <https://doi.org/10.3389/fpls.2024.1507628>
 31. Estell RE, Fredrickson EL, Havstad KM. Chemical composition of *Flourensia cernua* at four growth stages. *Grass Forage Sci.* 1996;51(4):434-41. <https://doi.org/10.1111/j.1365-2494.1996.tb02078.x>
 32. Afolabi QO, Shorinmade AY, Obero OJ. Proximate composition of *Jatropha curcas* leaves, phytochemical and antibacterial analysis of its ethyl acetate fraction. *Asian J Phys Chem Sci.* 2017;4(1):1-8. <https://doi.org/10.9734/AJOPACS/2017/38037>
 33. Djengue HW, Adjatin A, Djehoue R, Bonou-gbo Z, Odjo CT, Koukpo ZC, et al. Assessment of proximal, mineral composition and content of vitamin A and C of leaves and flowers from *Lippia multiflora* vegetable in Benin. *Afr J Biotechnol.* 2022;21(3):95-105. <https://doi.org/10.5897/AJB2021.17414>
 34. Itam A, Wati MS, Agustin V, Sabri N, Jumanah RA, Efdi M. Comparative study of phytochemical, antioxidant and cytotoxic activities and phenolic content of *Syzygium aqueum* (Burm. f. Alston f.) extracts growing in west Sumatera Indonesia. *Sci World J.* 2021;2021:5537597. <https://doi.org/10.1155/2021/5537597>
 35. Munteanu IG, Apetrei C. Analytical methods used in determining antioxidant activity: A review. *Int J Mol Sci.* 2021;22(7):3380. <https://doi.org/10.3390/ijms22073380>
 36. Vuolo MM, Lima VS, Maróstica-Junior MR. Phenolic compounds: Structure, classification and antioxidant power. In: Segura-Campos MR, editor. *Bioactive compounds: Health benefits and potential applications*. Netherlands: Elsevier; 2019. p. 33-50.
 37. Lang Y, Gao N, Zang Z, Meng X, Lin Y, Yang S, et al. Classification and antioxidant assays of polyphenols: A review. *J Future Foods.* 2024;4(3):193-204. <https://doi.org/10.1016/j.jfutfo.2023.07.002>
 38. Papalia T, Barreca D, Panuccio MR. Assessment of antioxidant and cytoprotective potential of *Jatropha* (*Jatropha curcas*) grown in southern Italy. *Int J Mol Sci.* 2017;18(3):660. <https://doi.org/10.3390/ijms18030660>
 39. Chies CE, Branco CS, Scola G, Agostini F, Gower AE, Salvador M. Antioxidant effect of *Lippia alba* (Miller) N. E. Brown. *Antioxidants.* 2013;2(4):194-205. <https://doi.org/10.3390/antiox2040194>
 40. Kumar S, Yadav M, Yadav A, Yadav JP. Impact of spatial and climatic conditions on phytochemical diversity and *in vitro* antioxidant activity of Indian *Aloe vera* (L.) Burm.f. *S Afr J Bot.* 2017;111:50-9. <https://doi.org/10.1016/j.sajb.2017.03.012>
 41. Umar AH, Syahrini R, Ranteta'dung I, Rafi M. FTIR-based fingerprinting combined with chemometrics method for rapid discrimination of *Jatropha* spp. (Euphorbiaceae) from different regions in South Sulawesi. *J Appl Pharm Sci.* 2023;13(1):139-49. <https://doi.org/10.7324/JAPS.2023.130113>
 42. de Sá-Filho JCF, de Castro-Nizio DA, Souza-de Oliveira AM, Freitas-Alves M, Camargos-de Oliveira R, Queiroz-Luz JM, et al. Geographic location and seasonality affect the chemical composition of essential oils of *Lippia alba* accessions. *Ind Crops Prod.* 2022;188(Part A):115602. <https://doi.org/10.1016/j.indcrop.2022.115602>
 43. Sukmawaty E, Ahmad A, Karim A, Dwyana Z, Karim H, Larekeng SH, et al. Effect of geographical and agroclimatic location on phytochemicals and antioxidant activity of *Moringa oleifera* leaves. *J Adv Biotechnol Exp Ther.* 2024;7(3):556-71. <https://doi.org/10.5455/jabet.2024.d48>
 44. Khan A, Kanwal F, Ullah S, Fahad M, Tariq L, Altaf MT, et al. Plant secondary metabolites-Central regulators against abiotic and biotic stresses. *Metabolites.* 2025;15(4):276. <https://doi.org/10.3390/metabo15040276>
 45. Yazıcı H, Çolak S, Duran U. Investigation geographic origin of *Laurus nobilis* L. leaves using FTIR, SEM-EDX and XRD analysis. *Spectrosc Lett.* 2025;58(2):149-60. <https://doi.org/10.1080/00387010.2024.2414053>
 46. Imelda E, Khairan K, Lubis RR, Karma T, Idroes R. Impact of environmental and geographical position on the chemometric classification of ethanol extracts from *Isotoma longiflora* leaves. *Glob J Environ Sci Manag.* 2024;10(1):155-68. <https://doi.org/10.22034/gjesm.2024.01.11>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc

See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.