



RESEARCH ARTICLE

Enhancing okra performance through osmo-priming: Effects on germination, yield and quality

Navneet Kaur¹, Dipika Mal^{1*} & Arshdeep Singh²

¹Department of Horticulture, School of Agriculture, Lovely Professional University, Phagwara 144 411, Punjab, India

²Department of Agronomy, School of Agriculture, Lovely Professional University, Phagwara 144 411, Punjab, India

*Correspondence email - dipika.21885@lpu.co.in, dipikaubkv@gmail.com

Received: 01 June 2025; Accepted: 19 July 2025; Available online: Version 1.0: 09 August 2025; Version 2.0: 22 August 2025

Cite this article: Navneet K, Dipika M, Arshdeep S. Enhancing okra performance through osmo-priming: Effects on germination, yield and quality. Plant Science Today. 2025; 12(3): 1-11. <https://doi.org/10.14719/pst.9755>

Abstract

To assess the impact of various seed priming techniques on okra (Punjab Suhawani variety) performance. Seeds were sourced from Punjab Agricultural University; Ludhiana and the study was structured using a randomized block design (RBD) with three replications and eight treatments with plot size of 2.25×3.75m and followed spacing was 45×15 cm. The treatments included: T₀ (un-primed) control, T₁ (hydro-priming), T₂ (osmo-priming with 5 % polyethylene glycol), T₃ (osmo-priming with 3 % potassium chloride), T₄ (halo-priming with 5 % potassium nitrate), T₅ (halo-priming with 1.5 % magnesium nitrate), T₆ (hormonal priming with 50 ppm gibberellic acid) and T₇ (hormonal priming with 50 ppm naphthalene acetic acid). For each treatment, seeds were soaked in the prepared solutions for 24 hrs before sowing under the shade. The study assessed germination percentage (92.17 %), number of days to germination (5.06 days), seedling dry weight (30.10 mg), seedling length (23.66 cm), seedling Vigor index (2166.95 I and 2627.83 II), root and shoot ratio (0.69), number of pods plant⁻¹ (12.28), individual pod weight (9.28 g), pod yield plot⁻¹ (14.24 kg), pod yield plant⁻¹ (14.29 g), pod yield tonnes hectare (12.89 t ha⁻¹), chlorophyll index (52.13) and mucilage content (5.57 %) and iodine content (mg/kg). The highest germination rate and seedling performance were recorded in T₂ (osmo-priming with 5 % PEG), followed closely by T₆ (hormonal priming with 50 ppm GA₃). In contrast, the unprimed seeds (T₀) control showed no significant improvement in any of the evaluated parameters.

Keywords: hormonal priming; okra; osmo-priming; responsible consumption and production; seed priming and quality

Introduction

Okra (*Abelmoschus esculentus*), a member of the Malvaceae family, as a popular vegetable of India. It is a staple summer vegetable of many countries (1, 2). It has a somatic chromosome number of 2n = 130. Commonly known as “lady finger,” okra is often called “a perfect villager’s vegetable” due to its widespread popularity and suitability for cultivation in rural areas (3), different species of the *Abelmoschus* genus are cultivated. Due to its significant economic value, particularly in nutrition, okra is extensively grown in Africa and America too. In India, okra cultivation spanned 526 thousand hectares in 2019-2020, yielding approximately 6,505 thousand metric tonnes (4). In Punjab, okra production stands at around 6.3 million tonnes from 0.51 million hectares. Okra seeds do not germinate at temperatures below 20 °C, with an optimal germination temperature of 29 °C. The fastest emergence rate occurs between 30°C and 35 °C; however, higher temperatures can delay flower initiation and blooming. Structurally, okra stems and leaves have pubescence, while its flowers are distinguishable by their yellow hue with a crimson-centered margin (5). The edible part of the plant is the capsule (pod), which typically measures 15-20 cm in length. Immature okra fruits are widely consumed as vegetables and used in various dishes such as salads, soups and stews, either dried or boiled

(6). Additionally, with the increasing demand for fresh vegetables in local and global markets, okra has gained significant popularity. Apart from its nutritional benefits, okra is also valued for its applications in oil production, biofuel and medicinal products (7, 8). Okra leaves are useful for many medicinal purposes. Nonetheless, okra leaves are thought to be effective in treating many illnesses, including diabetes, heart disease, digestive disorders and more. Mineral enrichment of okra is necessary to maintain normal cellular homeostasis. It is often foreseen that okra fruits can help prevent diabetes and obesity (9,10). After harvesting the okra fruit, the green stems are used to extract a crude mucilaginous substance, which finds applications in both the food industry and the pharmaceutical industry (11). Longitudinal okra flesh is rich in minerals and has significant levels of these minerals i.e. 0.6 to 1 mg iron (Fe), 81 to 82 mg calcium (Ca) and 61 to 63 mg phosphorus (P) respectively. Also present in okra is pro-vitamin A (185 0g), vitamin B2 (0.08 mg) and vitamin B complex (0.08 mg) and vitamin B complex (0.04 mg) respectively (12). Okra can be recognized as an abundant source of nutrients with advantageous health benefits. Consequently, because of the magnificence of polyphenols, they could serve as an extremely fruitful antioxidant source. Additionally, studies have confirmed that the fruit extract of okra contains several phenolic compounds, including protocatechuic acid, rutin,

catechin, quercetin and quercetin-3-O-gentiobioside (13). It has been reported that seed emergence and seedling growth, seed priming plays a vital role by initiating the hydration process, which activates the α -amylase enzyme. This enzyme facilitates the breakdown of starch reserves into simpler carbohydrates, providing essential energy to the embryo for respiration and supporting vigorous seedling development. Through priming, seeds become better prepared for rapid and uniform germination, enhancing overall plant establishment (14). Hydro-priming can improve the percentage of seeds that germinate and the emergence of seedlings in both saline and non-saline environments (15, 16). This controlled imbibition enables the embryo to activate its metabolic processes without radicle protrusion, thereby maintaining seed viability while enhancing germination (17). Generally, PEG (polyethylene glycol) is used for osmo-priming, but there are many other osmotic solutions involved, including potassium dihydrogen phosphate, magnesium sulfate, potassium nitrate, calcium chloride, potassium chloride, polyethylene glycol and mannitol are involved (18). When growing okra, seed hardness is a major issue since it hinders rapid germination and uniform field stands due to diminished, delayed and irregular emergence (3). Additionally, because okra seeds can be hard seeded, they may exhibit poor seedling emergence and vigour, resulting in a generally low germination rate (19). In seed priming, seeds are initially soaked in a pre-sowing treatment that allows them to absorb the necessary amount of water, triggering pre-germinative metabolic activities. However, this process can cause the seeds to lose their desiccation tolerance if the radicle begins to protrude prematurely. In addition to this, osmo-priming uses osmotic solutions with low water potential to slow down water uptake.

Seed priming treatments will have synergistic effects on the okra seed germination and the objectives of the study were, to determine the effects of seed priming on seed germination, yield and quality attributes and evaluate and identify the best priming treatment out of various methods of osmo-priming.

Materials and Methods

The field experiment was conducted at the experimental farm of the School of Agriculture, Lovely Professional University, Phagwara, Punjab, to evaluate the effects of various seed priming treatments on the emergence, yield and quality of okra. Seed priming, which involves soaking seeds in specific solutions before sowing, has been shown to significantly enhance Germination percentage (%), shoot length (cm), root length (cm), seedling length (cm), seedling dry weight (mg), seedling vigour index (i) and seedling vigour index (ii), root and shoot ratio, number of pods plant⁻¹, individual pod weight (g), pod yield plot⁻¹ (kg), yield plant⁻¹ (g), pod yield tonnes hectare (t ha⁻¹), chlorophyll index (SPAD) and mucilage content (%) and iodine content (mg kg⁻¹) of okra. The research trial utilized the okra variety Punjab-Suhawani. The Priming treatments applied were as follows: T₀- Control (un-primed seeds), T₁- Hydro-priming, T₂- Osmo-priming with 5 % polyethylene glycol, T₃- Osmo-priming with 3 % KCl, T₄- Halo-priming with 5 % KNO₃, T₅- Halo-priming with 1.5 % Mg (NO₃)₂, T₆- Hormonal-

priming with 50 ppm GA₃, T₇- Hormonal-priming with 50 ppm NAA. The site of the experimental plot falls under the subtropical category and the rainfall during the crop growing season was recorded as 486.1 mm. Seeds were fully submerged in the respective priming solutions for 24 hrs under the shade. After priming, the seeds were air-dried at room temperature for approximately 3 hrs to restore moisture content to normal levels.

Germination test

The rolled towel method was used for seeds in each priming treatment, which were tested for germination later. From each treatment, at least 25 seeds were taken and seeds were placed uniformly on the germination paper. Moreover, with the help of distilled water paper was moistened. Eventually, in the seed germinator, rolled towel papers were retained at a constant temperature of 25±1 °C and 95 % (RH).

Germination percentage

This formula was used to calculate the germination percentage (%).

Germination percentage =

$$\frac{\text{Total number of germination seeds}}{\text{Total number of seeds sown}} \times 100 \quad \text{Eq. 1}$$

Number of days to germination

The number of days from sowing until the emergence of the first leaf is recorded to determine the days to germination.

Seedling growth

To evaluate their growth response, seedlings were cultivated under standard settings. The length of the seedling (branch plus root) was measured in centimeters at 11 days after emergence (DAE). From the collar area to the cotyledon attachment site, the length of the shoot was measured. By measuring from the collar to the root tip, the length of the roots was ascertained. To assess the growth response, seedling growth was tracked.

Dry weight of seedling (mg)

The seedlings used for measuring root and shoot lengths were the same ones from the germination test. These seedlings were dried in a hot air oven at 60 °C for 24 hrs. Afterward, they were placed in desiccators for 30 min to complete the drying process. The dry weight of the seedlings was recorded in mg per ten seedlings. An electronic balance was utilized to obtain the precise weight of the dried seedlings.

Seedling length (cm)

Ten healthy seedlings were randomly selected during the first count to measure seedling length. The length was determined by measuring from the tip of the primary leaf to the tip of the primary root using a scale. The average length was then calculated and expressed in cm.

Seedling vigour index

Seedling vigour index (SVI) is commonly calculated using two formulas, known as SVI-I and SVI-II and was calculated by following the modified formula.

Seed Vigour index (i) =

$$\text{Germination (\%)} \times \text{Seedling length (cm)} \quad \text{Eq. 2}$$

Seed Vigour index (ii) =

$$\text{Germination (\%)} \times \text{Seedling dry weight (mg)} \quad \text{Eq. 3}$$

Root-Shoot ratio

From each treatment, 10 healthy plants were selected. The roots were carefully separated from the shoots by cutting at the soil line. Each root and shoot sample were weighed individually and the data for both parts of every plant were recorded. Using this information, the root-to-shoot ratio can be calculated.

$$\text{Root - Shoot ratio} = \frac{\text{Dry wt. of roots}}{\text{Dry wt. of top of plant}}$$

Eq. 4

Number of pods plant⁻¹

Pods were harvested from ten randomly tagged plants and after the counting, the average number of pods plant⁻¹ was calculated and then it will record the data.

Individual weight of the pods (g)

For each plot, ten healthy plants were selected and the weight of ten healthy pods from these plants was recorded. The individual weight of each pod was measured using a digital balance.

Pod yield plant⁻¹ (g)

At harvest, fresh fruits from the tagged plants were collected separately in polythene bags. The total average yield plant⁻¹ was then calculated for each plot.

Pod yield plot⁻¹ (kg) & Pod yield (t ha⁻¹)

During the harvest, fresh pods were picked from the tagged plants in all plots. The pods were collected separately in polythene bags throughout the harvesting period. Afterwards, the total average yield plant⁻¹ was calculated for each plot. Fresh pods were harvested from the tagged plants across all plots. Throughout the harvesting period, the pods were collected separately in polythene bags. Finally, the total average yield in tonnes ha⁻¹ was calculated for each plot.

Chlorophyll index

Chlorophyll index is computed after ten plants that are tagged are drawn randomly in each plot. The most common technique of measuring chlorophyll concentration is a SPAD meter (SPAD-502) and it was what was used to calculate the chlorophyll in the okra leaves.

Mucilage content (%)

A 25 g sample was mixed with 100 mL of distilled water and left to soak for 24 hrs. The resulting suspension was then filtered using a muslin cloth. After filtration, 50 mL of ethanol was added to the filtrate. The mixture was blended for at least 15 min using a magnetic stirrer. Next, the sample was filtered again using pre-weighed filter paper and then dried in an oven. Once dried, the filter paper with the residue was weighed again. The viscous matter is shown in the Fig. 1 (a, b, c, d). The mucilage content was calculated using the following formula (20).

$$\text{Mucilage content (\%)} = \frac{w_2 - w_1}{w} \quad \text{Eq. 5}$$

w_2 = Weight of filter paper along with material after drying, w_1 =




Fig. 1 (a, b, c, d). Illustrating the mucilage of okra.

Weight of pre-weighed filter paper, W = Weight of sample.

Iodine content (mg)

By using the Teepol detergent, the glassware was thoroughly cleaned, followed closely by washing with distilled water. After washing with distilled water, it was soaked overnight in a 10 % hydrochloric acid solution. The glassware was rinsed with tap water as well as thoroughly rinsed again with distilled water, after the soaking. Finally, before use, the glassware was dried in a hot air oven at a temperature of at least 60 °C.

For sample preparation, by using an electric mixer-cum-grinder the dried samples were ground into a fine powder. To maintain the quality, the powdered samples were then stored in airtight polyethylene bags. One gram of the dried and finely ground sample was accurately weighed and transferred into a clean test tube for the process of analysis. The sample was added with 1 mL of a 10 times diluted 6 M KOH solution. The sample was then placed in an oven and dried overnight at 95 ± 1 °C for the blank. Subsequently, the test tubes were transferred to a muffle furnace preheated to 100 °C. The samples were burnt for 1 hr, when the furnace temperature was then raised to 600 °C. Air was circulated by moving the tubes every 15 min for 15 seconds to ensure proper combustion during this period. After the ashing process, the tubes were transferred to a desiccator and allowed to cool to room temperature. If ashing was incomplete, 1 mL of a 10 times diluted zinc sulfate solution was added to all tubes and the drying and ashing process was repeated as described above.

Whenever ashing was complete, added the 0.2 mL of water to each test tube, followed by thorough stirring and vortexing. The volume was then adjusted to 5 mL with water. Finally, the tubes were centrifuged at 3500 rpm for 30 min and the supernatant was collected for the final assay. Into a test tube, the following reagents were added in sequence: 0.5 mL of double-distilled water, 0.5 mL of a H_2SO_4 : HCl solution, 0.5 mL of cerate reagent (prepared by dissolving 0.316 g of ammonium ceric (IV) sulfate in 15 mL of water, to which 40 mL of concentrated nitric acid was added dropwise), 5 mL of concentrated H_2SO_4 . The mixture was then made up to a final volume of 100 mL with distilled water. After that, 0.5 mL of arsenic reagent was added. (The arsenic reagent was prepared by dissolving 0.593 g of arsenic trioxide and 0.6 g of potassium hydroxide in 30 mL of water, then adding 0.1 mL of concentrated HCl and finally making the volume up to 100 mL with distilled water.)

The combined solution was thoroughly mixed and incubated at room temperature for 1 min. After incubation, 0.5 mL of blank, standard, or sample was added, mixed well and the reaction was allowed to proceed for 1 minute. The optical density (OD) was then measured at 400 nm. The slope was calculated from the standard curve using pure potassium iodide (KI) (21).

$$\text{Dry matter content (\%)} = \frac{\text{Final dry wt. of the sample}}{\text{Initial fresh wt. of the sample}} \times 100$$

Eq. 6

Statistical analysis

Data was organized and analyzed using one-way ANOVA through SPSS software. Treatment means were compared

using the Least Significant Difference (LSD) test at a significance level of $p < 0.05$.

Results and Discussion

Effect of different seed priming with osmo-priming on Okra crop

Germination percentage and number of days to germination

The seed pre-treatment with different chemicals had a significant effect ($p < 0.05$) on germination percentage (%). However, it showed no significant effect on the number of days to emergence. The pooled data indicated that PEG is a priming agent that germinates more okra seeds than un-primed seeds. The results indicated that T_2 (92.17 %) showed maximum germination percentage on the different sources of priming like T_6 (89.50 %) table 1. Conversely, T_0 (62.33 %) that contained unprimed seeds recorded low percentage of germination. Nonetheless, the overall combined data indicated that it has not been significantly influenced by various priming treatments on the number of days to germination ($p < 0.05$). The time to germination was highest in treatment T_6 with 5.53 days of germination and secondly in treatment T_2 which had 5.06 days. Similar results were found with T_2 and T_6 however, on 5.06 and 5.53 days. Similarly, T_6 and T_7 recorded the same trends with 5.06 and 7.81 days respectively. Conversely, T_0 (8.22) days that involved unprimed seeds had the least number of days to germination. All pre-germination metabolic functions are believed to require a particular amount of water to activate and enter into the lag phase before germination. This optimum water availability is offered by priming treatments and facilitates the occurrence of such metabolic activities successfully. As a result, different priming agents not only stimulate pre-germination metabolism but also positively influence the emergence percentage of seeds (17).

Shoot and root length (cm)

The combined data identified a significant difference in the shoot and root lengths of different kind of priming treatment ($p < 0.05$). This phenomenon of growth increase in shoot length after priming of seeds can be explained by increasing the cell division, meristem activity and the stimulation of the elongation of cells. All these physiological processes have been found to enhance seedling growth and vigour after being subjected to priming treatments (22) table 2. The finding was that regarding all the priming treatments, T_6 (15.83 cm) with GA_3 50 ppm had the longest shoot recorded as being significantly at par with T_2 which was 15.50 cm. Conversely, the bottom sample T_0 (7.55 cm) comprised of un-primed seeds that recorded the fewest shoots.

Conversely, the pooled data indicated that, the various priming treatments influenced the length of the longest root substantially ($p < 0.05$). Thus, T_2 with PEG 5 % had the maximum root length of 8.16 cm, followed by T_6 with GA_3 50 ppm having root length of 7.47 cm. Also, they were quite equal with T_2 and T_7 i.e., 8.16 cm and 7.63 cm. Likewise T_7 and T_6 are also very much at par with the T_6 being (7.47 cm) and the T_7 being (7.63 cm). Conversely, the minimum root length was observed in T_0 (3.57 cm), which is the value of the un-primed seeds table 2. High germination percent synchronized seedling emergence and better shoot and root length of primed seeds

Table 1. Effect of various priming treatments on the germination percentage (%) and number of days to germination in Okra

Treatment's detail	Germination percentage (%)			Number of days for germination Days		
	2019	2020	Pooled	2019	2020	Pooled
T ₀ (un-primed)	62.00 ^a ±1.00	62.67 ^a ±1.53	62.33 ^a ±1.15	8.00 ^a ±0.77	8.45 ^a ±0.77	8.22 ^a ±1.12
T ₁ (hydropriming)	77.33 ^d ±2.08	76.67 ^d ±1.53	77.00 ^d ±1.32	6.63 ^a ±0.83	6.55 ^{bcd} ±0.83	6.59 ^{bcd} ±0.96
T ₂ (PEG 5 %)	92.33 ^a ±1.53	92.00 ^a ±1.00	92.17 ^a ±0.29	5.00 ^b ±0.87	5.13 ^d ±0.87	5.06 ^d ±0.81
T ₃ (KCL) 3 %	72.33 ^e ±1.53	71.00 ^e ±1.00	71.67 ^e ±0.29	7.11 ^{ab} ±1.12	7.19 ^{abc} ±1.12	7.15 ^{abc} ±1.40
T ₄ (KNO ₃) 5 %	81.00 ^c ±1.00	82.00 ^c ±1.00	81.50 ^c ±0.50	6.33 ^{ab} ±1.58	6.67 ^{bcd} ±0.57	6.50 ^{bcd} ±0.87
T ₅ (MgNO ₃) ₂ 1.5 %	66.67 ^f ±1.53	67.33 ^f ±1.53	67.00 ^f ±0.50	7.77 ^a ±1.02	7.84 ^{ab} ±1.01	7.81 ^{ab} ±0.47
T ₆ (GA ₃) 50ppm	90.00 ^a ±1.00	89.00 ^a ±1.00	89.50 ^a ±1.00	5.43 ^b ±0.63	5.63 ^{cd} ±0.63	5.53 ^d ±0.51
T ₇ (NAA) 50ppm	85.00 ^b ±1.00	86.00 ^b ±1.00	85.50 ^b ±0.87	6.18 ^{ab} ±0.60	6.31 ^{bcd} ±0.60	6.25 ^{cd} ±0.74
CD (<i>p</i> <0.05)	2.50	1.69	1.51	1.92	1.51	1.41

Each value in the treatment is a means of three replicates ± SD. Means followed by the same letter (s), in each row, are not significantly different (ANOVA, LSD test *p*<0.05).

Table 2. Effect of various priming treatments shoot length and root length (cm) in Okra

Treatment's detail	Shoot length (cm)			Root length (cm)		
	2019	2020	Pooled	2019	2020	Pooled
T ₀ (un-primed)	7.28 ^f ±1.23	7.82 ^e ±0.55	7.55 ^e ±0.79	3.55 ^c ±0.95	3.59 ^c ±0.89	3.57 ^c ±0.92
T ₁ (hydropriming)	10.69 ^{de} ±1.05	11.36 ^{cd} ±1.12	11.02 ^{ef} ±0.58	7.46 ^{ab} ±1.11	6.46 ^a ±1.04	6.53 ^b ±1.00
T ₂ (PEG 5 %)	15.94 ^a ±0.99	15.06 ^{ab} ±1.42	15.50 ^{ab} ±1.20	8.84 ^a ±0.96	7.48 ^a ±1.03	8.16 ^a ±0.99
T ₃ (KCL) 3 %	11.55 ^{cde} ±0.94	12.74 ^c ±0.95	12.15 ^{de} ±0.56	6.32 ^b ±0.90	6.34 ^{ab} ±0.87	6.33 ^b ±0.88
T ₄ (KNO ₃) 5 %	12.54 ^{cd} ±1.04	13.15 ^{bc} ±1.03	12.85 ^{cd} ±0.87	7.41 ^{ab} ±0.09	7.49 ^a ±1.17	7.45 ^{ab} ±0.56
T ₅ (MgNO ₃) ₂ 1.5 %	9.66 ^e ±0.89	9.99 ^d ±0.93	9.83 ^f ±0.70	4.11 ^c ±1.02	5.21 ^b ±1.01	4.66 ^c ±0.52
T ₆ (GA ₃) 50ppm	14.74 ^{ab} ±0.88	16.92 ^a ±1.04	15.83 ^a ±0.83	6.59 ^b ±0.97	8.78 ^a ±0.96	7.47 ^{ab} ±1.07
T ₇ (NAA) 50ppm	13.13 ^{bc} ±1.44	14.84 ^b ±0.90	13.99 ^{bc} ±0.96	7.69 ^{ab} ±0.76	7.56 ^a ±1.28	7.63 ^{ab} ±1.01
CD (<i>p</i> <0.05)	1.91	1.92	1.54	1.48	1.19	1.91

Each value in the treatment is a means of three replicates ± SD. Means followed by the same letter (s), in each row, are not significantly different (ANOVA, LSD test *p*<0.05).

can be possibly explained by initiation of certain metabolic processes in the seed embryo (23, 17). So-called soybean and cumin have also been used to depict that osmo-priming as well as hydropriming can also enhance root and shoot growth (24).

Seedling length (cm) and dry weight of seedling (mg)

The pooled data in seedling length (root length+ shoot length) as recommended that numerous priming sources affected seedling length significantly (*p*<0.05) table 3. The result showed that among various priming sources, maximum seedling length was recorded in T₂ (23.66 cm), which is PEG 5 %, followed closely by T₆ (23.52 cm), which is GA₃ 50 ppm. Additionally, there were *at par* shows between T₂, T₇ and T₆ which were (23.66 cm, 23.61 cm and 21.52 cm, respectively). The minimum seedling length was observed in T₀ (13.60 cm), un-primed seeds, followed by T₅ (14.49 cm), which is Magnesium nitrate 1.5 %. On the opposite side, as shown in table no. 3, the pooled data revealed that different priming treatments had a significant impact on the dry weight of seedlings (*p*<0.05). The highest seedling dry weight was observed in T₂ at 30.10 mg, followed closely by T₇ at 28.46 mg and T₆ at 28.44 mg. Additionally, the dry weight of seedling values T₂, T₇ and T₆ are significantly at par with each other, which are 30.10 mg, 28.46 mg and 28.44 mg, respectively. In contrast, the lowest dry weight of seedlings of okra (23.59 mg) was observed in T₀, the unprimed control. Multiple studies

confirm that seed priming leads to a stronger and more effective root and shoot system, as well as a reduction in physiological deterioration, due to a series of physiological and biochemical changes within the seed. Seed priming initiates early metabolic activities-such as enzyme activation, protein and nucleic acid synthesis and membrane repair before radicle emergence, which increases the physiological activity of the embryo and supports the mobilization of food reserves into developing seedlings. This could have led to DNA repair because the imbibition process creates enzymes and protein membranes (25, 26). Osmo-priming treatments increased the dry weight of seedlings in all the investigated Berseem clover species, both primed and non-primed (27). When the seeds sprout, it reduces the seed coat's adherence. Seed germination and increased vigour during osmotic priming may be due to active nutrition and the resynthesis of certain enzymes. Furthermore, the production of DNA and RNA might have started. Eliminating obstacles causes seeds to germinate more quickly and eventually grow into a sturdy shoot with a higher dry weight (28).

Seedling vigour index (I) and (II)

Seed priming treatments significantly influenced both Seedling Vigour Index I and II, highlighting the impact of various priming methods on enhancing seedling growth and development. Priming can conceivably increase the competitive ability of the plant for the available resources such as nutrients, water and

Table 3. Effect of various priming treatments on seedling length (cm) and dry weight of seedlings (mg) in Okra

Treatment's detail	Seedling length (cm)			Dry weight of seedlings (mg)		
	2019	2020	Pooled	2019	2020	Pooled
T ₀ (un-primed)	10.83 ^d ±2.18	11.39 ^e ±1.26	13.60 ^e ±4.34	23.64 ^d ±0.89	23.54 ^d ±1.07	23.59 ^d ±0.98
T ₁ (hydropriming)	18.15 ^{bc} ±0.90	17.82 ^c ±0.11	17.98 ^d ±0.09	26.27 ^c ±1.05	27.42 ^{bc} ±0.96	26.85 ^{bc} ±0.83
T ₂ (PEG 5 %)	24.78 ^a ±0.02	22.54 ^a ±1.07	23.66 ^a ±1.03	29.84 ^a ±0.95	30.35 ^a ±1.37	30.10 ^a ±1.15
T ₃ (KCL) 3 %	17.87 ^{bc} ±0.81	19.08 ^c ±0.83	18.48 ^{cd} ±0.32	26.42 ^c ±0.84	26.54 ^c ±1.14	26.48 ^c ±0.28
T ₄ (KNO ₃) 5 %	19.95 ^{abc} ±1.11	20.64 ^{ab} ±1.01	20.30 ^{bcd} ±0.38	27.74 ^{bc} ±0.90	28.51 ^{abc} ±1.52	28.13 ^{ab} ±0.56
T ₅ (MgNO ₃) ₂ 1.5 %	13.77 ^d ±1.72	15.20 ^d ±1.93	14.49 ^e ±1.17	24.16 ^d ±0.92	24.61 ^d ±1.30	24.39 ^{de} ±0.19
T ₆ (GA ₃) 50ppm	21.33 ^a ±1.69	25.70 ^a ±0.84	23.52 ^{ab} ±0.55	28.73 ^{ab} ±0.81	28.15 ^{bc} ±0.95	28.44 ^a ±0.88
T ₇ (NAA) 50ppm	20.82 ^{ab} ±1.86	22.40 ^a ±2.16	21.61 ^{abc} ±1.77	28.06 ^{abc} ±1.55	28.85 ^{ab} ±1.08	28.46 ^a ±1.16
CD (<i>p</i> <0.05)	2.69	1.87	3.11	1.86	1.88	1.34

Each value in the treatment is a means of three replicates ± SD. Means followed by the same letter (s), in each row, are not significantly different (ANOVA, LSD test *p*<0.05).

Table 4. Effect of various priming treatments on seedling vigour index-i (SVI) and seedling vigour index-ii (SVI) in Okra

Treatment's detail	Seedling vigour index-i			Seedling vigour index-ii		
	2019	2020	Pooled	2019	2020	Pooled
T ₀ (un-primed)	687.96 ^e ±105.49	755.04 ^e ±93.96	721.50 ^e ±53.38	1374.28 ^e ±109.93	1342.86 ^e ±88.16	1358.57 ^e ±87.52
T ₁ (hydropriming)	1274.26 ^c ±95.08	1366.06 ^c ±91.03	1320.16 ^d ±43.26	2152.51 ^c ±84.50	2027.99 ^b ±137.75	2090.25 ^c ±88.59
T ₂ (PEG 5 %)	2260.81 ^a ±95.75	2073.08 ^a ±82.17	2166.95 ^a ±56.76	2721.48 ^a ±145.82	2534.18 ^a ±94.28	2627.83 ^a ±26.01
T ₃ (KCL) 3 %	1146.42 ^c ±141.03	1241.42 ^c ±94.43	1193.92 ^e ±72.89	1703.59 ^d ±157.18	1756.96 ^c ±103.35	1730.28 ^d ±81.88
T ₄ (KNO ₃) 5 %	1655.93 ^b ±83.14	1670.07 ^b ±107.78	1663.00 ^c ±93.55	2212.87 ^{bc} ±89.66	2116.53 ^b ±58.79	2164.70 ^c ±16.01
T ₅ (MgNO ₃) ₂ 1.5 %	983.15 ^d ±95.98	973.79 ^d ±94.86	978.47 ^f ±95.00	1673.48 ^d ±85.47	1573.84 ^d ±117.09	1623.66 ^d ±49.80
T ₆ (GA ₃) 50ppm	1955.36 ^a ±108.58	2263.46 ^a ±110.52	2109.41 ^a ±94.75	2473.04 ^a ±94.96	2736.82 ^a ±94.32	2604.93 ^a ±94.63
T ₇ (NAA) 50ppm	1728.08 ^b ±135.09	1770.72 ^b ±115.03	1749.4 ^b ±57.54	2365.17 ^{ab} ±113.86	2365.17 ^a ±108.06	2365.17 ^b ±110.86
CD (<i>p</i> <0.05)	138.30	171.54	82.25	186.50	185.22	119.97

Each value in the treatment is a means of three replicates ± SD. Means followed by the same letter (s), in each row, are not significantly different (ANOVA, LSD test *p*<0.05).

sunlight and thus, can affect seedling vigour (29). The pooled data indicated that different priming treatments had a significant effect on both Seedling Vigour Index-I and Seedling Vigour Index-II (*p*<0.05). Table 4 shows that among the several priming sources, the highest seedling vigour (i) was recorded in T₂ (2166.95) followed closely by T₆ (2109.41). In contrast, the lowest seedling vigour index (i) was observed in T₀ (721.50), the untreated control, followed by T₅ (978.47). On the other hand, table 4, the pooled data revealed that different priming treatments had a significant impact on seedling vigour index II (*p*<0.05). Therefore, seedling vigour index (ii) was recorded in T₂ (2627.83), which is PEG 5 %, followed by T₆ (2604.93). Whereas they were significantly at par shows between T₂ and T₆ which were 2627.83 and 2604.93. In addition, the minimum seedling vigour index (ii) was observed in T₀ (1358.57), which is un-primed seeds, followed by T₅ (1623.66), which is Magnesium nitrate 1.5 %. Consistent with this result is the finding of (30) that seed priming raised antioxidant levels in seeds, including glutathione and ascorbate. Through a reduction in lipid peroxidation activity, these enzymes speed up germination. A study was done on Osmo and halo-priming of okra showed that seed priming holds close conformity in the enhancement of seed vigour (17).

Root-shoot ratio

The pooled data indicated that various priming sources significantly affected shoot length and root length (*p*<0.05). Table 5 shows that among the various priming treatments, T₂ recorded the highest root-shoot ratio of okra at 0.69, followed by T₆ with a ratio of 0.65. Moreover, the minimum (0.21) root/shoot ratio of okra was found in T₀. (31) observed physiological and biochemical alterations that enhanced the physiological activity of the embryo and facilitated the mobilization of food stores into the developing seedlings. (32) that resulted in the establishment of a more resilient and efficient root and shoot system, thereby significantly mitigating physiological degradation (33). This process may have accelerated the repair of DNA, protein membranes and enzymes during imbibition (34). The priming-induced increase in the root to shoot ratio may be attributed to increased metabolic activities and greater cell wall extensibility. These physiological changes are possibly activated by the slower imbibition rate that occurs during hydro or matrix priming.

Effect of different seed priming with osmo-priming on yield attributes characters

Number of pods plant⁻¹ and individual pod weight (g)

The combination of data showed that there was a significant influence of different priming sources on the number of pods plant⁻¹ and individual pod weight (*p*<0.05). Table 6 indicates that

Table 5. Effect of various priming treatments on root-shoot ratio in Okra

Treatment's detail	Root-shoot ratio		
	2019	2020	Pooled
T ₀ (un-primed)	0.23 ^f ±0.04	0.19 ^f ±0.03	0.21 ^e ±0.03
T ₁ (hydropriming)	0.44 ^{de} ±0.10	0.38 ^{cd} ±0.04	0.41 ^{cd} ±0.06
T ₂ (PEG 5 %)	0.69 ^a ±0.06	0.62 ^{ab} ±0.05	0.65 ^{ab} ±0.03
T ₃ (KCL) 3 %	0.52 ^{cd} ±0.04	0.31 ^{de} ±0.04	0.42 ^{cd} ±0.02
T ₄ (KNO ₃) 5 %	0.48 ^d ±0.10	0.43 ^c ±0.06	0.45 ^c ±0.08
T ₅ (MgNO ₃) ₂ 1.5 %	0.38 ^e ±0.05	0.25 ^{ef} ±0.04	0.32 ^{de} ±0.01
T ₆ (GA ₃) 50ppm	0.72 ^{ab} ±0.10	0.66 ^a ±0.07	0.69 ^a ±0.08
T ₇ (NAA) 50ppm	0.59 ^{bc} ±0.10	0.55 ^b ±0.06	0.57 ^b ±0.08
CD ($p<0.05$)	0.12	0.09	0.09

Each value in the treatment is a means of three replicates $\pm$ SD. Means followed by the same letter (s), in each row, are not significantly different (ANOVA, LSD test $p<0.05$).

Table 6. Effect of different seed priming treatments on the number of pods plant⁻¹ and individual pod weight (g) in Okra

Treatment's detail	Number of pods plant ⁻¹			Individual pod weight (g)		
	2019	2020	Pooled	2019	2020	Pooled
T ₀ (un-primed)	7.00 ^e ±1.00	6.43 ^c ±0.59	6.72 ^c ±0.53	7.92 ^d ±0.54	7.20 ^d ±0.08	7.56 ^d ±0.23
T ₁ (hydropriming)	10.33 ^{cd} ±0.58	9.07 ^{ab} ±0.90	9.70 ^{ab} ±0.72	8.42 ^c ±0.06	8.40 ^c ±0.08	8.41 ^c ±0.06
T ₂ (PEG 5 %)	12.67 ^a ±0.58	11.90 ^a ±0.10	12.28 ^a ±0.29	9.02 ^a ±0.15	9.55 ^a ±0.011	9.28 ^a ±0.11
T ₃ (KCL) 3 %	10.00 ^{cd} ±1.00	8.00 ^b ±1.00	9.00 ^b ±0.50	8.48 ^{bc} ±0.03	8.45 ^c ±0.04	8.47 ^c ±0.02
T ₄ (KNO ₃) 5 %	12.00 ^{ab} ±1.00	9.10 ^{ab} ±0.95	10.55 ^a ±0.53	8.54 ^{bc} ±0.04	8.56 ^b ±0.03	8.55 ^{bc} ±0.03
T ₅ (MgNO ₃) ₂ 1.5 %	9.00 ^d ±1.00	9.33 ^{ab} ±0.58	9.17 ^b ±0.58	8.34 ^{cd} ±0.03	8.37 ^c ±0.05	8.36 ^c ±0.02
T ₆ (GA ₃) 50ppm	11.00 ^{bc} ±1.00	9.63 ^a ±0.55	10.32 ^a ±0.55	8.87 ^{ab} ±0.20	8.75 ^a ±0.03	8.81 ^a ±0.09
T ₇ (NAA) 50ppm	12.33 ^{ab} ±0.58	8.80 ^{ab} ±0.95	10.57 ^a ±0.73	8.65 ^{abc} ±0.04	8.68 ^a ±0.04	8.67 ^{ab} ±0.02
CD ($p<0.05$)	1.52	1.25	0.86	0.38	0.09	0.18

Each value in the treatment is a means of three replicates $\pm$ SD. Means followed by the same letter (s), in each row, are not significantly different (ANOVA, LSD test $p<0.05$).

plant⁻¹ had the highest number of pods number namely 12.28 which was under treatment T₂. And the second highest number of pods plant⁻¹ was 10.57 which represented in treatment T₇. However, they are significantly at par between the T₂ and T₇ of 12.28 and 10.57 and T₇ and T₆ are at par with each other (10.57 and 10.32). Similarly, T₆ and T₄ are in par with each other which is (10.32 and 10.55). Also, minimum (6.72) number of pods plant⁻¹ was obtained on treatment of T₀. On the other hand, pooled data revealed that the sources of priming had significant effect on the individual pod weight (g) of okra ($p<0.05$) table 6 shows that treatment T₂ had the highest individual pod weight of okra which was 9.28 g.

Individual pod weight values of T₂, T₆ and T₇ (9.28g, 8.81g and 8.67g, correspondingly) were however proved to be significantly at par with one another, priming treatments. These were opposed by the lowest individual pod weight of okra (7.56 g) observed in T₀ treatment. Certainly, the increase in pods plant⁻¹ can be characterized by the number of increases in leaves. According to (35), seed priming influenced seed emergence, increase seedling growth and Vigour has consequence resulted in showing an increased number of leaves and in the course of time, it will bear more pods. In addition, the number of pods plant⁻¹ is the outcome of the final output, which is one of the significant factors. The number of pods will also increase in the same

fashion with the leaves number. The correlation between the pods and leaves are coherent. Polyethylene glycol solution help in the enhancement of the yield of wheat which is noted by (36) and agrees with the current findings (37). As primed seed plots can possibly yield more grain since primed seeds germinate faster and more evenly and seedlings are avidly grown. All these leads to numerous phonological and yield.

Effect of different seed priming with osmo-priming on okra yield

Pod yield plant⁻¹ (g) and pod yield plot⁻¹ (Kg)

The pooled data showed that different priming sources had a significant impact on both pod yield plant⁻¹ (g) and pod yield plot⁻¹ (Kg) ($p<0.05$). Based on table 7, treatment T₂ recorded the highest pod yield plant⁻¹, with a value of 113.96 g. However, several pod yield plant⁻¹ (g) values of T₂ and T₆ are significantly at par with each which are (113.96 g and 90.99 g). Similarly, T₄ and T₇ significantly at par with each other (90.25 g and 91.51g, respectively). The minimum pod yield plant⁻¹ (50.90 g) was found from the T₀. On the other hand, the pooled data demonstrated that different priming sources had a significant effect on pod yield plot⁻¹ (Kg) of okra ($p<0.05$). Treatment T₂ produced the highest pod yield plot⁻¹ at 14.24 Kg, followed closely by T₇ at 11.73 Kg table 7. The number of pod yield plot⁻¹

Table 7. Effect of different seed priming treatments on the Pod yield plant⁻¹ (g) and Pod yield plot⁻¹ (Kg) in Okra

Treatment's detail	Pod yield plant ⁻¹ (g)			Pod yield plot ⁻¹ (Kg)		
	2019	2020	Pooled	2019	2020	Pooled
T ₀ (un-primed)	55.44 ^e ±9.26	46.37 ^c ±4.71	50.90 ^c ±3.92	6.93 ^e ±1.16	5.80 ^c ±0.59	6.17 ^d ±0.56
T ₁ (hydropriming)	86.76 ^{cd} ±4.70	76.14 ^{ab} ±7.26	81.45 ^b ±5.79	10.87 ^{cd} ±0.57	9.52 ^{ab} ±0.91	10.19 ^b ±0.72
T ₂ (PEG 5 %)	114.30 ^a ±6.86	113.61 ^a ±1.90	113.96 ^a ±3.97	14.29 ^a ±0.86	14.20 ^a ±0.24	14.24 ^a ±0.50
T ₃ (KCL) 3 %	84.79 ^{cd} ±8.27	68.35 ^b ±7.71	76.57 ^b ±3.63	10.60 ^{cd} ±1.03	9.21 ^{ab} ±0.97	9.90 ^b ±0.52
T ₄ (KNO ₃) 5 %	102.53 ^{ab} ±8.65	77.97 ^{ab} ±8.04	90.25 ^a ±4.28	12.82 ^{ab} ±1.08	9.74 ^{ab} ±1.00	11.28 ^a ±0.53
T ₅ (MgNO ₃) ₂ 1.5 %	75.02 ^d ±8.23	78.15 ^{ab} ±4.82	76.59 ^b ±5.01	9.38 ^d ±1.03	8.46 ^b ±1.09	8.92 ^c ±0.55
T ₆ (GA ₃) 50ppm	97.72 ^{bc} ±10.90	84.25 ^a ±4.64	90.99 ^a ±5.87	12.22 ^{bc} ±1.36	10.53 ^a ±0.58	11.37 ^a ±0.73
T ₇ (NAA) 50ppm	106.32 ^{ab} ±4.94	76.38 ^{ab} ±7.94	91.51 ^a ±6.23	13.33 ^{ab} ±0.62	10.13 ^a ±0.58	11.73 ^a ±0.60
CD (<i>p</i> <0.05)	4.84	10.45	7.00	1.76	1.33	0.87

Each value in the treatment is a means of three replicates ± SD. Means followed by the same letter (s), in each row, are not significantly different (ANOVA, LSD test *p*<0.05).

(Kg) values of T₂ and T₇, are significantly at par with each other, which is (14.24 Kg and 11.73 Kg. Likewise, T₆ and T₄ significantly at par with each other (11.37 Kg and 11.28 Kg respectively). The minimum pod yield plot⁻¹ (6.17 Kg) was found from T₀. Seeds primed plots have contributed to the best performance in the field (38). Uniform and improved germination, strong seedling growth, a well-developed root system and efficient subsequent growth may be the biggest reasons to increase the final yield. improved vigor of the seedlings and improved vegetative and reproductive characteristics may have come up with higher pod production. A typical limiting variable, likely crust development, dwindling soil moisture and high salt, hinder effective emergence, has been improved because of the head start of germination considering that primed seeds are in a different developmental stage than dry seeds (39).

Pod yield (t ha⁻¹)

The pooled data revealed that different priming sources had a significant impact on pod yield measured in t ha⁻¹ (*p*<0.05). Based on table 8, treatment T₂ recorded the highest pod yield of okra, producing 12.89 (t ha⁻¹). The number of pod yield tonnes (t ha⁻¹) values of T₂ and T₆ are significantly at par with each other (12.89 t ha⁻¹ and 12.36 t ha⁻¹). Similarly, T₆ and were shown to be significantly at par with each other 12.36 t ha⁻¹ and 12.10 t ha⁻¹ respectively. The minimum (9.92 t ha⁻¹) pod yield of okra was found from T₀ treatment. The priming treatment greatly raised overall biomass and raised wheat and soybean

yields (40, 41). For any crop being studied, yield is the final product and is influenced by several variables, including soil types, environmental conditions and genetic composition. Priming may boost biological yield because it improves early seedling growth and plant nutrition (42).

Chlorophyll index and Mucilage content (%)

The pooled data noticed that the source of priming etiologically influenced value of SPAD of okra was significant (*p*<0.05). Table 9 reveals that okra showed a maximum value of SPAD in treatment T₂ which was 52.13. Nonetheless, this was not statistically different with treatment T₆ that recorded a SPAD value of 50.44. In T₀, the lowest value of SPAD of okra (40.46) was obtained. Conversely, the pooled data revealed that all the sources of priming had significant effect on the content of mucilage in okra at *p*<0.05 table 9 reveals that T₆ priming recorded the highest value of the mucilage content in okra at 5.64, which is closely followed by treatment T₂ at 5.57. There were however shows between T₂ and T₆ that were at par which were 5.57 and 5.64 %. The lowest (3.68 %) mucilage content of okra was seen in the T₀ and T₇. The assistance of Chlorophylls, plants have the capability of capturing a large proportion of the light energy besides the fact that the performance of chlorophyll in the plants can result into photosynthesis reactions in plants hence, chlorophyll is highly important pigment. But there might be a possibility that chlorophyll is particularly sensitive to several environmental stresses, as a consequence of which, it may lead to

Table 8. Effect of different seed priming treatments on the Pod yield (t ha⁻¹) in okra

Treatment's detail	Pod yield (t ha ⁻¹)		
	2019	2020	Pooled
T ₀ (un-primed)	9.88 ^e ±0.17	9.95 ^c ±0.23	9.92 ^e ±0.03
T ₁ (hydropriming)	10.86 ^c ±0.27	10.28 ^c ±0.18	10.57 ^{cd} ±0.06
T ₂ (PEG 5 %)	12.75 ^a ±0.23	13.02 ^a ±1.80	12.89 ^a ±0.20
T ₃ (KCL) 3 %	10.54 ^{cd} ±0.34	10.86 ^b ±0.30	10.70 ^c ±0.32
T ₄ (KNO ₃) 5 %	11.61 ^b ±0.27	11.74 ^a ±0.19	11.68 ^b ±0.21
T ₅ (MgNO ₃) ₂ 1.5 %	10.13 ^{de} ±0.14	10.36 ^c ±0.23	10.24 ^{de} ±0.16
T ₆ (GA ₃) 50ppm	12.51 ^a ±0.24	12.21 ^a ±0.21	12.36 ^a ±0.18
T ₇ (NAA) 50ppm	12.05 ^b ±0.44	12.15 ^a ±0.40	12.10 ^a ±0.34
CD (<i>p</i> <0.05)	0.46	0.45	0.39

Each value in the treatment is a means of three replicates ± SD. Means followed by the same letter (s), in each row, are not significantly different (ANOVA, LSD test *p*<0.05).

Table 9. Effect of different seed priming treatments on okra's chlorophyll index (SPAD) and mucilage content (%)

Treatment's detail	Chlorophyll Index (SPAD)			Mucilage content (%)		
	2019	2020	Pooled	2019	2020	Pooled
T ₀ (un-primed)	40.06 ^f ±1.78	40.85 ^e ±1.52	40.46 ^d ±1.15	3.66 ^e ±0.11	3.70 ^e ±0.06	3.68 ^e ±0.08
T ₁ (hydropriming)	44.81 ^{de} ±1.18	43.58 ^{de} ±1.47	44.19 ^c ±0.96	4.42 ^{cd} ±0.38	4.58 ^{cd} ±0.17	4.50 ^{cd} ±0.26
T ₂ (PEG 5 %)	51.38 ^a ±1.21	52.88 ^a ±1.73	52.13 ^a ±0.74	5.82 ^a ±0.15	5.33 ^a ±0.32	5.57 ^a ±0.20
T ₃ (KCL) 3 %	46.34 ^{cd} ±1.96	46.01 ^{cd} ±3.08	46.18 ^c ±1.81	4.44 ^{cd} ±0.12	4.41 ^{cd} ±0.08	4.43 ^{cd} ±0.10
T ₄ (KNO ₃) 5 %	48.27 ^{bc} ±1.43	49.69 ^{ab} ±0.94	48.98 ^b ±1.16	4.84 ^{bc} ±0.11	4.83 ^{bc} ±0.10	4.84 ^{bc} ±0.10
T ₅ (MgNO ₃) ₂ 1.5 %	42.58 ^{ef} ±1.92	44.98 ^d ±1.43	43.78 ^c ±0.74	4.22 ^{de} ±0.03	4.19 ^{de} ±0.02	4.21 ^{de} ±0.01
T ₆ (GA ₃) 50ppm	50.42 ^{ab} ±0.41	50.45 ^{ab} ±0.98	50.44 ^{ab} ±0.67	5.39 ^{ab} ±0.40	5.88 ^a ±0.10	5.64 ^a ±0.25
T ₇ (NAA) 50ppm	49.01 ^{ab} ±2.41	49.06 ^{bc} ±2.35	49.04 ^b ±2.38	5.16 ^b ±0.56	5.12 ^{ab} ±0.60	5.14 ^{ab} ±0.58
CD (<i>p</i> <0.05)	2.38	3.27	2.36	0.53	0.47	0.49

Each value in the treatment is a means of three replicates ± SD. Means followed by the same letter (s), in each row, are not significantly different (ANOVA, LSD test *p*<0.05).

an outstanding decline of SPAD value and biosynthesis, consequently it may be subject to the growth and yield of plants (43, 44). Moreover, osmo-priming resulted in enhancement of photosynthetic processes in cucumber seedlings, such as those involving stomatal conductance, transpiration rate and photosynthesis rate when the seeds were primed (45). whereas the same finding is also comparable in the okra study in the sense that indubitably, this attribute also rises in photosynthetic activities to increased chlorophyll index as well in this okra study. However (46) depicted the same result that when the seeds were primed that has increased mucilage content it lies between 3.40 to 5.93 %. Besides that, okra mucilage has the capacity to hold more zinc, calcium and minerals which are also present in entire fruit of okra and okra mucilage on the whole are natural polysaccharides, which contain L-rhamnose and monosaccharide D-galactose, galacturonic acid combined with proteins and minerals. Polysaccharides that are available in mucilage can provide the biological significance like immunomodulators and anti-inflammatories that can be obtained by polysaccharides (47, 48).

Iodine content

The pooled data observed that several priming sources significantly affected the iodine content of okra (*p*<0.05). Table 10 shows that treatment T₂ recorded the highest iodine content in okra at 13.14 mg kg⁻¹, which was statistically at par with treatment T₆ that recorded 13.08 mg kg⁻¹. The minimum

iodine content of okra (6.71 mg kg⁻¹) was found in T₀ treatment. This result is supported by (49, 50).

Conclusion

The present research advises that Osmo-priming is a viable method of improving germination in okra. Osmo-priming using 5 % polyethylene glycol (PEG) as seed priming can be utilized to increase the percentage germination (%), a number of days to germinate, shoot length (cm), root length (cm), length of seedling (cm) and dry weight of seedling (mg) vigour index (i), seedling vigour index (ii), root-shoot ratio, pods plant⁻¹, individual pod weight (g), pod yield plot⁻¹ (kg), yield plant⁻¹ (g), pod yield (t ha⁻¹). It was also seen that osmo-priming seeds have the best degree of tolerance to biotic and abiotic factors, probably because of the enhanced functioning of antioxidant enzymes. Hence, it is possible to recommend osmo priming to farmers to be used in the future.

Acknowledgements

Authors are highly thankful for the Lovely Professional University for funding to conduct the research and all the authors are showing no conflict interest.

Table 10. Effect of different seed priming treatments on okra's iodine content (mg kg⁻¹)

Treatments	Iodine content (mg kg ⁻¹)		
	2019	2020	Pooled
T ₀ (un-primed)	6.75 ^e ±0.95	6.69 ^e ±1.07	6.71 ^f ±1.00
T ₁ (hydropriming)	9.64 ^{cd} ±0.96	9.75 ^{cd} ±1.12	9.69 ^d ±1.04
T ₂ (PEG 5 %)	13.44 ^a ±1.11	12.84 ^a ±0.91	13.14 ^a ±0.91
T ₃ (KCL) 3 %	10.31 ^{bc} ±0.95	10.41 ^{bc} ±0.84	10.36 ^{cd} ±0.89
T ₄ (KNO ₃) 5 %	11.01 ^b ±1.43	11.42 ^b ±1.43	11.22 ^b ±1.43
T ₅ (MgNO ₃) ₂ 1.5 %	8.76 ^d ±1.07	8.79 ^d ±1.03	8.77 ^e ±1.05
T ₆ (GA ₃) 50ppm	12.82 ^a ±1.17	13.354 ^a ±1.06	13.08 ^a ±0.93
T ₇ (NAA) 50ppm	10.47 ^{bc} ±0.90	10.67 ^{bc} ±1.16	10.57 ^{bc} ±1.03
CD (<i>p</i> <0.05)	0.90	1.03	0.81

Each value in the treatment is a means of three replicates ± SD. Means followed by the same letter (s), in each row, are not significantly different (ANOVA, LSD test *p*<0.05).

Authors' contributions

NK did field work and wrote the paper. DM carried out the analysis and AS did the corrections.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

References

- Dhakal S, Hassan J, Rajib MM, Ghosh TK, Gomasta J, Biswas MS, et al. Seed priming and GA3 field application enhanced growth, yield and postharvest quality of okra. *Trends Hortic.* 2023;6:3578. <https://doi.org/10.24294/th.v6i2.3578>
- Pandey A, Nivedhitha S, Sagar V, Malav PK, Sharma S, Ahlawat SP. Notes on diversity distribution and systematics study of *Abelmoschus tuberculatus* Pal & Har B. Singh: a close wild relative of okra from India. *Indian J Plant Genet Resour.* 2020;33(1):77-84. <https://doi.org/10.5958/0976-1926.2020.00011.x>
- Lamichhane A, Mamata KC, Shrestha M. Effect of seed priming on germination of okra (*Abelmoschus esculentus* var. Arka Anamika). *Malays J Sustain Agric.* 2021;5(2):111-4. <https://doi.org/10.26480/mjsa.02.2021.111.114>
- Nhb, D. A. C. National horticulture board. 2018.
- Islam MT. Phytochemical information and pharmacological activities of okra (*Abelmoschus esculentus*): a literature-based review. *Phytother Res.* 2019;33(1):72-80. <https://doi.org/10.1002/ptr.6212>
- Elkhalifa AE, Alshammari E, Adnan M, Alcantara JC, Awadelkareem AM, Eltoum NE, et al. Okra (*Abelmoschus esculentus*) as a potential dietary medicine with nutraceutical importance for sustainable health applications. *Molecules.* 2021;26(3):696. <https://doi.org/10.3390/molecules26030696>
- Anwar F, Qadir R, Ahmad N. Cold pressed okra (*Abelmoschus esculentus*) seed oil. In: *Cold pressed oils*. Academic Press; 2020. p. 309-14. <https://doi.org/10.1016/B978-0-12-818188-1.00027-X>
- Schaffleitner R, Lin CY, Lin YP, Wu TH, Hung CH, Phooi CL, et al. The world vegetable center okra (*Abelmoschus esculentus*) core collection as a source for flooding stress tolerance traits for breeding. *Agriculture.* 2021;11(2):165. <https://doi.org/10.3390/agriculture11020165>
- Romdhane MH, Chahdoura H, Barros L, Dias MI, Corrêa RC, Morales P, et al. Chemical composition, nutritional value and biological evaluation of Tunisian okra pods (*Abelmoschus esculentus* L. Moench). *Molecules.* 2020;25(20):4739. <https://doi.org/10.3390/molecules25204739>
- Olawuyi IF, Lee WY. Structural characterization, functional properties and antioxidant activities of polysaccharide extract obtained from okra leaves (*Abelmoschus esculentus*). *Food Chem.* 2021;354:129437. <https://doi.org/10.1016/j.foodchem.2021.129437>
- Gemedede HF, Haki GD, Beyene F, Rakshit SK, Woldegiorgis AZ. Indigenous Ethiopian okra (*Abelmoschus esculentus*) mucilage: a novel ingredient with functional and antioxidant properties. *Food Sci Nutr.* 2018;6(3):563-71. <https://doi.org/10.1002/fsn3.596>
- Sharma K, Gupta A, Kumar M, Singh MK, Malik S, Singh B, et al. Effect of integrated nutrient management and foliar spray of bioregulators on growth and yield of okra. *Int J Curr Microbiol Appl Sci.* 2020;9:344-54. <https://doi.org/10.20546/ijcmas.2020.912.044>
- Wu DT, Nie XR, Shen DD, Li HY, Zhao L, Zhang Q, et al. Phenolic compounds, antioxidant activities and inhibitory effects on digestive enzymes of different cultivars of okra (*Abelmoschus esculentus*). *Molecules.* 2020;25(6):1276. <https://doi.org/10.3390/molecules25061276>
- Mabuza M, Tana T. Effects of osmo-priming on germination, growth and green pod yield of okra [*Abelmoschus esculentus* (L.) Moench] at Luyengo, Middleveld of Eswatini. *World J Adv Res Rev.* 2021;11(1):29-38. <https://doi.org/10.30574/wjarr.2021.11.1.0248>
- Wang C, Yu YB, Chen TT, Wang ZW, Yan JK. Innovative preparation, physicochemical characteristics and functional properties of bioactive polysaccharides from fresh okra (*Abelmoschus esculentus* (L.) Moench). *Food Chem.* 2020;320:126647. <https://doi.org/10.1016/j.foodchem.2020.126647>
- Kaur N, Gandhi A, Mal D, Gharde SK. Impact of seed hydro priming duration on germination and seedling performance of okra (*Abelmoschus esculentus* L. var. Punjab Suhawani). *Ecol Environ Conserv.* 2022;29(2):718-25. <https://doi.org/10.53550/EEC.2023.v29i02.028>
- Tania SS, Rhaman MS, Hossain MM. Hydro-priming and halo-priming improve seed germination, yield and yield contributing characters of okra (*Abelmoschus esculentus* L.). *Trop Plant Res.* 2020;7(1):86-93. <https://doi.org/10.22271/tpr.2020.v7.i1.012>
- Marthandan V, Geetha R, Kumutha K, Renganathan VG, Karthikeyan A, Ramalingam J. Seed priming: a feasible strategy to enhance drought tolerance in crop plants. *Int J Mol Sci.* 2020;21(21):8258. <https://doi.org/10.3390/ijms21218258>
- Mukhtar NK, Shamsudin NN, Zain NM, Naher L, Rahman KA, Sidek N. Enhancing okra (*Abelmoschus esculentus*) growth performance through seed priming. In: *BIO Web of Conferences 2024* (Vol. 131, p. 05010). EDP Sciences. <https://doi.org/10.1051/bioconf/202413105010>
- Rao KG, Sulladmath UV. Changes in certain chemical constituents associated with maturation of okra (*Abelmoschus esculentus* (L.) Moench), pods. 1977;4(1) 37-42. 10.5555/19790370852
- Kaur H, Chawla N, Pathak M. Effect of different seed priming treatments and priming duration on biochemical parameters and agronomic characters of okra (*Abelmoschus esculentus* L.). *International Journal of Plant Physiology and Biochemistry.* 2015;7(1):1-1.10.5897/IJPPB2015.0224
- Balchhaudi A. Enhancement of okra (*Abelmoschus esculentus* L.) germination through seed priming techniques. *Indones J Agric Res.* 2023;6(2):137-46. <https://doi.org/10.32734/injar.v6i2.13660>
- Hussain S, Hussain S, Khaliq A, Ali S, Khan I. Physiological, biochemical and molecular aspects of seed priming. In: *Priming and pretreatment of seeds and seedlings: implication in plant stress tolerance and enhancing productivity in crop plants.* 2019. p. 43-62. https://doi.org/10.1007/978-981-13-8625-1_3
- Yuan-Yuan SU, Yong-Jian SU, Ming-Tian WA, Xu-Yi LI, Xiang GU, Rong HU, Jun MA. Effects of seed priming on germination and seedling growth under water stress in rice. *Acta Agronomica Sinica.* 2010 Nov 1;36(11):1931-40. [https://doi.org/10.1016/S1875-2780\(09\)60085-7](https://doi.org/10.1016/S1875-2780(09)60085-7)
- Thakur V, Mal D, Gharde SK, Durlabh DA. Effect of different seed priming treatments on seed germination and seedling vigor of okra (*Abelmoschus esculentus* L. var. Punjab Suhawani). *J Food Chem Nanotechnol.* 2023;9(S1):S271-6. <https://doi.org/10.17756/jfcn.2023-s1-035>
- Soga K, Gandhi A, Mal D, Gharde SK. Impact of seed hydropriming duration on growth and yield parameter of okra (*Abelmoschus esculentus* L. var. Punjab Suhawani). *J Food Chem Nanotechnol.* 2023;9(S1):S36-43. <https://doi.org/10.17756/jfcn.2023-S1-007>
- Khanal S, Khanal S, Koirala S, Katel S, Mandal HR. Seed priming's effectiveness in improving okra germination and seedling growth. *Asian J Adv Agric Res.* 2022;18(2):29-34. <https://doi.org/10.9734/AJAAR/2022/v18i230215>
- Adhikari A, Shrestha A. Effect of primed and un-primed seeds on germination, growth performance and yield in okra (*Abelmoschus*

- esculentus* (L.) Moench). Turk J Agric Food Sci Technol. 2020;8(8):1686-91. <https://doi.org/10.24925/turjaf.v8i8.1686-1691.3460>
29. Adhikari A, Shrestha A. Effect of primed and un-primed seeds on germination, growth performance and yield in okra (*Abelmoschus esculentus* (L.) Moench). Turk J Agric Food Sci Technol. 2020;8(8):1686-91. <https://doi.org/10.24925/turjaf.v8i8.1686-1691.3460>
 30. Tania SS, Rhaman MS, Hossain MM. Hydro-priming and halo-priming improve seed germination, yield and yield contributing characters of okra (*Abelmoschus esculentus* L.). Trop Plant Res. 2020;7(1):86-93. <https://doi.org/10.22271/tpr.2020.v7.i1.012>
 31. Bereded Sheferie M. Effect of seed priming methods on seed quality of okra (*Abelmoschus esculentus* (L.) Moench) genotypes. Adv Agric. 2023;2023:3951752. <https://doi.org/10.1155/2023/3951752>
 32. Lakmali KN, Seran TH. Impact of seed priming with king coconut water on growth and yield of okra (*Abelmoschus esculentus* L.). J Food Agric. 2022;15(2). <https://doi.org/10.4038/jfa.v15i2.5271>
 33. Zhang K, Zhang Y, Sun J, Meng J, Tao J. Deterioration of orthodox seeds during ageing: influencing factors, physiological alterations and the role of reactive oxygen species. Plant Physiol Biochem. 2021;158:475-85. <https://doi.org/10.1016/j.plaphy.2020.11.031>
 34. Tiwari TN, Prasad SR, Agrawal DK. Seed priming improves crop growth and yield performance of pigeonpea (*Cajanus cajan* L.). J Food Legumes. 2019;32(1):9-12. <https://doi.org/10.59797/jfl.v32i1.692>
 35. Devi LA, Selvam S. Efficacy of seed priming using potassium silicate on okra's (*Abelmoschus esculentus* var. Raadhika) germination performance. Biochem Cell Arch. 2022;22(2). <https://doi.org/10.51470/bca.2022.22.2.4321>
 36. Anilkumar L, Malarkodi K. Combined seed enhancement technique involving seed priming and coating for improvised anatomical potential and vigour of okra seeds (*Abelmoschus esculentus* L.). J Phytol. 2019;11:25-30. <https://doi.org/10.25081/jp.2019.v11.3857>
 37. Alam MM, Khan MR, Hossain MS, Sultana MS, Anjum MT. Effects of different priming treatments on the germination and growth of okra (*Abelmoschus esculentus* L.). J Trop Crop Sci. 2024;11(1). <https://doi.org/10.29244/jtcs.11.01.91-96>
 38. Namboodiri RV, Bastian D, Anitha P, Rashmi CR. Enhancing okra (*Abelmoschus esculentus* L. (Moench)) performance: a comparative study of priming across varieties. Int J Plant Soil Sci. 2024;36(12):280-91. <https://doi.org/10.9734/ijpss/2024/v36i125202>
 39. Bereded Sheferie M. Effect of seed priming methods on seed quality of okra (*Abelmoschus esculentus* (L.) Moench) genotypes. Adv Agric. 2023;2023:3951752. <https://doi.org/10.1155/2023/3951752>
 40. Kamboj A, Sangha MK, Devi V, Kumar P, Pathak M, Singla D. Enhancing germination and growth in wild okra genotypes through gibberellic acid priming. Sci Hortic. 2024;334:113332. <https://doi.org/10.1016/j.scienta.2024.113332>
 41. Rhaman MS, Rauf F, Tania SS, Karim MM, Sagar A, Robin AH, et al. Seed priming and exogenous application of salicylic acid enhance growth and productivity of okra (*Abelmoschus esculentus* L.) by regulating photosynthetic attributes. J Exp Biol Agric Sci. 2021;9(6):759-69. [https://doi.org/10.18006/2021.9\(6\).759.769](https://doi.org/10.18006/2021.9(6).759.769)
 42. Fomekong MK, Tetang EF, Temegne CN, Atabong PA, Ntsomboh GN, Mir BA, et al. Effect of halopriming on seed germination performance and early seedling establishment in okra (*Abelmoschus esculentus* (L.) Moench) under saline conditions. Plant Physiol Rep. 2025;30(1):146-59. <https://doi.org/10.1007/s40502-024-00818-z>
 43. Devkota P, Jagadish Joshi PR, Dhital PR. Efficacy of priming on germination and seedling growth of okra in field condition in Khajura, Nepal. 2025;5(1):35-42. <http://doi.org/10.26480/ppsc.01.2025.35.42>
 44. Adheena P, Namboodiri RV, Bastian D, Anitha P, Rashmi CR. Optimizing pre-sowing treatments for the enhanced growth, fruit yield and seed quality in *Abelmoschus esculentus* L. Moench. Int J Plant Soil Sci. 2024;36(2):53-62. <https://doi.org/10.9734/IJPSS/2024/v36i24363>
 45. Khaskheli GN, Talpur A, Jamali MF, Jamali AR, Lund MM, Khaskheli M. Growth and yield response of okra (*Abelmoschus esculentus* L.) to various levels of biostimulant. J Appl Res Plant Sci. 2024;5(2):160-9. <https://doi.org/10.38211/joarps.2024.05.247>
 46. Akpor O, Ajinde A, Ogunnusi T. Effects of priming duration and rhizosphere bacteria metabolite concentration on the germinability of cowpea, soybean, sesame and okra seeds. F1000Res. 2024;12:781. <https://doi.org/10.12688/f1000research.137322.5>
 47. Mahood HE, Sarropoulou V. Growth, yield and phenolic compounds in okra (*Abelmoschus esculentus*) with culture condition and germination of seeds elicited by gamma radiation. Int J Veg Sci. 2025;31(3):376-408. <https://doi.org/10.1080/19315260.2025.2475832>
 48. Zhu XM, Xu R, Wang H, Chen JY, Tu ZC. Structural properties, bioactivities and applications of polysaccharides from okra (*Abelmoschus esculentus* (L.) Moench): a review. J Agric Food Chem. 2020;68(48):14091-103. <https://doi.org/10.1021/acs.jafc.0c04475>
 49. Srivastava M, Kathayat K, Mashkey VK. Performance of okra genotypes for different quality parameters. J Food Chem Nanotechnol. 2023;9(S1):S285-90. <https://doi.org/10.17756/jfcn.2023-s1-037>
 50. Khan FA, Narayan S, Bhat SA, Ashraf S, Fayaz F, Aamir M. Physiological potential of seed germination and seedling vigour of okra and parsley as influenced by seed priming treatments. SKUAST J Res. 2023;25(1):34-42. <https://doi.org/10.5958/2349-297X.2023.00004.1>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.