



RESEARCH ARTICLE

The effect of potassium fertilizer and planting dates on the growth and production of maize (*Zea mays* L.)

Fadhil Hussein Mukhlif¹, H I H El-Hamdani^{1*}, Noor M A Fadama², Suad Shallal Shahatha¹ & Mohammed Othman Mousa¹

¹Center of Desert Studies, University of Anbar, Ramadi 31001, Iraq

²General Directorate of Education, Ministry of Education, Ramadi 31001, Anbar, Iraq

*Correspondence email - hajer.i.hashash@uoanbar.edu.iq

Received: 04 June 2025; Accepted: 15 September 2025; Available online: Version 1.0: 22 September 2025

Cite this article: Fadhil HM, El-Hamdani HIH, Noor FMA, Suad SS, Mohammed OM. The effect of potassium fertilizer and planting dates on the growth and production of maize (*Zea mays* L.). Plant Science Today (Early Access). <https://doi.org/10.14719/pst.9827>

Abstract

An agricultural field study was conducted in Ramadi City, located in the Western region of Iraq, during the fall season of 2023, with the objective of examining the effect of three planting dates. viz 25 March, 5 and 15 April, and three levels of potassium fertilizer, including 100, 150 and 200 kg K₂O per hectare, on maize, variety Buhuth 106. Variations in planting dates exerted a notable impact on traits, including plant height, leaf area, ear length, number of ears per plant, weight of 300 grain total grain yield. The planting date of 25 March gave the highest rate of the studied traits, reaching 202.00 cm, 3990 cm², 15.138 cm, 1.824, 73.82 g and 4.372 tons. ha⁻¹ respectively. Significant differences were observed between the levels of potassium fertilizer for the same traits above. the potassium fertilizer level of 100 kg K₂O. ha⁻¹ was associated with the highest average values observed across all of the measured traits, which amounted to 187.156 cm, 3490 cm², 13.413 cm, 1.752 cm, 71.15 g and 3.301 tons. ha⁻¹ respectively. The combined effect of the experimental variables also demonstrated a clear influence on grain yield, as planting on 25 March combined with the application of 100 kg K₂O per hectare resulted in the highest recorded grain yield of 5.210 tons.

Keywords: grain yield planting dates; growth characteristics; maize (*Zea mays* L.); potassium fertilization

Introduction

Maize (*Zea mays* L.) occupies the third position worldwide, following wheat and rice, in terms of cultivated area and total production (1). Its grains are used in the manufacture of pastries and in the composition of concentrated poultry and animal feed, in addition to being used as human food in poor countries due to its high content of proteins, carbohydrates, minerals, and oil, as well as some vitamins (1). Although this crop holds significant importance, its productivity per unit area remains low. In Iraq, the cultivated area of maize in 2017 reached approximately 57.2 thousand hectares, yielding a total production of 185.3 thousand tons. (2). The significant gap in the productivity rate necessitates conducting scientific and applied studies to address this issue. Research has shown that potassium fertilization contributes significantly to enhancing plant growth and productivity, as potassium is one of the major nutrients that plants require in large quantities-second only to nitrogen-as the average content of plant tissues reaches 1.5% and may rise to 8% of the dry weight, as seen in tobacco tissues (3). Potassium additionally It has an effective role in improving water use efficiency required to produce one gram of yield and in reducing overall water consumption (4). In addition, it enhances the plant's ability to tolerate drought conditions and increases its resistance to extreme cold, without negatively affecting growth, productivity, cell division,

resistance to lodging, or susceptibility to plant diseases. Pollen grains are greatly affected by high temperatures and lose their ability to pollinate within a short period at temperatures above 35 °C. Therefore, we see that the yield of maize in spring cultivation is less than that of autumn cultivation. It is known that spring cultivation begins in the second half of March, when the temperature ranges between 22-28 °C. Since the male organs mature within two months of cultivation, i.e., around mid-May, when the temperature has risen to between 35-40 °C. Therefore, we see that the pollination rate in spring cultivation is low and ranges between 50-70% approximately, depending on the air temperature. Accordingly, the yield is less than that of the autumn crop (5). Therefore, this study aimed to identify the optimal potassium fertilizer level and the most suitable planting date to achieve maximum vegetative development and grain yield.

Materials and Methods

An agricultural field experiment was carried out during the 2023 spring growing season in a farmer's field located in Ramadi City, the administrative center of Al-Anbar Province Iraq. Three levels of potassium fertilizer were used in the experiment: 100, 150 and 200 kg K₂O. ha⁻¹, all of which were added mixed as basal. The plants were planted on three planting dates: March 25, April 5 and April 15. The experimental land was plowed, smoothed and

leveled and then divided into experimental units with dimensions of 2x3 m. Each experimental unit was composed of four planting rows, with each row measuring 3 meters in length, the spacing between rows was 50 cm (6) and the distance between planting holes along each row was 25 cm. A spacing of 1.5 m was maintained between each experimental unit to ensure proper isolation and to prevent overlap or interference between treatments. Random soil samples were collected from the study site prior to planting, at a depth of 0-30 cm, in order to assess some of its physicochemical properties. Then, the phosphate fertilizer was incorporated into the soil in a single dose at a rate of 100 kg. ha⁻¹ in the form of triple superphosphate. The nitrogen fertilizer was uniformly added at a rate of 200 kg N. ha⁻¹ in the form of urea (46% N) in two batches, the first immediately after germination and the second at flowering. The experiment was conducted following a randomized complete block design (RCBD) to ensure the accurate evaluation of treatments under controlled field conditions. Immediately after planting, the experimental field was initially irrigated, and subsequent irrigation was carried out as needed throughout the growing period, periodically depending on the soil moisture and plant condition. The experimental land was tilled twice during the growing season, the corn stalk borer (*Sesamia cretica*) was managed through the application of granular diazinon insecticide (6).

Ten randomly selected plants were taken from each experimental unit to assess the following traits:

1. Plant height cm:
2. Leaf Area (cm²): Calculated using the following formula: (Leaf length below the upper ear leaf)² × 0.75 (7).
3. Ear length:
4. was recorded by counting ears per individual plant

Table 1. It contains the physical and chemical properties of the experimental soil during the 2023 agricultural season.

traits	Value
PH	7.91
Ece	2.38
Ds.m ⁻¹	
Ready Potassium PPM	132
Ready Phosphorus PPM	8.1
Soluble Nitrate PPM	19.01
Lime (%)	23.2
Gypsum (%)	8.7
Organic Matter (%)	1.04
Sand	93
g. kg ⁻¹ . Soil	
Clay g. kg ⁻¹ . Soil	386
Green g. kg ⁻¹ . Soil	515
Tissue	Silty clay mixture

Table 2. Effect of Planting Dates and Potassium Fertilizer Levels on Maize Plant Height

Sowing times	Potassium Fertilizer			Mean Sowing times
	100 K ₂ O kg. ha ⁻¹	150 K ₂ O kg. ha ⁻¹	200 K ₂ O kg. ha ⁻¹	
March 25	201.177	200.047	204.013	202.00
April 5	192.183	184.373	181.053	186.00
April 15	168.107	163.027	160.013	164.00
Mean Potassium Fertilizer	187.00	182.482	182.00	
L.S. D 0.05	potassium 0.213	Sowing Times 0.2138	Sowing Times × Potassium 0.3703	

5. Weight of 300 Grains (g): A total of 300 grains were counted and weighed, and the final weight was adjusted to a standard grain moisture level of 15.5% (8).
6. Kernel yield tons. ha⁻¹:

The data for the studied traits were statistically analyzed based on the experimental design using the Gen Stat software. The minimum significant difference (MSD) test was employed to determine statistically significant differences between means at the 5% probability level.

Results and Discussion

Plant Height

It is evident from Table 2 that; a significant increase was observed in shoot height values with increasing fertilizer levels of potassium addition. The application rate of 100 kg K₂O. ha⁻¹ gave the highest rate for this trait, reaching 187.00 cm. In comparison with the application rate of 200 kg K₂O. ha⁻¹, which achieved 182.00 cm. This increase in the trait may be due to the activation of factors growth This increase in plant height may be attributed to enhanced cell multiplication and extension, processes stimulated by the role of potassium in activating effective plant hormones, especially gibberellins and auxins (9). Regarding planting dates, the results of the same table showed that the difference in planting dates had a significant effect on the average of this trait, as planting on the date of March 25 Recorded the highest mean plant height, reaching 202.00 cm when compared with the dates of April 5 and April 15, which gave 186.00 cm and 164.00 cm, respectively. This could be due to a reduction in temperature rates on the first date and the increase in relative humidity, which led to stimulating the division of plant cells and the elongation of internodes (6). As for the interaction of the studied factors, the treatment (100 kg K₂O. ha⁻¹ in addition the date of March 25) was characterized by the highest value for the same trait, and perhaps the reason for this is the combined efforts of the individual factors with each other. Similar results were indicated by (9) and (10).

Leaf area

Table (3) clearly shows a significant increase in the average leaf area values, which is directly proportional to the amount of potassium fertilizer applied, as the concentration exceeded 100 kg K₂O. h⁻¹ showing the maximum value of 3490 cm² for the trait, in comparison with the high concentration that gave 3303 cm². This is attributed to the role of potassium in increasing the division of leaf cells and the production of gibberellins and auxins hormones and its work with these hormones in increasing the flexibility and elasticity of the cell walls, which encourages the withdrawal of water and the formation of high swelling pressure that contributes to the elongation of leaf cells and thus

increasing the leaf area (6). As for the planting dates, they were also significantly different from each other in terms of their effect on the rate of this trait. Planting on the first date yielded the greatest average leaf area, which reached 3990 cm², while a reduction in the average value of this trait was observed when the planting date was delayed to 2803 cm². This may be due to the effective role of low temperatures in enhancing the activity of enzymes that stimulate leaf cell division. The interaction between the individual factors exerted a significant influence on increasing leaf area, as the combination of the first fertilizer level with the early planting date resulted in the highest average value for this trait, reaching 4167 cm². These results are consistent with the findings of (11) and (12).

Ear length

Experimental results presented in Table (4) indicated that the study factors exerted a significant influence on the average ear length of maize plant. The potassium fertilizer levels differed in their impact on this trait, with the highest ear length reaching 13.413 cm when fertilized with 100 kg K₂O. h⁻¹, while the remaining levels of 150 and 200 kg K₂O. h⁻¹ gave 12.858 cm and 12.143 cm, respectively.

As for planting dates, the early planting date recorded the highest mean ear length of 15.138 cm, compared to the late planting date, which recorded 11.205 cm. The combined effect of the studied factors had a meaningful effect on the value of the trait as the highest average was 16.223 cm when fertilized with 100 kg K₂O. h⁻¹ and planted on 25/3. Both (13) and (14) reached similar results.

Number of ears per plant

As shown in Table (5) potassium fertilizer levels had a significant effect on the number of ears per plant, as the application of 100 kg K₂O. Ha⁻¹ resulted in the highest value for this trait, recording 1.752 ears per plant.

Compared to 200 kg K₂O. ha⁻¹, which gave 1.666 ears per plant¹. This may be attributed to the role of potassium in enhancing the soil's physical and chemical properties, which helps increase the penetration of roots into the soil, thus making the plant obtain a larger amount of its nutritional needs (15), thus

increasing the dry matter of the plant, which in turn leads to stimulating the growth of secondary ears and reaching the stages of growth of full ears (16). Also, planting dates differed significantly in their effect on the trait rate, as the date 25/3 recorded the highest mean value for the trait under investigation, reaching 1.824 ears per plant¹, when compared to the dates 5/4 and 15/4, which gave 1.686 ears plant¹ and 1.608 ears per plant¹, respectively. The superiority of the first planting date in the number of ears per plant may be attributed to the fact that early planting promotes the development of more ears per plant and late flowering, while the plants are still small due to the incompleteness of their natural physiological growth (17). Significant differences in the trait value were observed due to the interaction between the study factors, as the highest number of ears per individual plant was 1.910 ears per plant¹, which was recorded by plants planted at the first date and fertilized with the first level of potassium fertilizer. These results were consistent with what was stated in this table.

Grain Weight per 300 Seeds

According to the findings in Table (6), yellow corn plants exhibited significant variation in grain weight in response to different potassium fertilizer levels, as the highest average was 71.15 g achieved by the 100 kg K₂O. ha⁻¹ treatment, while the average value of the trait decreased with increasing the level of potassium fertilizer to reach the lowest 68.62 g when the plant was treated with 200 kg K₂O. ha⁻¹.

This may be attributed to the role of potassium in transferring carbon metabolism products from the source to the sink, with plants sown on 25 March recording the highest mean value for the grain weight per plant, reaching 73.82 g, while delaying the planting date led to a decrease in the average of the trait to 66.00 g.

The interactions between the study factors also followed a similar trend to that of the individual factors, as the first fertilizer level with the earliest planting date gave the highest recorded value for the studied trait (75.80 g). This finding aligns with the results reported by previous researchers (13-15).

Table 3. Impact of Planting Times and Potassium Fertilizer Levels on Leaf Area of Maize Plants

Sowing times	Potassium Fertilizer			Mean Sowing times
	100 K ₂ O kg. ha ⁻¹	150 K ₂ O kg. ha ⁻¹	200 K ₂ O kg. ha ⁻¹	
March 25	4167	3700	4103	3990
April 5	3400.	3200.	3100	3233
April 15	2904	2800	2705.	2803
Mean Potassium Fertilizer	3490	3233	3303	
L.S. D 0.05	potassium 67.3	Sowing Times 67.3	Sowing Times × Potassium 116.5	

Table 4. Impact of Planting Dates and Potassium Fertilizer Levels on Ear Length of Maize Plants

Sowing times	Potassium Fertilizer			Mean Sowing times
	100 K ₂ O kg. ha ⁻¹	150 K ₂ O kg. ha ⁻¹	200 K ₂ O kg. ha ⁻¹	
March 25	16.223	15.130	14.063	15.138
April 5	13.013	12.000	11.200	12.071
April 15	11.003	11.446	11.166	11.205
Mean Potassium Fertilizer	13.413	12.858	12.143	
L.S. D 0.05	potassium 0.040	Sowing Times 0.040	Sowing Times × Potassium 0.070	

Table 5. Impact of Planting Dates and potassium fertilizer levels on the Number of Ears per Maize Plant

Sowing times	Potassium Fertilizer			Mean Sowing times
	100 K ₂ O kg. ha ⁻¹	150 K ₂ O kg. ha ⁻¹	200 K ₂ O kg. ha ⁻¹	
March 25	1.910	1.803	1.7600	1.824
April 5	1.716	1.686	1.656	1.686
April 15	1.630	1.613	1.583	1.608
Mean Potassium Fertilizer	1.752	1.701	1.666	
L.S. D 0.05	potassium 0.005	Sowing Times 0.005	Sowing Times × Potassium 0.009	

Table 6. Impact of Planting Dates and Potassium fertilizer levels on the Grain Weight per 300 of maize

Sowing times	Potassium Fertilizer			Mean Sowing times
	100 K ₂ O kg. ha ⁻¹	150 K ₂ O kg. ha ⁻¹	200 K ₂ O kg. ha ⁻¹	
March 25	75.80	73.45	72.21	73.82
April 5	70.28	69.21	69.00	69.50
April 15	67.36	66.00	64.64	66.00
Mean Potassium Fertilizer	71.15	69.55	68.62	
L.S. D 0.05	Potassium 0.572	Sowing Times 0.572	Sowing Times × Potassium 0.990	

Grain yield

Table (7) indicates the significant impact of potassium fertilization levels on grain yield. The increase in yield was more evident under the application rate of 100 kg K₂O. ha⁻¹, which gave 3.301 tons. ha⁻¹.

While the trait rate decreased when plants were treated with 200 kg K₂O. ha⁻¹, which reached 2.296 tons. ha⁻¹. Potassium contributes significantly to enhancing cell division and elongation and it also facilitates the translocation of carbon metabolism products to the sites of leaf emergence, this was reflected in the expansion of the leaf area, which subsequently enhanced the supply of assimilates to the developing flowers, leading to an increase in grain weight and fullness, in addition to its contribution to increasing the number of ears per plant (6).

From the same table, it is noted that a significant variation exists among planting dates regarding the rate of the studied trait. Early planting led to achieving the maximum grain yield of 4.372 tons. ha⁻¹, while delaying the planting date contributed to a decrease in the grain yield to 1.482 tons. ha⁻¹. The reason is due to the favourable thermal conditions and elevated relative humidity during the flowering period, which positively influenced pollen viability and consequently enhanced the fertilization success, allowing for an extended grain filling period (17). The significance of the interaction in the trait rate shows the synergy of the two factors in increasing the grain yield rate, as the treatment 100 kg K₂O. ha⁻¹ in along with planting on 25 March resulted in the maximum grain yield of 5.210 tons. ha⁻¹ (18). The results were consistent with what was found by (19).

Conclusion

The results showed that planting dates and potassium fertilizer levels have two major effects on plant growth and yield. It was found that early planting allows the plant to benefit from the photoperiod, which contributes to the plant benefiting as much as possible from suitable climatic conditions, increasing flower setting and avoiding high temperatures that kill pollen grains. Also, low potassium fertilizer levels gave the best results. This indicates that low levels spare the plant the risk of toxicity and provide economic feasibility.

Acknowledgements

The authors would like to express their sincere appreciation to the Center of Desert Studies - Anbar University for providing support and facilities to complete this research.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

FHM conceived the study, designed the experiment and supervised the manuscript preparation. MOM was responsible for field management and plant data collection. HHH contributed to the assessment of plant physiological traits. NMAF assisted in agronomic evaluations and data analysis. SSS contributed to reviewing the manuscript and coordinating logistics. All authors read and approved the final manuscript.

Table 7. Impact of Planting Dates and Potassium fertilizer levels on Grain yield

Sowing times	Potassium Fertilizer			Mean Sowing times
	100 K ₂ O kg. ha ⁻¹	150 K ₂ O kg. ha ⁻¹	200 K ₂ O kg. ha ⁻¹	
March 25	5.210	4.203	3.703	4.372
April 5	2.896	2.706	2.103	2.568
April 15	1.796	1.566	1.083	1.482
Mean Potassium Fertilizer	3.301	2.825	2.296	
L.S. D 0.05	potassium 0.0135	Sowing Times 0.0135	Sowing Times × Potassium 0.023	

Compliance with ethical standards

Conflict of Interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

References

- Mukhlif FH, Ramadan ASA, Hammody DT, Mousa MO, Shahatha SS. Molecular assessment of genetic divergence among maize genotypes. *SABRAO J. Breed. Genet.* 2023;55(3):739–48. <http://doi.org/10.54910/sabrao2023.55.3.12>
- Elsahookie MM. Corn, production and improvement. Baghdad: Ministry of Higher Education and Scientific Research, University of Baghdad; 1990.
- Directorate of Agricultural Statistics. Production of cotton, yellow corn and potatoes. Baghdad: Central Statistical Organization, Ministry of Planning – Iraq; 2017.
- Adrian J. Potassium nutrition in North Great Plains: new insights and perspectives. Potash and Phosphate Institute; 2004.
- Marschner H. Mineral nutrition of higher plants. 2nd ed. London: Academic Press; 1995.
- Barzaq J. Planting yellow corn in the spring season [Internet]. Iraq-datepalms.net; 2012 [cited 2025 Jun 4]. Available from: <https://www.iraq-datepalms.net>
- AL-Dulami BHA, AL-Hadethi NDH. Response of maize to potassium fertilizer and boron leaves nutrition. *Anbar J Agric Sci.* 2015;13(1):213–25.
- Elsahookie MM. Shortcut method for estimating plant leaf area in maize. *Agron J Crop Sci.* 1985;154:157–60.
- Al-Kazaali HA, Bahtash FY. Response of corn grain traits to harvesting moisture. *Iraqi J Agric Sci.* 2017;48:12–17. <http://doi.org/10.36103/ijas.v48iSpecial.239>
- Farhan EA. Effect of foliar application of potassium on growth characters and NPK content of popcorn *Zea mays* L. var. everta. *Iraqi J Desert Stud.* 2020;10(2):20–32.
- Ngosong CA, EnowATDVV, TeningMNEOAS. Optimising potassium fertilizer rates for sustainable maize (*Zea mays* L.) production on the volcanic soils of Buea, Cameroon. *Fundam Appl Agric.* 2022;7(1):11–20. <http://doi.org/10.5455/faa.96971>
- Liagat W, Akmal M, Ali J. Sowing dates effect on production of high yielding maize varieties. *Sarhad J Agric.* 2018;34(1):102–13. <http://doi.org/10.17582/journal.sja/2018/34.1.102.113>
- Mukhlif FH, AlKaisy AMA, Hammody DT, Mousa MO, Shahatha SS. The effect of potassium fertilizer and planting dates on growth, yield and quality of wheat (*Triticum aestivum* L.). *Kirkuk Univ J Agric Sci.* 2023;14(3):1–10. <http://doi.org/10.58928/ku23.14301>
- He B, Ji ZSQ, Wei J, Ma W. Effect of different long-term potassium dosages on crop yield and potassium use efficiency in the maize-wheat rotation system. *Agronomy.* 2022;12:2565. <http://doi.org/10.3390/agronomy12102565>
- Mohammed KAS, Eid MSM, Usman ARA. Enhancing the agronomic performance of potassium fertilizer and potassium-bearing minerals in sandy loam soil by adding humic acids and mycorrhiza. *Assiut J Agric Sci.* 2024;55(1):236–54. <http://doi.org/10.21608/ajas.2023.243994.1303>
- Joy B, Yuniarti A, Priyanka, Hakim FK, Taufik I, Wibawa RJ, Wahyu GT. Growth and yield of maize by nitrogen and potassium inorganic fertilizers application to Fluventic Eutrodepts. *E3S Web Conf.* 2023;444:04026. <http://doi.org/10.1051/e3sconf/202344404026>
- Al-Hilfy IHH, Al-Temimi AHM. Response of some synthetic maize cultivars to mineral, organic and bio fertilizer. *Iraqi J Agric Sci.* 2017;48(7):2763–71.
- Regab KH, Jassim WM. The effect of sowing dates on grains yield and components for some crosses corn (*Zea mays* L.). *Tikrit J Agric Sci.* 2016;16(1):21–29.
- Zhang Q, Li G, Lu WLD. Interactive effects of nitrogen and potassium on grain yield and quality of waxy maize. *Plants.* 2022;11:2528. <http://doi.org/10.3390/plants11192528>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc. See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.