



RESEARCH ARTICLE

# Antibiotic sensitivity of phosphate-mobilizing rhizobacteria of wheat rhizosphere

Saidakhon I Zakiryeva<sup>1</sup>, Zair S Shakirov<sup>1</sup>, Khursheda M Khamidova<sup>1</sup>, Husniddin Kh Karimov<sup>1</sup>,  
Ogiljon U Jamalova<sup>2</sup>, Nilufar A Elova<sup>3</sup> & Bakhtiyor K Rakhmanov<sup>4\*</sup>

<sup>1</sup>Soil Microbiology and Biotechnology Laboratory, The Institute of Microbiology of the Academy of Sciences of the Republic of Uzbekistan, Tashkent 100 128, Uzbekistan

<sup>2</sup>Department of Biotechnology and Microbiology, National University of Uzbekistan, Tashkent 100 174, Uzbekistan

<sup>3</sup>Probiotics Microbiology and Biotechnology Laboratory, The Institute of Microbiology of the Academy of Sciences of the Republic of Uzbekistan, Tashkent 100 128, Uzbekistan

<sup>4</sup>Structural and Functional Genomics Laboratory, Center of Genomics and Bioinformatics of the Academy of Sciences of the Republic of Uzbekistan, Tashkent 111 215, Uzbekistan

\*Correspondence email - [bakhtiyor.rakhmanov@gmail.com](mailto:bakhtiyor.rakhmanov@gmail.com)

Received: 14 June 2025; Accepted: 29 September 2025; Available online: Version 1.0: 27 November 2025

**Cite this article:** Zakiryeva SI, Shakirov ZS, Khamidova KHM, Karimov HKH, Jamalova OU, Elova NA, Rakhmanov BK. Antibiotic sensitivity of phosphate-mobilizing rhizobacteria of wheat rhizosphere. *Plant Science Today* (Early Access). <https://doi.org/10.14719/pst.10048>

## Abstract

Antibiotic sensitivity is a serious problem, especially relevant when using bacterial biofertilizers. Many strains of rhizobacteria used as bacterial biofertilizers and biofungicides demonstrate resistance to several antibiotics or contain antibiotic resistance genes. In this study, we investigated the antibiotic sensitivity of 12 strains of phosphate-mobilizing rhizobacteria isolated from the rhizosphere of wheat grown on irrigated lands of the Sirdarya, Tashkent andijan and Kashkadarya regions of Uzbekistan in 2021, belonging to the genera *Enterobacter*, *Rahnella*, *Bacillus*, *Pantoea* and *Pseudomonas*, to seven antibiotics of different classes: erythromycin, streptomycin, gentamicin, chloramphenicol, amikacin, tetracycline and cephalixin. Most strains showed high sensitivity to aminoglycosides, chloramphenicol and tetracycline, while some strains showed resistance to erythromycin, streptomycin and cephalixin. Moderate resistance to streptomycin, tetracycline and cephalixin was noted, especially among strains of the genus *Enterobacter*. Strains of the genus *Rahnella* showed moderate resistance to streptomycin, gentamicin, tetracycline and cephalixin. Strains of the genus *Bacillus* showed moderate resistance to chloramphenicol, tetracycline and cephalixin; the diameter of the zones of growth inhibition was 15 mm to 17 mm. The strain *P. agglomerans* 19 showed moderate resistance only to tetracycline, while remaining sensitive to other antibiotics, but this result requires confirmation through molecular genetic studies in the future. The data revealed are important for assessing the safety of phosphate-mobilising rhizobacteria when used as bacterial fertilisers, taking into account their resistance, to minimise the risk of spreading antibiotic resistance genes (ARGs) in agroecosystems.

**Keywords:** agroecosystems; antibiotic resistance; microbial biofertilizers; phosphate-mobilising rhizobacteria; wheat

## Introduction

Plant growth-promoting bacteria (PGPB) have a number of beneficial properties that make them widely used in sustainable agriculture. The use of these microorganisms helps to reduce dependence on chemical fertilisers, increases soil fertility and improves overall plant health (1-3). An important concern is the presence of antibiotic resistance genes (ARGs) in many PGPB strains. Introducing multiresistant bioinoculants into agroecosystems may facilitate gene transfer to soil microbes and, ultimately, to animals and humans. Large-scale use of those strains could therefore contribute to the global antibiotic resistance crisis, as ARGs may persist in soils and even enter the food chain (4). In the context of bacterial biofertilizer applications, the presence of antibiotic resistance genes in PGPB represents a potential risk, as these genes may be innate or acquired through horizontal transfer (5). Careful monitoring of agricultural microbial strains is essential to assess their

safety and role in resistance spread. In PGPR, resistance has a dual role, serving both as a biosafety concern and as a marker for tracking strain viability in laboratory and field conditions, while also giving strains a competitive advantage in complex microbial communities (6, 7). The application of these strains as bioinoculants in soil systems may pose a risk of horizontal transfer of antibiotic resistance genes to coexisting microorganisms (3). This requires a balanced approach to the selection of bacterial agents for use in agriculture.

Humans may directly or indirectly introduce antibiotic resistance genes into plant-associated microbial communities. Risk assessment of PGPB strains carrying ARGs remains limited, despite their continued promotion as biofertilizers. The presence of ARGs may further enhance their survival in antibiotic-contaminated soils (8). The large-scale use of biofertilizers carrying ARGs may extend antibiotic resistance from agricultural soils to natural habitats, aquatic systems, animals and humans. Thus, while resistant strains

can support sustainable farming, they also raise serious concerns about the global spread of antimicrobial resistance and its threat to public health (4).

Although there is evidence for the presence of antibiotic resistance genes in some PGPB strains, the extent of the risk of their transmission to other soil microorganisms and their possible impact on human pathogenic bacteria are still poorly understood (4). Antibiotics create selection pressure by promoting the spread of resistant forms in the environment. This problem is of particular relevance due to the uncontrolled use of antibiotics in medicine and veterinary medicine, which, in turn, leads to an increase in the number of multidrug-resistant strains and a decrease in the effectiveness of antibiotic therapy, increasing the frequency of severe and fatal infections (9-11). Antibiotics enter the environment in various ways, the main ones being wastewater from medical institutions, household and animal waste and drug residues in food and feed (12, 13). The agricultural sector, where antibiotics are widely used as growth stimulants in farm animals, creates a particularly significant load on ecosystems. As a result, significant amounts of antibiotics enter the soil, forming selective pressure and promoting the spread of resistance genes in agrocenoses (14).

The large-scale use of PGPB strains requires careful evaluation of their biological and environmental traits, particularly antibiotic sensitivity. Assessing the antibiotic response of phosphate-mobilizing bacteria is therefore an essential step in determining their safety and effectiveness. This study examined the sensitivity of wheat-associated phosphate-mobilising rhizobacteria to several commonly used antibiotics.

## Materials and Methods

The work studied twelve strains of phosphate-mobilizing rhizobacteria isolated from the rhizosphere of wheat and identified on the basis of morphological and physiological-biochemical characteristics. The identification was additionally confirmed using MALDI-TOF mass spectrometry. The isolates belonged to the genera *Enterobacter*, *Rahnella*, *Bacillus*, *Pantoea* and *Pseudomonas* (15-17). *Rahnella* strains were further identified based on molecular genetic characteristics as described in our previous study. The 16S rRNA gene of *Rahnella aquatilis* strain UT4 was sequenced and compared with sequences available in the National Center for Biotechnology Information database. The analysis showed 99 % identity with several *Rahnella aquatilis* sequences, including accession numbers MN826597.1, MN826570.1, MT256284.1, CP036490.1 and CP034483.1 (17).

### Cultivation of rhizobacteria

Rhizobacterial strains were cultivated on a shaker at 220 rpm at a temperature of  $28 \pm 2$  °C for 24 hours under aerobic conditions in a peptone liquid nutrient medium with the addition of glucose and NaCl, pH  $7.0 \pm 0.1$ . (18).

### Study the sensitivity of rhizobacteria to antibiotics

Antibiotic sensitivity of rhizobacteria was evaluated by the disk-diffusion method according to the standard of Clinical and Laboratory Standards Institute (CLSI), 30th ed. Wayne, PA: Clinical and Laboratory Standards Institute; 2020 (19). The antibiotics used were erythromycin (Ery, 15 µg), streptomycin (Str, 10 µg), gentamicin (Gen, 10 µg), chloramphenicol (Chl, 30 µg), amikacin (Amk, 30 and 50 µg), tetracycline (Tet, 30 µg) and cephalexin (Cep, 75 µg). The

diameter of growth inhibition zones was measured in millimetres and interpreted as sensitive (19 mm-36 mm), moderately sensitive (14 mm-18 mm) or resistant (0 mm-13 mm). The diameters of growth inhibition zones around the disks were measured after 24 hr of incubation at  $28 \pm 2$  °C.

### Statistical analysis

All experiments were performed three times to ensure the reliability and reproducibility of the results. Statistical analysis was performed using a standard licensed ANOVA software package. Differences between groups were considered statistically significant at  $p \leq 0.001$ .

## Results and Discussion


The results of the study showed significant variability in the sensitivity of phosphate-mobilising rhizobacteria to the antibiotics studied. The data are presented in Table 1. Overall, 75 % of strains showed sensitivity to chloramphenicol and aminoglycosides (amikacin), indicating the potential for the use of these bacteria without the risk of spreading resistance to these classes of antibiotics. Moderate resistance to streptomycin, tetracycline and cephalexin was noted, especially among strains of the genus *Enterobacter*. Strain of *E. cloacae* 7 is resistant to erythromycin, tetracycline and cephalexin. Strains of the genus *Rahnella* showed moderate resistance to streptomycin, gentamicin, tetracycline and cephalexin. Strain of *R. aquatilis* UT4 showed resistance to cephalexin and the diameter of the growth inhibition zone was  $5.7 \pm 2.5$  mm, respectively. Strains of the genus *Bacillus* showed moderate resistance to chloramphenicol, tetracycline and cephalexin; the diameter of growth inhibition zones was 15 mm to 17 mm. Strain of *B. subtilis* 24 showed resistance to gentamicin. Strain of *P. agglomerans* 19 showed moderate resistance only to tetracycline, while they were sensitive to the other antibiotics, but this result requires confirmation through molecular genetic studies in the future. The diameter of growth inhibition zones was 19 mm-30 mm. Among all the strains studied, *Pseudomonas antarctica* 23 depicted the highest level of multiple resistance. This exhibited resistance to gentamicin, amikacin, tetracycline and cephalexin (Table 1 and Fig. 1).

Antibiotic-resistant bacteria in biofertilizers may reach surface and groundwater after soil application, transferring resistance genes to local microbial communities and contributing to the accumulation of ARGs in environmental reservoirs (20). Moreover, strains able to colonize plants as endophytes can potentially transmit resistance genes to consumers through plant-derived products (21). The results obtained are consistent with the literature data, according to which Gram-positive bacteria of the genus *Bacillus* usually show high sensitivity to most antibiotics, while Gram-negative representatives, especially *Pseudomonas*, are more resistant due to their multistage defense mechanisms, notably active release of antibiotics (22, 23).

Many strains of rhizobacteria used as PGPB inoculants and biocontrol agents, in particular representatives of the genera *Pseudomonas* and *Bacillus*, demonstrate resistance to several antibiotics and/or contain antibiotic resistance genes (24-26). Thus, research indicates that phosphate-mobilising bacteria were isolated and their resistance was tested against various antibiotics, including kanamycin, streptomycin, tetracycline, ampicillin, chloramphenicol and spectinomycin (13). As a result, 11 out of 13 strains were found to be resistant to at least one of these drugs (27). Similarly, research

**Table 1.** Sensitivity of phosphate-mobilising rhizobacteria to antibiotics inhibitory zone diameter (mm)

Shtamm	Ery 15	Str 10	Gen 10	Chl 30	Amk 30	Amk 50	Tet 30	Cep 75
<i>E. cloacae</i> 7	12.0 ± 1.5	17.3 ± 3.5	17.7 ± 2.5	18.0 ± 2.0	18.7 ± 2.5	18.3 ± 2.1	0.0 ± 0.0	12.3 ± 2.5
<i>E. cloacae</i> 12	30.3 ± 2.5	28.7 ± 4.6	29.7 ± 3.1	21.7 ± 3.1	31.0 ± 3.6	29.7 ± 3.1	16.7 ± 2.1	14.3 ± 2.5
<i>E. cloacae</i> 18	17.7 ± 2.5	17.0 ± 2.0	19.7 ± 3.1	25.3 ± 3.1	22.0 ± 2.0	20.0 ± 3.0	20.3 ± 2.5	17.7 ± 1.5
<i>R. aquatilis</i> UT4	35.0 ± 2.6	17.0 ± 2.0	17.0 ± 2.0	20.3 ± 2.5	20.0 ± 2.0	20.3 ± 2.5	15.7 ± 4.7	5.7 ± 2.5
<i>R. aquatilis</i> 14	17.0 ± 1.0	21.0 ± 2.0	21.3 ± 2.1	23.0 ± 2.6	28.0 ± 2.6	23.0 ± 3.6	18.0 ± 3.0	13.3 ± 1.5
<i>B. cereus</i> 22	27.3 ± 2.1	30.3 ± 3.1	30.3 ± 2.5	16.0 ± 1.5	32.0 ± 3.0	29.3 ± 4.0	19.0 ± 2.0	17.1 ± 2.1
<i>B. subtilis</i> 24	32.3 ± 2.1	31.0 ± 2.0	0.0 ± 0.0	32.7 ± 2.9	27.3 ± 2.1	27.0 ± 2.6	16.0 ± 2.0	20.0 ± 5.0
<i>B. megaterium</i> 29	32.3 ± 2.5	30.0 ± 3.6	32.7 ± 2.5	30.3 ± 2.5	30.0 ± 3.6	26.7 ± 8.1	15.0 ± 1.5	17.0 ± 2.0
<i>P. agglomerans</i> 19	19.0 ± 2.0	20.3 ± 2.5	24.0 ± 3.0	30.0 ± 2.5	24.0 ± 2.0	22.0 ± 3.0	14.0 ± 2.5	20.7 ± 3.5
<i>P. agglomerans</i> 20	22.7 ± 2.5	20.3 ± 4.0	20.7 ± 2.5	32.0 ± 3.0	26.0 ± 3.0	22.3 ± 3.2	22.0 ± 3.0	20.7 ± 2.1
<i>P. kilonensis</i> 30	36.0 ± 3.0	32.7 ± 2.5	32.3 ± 2.5	27.0 ± 2.0	33.0 ± 2.6	32.3 ± 3.5	17.0 ± 2.0	18.0 ± 3.0
<i>P. antarctica</i> 23	15.3 ± 2.5	18.0 ± 3.0	0.0 ± 0.0	27.7 ± 1.5	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0



**Fig. 1.** Antibiotic sensitivity of phosphate-mobilising rhizobacteria *Pantoaea agglomerans* 20. **A.** And *Bacillus megaterium* 29; **B.** 1 - erythromycin (Ery, 15 µg); 2 - streptomycin (Str, 10 µg); 3 - gentamicin (Gen, 10 µg); 4 - chloramphenicol (Chl, 30 µg); 5 - amikacin (Amk, 30 µg); 6 - amikacin (50 µg), 7 - tetracycline (Tet, 30 µg); 8 - cefalexin (Cep, 75 µg).

indicates that among 11 phosphate-mobilising bacteria isolated from quinoa rhizosphere, five strains exhibited multiple resistance to at least two antibiotics. It is noteworthy that even widely used biocontrol agents like *Bacillus subtilis*, officially registered in Europe for plant protection against phytophthorosis, may contain ARGs, which require special attention in the development and introduction of microbial preparations in agriculture (28).

*Pseudomonas aeruginosa*, frequently found in agricultural soils and belonging to the group of plant growth-promoting rhizobacteria, is a multidrug-resistant opportunistic pathogen. This

bacterium is capable of carrying antibiotic resistance genes located on both the chromosome and plasmids. The resistance of *P. aeruginosa* to various antibiotics is provided due to the enhanced expression of  $\beta$ -lactamases, the work of efflux systems, changes in the structure of the cell membrane and the ability to modify antibiotic molecules (in particular, aminoglycosides) by chemical means (29, 30). *Pseudomonas chlororaphis* GP72, isolated from the rhizosphere of green pepper, shows resistance to antibiotics, particularly chloramphenicol and bleomycin. The resistance was noticed in the presence of *emrB* and *rarD* genes encoding proteins

involved in efflux systems (31, 32). These data indicate the ability of even environmentally relevant rhizobacteria, such as *P. chlororaphis*, to possess multidrug resistance mechanisms, which calls for careful regulation of their use in agricultural practice.

Strains of *Bacillus cereus* isolated from sunflower rhizosphere also show resistance to a wide range of antibiotics such as hygromycin B, chloramphenicol, streptomycin, penicillin, fosfomycin, bacitracin, teicoplanin and vancomycin. The resistance of these strains is due to the presence of several antibiotic resistance genes like *bl2a*, *bacA*, *bcrA*, *fos* and *vanSA* (33-35). The commercial strain of *Pantoea agglomerans* P10c used as a biocontrol agent exhibited resistance to a wide range of antibiotics, namely chloramphenicol, streptomycin, polymyxin, nalidixic acid, penicillin, bacitracin, novobiocin, fosfomycin, bicyclomycin, kasugamycin, cloxacillin, spectinomycin and fluoroquinolones. Research revealed that there was the presence of multiple antibiotic resistance genes like *acrD*, *ant3IA*, *bacA*, *bl2BE*, *catB3*, *emrD*, *emrA*, *mdtC*, *mdtB*, *mdtA*, *ksgA*, *rarD*, *tolC* and *qnrB* in genomic analysis (36). In contrast to the commercial strain of *P. agglomerans* P10c, which has multiple resistance to a wide range of antibiotics and carries a significant number of antibiotic resistance genes, our isolated strain of *P. agglomerans* 19 showed a limited resistance profile. It was found to be moderately resistant only to tetracycline, maintaining sensitivity to the other antibiotics tested. However, this result required confirmation in our future molecular genetic studies. These differences indicate the potential environmental safety of our strain when used in agroecosystems, as the risk of ARG spread into the environment is significantly lower compared to industrial strains.

Strain of *Enterobacter cloacae* 13047 shows resistance to a wide range of antibiotics, including tetracyclines, chloramphenicol, aminoglycosides, beta-lactams, glycolcyclines, macrolides, cephalosporins, sisomicin, dibecacin, penicillin, tobramycin, gentamicin, netilmicin, acriflavine, carbapenems, fluoroquinolones, trimethoprim and sulfonamides. Resistance is conferred by the presence of multiple ARGs, namely *aac3IIA*, *aac6IB*, *acrA*, *ant2IA*, *bl2F*, *bl3IMP*, *bl1AMP*, *bl3VIM*, *dfrA17*, *catB3*, *catA2*, *qnrA*, *sul1*, *sul2* and *tetD* (37, 38). The broad spectrum of antibiotic resistance of this strain is of concern, especially given its presence in both natural, clinical and agrarian environments, indicating a high risk of ARG transmission between different ecosystems. Our strain of *E. cloacae* 7 was resistant to erythromycin, tetracycline and cephalixin and showed only moderate resistance to streptomycin. These findings highlight the importance of evaluating the safety and genetic traits of bioagents before their application in agroecosystems. Monitoring antibiotic sensitivity is essential to minimize the risk of resistance gene transfer and to ensure the safe development of rhizobacterial inoculants.

## Conclusion

This study underscores the importance of evaluating phosphate-mobilizing rhizobacteria not only for their plant growth-promoting traits but also for their antibiotic resistance profiles. The observed differences among strains emphasize their genetic diversity and the potential risk of resistance gene transfer within soil microbial communities. Strains demonstrating strong phosphate-mobilizing capacity alongside antibiotic sensitivity are promising candidates for the development of safer biofertilizers, whereas moderately resistant isolates, such as *Pantoea agglomerans* 19, require further

molecular genetic studies to verify the absence of ARGs. Moreover, the sequencing of *Rahnella aquatilis* strain UT4 contributes additional insights into the diversity of phosphate-mobilizing bacteria and will serve as a reference point in forthcoming comparative analyses. Future work will expand the screening to a broader set of strains and apply molecular approaches, including PCR and sequencing, to characterize resistance determinants more precisely and evaluate their biosafety for agricultural applications.

## Acknowledgements

This study was funded by the State Budget of the Republic of Uzbekistan with the support of the Academy of Sciences and the Ministry of Higher Education, Science and Innovation. The work was carried out within the framework of the state research program titled "Development of a new microbial technology that increases the assimilable forms of phosphorus in the soil". We thank the scientific group for their assistance in conducting the study.

## Authors' contributions

SIZ designed and carried out experiments, cultivated phosphate-mobilising bacteria and prepared the manuscript. KMK, HKK and NAE participated in the experiments and drafted the manuscript. ZSS coordinated the study. OUJ collected the literature and participated in drafting the manuscript. BKR performed the processing of data and revised the manuscript. All authors read and approved the final manuscript.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** None

## References

- Glick BR. Plant growth-promoting bacteria: Mechanisms and applications. Scientifica (Cairo). 2012;963401. <https://doi.org/10.6064/2012/963401>
- Souza RD, Ambrosini A, Passaglia LM. Plant growth-promoting bacteria as inoculants in agricultural soils. Genet Mol Biol. 2015;38:401-419. <https://doi.org/10.1590/S1415-475738420150053>
- Ramakrishna W, Yadav R, Li K. Plant growth promoting bacteria in agriculture: Two sides of a coin. Appl Soil Ecol. 2019;138:10-18. <https://doi.org/10.1016/j.apsoil.2019.02.019>
- Mahdi I, Fahsi N, Hijri M, Sobeh M. Antibiotic resistance in plant growth promoting bacteria: A comprehensive review and future perspectives to mitigate potential gene invasion risks. Front Microbiol. 2022;13:999988. <https://doi.org/10.3389/fmicb.2022.999988>
- Kang Y, Shen M, Xia D, Ye K, Zhao Q, Hu J. Caution of intensified spread of antibiotic resistance genes by inadvertent introduction of beneficial bacteria into soil. Acta Agric Scand Sect B Soil Plant Sci. 2017;67:576-82. <https://doi.org/10.1080/09064710.2017.1314548>
- Trivedi P, Pandey A, Palni LMS, Bag N, Tamang M. Colonization of rhizosphere of tea by growth promoting bacteria. Int J Tea Sci. 2005;4(1&2):19-25.
- Gamalero E, Glick BR. Mechanisms used by plant growth-promoting bacteria. In: Bacteria in Agrobiolgy: Plant Nutrient Management. Berlin Heidelberg: Springer; 2011. p. 17-46. [https://doi.org/10.1007/978-3-642-11111-1\\_2](https://doi.org/10.1007/978-3-642-11111-1_2)

- [doi.org/10.1007/978-3-642-21061-7\\_2](https://doi.org/10.1007/978-3-642-21061-7_2)
8. Cray JA, Bell AN, Bhaganna P, Mswaka AY, Timson DJ, Hallsworth JE. The biology of habitat dominance; Can microbes behave as weeds? *Microb Biotechnol.* 2013;6:453–92. <https://doi.org/10.1111/1751-7915.12027>
  9. Hawkey PM, Jones AM. The changing epidemiology of resistance. *J Antimicrob Chemother.* 2009;64:i3–i10. <https://doi.org/10.1093/jac/dkp256>
  10. Paul M, Shani V, Muchtar E, Kariv G, Robenshtok E, Leibovici L. Systematic review and meta-analysis of the efficacy of appropriate empiric antibiotic therapy for sepsis. *Antimicrob Agents Chemother.* 2010;54:4851–63. <https://doi.org/10.1128/AAC.00627-10>
  11. Riber L, Poulsen PH, Al-Soud WA, Skov Hansen LB, Bergmark L, Brejnrod A, et al. Exploring the immediate and long-term impact on bacterial communities in soil amended with animal and urban organic waste fertilizers using pyrosequencing and screening for horizontal transfer of antibiotic resistance. *FEMS Microbiol Ecol.* 2014;90:206–24. <https://doi.org/10.1111/1574-6941.12403>
  12. Keswani C, Prakash O, Bharti N, Vílchez JI, Sansinenea E, Lally RD, et al. Re-addressing the biosafety issues of plant growth promoting rhizobacteria. *Sci Total Environ.* 2019;690:841–52. <https://doi.org/10.1016/j.scitotenv.2019.07.046>
  13. Xiao R, Huang D, Du L, Song B, Yin L, Chen Y, et al. Antibiotic resistance in soil-plant systems: A review of the source, dissemination, influence factors and potential exposure risks. *Sci Total Environ.* 2023;869:161855. <https://doi.org/10.1016/j.scitotenv.2023.161855>
  14. Hung W. Prevalence, fate and co-selection of heavy metals and antibiotic resistance genes in urban and agricultural soils [dissertation]. Los Angeles (CA): University of California, Los Angeles; 2020. <https://escholarship.org/uc/item/5v4895hs>
  15. Zakiryeva SI, Shakirov ZS, Khamidova HM, Normuminov AA, Atajanova ShSh, Azatov FR. Search for phosphate-mobilizing bacteria in the soils of Uzbekistan. *Universum: Chem Biol.* 2021;9(87):5–11. <https://7universum.com/ru/nature/archive/item/12238>
  16. Zakiryeva SI. Study of rhizosphere bacteria ability to mobilize P<sub>2</sub>O<sub>5</sub> from hard-to-access soil phosphates. *Web Agric J Agric Biol Sci.* 2024;2(10):90–5. <https://webofjournals.com/index.php/8/article/view/1994>
  17. Shakirov ZS, Mamanazarova KS, Yakubov IT, Zakiryeva SI, Khamidova KM. Nitrogen-fixing, phosphate–potassium–mobilizing ability of *Rahnella* bacteria isolated from wheat roots. *Regul Mech Biosyst.* 2022;13(4):379–384. <https://medicine.dp.ua/index.php/med/article/view/838/851>
  18. Zvyagintsev DG. *Methods of soil microbiology and biochemistry.* Moscow; 1991. 350 p. <https://www.scirp.org/reference/referencespapers?referenceid=2564001>
  19. CLSI. *Performance standards for antimicrobial susceptibility testing.* 30th ed. Wayne (PA): Clinical and Laboratory Standards Institute; 2020. <https://clsi.org>
  20. Topp E, Monteiro SC, Beck A, Coelho BB, Boxall ABA, Duenk PW, et al. Runoff of pharmaceuticals and personal care products following application of biosolids to an agricultural field. *Sci Total Environ.* 2008;396:52–9. <https://doi.org/10.1016/j.scitotenv.2008.02.011>
  21. Scaccia N, Vaz-Moreira I, Manaia CM. The risk of transmitting antibiotic resistance through endophytic bacteria. *Trends Plant Sci.* 2021;26:1213–226. <https://doi.org/10.1016/j.tplants.2021.09.001>
  22. Singh RP, Jha PN. The PGPR *Stenotrophomonas maltophilia* SBP-9 augments resistance against biotic and abiotic stress in wheat plants. *Front Microbiol.* 2017;8:1945. <https://doi.org/10.3389/fmicb.2017.01945>
  23. Blair JM, Webber MA, Baylay AJ, Ogbolu DO, Piddock LJ. Molecular mechanisms of antibiotic resistance. *Nat Rev Microbiol.* 2015;13(1):42–51. <https://doi.org/10.1038/nrmicro3380>
  24. Steil L, Hoffmann T, Budde I, Völker U, Bremer E. Genome-wide transcriptional profiling analysis of adaptation of *Bacillus subtilis* to high salinity. *J Bacteriol.* 2003;185:6358–370. <https://doi.org/10.1128/JB.185.21.6358-6370.2003>
  25. Santoyo G, Orozco-Mosqueda MD, Govindappa M. Mechanisms of biocontrol and plant growth-promoting activity in soil bacterial species of *Bacillus* and *Pseudomonas*: A review. *Biocontrol Sci Technol.* 2012;22:855–72. <https://doi.org/10.1080/09583157.2012.694413>
  26. Wellington EMH, Boxall ABA, Cross P, Feil EJ, Gaze WH, Hawkey PM, et al. The role of the natural environment in the emergence of antibiotic resistance in Gram-negative bacteria. *Lancet Infect Dis.* 2013;13:155–65. [https://doi.org/10.1016/S1473-3099\(12\)70317-1](https://doi.org/10.1016/S1473-3099(12)70317-1)
  27. Fahsi N, Mahdi I, Mesfioui A, Biskri L, Allaoui A. Phosphate solubilizing rhizobacteria isolated from *Jujube ziziphus lotus* plant stimulate wheat germination rate and seedling growth. *Peer J.* 2021;9:e11583. <https://doi.org/10.7717/peerj.11583>
  28. Mahdi I, Fahsi N, Hafidi M, Benjelloun S, Allaoui A, Biskri L. Rhizospheric phosphate-solubilizing *Bacillus atrophaeus* GQJK17 S8 increases quinoa seedling growth, withstands heavy metals and mitigates salt stress. *Sustainability.* 2021;13:3307. <https://doi.org/10.3390/su13063307>
  29. Lister PD, Wolter DJ, Hanson ND. Antibacterial-resistant *Pseudomonas aeruginosa*: Clinical impact and complex regulation of chromosomally encoded resistance mechanisms. *Clin Microbiol Rev.* 2009;22(4):582–610. <https://doi.org/10.1128/CMR.00040-09>
  30. Poole K. *Pseudomonas aeruginosa*: Resistance to the max. *Front Microbiol.* 2011;2:65. <https://doi.org/10.3389/fmicb.2011.00065>
  31. Ma M, Wang C, Ding Y, Li L, Shen D, Jiang X, et al. Complete genome sequence of *Paenibacillus polymyxa* SC2, a strain of plant growth-promoting rhizobacterium with broad-spectrum antimicrobial activity. *J Bacteriol.* 2011;193(1):311–12. <https://doi.org/10.1128/JB.01234-10>
  32. Kang Y, Shen M, Wang H, Zhao Q. A possible mechanism of action of plant growth-promoting rhizobacteria (PGPR) strain *Bacillus pumilus* WP8 via regulation of soil bacterial community structure. *J Gen Appl Microbiol.* 2013;59(4):267–77. <https://doi.org/10.2323/jgam.59.267>
  33. Ivanova N, Sorokin A, Anderson I, Galleron N, Candelon B, Kapatral V, et al. Genome sequence of *Bacillus cereus* and comparative analysis with *Bacillus anthracis*. *Nature.* 2003;423(6935):87–91. <https://doi.org/10.1038/nature01582>
  34. Joo G-J, Kim Y-M, Lee I-J, Song K-S, Rhee I-K. Growth promotion of red pepper plug seedlings and the production of gibberellins by *Bacillus cereus*, *Bacillus macroides* and *Bacillus pumilus*. *Biotechnol Lett.* 2004;26(6):487–91. <https://doi.org/10.1023/B:BILE.0000019555.87121.34>
  35. Khan N, Zandi P, Ali S, Mehmood A, Shahid MA, Yang J. Impact of salicylic acid and PGPR on the drought tolerance and phytoremediation potential of *Helianthus annuus*. *Front Microbiol.* 2018;9:2507. <https://doi.org/10.3389/fmicb.2018.02507>
  36. Smits TH, Rezzonico F, Blom J, Goesmann A, Abelli A, Morelli RK, et al. Draft genome sequence of the commercial biocontrol strain *Pantoea agglomerans* P10c. *Genome Announc.* 2015;3(6):e01448–15. <https://doi.org/10.1128/genomeA.01448-15>
  37. English M, Coulson T, Horsman S, Patten C. Overexpression of *hns* in the plant growth-promoting bacterium *Enterobacter cloacae* UW5 increases root colonization. *J Appl Microbiol.* 2010;108(6):2180–190. <https://doi.org/10.1111/j.1365-2672.2009.04620.x>

38. Ren Y, Ren Y, Zhou Z, Guo X, Li Y, Feng L, et al. Complete genome sequence of *Enterobacter cloacae* subsp. *cloacae* type strain ATCC 13047. J Bacteriol. 2010;192(9):2463–464. <https://doi.org/10.1128/JB.00067-10>

#### Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonpublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonpublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonpublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

**Publisher information:** Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.