



RESEARCH ARTICLE

# Morpho-anatomical investigation and metabolic profiling of *Gastrochilus calceolaris* (Buch.-Ham. ex Sm.) D.Don: A critically endangered epiphytic orchid

Anukriti<sup>1</sup>, Nainsy Gill<sup>1</sup>, Kajal Choudhary<sup>1</sup>, Mukesh Verma<sup>1</sup>, Shelja<sup>1</sup>, Jitender Kumar<sup>1\*</sup>, Harsharan Singh<sup>1</sup> & Pankaj Bagga<sup>2</sup>

<sup>1</sup>Department of Plant Sciences, Central University of Himachal Pradesh, Kangra 176 206, Himachal Pradesh, India

<sup>2</sup>Department of Zoology, DAV College, Jalandhar 144 008, Chandigarh, India

\*Correspondence email - [jitenderkumarcuhp@gmail.com](mailto:jitenderkumarcuhp@gmail.com)

Received: 31 August 2025; Accepted: 21 November 2025; Available online: Version 1.0: 16 December 2025; Version 2.0: 01 January 2026

**Cite this article:** Anukriti, Nainsy G, Kajal C, Mukesh V, Shelja, Jitender K, Harsharan S, Pankaj B. Morpho-anatomical investigation and metabolic profiling of *Gastrochilus calceolaris* (Buch.-Ham. ex Sm.) D.Don: A critically endangered epiphytic orchid. Plant Science Today. 2026; 13(1): 1-10. <https://doi.org/10.14719/pst.11555>

## Abstract

*Gastrochilus calceolaris* (Buch.-Ham. ex Sm.) D.Don, commonly known as the shoe-shaped belly-lip orchid, is a rare monopodial epiphytic orchid of high medicinal value. The present study provides the first integrated morpho-anatomical and metabolic profiling of this critically endangered species. Morphological and anatomical observations revealed distinct xerophytic adaptations, including a thick cuticle, well-developed velamen and prominent vascular bundles, which enable drought tolerance. Preliminary phytochemical screening of the methanolic leaf extract confirmed the presence of diverse secondary metabolites, including alkaloids, flavonoids, tannins and phenols. Gas Chromatography-Mass Spectrometry (GC-MS) identified 43 phytoconstituents, of which major compounds such as 3-Heptyn-1-ol and 3-heptyn-1-ol are known for their antioxidant, antifungal and anti-inflammatory activities. These findings highlight the pharmacological and conservation significance of *G. calceolaris*, providing a baseline for future bioprospecting and *in vitro* conservation strategies.

**Keywords:** anatomy; *Gastrochilus calceolaris*; GC-MS; secondary metabolites

## Introduction

Medicinal plants have served as a vital source of therapeutic compounds since ancient times. Their secondary metabolites, such as flavonoids, saponins, alkaloids, terpenoids, steroids, etc., play key roles in protecting plants from environmental stress and are extensively utilized in modern drug development. Globally, these secondary metabolites form the basis of major pharmaceutical, cosmetic and nutraceutical products due to their diverse pharmacological activities. (1, 2).

The family Orchidaceae, one of the largest among the flowering plants, includes approximately 28000 species within 763 genera (3). Members of this family exhibit remarkable ecological and morphological diversity occurring as epiphytes, terrestrials, or lithophytes (4). The anatomical features correlate with ecological adaptations and can serve as an important marker for distinguishing and authenticating species within a genus (5). Epiphytic orchids in particular show xeromorphic adaptations such as thick cuticle, well-developed velamen and reduced stomata to withstand intermittent water availability (6-8).

*Gastrochilus calceolaris* (Buch.-Ham. ex Sm.) D.Don is a small, pendulous, monopodial epiphytic orchid renowned for its attractive yellowish-green flowers with purple specks. It commonly grows on the tree trunks of *Ficus glomerata* and *Mangifera indica*

across the Eastern Himalayas, extending through India, Nepal, Bhutan, China, Thailand, Vietnam and Malaysia (9, 10). Despite its horticultural and medicinal values, *G. calceolaris* faces severe threats from habitat destruction and overcollection.

Metabolic profiling is a widely used technique to identify and analyze various bioactive compounds present within plant species. Previously, Gas Chromatography-Mass Spectrometry (GC-MS) of *Dendrobium amoenum* and *Dendrobium anceps* has revealed a variety of phytoconstituents with antioxidant, antimicrobial and anticancer potential (11, 12). As there are no previous studies, to the best of our knowledge, investigating the bioactive compounds in *G. calceolaris*, the present study aims to explore its morpho-anatomical features and metabolic profile.

## Materials and Methods

### Collection of plant material

The plant samples were collected from Sakoh, Kunal Pathri near Dharamshala, District Kangra, Himachal Pradesh, India (Latitude 32.175036°, Longitude 76.310357°). The collection was carried out during the flowering season (March-April 2024). The collected plants were authenticated at the Herbarium, Department of Botany, Panjab University, Chandigarh and recorded under

accession number 22974. Fresh leaves and root tissues were used for morpho-anatomical and metabolic investigation. All experiments were performed in triplicate to ensure reproducibility.

### Morpho-anatomical investigation

#### Morphological studies

The morphological study of *G. calceolaris* was conducted through detailed field observations and examination of freshly collected specimens from their natural habitat. Morphological assessments were performed on vegetative and floral parts, including leaves, stems and flowers. Key parameters such as length, breadth, shape, size and color were recorded following standard orchid terminology (13). Measurements were taken in triplicate and representative photographs were captured to document diagnostic features.

#### Anatomical study

For anatomical investigations, healthy leaf and root samples of *G. calceolaris* were collected from naturally growing individuals. The samples were washed thoroughly to remove surface debris and sectioned manually using a sharp razor blade. Thin transverse sections (TS) were stained with safranin to enhance the visibility of cellular structures, mounted on glass slides using glycerine and observed under a binocular light microscope (Model CH20iLED, Olympus, Japan). Photomicrographs (40X) were captured to document and analyze the anatomical features qualitatively. 10-15 manually sectioned leaf and root segments were analyzed.

### Preparation of plant extract

The leaves and roots of *G. calceolaris* were first washed thoroughly to remove any dirt or debris. After washing, the plant material was weighed and left to dry naturally in the shade at room temperature to preserve its chemical constituents. Once completely dried, the leaves and roots were ground separately into a fine powder using a mechanical grinder.

To prepare the methanolic extract, 10 g of each powdered sample (leaves and roots) were placed in a Soxhlet apparatus and extracted with methanol at 60 °C for about 6 hr. This process allowed the active compounds to be thoroughly extracted from the plant material. The obtained extracts were then concentrated using a rotary evaporator to remove the solvent and yield a thick, crude extract. These extracts were stored in clean, airtight containers at 4 °C until further analysis.

### Qualitative test for the phytochemical analysis

Prepared plant extracts were investigated for several qualitative tests using standard procedures (Table 1) such as alkaloids, proteins, carbohydrates, flavonoids, phenolics, etc. Each test was conducted in triplicate and the presence of various secondary metabolites was indicated by characteristic colour changes or precipitate formation.

### Gas chromatography-mass spectroscopic analysis of *G. calceolaris*

To determine the existence of active constituents and the chemical composition of *G. calceolaris* extract were analyzed using gas chromatography and mass spectrometry (GC-MS). GC-MS analysis of the whole plant extract of *G. calceolaris* was characterized using the equipment Shimadzu QP 2010 Ultra GC-

**Table 1.** Tests for qualitative phytochemical analysis

| Phytochemical class | Name of the tests                               | References |
|---------------------|---|------------|
| Alkaloids           | Dragendorff's test                              | (2,14)     |
|                     | Wagner's test                                   | (2,14)     |
|                     | Picric acid test                                | (14)       |
|                     | Iodine test                                     | (14)       |
|                     | Benedict's test                                 | (14)       |
| Carbohydrates       | Fehling's test                                  | (14)       |
|                     | Resorcinol test                                 | (14)       |
|                     | Starch test                                     | (14)       |
|                     | Million test                                    | (14)       |
| Proteins            | Ninhydrin test                                  | (14)       |
|                     | Xanthoproteic test                              | (14)       |
|                     | Lead acetate test                               | (2,14)     |
| Flavonoids          | Ferric chloride test                            | (2,14)     |
|                     | Conc. H <sub>2</sub> SO <sub>4</sub> test       | (2,14)     |
|                     | Shinoda's test/ Mg-hydrochloride reduction test | (2,14)     |
|                     | Lead acetate test                               | (2,14)     |
| Phenolic            | Iodine test                                     | (2,14)     |
|                     | Ferric chloride test                            | (2,14)     |
| Tannins             | 10 %NaOH test                                   | (2,14)     |
| Phylobatannins      | HCL test  | (2,14)     |
| Saponins            | Foam Test                                       | (2,14)     |
| Steroids            | Liebermann-Burchard test                        | (2)        |
| Phytosterols        | Salkowski Test                                  | (2,14)     |
| Terpenoids          | Sulphuric acid test                             | (2,14)     |
| Quinones            | Sulphuric acid test                             | (2,14)     |
| Anthroquinones      | Alcoholic KOH test                              | (2,14)     |
| Coumarin            | Ammonium hydroxide test                         | (2,14)     |
| Fats and oils       | NaOH test                                       | (2,14)     |
| Lignin              | Spot/stain test                                 | (14)       |
|                     | Labat test                                      | (14)       |

MS. An injection volume of 1  $\mu$ L was employed through splitless mode at 280 °C injection temperature and 200 °C ion source temperature. Three replicates were used in the study. The oven was programmed to 70 °C for a duration of 5 min, after which the temperature was to rise by 5 °C min<sup>-1</sup> until it attained 310 °C for 10 min. The identification of constituents was done by the National Institute of Standards and Technology Library (NIST20) as well as comparison of their retention indices. The constituents were characterized after comparison with those accessible within the computer library (NIST) attached to the GC-MS instrument and the gathered results have been tabulated (Table 2). Peaks identified in the GC-MS chromatogram serve as an indicative of the presence of secondary metabolites.

## Results

### Morphology

*G. calceolaris* is an epiphytic species with a slender stem bearing 8-9 distichously arranged, fleshy leaves that are oblong-lanceolate to linear-lanceolate. Inflorescences are lateral, short and bear 4-10 crowded flowers, faintly fragrant and typically green with reddish-brown blotches. The labellum is distinctly divided into a cup shaped hypochile and a broader epichile with white-papillose surface and purplish-red blotches along the margins. The column is short and carries 2 ovoid-elliptic pollinia with a linear stipe and oblong-elliptic viscidium. The detailed morphological characters are mentioned in Table 3 and the digital diagrams in Fig. 1.

### Anatomy

The TS of leaf of the *G. calceolaris* showed the hexagonal-shaped epidermis covered with a thick layer of cuticle; mesophyll cells were filled with green pigment called chloroplast. Vascular bundles were packed within the bundle sheath layer and

**Table 2.** Phytoconstituents identified in methanolic leaves extract of *G. calceolaris*


| Peak No. | Compound name                          | Retention time | Peak area % | Molecular weight (g mol <sup>-1</sup> ) | Molecular formula                              | Chemical structure |
|----------|--|----------------|-------------|---|--|--------------------|
| 1.       | 3-Heptyn-1-ol                          | 10.518         | 31.07       | 112.17                                  | C <sub>7</sub> H <sub>12</sub> O               |                    |
| 2.       | 2-Buten-1-ol, 2-methyl                 | 14.327         | 5.94        | 86.13                                   | C <sub>5</sub> H <sub>10</sub> O               |                    |
| 3.       | 4-Vinylphenol                          | 15.608         | 0.49        | 120.15                                  | C <sub>8</sub> H <sub>8</sub> O                |                    |
| 4.       | 4-Oxepincarboxylic acid, 2,3,6,7-tetra | 17.677         | 7.63        | 170.21                                  | C <sub>9</sub> H <sub>14</sub> O <sub>3</sub>  |                    |
| 5.       | Phenol,4-(methoxymethyl)-              | 18.047         | 3.23        | 138.16                                  | C <sub>8</sub> H <sub>10</sub> O <sub>2</sub>  |                    |
| 6.       | 2-Methoxy-4-vinylphenol                | 18.304         | 1.24        | 150.170                                 | C <sub>9</sub> H <sub>10</sub> O <sub>2</sub>  |                    |
| 7.       | Benzenemethanol, 4-hydroxy-            | 19.683         | 0.32        | 124.14                                  | C <sub>7</sub> H <sub>8</sub> O <sub>2</sub>   |                    |
| 8.       | Heptanoic acid, 1-methylethyl ester    | 19.825         | 1.34        | 172.26                                  | C <sub>10</sub> H <sub>20</sub> O <sub>2</sub> |                    |
| 9.       | Megastigmatrienone                     | 25.600         | 0.16        | 190.280                                 | C <sub>13</sub> H <sub>18</sub> O              |                    |
| 10.      | Megastigmatrienone                     | 26.754         | 0.23        | 190.280                                 | C <sub>13</sub> H <sub>18</sub> O              |                    |
| 11.      | beta-(4-Hydroxy-3-methoxyphenyl)pr     | 29.483         | 0.82        | 196.20                                  | C <sub>10</sub> H <sub>12</sub> O <sub>4</sub> |                    |
| 12.      | Tetradecanoic acid                     | 29.940         | 0.36        | 228.370                                 | C <sub>14</sub> H <sub>28</sub> O <sub>2</sub> |                    |
| 13.      | 2,3-Bis(1-methylallyl)pyrrolidine      | 30.653         | 0.91        | 179.30                                  | C <sub>12</sub> H <sub>21</sub> N              |                    |
| 14.      | Octadecanal                            | 31.184         | 0.25        | 268.500                                 | C <sub>18</sub> H <sub>36</sub> O              |                    |


|     |   |        |      |          |  |  |
|-----|---|--------|------|----------|--|--|
| 15. | Pentadecanoic acid, methyl ester  | 31.327 | 0.13 | 256.420  | C <sub>16</sub> H <sub>32</sub> O <sub>2</sub> |  |
| 16. | Neophytadiene   | 31.590 | 1.42 | 278.500  | C <sub>20</sub> H <sub>38</sub>                |  |
| 17. | 2-Pentadecanone, 6,10,14-trimethyl  | 31.707 | 0.17 | 268.5    | C <sub>18</sub> H <sub>36</sub> O              |  |
| 18. | Pentadecanoic acid  | 32.140 | 0.76 | 242.400  | C <sub>15</sub> H <sub>30</sub> O <sub>2</sub> |  |
| 19. | Neophytadiene   | 32.520 | 0.57 | 278.500  | C <sub>20</sub> H <sub>38</sub>                |  |
| 20. | Hexadecanoic acid, methyl ester   | 33.492 | 1.80 | 270.500  | C <sub>17</sub> H <sub>34</sub> O <sub>2</sub> |  |
| 21. | Benzene propanoic acid, 3,5-bis(1,1-dimethylethyl)-4-hydroxy-, ethylester | 33.586 | 0.20 | 292.4131 | C <sub>18</sub> H <sub>28</sub> O <sub>3</sub> |  |
| 22. | n-Hexadecanoic acid   | 34.365 | 7.11 | 256.420  | C <sub>16</sub> H <sub>32</sub> O <sub>2</sub> |  |
| 23. | Hexadecanoic acid, 2-oxo-, methyl ester                                   | 34.874 | 0.21 | 284.4    | C <sub>17</sub> H <sub>32</sub> O <sub>3</sub> |  |
| 24. | Heptadecanoic acid, methyl ester  | 35.556 | 0.23 | 284.500  | C <sub>18</sub> H <sub>36</sub> O <sub>2</sub> |  |
| 25. | Heptadecanoic acid  | 36.304 | 0.80 | 270.500  | C <sub>17</sub> H <sub>34</sub> O <sub>2</sub> |  |
| 26. | 9,12-Octadecadienoic acid (Z,Z)-, methyl ester                            | 36.888 | 3.40 | 294.472  | C <sub>19</sub> H <sub>34</sub> O <sub>2</sub> |  |
| 27. | 11,14,17-Eicosatrienoic acid, methyl ester                                | 37.010 | 4.28 | 320.509  | C <sub>21</sub> H <sub>36</sub> O <sub>2</sub> |  |
| 28. | Phytol  | 37.215 | 1.46 | 296.500  | C <sub>20</sub> H <sub>40</sub> O              |  |
| 29. | Methyl stearate   | 37.530 | 0.27 | 298.500  | C <sub>19</sub> H <sub>38</sub> O <sub>2</sub> |  |
| 30. | 10E,12Z-Octadecadienoic acid  | 37.770 | 4.93 | 280.400  | C <sub>18</sub> H <sub>32</sub> O <sub>2</sub> |  |
| 31. | cis,cis,cis-7,10,13-Hexadecatrienal                                       | 37.895 | 6.80 | 234.38   | C <sub>16</sub> H <sub>26</sub> O              |  |

|     |                                     |        |      |         |  |  |
|-----|-------------------------------------|--------|------|---------|--|--|
| 32. | Octadecanoic acid                   | 38.281 | 0.98 | 284.500 | C <sub>18</sub> H <sub>36</sub> O <sub>2</sub> |  |
| 33. | Heptadecyl acetate                  | 40.555 | 0.25 | 298.500 | C <sub>19</sub> H <sub>38</sub> O <sub>2</sub> |  |
| 34. | Eicosanoic acid, methyl ester       | 41.225 | 0.06 | 326.600 | C <sub>21</sub> H <sub>42</sub> O <sub>2</sub> |  |
| 35. | Tetradecyl cyclohexane              | 41.502 | 0.32 | 280.500 | C <sub>20</sub> H <sub>40</sub>                |  |
| 36. | Tetramethylheptadecan-4-ol          | 41.615 | 0.14 | 324.500 | C <sub>21</sub> H <sub>40</sub> O <sub>2</sub> |  |
| 37. | 3-Allyl-6-methoxyphenol             | 50.489 | 1.16 | 164.20  | C <sub>10</sub> H <sub>12</sub> O <sub>2</sub> |  |
| 38. | Delta-Tocopherol                    | 50.957 | 0.42 | 402.700 | C <sub>27</sub> H <sub>46</sub> O <sub>2</sub> |  |
| 39. | (R)-6-Methoxy-2,8-dimethyl-2-(4R,8R | 52.200 | 2.50 | 416.679 | C <sub>28</sub> H <sub>48</sub> O <sub>2</sub> |  |
| 40. | DL-alpha-Tocopherol                 | 53.504 | 1.13 | 430.700 | C <sub>29</sub> H <sub>50</sub> O <sub>2</sub> |  |
| 41. | Stigmasterol                        | 55.547 | 0.66 | 412.7   | C <sub>29</sub> H <sub>48</sub> O              |  |
| 42. | gamma-Sitosterol                    | 56.626 | 2.56 | 414.700 | C <sub>29</sub> H <sub>50</sub> O              |  |
| 43. | Stigmast-4-en-3-one                 | 59.197 | 1.30 | 412.700 | C <sub>29</sub> H <sub>48</sub> O              |  |

**Table 3.** Morphological description of *G. calceolaris*

| Features          | Description  |
|-------------------|--|
| Habit             | Epiphytic; plant length ranges from 10-30 cm (root to stem)  |
| Stem              | Length: 12 cm; diameter: 0.4-1.5 cm; internode enclosed by sheathing leaf bases  |
| Leaves            | Number: 8-9; shape: oblong-lanceolate to narrowly oblong or linear-lanceolate; size: 6-30 × 0.8-3.0 cm; arrangement: distichous; texture: fleshy; structure: carinate or nearly flat; base: articulate; apex: acute to unevenly bifid    |
| Inflorescence     | Type: lateral; number: 1-4; length: 3.5-5.0 cm; shape: corymbose to umbellate; flowers per inflorescence: 4-10   |
| Floral bracts     | Shape: broadly ovate-triangular; size: 4-5×3-3.5 mm; apex: acute   |
| Flowers           | Size: 1.5-3.0 cm long × 1.3-2.0 cm wide; arrangement: crowded; fragrance: faint  |
| Coloration        | Sepals and petals: soft green with reddish-brown blotches (variable); hypochile: light greenish yellow (base) to yellowish-white (mouth), purplish-red spots; epichile: white with yellow subtriangular disc blotched with purplish-red  |
| Pedicel and ovary | Length: 1.2-2.8 cm   |
| Sepals            | Shape: obovate-oblong to obovate-lanceolate; Size: 5.5-12.0 × 3.5-7.0 mm; Apex: obtuse; arrangement: spreading; laterals: slightly narrower and shorter  |
| Petals            | Shape: obliquely oblong-spathulate to oblong-obovate or elliptic-obovate; size: 5.5-11.0 × 3-6 mm; apex: obtuse  |
| Labellum          | Attachment: to the lower half of the column  |
| Hypochile         | Shape: cup-shaped; size: 6-7 × 5-6 mm; features: may have a truncate upper margin and vertical front edges, or lack vertical edges   |
| Epichile          | Shape: suborbicular to subreniform or subovate; size: 3.2-4.5 × 6.0-8.5 mm; surface: white-papillose adaxially (except central yellow cushion); cushion minutely hairy or rough, with a basal cavity; margins: lacerate-fringed to erose |
| Column            | Length: 2.5-4.0 mm   |
| Anthers           | Size: 1.5-2.0 × 2.0-2.4 mm   |
| Pollinia          | Number: 2; shape: ovoid-elliptic; size 0.7-1.1 × 0.4-0.8 mm  |
| Stipe             | Shape: linear; length: 1.6-2.5 mm; texture: hyaline  |
| Viscidium         | Shape: oblong-elliptic; size: 0.5-0.8 × 0.4-0.6 mm   |

**Fig. 1.** A. Front view of inflorescence b. whole plant with young inflorescence; c. front view of flower; d. floral perigone showing: a. dorsal sepal, b. lateral sepal, c. petals; e. lip; f. pedicellate ovary, column and lip with mentum; g. pollinia; h. anther cap.



**Fig. 2.** Transverse section of *G. calceolaris* leaf- (A): shows mesophyll with chloroplasts in green color (c), mid rib vascular bundles (a), side vein vascular bundles (b); (B): shows vascular bundles xylem tissues (d) and phloem tissues (e); (C): shows the epidermis covered with thick layer of cuticle (f).

differentiated the phloem and xylem tissues (Fig. 2). The TS of the root constituted a special type of multilayer before the epidermis called velamen. The vascular bundles are radial in nature with well-developed pith in the center (Fig. 3).


#### Phytochemical analysis

The qualitative phytochemical analysis (Table 4) revealed that roots contained a higher accumulation of alkaloids compared to the leaves. In contrast, leaves exhibited a greater presence of tannins, steroids and phytosterols. Terpenoids, anthraquinone and coumarin were absent in both leaves and roots. Carbohydrates, proteins, flavonoids, phenolics, saponins, phytobatannin, quinone, fats and oils and lignin were present in both leaves and roots.


#### GC-MS analysis

The methanolic leaves extract of *G. calceolaris* had revealed the presence of 43 different phytoconstituents through GC-MS chromatogram identified by the National Institute of Standards and Technology Library (Fig. 4). A major percentage of

phytoconstituents (Table 4) is 3-heptyn-1-ol (31.07 %), 4-oxepincarboxylic acid, 2,3,6,7-tetra (7.63 %), n-hexadecanoic acid (7.11 %) and the least percentage of phytoconstituents is eicosanoic acid, methyl ester (0.06 %). Among these, 4-oxepincarboxylic acid, 2,3,6,7-tetra is reported to possess antihistamine activity (15). n-hexadecanoic acid exhibits several pharmacologically relevant activities such as anti-androgenic, antioxidant, hypcholesterolemic, haemolytic, nematicidal, pesticide and 5- $\alpha$ -reductase inhibitory properties (16). Additionally, minor constituents detected in the extract also contribute to significant activities such as anti-inflammatory properties of 11,14,17-eicosatrienoic acid methyl ester (17), hepatoprotective, anticancer, antiacne, antihistaminic and anticoronal activities of 9,12-octadecadienoic acid (Z,Z) methyl ester (18) and analgesic and antioxidant potential of 10E,12Z-octadecadienoic acid (19). The presence of cis-cis-cis-7,10,13-hexadecatrienal associated with neurological benefits such as in Alzheimer's disease, epilepsy, depression and stroke (20), further highlights the therapeutic



**Fig. 3.** Transverse section of *G. calceolaris* root - (A): shows cortex region (a); (B): shows endodermis (e), well developed pith (d), radial vascular bundles- phloem(h)and xylem is differentiated into metaxylem (g) and protoxylem (f); (C): shows a special layer called velamen (b) and epidermis (c).



**Fig. 4.** GC MS chromatogram of *G. calceolaris*.

**Table 4.** Qualitative test for phytochemicals of *G. calceolaris*

| Phytochemicals | Tests                                    | Roots | Leaves |
|----------------|--|-------|--------|
| Alkaloids      | Dragendorff` s test                      | +++   | ++     |
|                | Wagner` s test                           | +++   | ++     |
|                | Picric acid test                         | +++   | ++     |
|                | Iodine test                              | ++    | +      |
| Carbohydrates  | Benedict` s test                         | +++   | +++    |
|                | Fehling` s test                          | +++   | +++    |
|                | Resorcinol test                          | +++   | +++    |
|                | Starch test                              | +++   | +++    |
| Protein        | Million test                             | +++   | +++    |
|                | Ninhydrin test                           | +++   | +++    |
|                | Xanthoproteic test                       | +++   | +++    |
|                | Lead acetate test                        | +++   | +++    |
| Flavonoids     | Ferric chloride test                     | +++   | +++    |
|                | Conc. Sulfuric acid ( $H_2SO_4$ ) test   | +++   | ++     |
|                | Shinoda` s test                          | +++   | +++    |
|                | Lead acetate test                        | +++   | +++    |
| Phenolics      | Ferric chloride test                     | +++   | +++    |
|                | Iodine test                              | +++   | +++    |
| Saponins       | Foam test                                | +     | +      |
| Tannin         | 10 % sodium hydroxide (NaOH) test        | +     | ++     |
| Steroids       | Liebermann-Burchard test                 | ++    | +++    |
| Phytosterol    | Salkowski` s test                        | +     | ++     |
| Phylobatannin  | Hydrochloric acid (HCl) test             | +     | +      |
| Quinone        | Sulphuric acid test                      | ++    | ++     |
|                | Alcoholic potassium hydroxide (KOH) test | ++    | ++     |
| Terpenoid      | Sulphuric acid test                      | -     | -      |
| Anthraquinone  | Ammonium hydroxide test                  | -     | -      |
| Coumarin       | NaOH test                                | -     | -      |
| Fats & oils    | Spot/stain test                          | +++   | +++    |
| Lignin         | Labat` s test                            | +++   | ++     |

**Note:** The '+++' sign indicates high presence, '++' sign indicates moderate presence, '+' sign indicates low presence and '-' sign indicates the absence of the phytochemical.

potential of the species.

## Discussion

In the present study, *G. calceolaris* showed several morphological and anatomical characters that differentiate it from other plant species and exhibited similarities with the family. The epidermis contained an extra layer made up of dead cells forming velamen, which prevents water loss and facilitates water absorption. This is consistent with earlier descriptions of epiphytic orchids, which are

characterized by velamen-covered roots, succulent leaves with thick cell walls, cuticles and specialized water storage structures that help them cope with drought and canopy microclimates (21).

The methanolic extract of leaves and roots showed the positive result for 13 secondary metabolites (proteins, carbohydrates, flavonoids, phenolics, tannin, lignin, etc.) and absence of 3 secondary metabolites (terpenoids, anthraquinone and coumarin). The methanolic extract of roots showed a more positive result than the leaves. The methanolic leaves extract of *G. calceolaris* has revealed 43 phytoconstituents through the GC-MS analysis, many of which are associated with diverse biological activities, including anti-inflammatory, antifungal and anti-diabetic properties. Several of these compounds also exhibited antioxidant, anticancer, anti-bacterial, antifungal, anti-microbial, nematicidal, pesticidal, hypocholesterolemic, anti-arthritis effects hepatoprotective activities and anti-microbial properties. Some of the phytoconstituents are adhesive in nature and contain bioactive chemicals effective against pests and nematodes.

Comparable GC-MS-based phytochemical studies in orchids further support the medical relevance of *G. calceolaris*. A previous study identified the major bioactive compounds such as 3-methoxy-hexane-1,6-diol, 1-methylene-2b-hydroxymethyl-3, 3-dimethyl-4b-(3-methylbut-2-enyl)-cyclohexane and butyl 9,12,15-octadecatrienoate in *Rhynchostylis retusa* leaf extract by GC-MS (22). The phytochemical examination of *Dendrobium candidum* by GC-MS reveals 21 bioactive compounds in the chromatogram and the major compound was heneicosane, showing insecticidal activity against mosquitoes (23). Another study identified a wide range of bioactive compounds by GC-MS in the stem and leaf methanolic extract of *Dendrobium jenkinsii* (24). The main constituents of stem were hexadecanoic acid, methyl ester, dotriacontane, gamma-sitosterol, mandelic acid, 3,4-dimethoxy-, methyl ester, whereas in leaf, the most prevalent compounds were 9,12,15-octadecatrienoic acid, methyl ester, (Z, Z, Z)-; gamma-sitosterol, hexadecanoic acid, methyl ester and n-hexadecanoic acid. Eight phytochemical compounds, namely phytol, 2-butyne, 2-cyclopenten-1-one, n-hexadecanoic acid, 9,12-octadecadienoic acid (Z,Z), 9,12,15-octadecatrienoic acid (Z,Z,Z), octadecanoic acid and 1,4-benzenedicarboxylic acid, were identified through GC-MS in

*Cymbidium aloifolium* (25).

The presence of these compounds marked it as a highly medicinal plant and has potential for new drug manufacturing in the future. It also holds promise to produce novel drugs with the isolation of specific compounds. *G. calceolaris* contains numerous biologically active compounds. Therefore, it is advised to consider this plant significant in the field of phytopharmaceuticals. However, additional research is recommended to thoroughly study the plant to fully evaluate its biological activity, pharmacological properties, toxicity profile, ecological impact and effects on agricultural products.

## Conclusion

The present study presents the first comprehensive morpho-anatomical and metabolic profiling of *G. calceolaris*, a critically endangered epiphytic orchid of the Western Himalaya. The xeromorphic anatomical traits identified, such as a multi-layered velamen, thick cuticle and compact mesophyll, highlight the species' ecological adaptation to intermittent water availability and arboreal habitats. GC-MS analysis revealed a rich profile of 43 phytoconstituents, including major bioactive compounds such as n-hexadecanoic acid, phytol, stigmasterol and  $\gamma$ -sitosterol, which are known for their antioxidant, anti-inflammatory and antimicrobial activities. These findings underscore the significant pharmacological potential of *G. calceolaris* and provide foundational data for its future bioprospecting. Given its threatened status and declining natural populations, the study also emphasizes the urgent need for conservation and further research, including *in vitro* propagation, targeted metabolomics and biological activity validation.

## Acknowledgements

The authors are grateful to the DST-SERB for providing funding (Project No. EEQ/2022/0009777) and to the Central University of Himachal Pradesh for providing the necessary facilities for the ongoing research.

## Authors' contributions

A and NG performed the study. A, NG, KC, MV, S, HS and PB wrote the manuscript. JK edited the final version of the manuscript. All authors read and approved the final manuscript.

## Compliance with ethical standards

**Conflict of interest:** The authors do not have any conflicts of interest to declare.

**Ethical issues:** None

## References

- Mathai K. Nutrition in the adult years. In: Mahan LK, Escott-Stump S, editors. Krause's food, nutrition and diet therapy. 10th ed. 2000. p. 271-5. <https://doi.org/10.1054/clnu.2000.0105>
- Akter M, Huda MK, Hoque MM. Investigation of secondary metabolites of nine medicinally important orchids of Bangladesh. J Pharmacogn Phytochem. 2018;7(5):602-6.
- Christenhusz M. The number of known plant species in the world and its annual increase. Phytotaxa. 2016;261:201-27. <https://doi.org/10.11646/phytotaxa.261.3.1>
- De LC. Approaches for natural farming in orchids. Vigyan Varta. 2022;3:28-33.
- Rindyastuti R, Nurfadilah S, Rahadiantoro A, Hapsari L, Abywijaya I. Leaf anatomical characters of four epiphytic orchids of Sempu Island, East Java, Indonesia. Biodiversitas. 2018;19(5):1906-18. <https://doi.org/10.13057/biodiv/d190543>
- Arditti J. Fundamentals of orchid biology. New York: John Wiley & Sons; 1992.
- Sailo N, Rai D, De LC. Physiology of temperate and tropical orchids-an overview. Int J Sci Res. 2014;3(7).
- Guan ZJ, Zhang SB, Guan KY, Li SY, Hu H. Leaf anatomical structures of Paphiopedilum and Cypripedium and their adaptive significance. J Plant Res. 2011;124(2):289-98. <https://doi.org/10.1007/s10265-010-0372-z>
- Pathak P, Piri H, Vij SP, Mahant KC, Chauhan S. *In vitro* propagation and mass scale multiplication of a critically endangered epiphytic orchid, *Gastrochilus calceolaris*. Indian J Exp Biol. 2011;49(9):711.
- Tang F, Song Y, Liu Q. The chloroplast genome of an endangered orchid species, *Gastrochilus calceolaris* (Orchidaceae: Aeridinae). Mitochondrial DNA B. 2018;3(2):988-9. <https://doi.org/10.1080/23802359.2018.1507646>
- Paudel MR, Pant B. Cytotoxic activity of crude extracts of *Dendrobium amoenum* and detection of bioactive compounds by GC-MS. Bot Orient. 2017;11:38-42. <https://doi.org/10.3126/botor.v11i0.21030>
- Noorjahan S, Rahamtulla M, Khasim SM. Phytochemical profiling and GC-MS analysis of leaf extracts of *Dendrobium anceps* Sw. J Med Pharm Allied Sci. 2023;12(6):6230-40. <https://doi.org/10.55522/jmpas.V12I6.5876>
- Dressler RL. Phylogeny and classification of the orchid family. Cambridge: Cambridge University Press; 1993.
- Shaikh JR, Patil M. Qualitative tests for preliminary phytochemical screening: An overview. Int J Chem Stud. 2020;8(2):603-8. <https://doi.org/10.22271/chemi.2020.v8.i2.8834>
- Higashiura K, Ogino T, Furukawa K, Yamazaki Y. Oxepin derivative. Eur Pat. 2009;EP 2067777 A1.
- Padmashree MS, Roopa B, Ashwathanarayana R, Naika R. Antibacterial properties of *Ipomoea staphylina* Roem & Schult. plant extracts: preliminary phytochemical and GC-MS analysis. Trends Pharmacol Res. 2018;5(3):349-69. <https://doi.org/10.22271/tpr.2018.v5.i3.044>
- Chen SJ, Chuang LT, Chen SN. Incorporation of eicosatrienoic acid exerts mild anti-inflammatory properties in murine RAW264.7 cells. Inflammation. 2015;38(2):534-45. <https://doi.org/10.1007/s10753-014-9960-8>
- Katta J, Rampilla V, Khasim SM. A study on phytochemical and anticancer activities of epiphytic orchid *Aerides odorata* Lour. Eur J Med Plants. 2019;28:1-21. <https://doi.org/10.9734/ejmp/2019/v28i30135>
- Könczöl Á, Engel R, Szabó K, Hornok K, Tóth S, Béni Z, et al. Topical analgesic, anti-inflammatory and antioxidant properties of *Oxybaphus nyctagineus*: phytochemical characterization of active fractions. J Ethnopharmacol. 2014;155(1):776-84. <https://doi.org/10.1016/j.jep.2014.06.020>
- Liu F, Zhao Q, Liu S, Xu Y, Zhou D, Gao Y, et al. Revealing the pharmacological mechanism of *Acorus tatarinowii* in the treatment of ischemic stroke based on network pharmacology. Evid Based Complement Altern Med. 2020;2020:3236768. <https://doi.org/10.1155/2020/3236768>

21. Zhang S, Yang Y, Li J, Qin J, Zhang W, Huang W, Hu H. Physiological diversity of orchids. *Plant Divers.* 2018;40(4):196-208. <https://doi.org/10.1016/j.pld.2018.06.003>
22. Kumar A, Mahanty B, Goswami RC, Barooah PK, Choudhury B. *In vitro* antidiabetic and antioxidant activities and GC-MS analysis of *Rhynchosystylis retusa* and *Euphorbia nerifolia* leaf extracts. *3 Biotech.* 2021;11(7):315. <https://doi.org/10.1007/s13205-021-02869-7>
23. Majumder S, Ghosh A, Chakraborty S, Bhattacharya M. GC-MS analysis reveals *Dendrobium candidum* is a mosquito-attractant orchid with mosquitocidal compounds. *Int J Mosq Res.* 2020;7(6):9-12. <https://doi.org/10.22271/23487941.2020.v7.i6a.483>
24. Borah O, Das AK, Vijayan D. Chemical composition, antioxidant potential and GC-MS analysis of methanolic extracts of leaves and stems of *Dendrobium jenkinsii* Wall. ex Lindl. *Vegetos.* 2025;1-10. <https://doi.org/10.1007/s42535-025-01260-8>
25. Rampilla V, Khasim SM. GC-MS analysis of organic extracts of *Cymbidium aloifolium* (L.) Sw. leaves from Eastern Ghats of India. In: *Orchid Biology: Recent Trends & Challenges.* Singapore: Springer; 2020. p. 507-17. [https://doi.org/10.1007/978-981-32-9456-1\\_26](https://doi.org/10.1007/978-981-32-9456-1_26)

#### Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonpublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonpublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc  
See [https://horizonpublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

**Publisher information:** Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.