



RESEARCH ARTICLE

Screening of salt stress in the overexpressed type of *Arabidopsis thaliana* (L.) Heynh. for the identification of significant hub genes using a systems biology approach

Elham Amjad¹, Babak Sokouti^{1*}, Solmaz Asnaashari^{1*} & Siavoush Dastmalchi^{1,2*}

¹Biotechnology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

²School of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran

*Email: b.sokouti@gmail.com, asnaasharisolmaz@gmail.com, dastmalchi.s@tbzmed.ac.ir



ARTICLE HISTORY

Received: 07 June 2021

Accepted: 09 December 2021

Available online

Version 1.0 : 23 July 2022

Version 2.0 : 01 October 2022



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS etc. See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Amjad E, Sokouti B, Asnaashari S, Dastmalchi S. Screening of salt stress in the overexpressed type of *Arabidopsis thaliana* (L.) Heynh. for the identification of significant hub genes using a systems biology approach. *Plant Science Today*. 2022;9(4):773–784. <https://doi.org/10.14719/pst.1301>

Abstract

Worldwide, it is known that abiotic and biotic stresses can affect the production of crops by a declining trend. To control the situation, SnRK2 (a sub-family 2 of SNF1-related protein kinase) overexpression levels can induce salt tolerance. This study used a dataset for 2 types of *Arabidopsis thaliana* including the wild and PtSnRK2.7 overexpressed in mock and salt conditions to compare and identify the salt stress-responsive genes. A computational systems biology approach was employed to identify the differentially expressed genes and determine their mechanisms in terms of molecular functionalities, cellular components, KEGG enrichment pathways and plant ontology analyses. The results indicate that the 15 genes identified for PtSnRK2.7 overexpressed type in mock against salt conditions were upregulated (AT1G19180 and AT2G23150 were downregulated) and related to various environmental stresses. Furthermore, 8 out of 15 identified genes were downregulated for the wild type exposed to salt stress and the rest were upregulated. And, the only upregulated gene found differentially expressed between wild and overexpressed types in salt stress conditions was AT4G15110. In contrast, the other two AT1G15010 and AT4G19430 were downregulated and involved in transient stress and inactivation of chloroplast, respectively. Taken together, it has been shown that *A. thaliana* PtSnRK2.7 overexpressed type can resist salt stress. Finally, more experimental studies and computational systems biology methodologies are needed to reveal and confirm the responsive gene for salt stress in *A. thaliana*.

Keywords

Systems biology, *Arabidopsis thaliana*, differentially expressed genes, function, plant ontology, gene ontology

Introduction

Globally, the agricultural products suffer from stressful environmental conditions such as high temperature, high soil salinity and drought, which hamper a lot of accounting for large cultivable areas in many countries like India, the US and China (1, 2). Among various types of abiotic stressors limiting the productivity of the crop, salt stress is considered plant toxicity that interferes significantly with cellular and physiological processes such as photosynthesis, chlorophyll contents, chlorosis and necrosis due to ion toxicity (3-5). So, the management of modern agriculture in the presence of salt stress, which increases annually due to the prolonged shortages in the wa-

ter supply, is essential. Hence, several biotechnological techniques (e.g., *Arbuscular mycorrhizal* fungi (AMF)) are needed to alleviate the adverse effects of salinity (6). The plants are structured as a nonlinear model such that any input can impose a complicated output. This property of plants' nature makes them be reprogrammed in physiological, molecular, and developmental processes to optimize salt stress response (7). One of the well-known plant model systems is *Arabidopsis thaliana* from the Brassicaceae family, with 120 Mbp of DNA organized into five chromosomes. That has specifically become the focus of molecular genome-based analysis research for over forty years to provide the most basic information for all eukaryotes (8, 9). Several studies were performed on inducing various stressors in *A. thaliana* to investigate the changes in the microarray gene expression levels (10-12).

SnRKs (sucrose non-fermenting related protein kinase) with 3 subfamilies of SnRK1, SnRK2 and SnRK3 are involved in associating abiotic stresses and their corresponding metabolic responses (13). Among these, members of SnRK2 (i.e., subfamily 2 of SNF1-related protein kinase) family, plant-specific serine/threonine kinases, have taken special interests of researchers while studying the plant responses to various stressors (e.g., abiotic stresses and abscisic acid-dependent signaling pathway) (13). Several genome-wide analysis studies were performed on the SnRK2 family using some plant models such as *Gossypium hirsutum* and *Populus trichocarpa* to appraise the overexpression levels of SnRK2 members *Arabidopsis thaliana* dealing with environmental stresses (14, 15).

Moreover, the rationale of the current systems biology research to study the overexpressed *Arabidopsis thaliana* with PtSnRK2.7 overexpression is to investigate the differentially expressed genes (DEGs) in the presence and absence of the overexpression in salt stress situations. However, the control of the position will bring the policy-makers many advantages in terms of commercial or economic purposes.

Materials and Methods

The overall flowchart of the current study, starting from the dataset selection, pre-processing steps, and post-processing to analysis procedures, were depicted in Fig. 1.

Data Source

The GSE79997 dataset was downloaded from the NCBI GEO database (i.e., <https://www.ncbi.nlm.nih.gov/geo/>) with twelve samples consisting of 4 groups of wild and SnRK2.7 overexpressed types in mock and salt treatment status, which was the only available GEO dataset. The platform of this microarray dataset was GPL198 [ATH1-121501] Affymetrix *Arabidopsis* ATH1 genome array which includes twelve CEL files (GSM2109727, GSM2109728, GSM2109729, GSM2109730, GSM2109731, GSM2109732, GSM2109733, GSM2109734, GSM2109735, GSM2109736, GSM2109737, GSM2109738). The four groups were denoted by *wild-mock-salt*, *wild-over-mock*, *wild-over-salt*, *over-mock-salt*. Con-




Fig. 1. Flowchart of step by step approaches to achieve the final validated genes in terms of *Arabidopsis* salt stress.

sidering the group naming, it should be noted that the italic format of the font represented the mock and salt stress conditions and the wild and SnRK2.7 overexpressed types used the normal font.

DEGs determination through pre-processing and statistical tests

The BRB-ArrayTools version 4.6.0, developed by Dr. Richard Simon and the BRB-ArrayTools Development Team, was used as a free and non-commercial integrated excel toolbox for microarray analyses. The Affymetrix *Arabidopsis* ATH1 genome array annotation data (chip ath1121501) "ath1121501.db"(16) developed for the R programming environment was also utilized for the corresponding gene symbols retrieval of probe ids. Moreover, for pre-processing procedures, the Microarray Suite version 5.0 (MAS 5.0) algorithm and pre-defined settings for enabling the spot filtering, quantile normalization and gene filtering were considered where the fold change parameter was set to less than 2. To identify the significant differentially expressed genes (DEGs), the "class comparison between groups" menu of BRB-Array Tools was selected by setting the univariate permutation tests and fold change thresholds as 10000 and 2. The additional outputs from the analyses include the illustrations of the volcano plot and box plot based on the microarray data.

Analyses of GO, PO and KEGG pathway and functional enrichment

To determine the molecular and cellular functional processes of the obtained significant gene lists as well as their enriched pathways, the online tool, PlantGSEA (Plant Gene Set Enrichment Analysis) website (i.e., <http://structuralbiology.cau.edu.cn/PlantGSEA>), was applied (17). This website took advantage of gene ontology (GO) annotation analysis, plant ontology (PO), and the Kyoto Encyclopedia of Genes and Genomes (KEGG) to cover the required assessment properties. Moreover, the results were tested and analyzed with hypergeometric statistical test and Bonferroni multi-test adjustment methods and a significant level of 0.05. The input queries consisted of two gene lists: (i) the list of probe ids retrieved from potential significant DEGs and (ii) customized background corresponding to anatomy type of TAIR (The *Arabidopsis* Information Resource) genes. However, the temporal type of TAIR gene list was not considered due to the full coverage of genes overlapped with anatomy type related to the salt stress (i.e., with "salt" keyword).

Construction of protein-protein interaction network and identification of significant genes

For further analysis of the identified DEGs in their intra-protein interactions, the STRING database (with 5090 organisms, over 24.6 million proteins and over 2000 million protein-protein interactions) plugin for Cytoscape 3.9.0 was used (updated on October 21, 2021). The confidence score cutoff default value was set to 0.4. After, the ClusterOne v.1.0 as one of the well-known graph clustering algorithms was used to generate overlapping clusters of PPI network (18) to determine the modules with p -value ≤ 0.05 where the genes with the highest degrees of connectivity were introduced as the potential biomarkers or the most involved gene in the expression of the differences between two defined groups (e.g., the wild types at mock and salt conditions).

Analysis of DEGs identified from this study and literature

This section includes the identified DEGs from BRB-array tools and those identified from the literature for further analysis and comparison (15). However, the significant DEGs from the literature were available for only 3 groups denoted by *wild-mock-salt*, *wild-over-mock*, *over-mock-salt*. And, the calculation for *wild-over-salt* was separately performed using the data obtained in the current investigation. For achieving a robust comparison procedure, the TAIR IDs were of interest as some of them might not have a corresponding official gene symbol. First, the target DEGs included in the groups mentioned above were used to construct the PPI network based on the STRING database using the Cytoscape 3.9.0. Then, the cytoHubba plugin was utilized to determine the most significant DEGs by performing eleven topological methodologies (19).

Moreover, Maximal Clique Centrality (MCC), Maximum Neighborhood Component (MNC), Degree, Edge Percolated Component (EPC) and EcCentricity were considered to obtain ten top-ranked genes for each ranking score distinctly. After overlapping the top 10 ranked list of genes,

5 ranking score algorithms were extracted for the 4 groups mentioned above. Finally, the shared genes between our study and the literature would be identified and were the target of interest for being analyzed for their functionality using the literature researches.

Results

The numbers of those genes that passed the filtering criteria after the microarray dataset import for *wild-mock-salt*, *wild-over-mock*, *wild-over-salt* and *over-mock-salt* groups are 2216, 86, 234 and 1573 respectively. Additionally, the outcomes of pre-processing procedure obtained from class comparison of BRB-ArrayTools for 4 groups mentioned above demonstrate the significant differentially expressed genes (DEGs) using the two-sample T-test. The numbers of DEGs for *wild-mock-salt*, *wild-over-mock*, *wild-over-salt* and *over-mock-salt* groups are 1288 (730 down-regulated genes and 558 upregulated genes), 33 (30 down-regulated genes and 3 upregulated genes), 3 (0 down-regulated genes and 3 upregulated genes) and 1525 (884 down-regulated genes and 641 upregulated genes) respectively. Fig. 2 shows the boxplot, a representation of gene expression levels and volcano plots (i.e., the fold change values) of the microarray data for four groups after pre-processing approach. The list of top upregulated and downregulated genes for 4 groups is shown in Supplementary Table 1.

The output of the plant GSEA online tool provides several gene ontology functional analyses and plant ontology assessments on the DEGs of four groups. The analyses results include gene ontology cellular components (GO-CC), gene ontology molecular functions (GO-MF), the KEGG pathway as well as the plant ontology (PO) assessments (Fig. 3).

According to *over-mock-salt*, several cellular components (e.g. cell, cell part), molecular functions (e.g. catalytic activity, binding), KEGG pathways (e.g. metabolic pathways, biosynthesis of phenylpropanoids) and plant ontology (e.g. fluorescence meristem, petiole) can be significantly enriched by DEGs based on the identified p -values. Moreover, the *wild-mock-salt* group demonstrates the DEGs enrichment by cellular components (e.g. cell, cell part), molecular functions (e.g. binding, catalytic activity), KEGG pathways (e.g. metabolic pathways, biosynthesis of plant hormones) and plant ontology (e.g. petiole, fluorescence meristem). Finally, the *wild-over-mock* set offers only the plant ontology assessment such as stamen and collective leaf structure.

The construction and visualization of protein-protein interaction networks for the 4 groups have resulted in PPI networks for *over-mock-salt* with 850 nodes and 4895 edges, for *wild-mock-salt* with 667 nodes and 3300 edges and *wild-over-mock* with 20 nodes and 23 edges. However, due to only one DEG for the *wild-over-salt* group, no PPI network has been determined. Supplementary Table 2 shows the list of the top 10 genes which have been ranked based on their interconnectivity degree considering the 4 constructed PPI networks. Among them, JAZ1




Fig. 2. Boxplot and volcano plots of four groups. (a) wild-mock-salt, (b) wild-over-mock, (c) over-mock-salt and (d) wild-over-salt.




Fig. 3. Gene ontology cellular components (GO-CC), gene ontology molecular functions (GO-MF), the KEGG pathway as well as the plant ontology (PO) assessments. (a), (b), (c) and (d) are related to over-mock-salt for CC, MF, KEGG and PO respectively. (e), (f), (g) and (h) are related to wild-mock-salt for CC, MF, KEGG and PO respectively. (i) is the PO analysis for wild-over-mock.




Fig. 4B. Wild-mock--salt (a)-(i).

lapped genes between current and earlier studies (15), including TAIR IDs, gene symbols, fold changes, up/down regulated status of genes, and the significant p values, are listed in Supplementary Table 3. And, the genes listed as the last group in Supplementary Table 3 are those only identified through the BRB-array tool as significant DEGs. However, the PPI network analysis is not applicable due to the low number of DEGs (with only three critical genes).

Discussion

The extensive need for food is directly linked with human life, which can be affected by various environmental risk

factors such as abiotic and biotic stresses (20). So, providing food production security for the whole world is of the top-most requirements that policymakers of the countries should be aware of and make thoughtful decisions for increasing the plants' stress tolerance. Because the plants are always immobile and affected mainly by stressful situations, they need more attention in manipulating mechanisms and functionalities through upregulating or down-regulating the gene expression levels to achieve optimal productions (21, 22). In the current study, salt stress as one of the substantial abiotic stresses in *Arabidopsis thaliana* was investigated for 2 types, including wild and PtSnRK2.7 over expression.




Fig. 4C. Wild-over-mock (a)-(d)




Fig. 5. The number of overlapping genes between current and Song et al. studies.

In the case of wild and PtSnRK2.7 overexpression types (7-day-old seedlings) by 200mM NaCl treatment for two days (wild-over-salt), the survival rate of the latter was improved by 2.5 as reported in the literature (15); while the

chlorophyll contents were also higher than that of wild type due to the effects of PtSnRK2.7 over expression on chlorophyll biosynthesis. And, this shows a direct relationship between chlorophyll contents and survival rate. The comparison outcomes of the current study in terms of wild and PtSnRK2.7 overexpression types showed that the only upregulated differentially expressed gene was CYP97B3 (i.e., AT4G15110 at chromosome 4). The 2 others were AT1G15010 (at chromosome 1 affected mostly in transient stress conditions (23)) and AT4G19430 (at chromosome 4 related to inactivation of the chloroplast (24)) without any annotated gene symbols. CYP97B3 as a member of plant cytochrome P450 involved in several biosynthetic reactions that can result in the production of defensive compounds (25). And, this can be almost regarded as no changes between wild and PtSnRK2.7 over expression types while treated by NaCl. Additionally, it induces the fact that the wild type will overexpress the level of PtSnRK2.7 gene to get adapted to the salt tolerance condition by consuming some resources resulting in a lower survival rate and chlorophyll contents.

It has been reported that in the absence of salt stress, significant changes were not observed in the chlorophyll contents of both types (15). This study also demonstrates that the 2 types have only changes in 8 downregulated genes with the highest interconnectivity degree, which can be the reason for the PtSnRK2.7 overexpression processes. It is worth noting that 5 out of 8 identified genes were also determined in a previous study for this condition (15). The other 3 were novel ones At1g73010, ATRNS1 and ATSPX1, which have been reported in the literature for stresses such as hypoxia and reoxygenation (23), plant defensive response (26) and transient stress (23).

Despite the existence of few conflicts in terms of the relationships between fold changes and the over/under-expression status of the overlapped genes, which may be directly related to the nature of the statistics-based algorithms used in the study for analyzing (15) and identifying the initial significant DEGs, the overlapping genes have substantial effects on the 3 compared groups *wild-mock-salt*, *wild-over-mock*, *over-mock-salt* (Supplementary Table 3). A variety of statistical tests are commonly used to identify differentially expressed genes (DEGs), including Welch's t-test, moderated t-test and permutation tests. For parametric tests, accurately estimating intra-sample-group variance is a critical issue; 2 improved variance estimation techniques are the locally pooled error and empirical Bayes methods. Because Omics data analysis typically involves tens of thousands of statistical tests, correcting multiple hypotheses is essential (27). To find DEGs, we usually use the t-test with the ordered set of P values converted to cumulative false discovery rate (FDR) estimates. A typical cutoff would be 10%. Both statistical functions are implemented in BRB-Array Tools (28).

Furthermore, the remaining group *wild-over-salt* was the only group being analyzed in this study and three essential genes (i.e., AT4G15110, AT1G15010, AT4G19430) were determined responsible for salt stress and chloroplast in similar conditions in such that the salt resistance

response of SnRK2.7 overexpression can be revealed (Supplementary Table 3). Finally, Supplementary Table 4 lists the overlapping genes, chromosome location and function in agreement with their corresponding literature studies.

Taking in to consideration the wild type exposed to salt condition (*wild-mock-salt*), thirteen unique genes were observed within nine statistically significant modules (p -value < 0.05) from the constructed PPI network. Supplementary Table 5 expounds the genes functions, chromosome position, accession no., as well as their cited references.

Moreover, fifteen genes with the highest interconnectivity with eleven identified gene modules were found statistically significant (p -value < 0.05), which are listed in Supplementary Table 6, along with their specific functionalities mostly related to several stresses such as salt stress.

A system biology approach was applied for studying the responses of *A. thaliana* to salt stress (abiotic stress) in wild and PtSnRK2.7 overexpression types. Furthermore, several analyses in terms of gene ontology (cellular component and molecular function), KEGG enrichment and plant ontology were performed. The outcomes were a set of significant genes identified from the PPI networks module clustering for four conditions mentioned above. Then, the identified genes were further validated through the literature findings considering their relations to abiotic and biotic stresses.

Conclusion

To provide salt tolerance in plants specifically *A. thaliana*, various mechanisms at the cellular, molecular, biological and whole-plant levels are involved. So, suitable strategies are needed to control the plant responses and adaptations to salinity stress. Although many studies have been carried out in this field, it is required to accumulate their results for the plant biologists to identify the responsible genomic modifications accounted for different corresponding mechanisms and critical pathways. Moreover, a gap exists as the investigations are routinely performed on one single gene rather than several potential genes. And, this urges future studies to take advantage of the retrospective computational-based studies to confirm the previously achieved outcomes possibly.

In conclusion, our findings suggest that the overexpressed PtSnRK2.7 gene primarily involved in resistive salt stress responses is, at least partially, responsible for molecular and functional mechanisms of the *A. thaliana* plant model. One of the significant outcomes of the current study, which was not mentioned in previous studies, is the effect of over expression in responsive genes related to unregulated chloroplast contents. Also, it was found out that the number of shared genes between the current study and the previous investigation for the 3 groups *wild-mock-salt*, *wild-over-mock*, *wild-over-salt* were 8, 9, 7. And, for the remaining group, only 3 significant genes were identified. Thus, additional investigation may be essential

to validate the critical role of PtSnRK2.7 overexpressed protein against stress responses to accommodate the required evidence in improving stress tolerance in several plants and trees.

Acknowledgements

The authors would like to thank the Research Office of Tabriz University of Medical Sciences for providing support under financial support scheme (Grant No. 64896).

Authors contributions

EA carried out the gathering and analyzing of studies and drafted the manuscript. SA participated in the design of the study and performed the statistical analysis. BS and SD conceived of the study and participated in its design and coordination as well as the analyses of data. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

Supplementary data

Table 1. Top upregulated and downregulated genes for four groups

Table 2. List of potential gene biomarkers using ClusterOne module analysis.

Table 3. Overlapping genes between current and Song et al. study along with their properties.

Table 4. Overlapping genes, chromosome location, and literature functional mechanisms.

Table 5. Overlapping genes, chromosome location, and literature functional mechanisms.

Table 6. The details of fifteen significant genes determined from the over-*mock-salt* type.

References

1. Yadav SP, Bharadwaj R, Nayak H, Mahto R, Singh RK, Prasad SK. Impact of salt stress on growth, productivity and physicochemical properties of plants: A Review. *IJCS*. 2019;7(2):1793-98.
2. Yamaguchi T, Blumwald E. Developing salt-tolerant crop plants: challenges and opportunities. *Trends in Plant Science*. 2005;10(12):615-20. <https://doi.org/10.1016/j.tplants.2005.10.002>
3. Morton MJ, Awlia M, Al-Tamimi N, Saade S, Pailles Y, Negrão S et al. Salt stress under the scalpel-dissecting the genetics of salt tolerance. *The Plant Journal*. 2019;97(1):148-63. <https://doi.org/10.1111/tpj.14189>
4. Acosta-Motos JR, Ortuño MF, Bernal-Vicente A, Diaz-Vivancos P, Sanchez-Blanco MJ, Hernandez JA. Plant Responses to Salt Stress: Adaptive Mechanisms. *Agronomy*. 2017;7(1):18. <https://doi.org/10.3390/agronomy7010018>
5. Pardo JM. *Biotechnology of water and salinity stress tolerance*. Current Opinion in Biotechnology. 2010;21(2):185-96. <https://doi.org/10.1016/j.copbio.2010.02.005>
6. Evelin H, Devi TS, Gupta S, Kapoor R. Mitigation of salinity stress in plants by arbuscular mycorrhizal symbiosis: current understanding and new challenges. *Frontiers in Plant Science*. 2019;10:470. <https://doi.org/10.3389/fpls.2019.00470>
7. Islam F, Wang J, Farooq MA, Yang C, Jan M, Mwamba TM et al. Rice responses and tolerance to salt stress: Deciphering the physiological and molecular mechanisms of salinity adaptation. *Advances in Rice Research for Abiotic Stress Tolerance*: Elsevier; 2019. p. 791-819. <https://doi.org/10.1016/B978-0-12-814332-2.00040-X>
8. Prasad B, Karodia S, Malviya J. Identifying genes responsible for shoot development in *Arabidopsis thaliana* using microarray technique. *Research Journal of Life Sciences, Bioinformatics, Pharmaceutical and Chemical Sciences*. 2019;5(1):262-82.
9. Meinke DW, Cherry JM, Dean C, Rounsley SD, Koornneef M. *Arabidopsis thaliana*: a model plant for genome analysis. *Science*. 1998;282(5389):662-82. <https://doi.org/10.1126/science.282.5389.662>
10. Liu H-H, Tian X, Li Y-J, Wu C-A, Zheng C-C. Microarray-based analysis of stress-regulated microRNAs in *Arabidopsis thaliana*. *Rna*. 2008;14(5):836-43. <https://doi.org/10.1261/rna.895308>
11. Patankar HV, Al-Harrasi I, Al Kharusi L, Jana GA, Al-Yahyai R, Sunkar R et al. Overexpression of a metallothionein 2A gene from date palm confers abiotic stress tolerance to yeast and *Arabidopsis thaliana*. *International Journal of Molecular Sciences*. 2019;20(12):2871. <https://doi.org/10.3390/ijms20122871>
12. Coolen S, Van Pelt JA, Van Wees SC, Pieterse CM. Mining the natural genetic variation in *Arabidopsis thaliana* for adaptation to sequential abiotic and biotic stresses. *Planta*. 2019;249(4):1087-105. <https://doi.org/10.1007/s00425-018-3065-9>
13. Kulik A, Wawer I, Krzywińska E, Bucholc M, Dobrowolska G. SnRK2 protein kinases-key regulators of plant response to abiotic stresses. *Omics: a Journal of Integrative Biology*. 2011;15(12):859-72. <https://doi.org/10.1089/omi.2011.0091>
14. Liu Z, Ge X, Yang Z, Zhang C, Zhao G, Chen E et al. Genome-wide identification and characterization of SnRK2 gene family in cotton (*Gossypium hirsutum* L.). *BMC Genetics*. 2017;18(1):54. <https://doi.org/10.1186/s12863-017-0517-3>
15. Song X, Yu X, Hori C, Demura T, Ohtani M, Zhuge Q. Heterologous overexpression of poplar SnRK2 genes enhanced salt stress tolerance in *Arabidopsis thaliana*. *Frontiers in Plant Science*. 2016;7:612. <https://doi.org/10.3389/fpls.2016.00612>
16. Carlson M. ath1121501.db: Affymetrix *Arabidopsis* ATH1 Genome Array annotation data (chip ath1121501). *Bioconductor.org*. 2016.
17. Yi X, Du Z, Su Z. PlantGSEA: a gene set enrichment analysis toolkit for plant community. *Nucleic Acids Research*. 2013;41(W1):W98-W103. <https://doi.org/10.1093/nar/gkt281>
18. Nepusz T, Yu H, Paccanaro A. Detecting overlapping protein complexes in protein-protein interaction networks. *Nature Methods*. 2012;9(5):471-72. <https://doi.org/10.1038/nmeth.1938>
19. Chin C-H, Chen S-H, Wu H-H, Ho C-W, Ko M-T, Lin C-Y. cytoHubba: identifying hub objects and sub-networks from complex interactome. *BMC Syst Biol*. 2014;8(4):S11. <https://doi.org/10.1186/1752-0509-8-S4-S11>
20. Teshome DT, Zharare GE, Naidoo S. The threat of the combined effect of biotic and abiotic stress factors in forestry under a changing climate. *Frontiers in Plant Science*. 2020;11(1874):601009. <https://doi.org/10.3389/fpls.2020.601009>
21. Zampieri M, Weissteiner CJ, Grizzetti B, Toreti A, van den Berg M, Dentener F. Estimating resilience of crop production systems: From theory to practice. *ScTEen*. 2020;735:139378. <https://doi.org/10.1016/j.scitotenv.2020.139378>

22. Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A et al. Crop production under drought and heat stress: Plant responses and management options. *Frontiers in Plant Science*. 2017;8:1147. <https://doi.org/10.3389/fpls.2017.01147>
23. Lee TA, Bailey-Serres J. Integrative analysis from the epigenome to translome uncovers patterns of dominant nuclear regulation during transient stress. *The Plant Cell*. 2019;31(11):2573-95. <https://doi.org/10.1105/tpc.19.00463>
24. Dal Bosco C, Lezhneva L, Biehl A, Leister D, Strotmann H, Wanner G, et al. Inactivation of the chloroplast ATP synthase γ subunit results in high non-photochemical fluorescence quenching and altered nuclear gene expression in *Arabidopsis thaliana*. *Journal of Biological Chemistry*. 2004;279(2):1060-69. <https://doi.org/10.1074/jbc.M308435200>
25. Schuler MA, Werck-Reichhart D. Functional genomics of P450s. *Annual Review of Plant Biology*. 2003;54(1):629-67. <https://doi.org/10.1146/annurev.arplant.54.031902.134840>
26. Kim K, Yadav D, Cho M. Multi-phased internalization of murine norovirus (MNV) in *Arabidopsis* seedlings and its potential correlation with plant defensive responses. *Microbial Pathogenesis*. 2019;135:103648. <https://doi.org/10.1016/j.micpath.2019.103648>
27. Ge Y, Sealson SC, Speed TP. Multiple testing and its applications to microarrays. *Stat Methods Med Res*. 2009;18(6):543-63. <https://doi.org/10.1177/0962280209351899>
28. Zhao Y, Simon R. BRB-ArrayTools Data Archive for human cancer gene expression: a unique and efficient data sharing resource. *Cancer Inform*. 2008;6:9-15. <https://doi.org/10.4137/CIN.S448>
29. Méteignier LV, Ghandour R, Meierhoff K, Zimmerman A, Chicher J, Baumberger N et al. The *Arabidopsis* mTERF-repeat MDA1 protein plays a dual function in transcription and stabilization of specific chloroplast transcripts within the psbE and ndhH operons. *New Phytol*. 2020;227(5):1376-91. <https://doi.org/10.1111/nph.16625>
30. Reinholdt O, Schwab S, Zhang Y, Reichheld JP, Fernie AR, Hagemann M et al. Redox-Regulation of Photorespiration through Mitochondrial Thioredoxin o1. *Plant Physiol*. 2019;181(2):442-57. <https://doi.org/10.1104/pp.19.00559>
31. Jin Y, Chen S, Fan X, Song H, Li X, Xu J et al. Diuron treatment reveals the different roles of two cyclic electron transfer pathways in photosystem II in *Arabidopsis thaliana*. *Pestic Biochem Physiol*. 2017;137:15-20. <https://doi.org/10.1016/j.pestbp.2016.09.002>
32. Qian H, Tsuji T, Endo T, Sato F. PGR5 and NDH pathways in photosynthetic cyclic electron transfer respond differently to sublethal treatment with photosystem-interfering herbicides. *J Agric Food Chem*. 2014;62(18):4083-89. <https://doi.org/10.1021/jf500143f>
33. Rawat R, Sandoval FJ, Wei Z, Winkler R, Roje S. An FMN hydroxylase of the haloacid dehalogenase superfamily is active in plant chloroplasts. *J Biol Chem*. 2011;286(49):42091-98. <https://doi.org/10.1074/jbc.M111.260885>
34. Ehrling J, Sauveplane V, Olry A, Ginglinger JF, Provart NJ, Werck-Reichhart D. An extensive (co-)expression analysis tool for the cytochrome P450 superfamily in *Arabidopsis thaliana*. *BMC Plant Biol*. 2008;8:47. <https://doi.org/10.1186/1471-2229-8-47>
35. Ascencio-Ibanez JT, Sozzani R, Lee T-J, Chu T-M, Wolfinger RD, Cella R et al. Global analysis of *Arabidopsis* gene expression uncovers a complex array of changes impacting pathogen response and cell cycle during geminivirus infection. *Plant Physiology*. 2008;148(1):436-54. <https://doi.org/10.1104/pp.108.121038>
36. Yoshida K, Hisabori T. Determining the Rate-Limiting Step for Light-Responsive Redox Regulation in Chloroplasts. *Antioxidants (Basel)*. 2018;7(11). <https://doi.org/10.3390/antiox7110153>
37. Simkin AJ, Lopez-Calcagno PE, Davey PA, Headland LR, Lawson T, Timm S et al. Simultaneous stimulation of sedoheptulose 1,7-bisphosphatase, fructose 1,6-bisphosphatase aldolase and the photorespiratory glycine decarboxylase-H protein increases CO₂ assimilation, vegetative biomass and seed yield in *Arabidopsis*. *Plant Biotechnol J*. 2017;15(7):805-16. <https://doi.org/10.1111/pbi.12676>
38. Bach-Pages M, Homma F, Kourelis J, Kaschani F, Mohammed S, Kaiser M et al. Discovering the RNA-binding proteome of plant leaves with an improved RNA interactome capture method. *Biomolecules*. 2020;10(4). <https://doi.org/10.3390/biom10040661>
39. Fares A, Rossignol M, Peltier JB. Proteomics investigation of endogenous S-nitrosylation in *Arabidopsis*. *Biochem Biophys Res Commun*. 2011;416(3-4):331-36. <https://doi.org/10.1016/j.bbrc.2011.11.036>
40. Rocha J, Nitenberg M, Girard-Egrot A, Jouhet J, Maréchal E, Block MA et al. Do galactolipid synthases play a key role in the biogenesis of chloroplast membranes of higher plants? *Front Plant Sci*. 2018;9:126. <https://doi.org/10.3389/fpls.2018.00126>
41. Botté CY, Deligny M, Rocchia A, Bonneau AL, Saïdani N, Hardré H et al. Chemical inhibitors of monogalactosyldiacylglycerol synthases in *Arabidopsis thaliana*. *Nat Chem Biol*. 2011;7(11):834-42. <https://doi.org/10.1038/nchembio.658>
42. Abercrombie JM, Halfhill MD, Ranjan P, Rao MR, Saxton AM, Yuan JS et al. Transcriptional responses of *Arabidopsis thaliana* plants to As (V) stress. *BMC Plant Biol*. 2008;8:87. <https://doi.org/10.1186/1471-2229-8-87>
43. Giraud E, Ho LH, Clifton R, Carroll A, Estavillo G, Tan YF et al. The absence of alternative oxidase1a in *Arabidopsis* results in acute sensitivity to combined light and drought stress. *Plant Physiol*. 2008;147(2):595-610. <https://doi.org/10.1104/pp.107.115121>
44. Sottosanto JB, Saranga Y, Blumwald E. Impact of AtNHX1, a vacuolar Na⁺/H⁺ antiporter, upon gene expression during short- and long-term salt stress in *Arabidopsis thaliana*. *BMC Plant Biol*. 2007;7:18. <https://doi.org/10.1186/1471-2229-7-18>
45. Dombrecht B, Xue GP, Sprague SJ, Kirkegaard JA, Ross JJ, Reid JB et al. MYC2 differentially modulates diverse jasmonate-dependent functions in *Arabidopsis*. *Plant Cell*. 2007;19(7):2225-45. <https://doi.org/10.1105/tpc.106.048017>
46. Tran LS, Nakashima K, Sakuma Y, Simpson SD, Fujita Y, Maruyama K et al. Isolation and functional analysis of *Arabidopsis* stress-inducible NAC transcription factors that bind to a drought-responsive cis-element in the early responsive to dehydration stress 1 promoter. *Plant Cell*. 2004;16(9):2481-98. <https://doi.org/10.1105/tpc.104.022699>
47. Bharti P, Mahajan M, Vishwakarma AK, Bhardwaj J, Yadav SK. AtROS1 overexpression provides evidence for epigenetic regulation of genes encoding enzymes of flavonoid biosynthesis and antioxidant pathways during salt stress in transgenic tobacco. *J Exp Bot*. 2015;66(19):5959-69. <https://doi.org/10.1093/jxb/erv304>
48. Mahmood K, Xu Z, El-Kereamy A, Casaretto JA, Rothstein SJ. The *Arabidopsis* transcription factor ANAC032 represses anthocyanin biosynthesis in response to high sucrose and oxidative and abiotic stresses. *Front Plant Sci*. 2016;7:1548. <https://doi.org/10.3389/fpls.2016.01548>
49. Huang KC, Lin WC, Cheng WH. Salt hypersensitive mutant 9, a nucleolar APUM23 protein, is essential for salt sensitivity in association with the ABA signaling pathway in *Arabidopsis*. *BMC Plant Biol*. 2018;18(1):40. <https://doi.org/10.1186/s12870-018-1255-z>
50. Wang L, Waters MT, Smith SM. Karrikin-KAI2 signalling provides *Arabidopsis* seeds with tolerance to abiotic stress and inhibits germination under conditions unfavourable to seedling estab-

- ishment. *New Phytologist*. 2018;219(2):605-18. <https://doi.org/10.1111/nph.15192>
51. Divi UK, Rahman T, Krishna P. Gene expression and functional analyses in brassinosteroid-mediated stress tolerance. *Plant Biotechnol J*. 2016;14(1):419-32. <https://doi.org/10.1111/pbi.12396>
 52. Turek I, Irving H, Gehring C. Dataset on interactors of the *Arabidopsis thaliana* Plant Natriuretic Peptide (AtPNP-A) determined by mass spectrometry. *Data Brief*. 2020;30:105606. <https://doi.org/10.1016/j.dib.2020.105606>
 53. Lee KP, Liu K, Kim EY, Medina-Puche L, Dong H, Duan J et al. Plant natriuretic peptide A and its putative receptor PNP-R2 antagonize salicylic acid-mediated signaling and cell death. *Plant Cell*. 2020;32(7):2237-50. <https://doi.org/10.1105/tpc.20.00018>
 54. Gandia-Herrero F, Lorenz A, Larson T, Graham IA, Bowles DJ, Rylott EL et al. Detoxification of the explosive 2,4,6-trinitrotoluene in *Arabidopsis*: discovery of bifunctional O- and C-glucosyltransferases. *Plant J*. 2008;56(6):963-74. <https://doi.org/10.1111/j.1365-313X.2008.03653.x>
 55. Ascencio-Ibáñez JT, Sozzani R, Lee TJ, Chu TM, Wolfinger RD, Cella R et al. Global analysis of *Arabidopsis* gene expression uncovers a complex array of changes impacting pathogen response and cell cycle during geminivirus infection. *Plant Physiol*. 2008;148(1):436-54. <https://doi.org/10.1104/pp.108.121038>
 56. Li B, Fang J, Singh RM, Zi H, Lv S, Liu R et al. Fatty acid desaturase 5 is required to induce autoimmune responses in gigantic chloroplast mutants of *Arabidopsis*. *Plant Cell*. 2020;32(10):3240-55. <https://doi.org/10.1105/tpc.20.00016>
 57. Liu F, Zhao Q, Jia Z, Song C, Huang Y, Ma H et al. N-3-oxo-octanoyl-homoserine lactone-mediated priming of resistance to *Pseudomonas syringae* requires the salicylic acid signaling pathway in *Arabidopsis thaliana*. *BMC Plant Biol*. 2020;20(1):38. <https://doi.org/10.1186/s12870-019-2228-6>
 58. Li LS, Ying J, Li E, Ma T, Li M, Gong LM et al. *Arabidopsis* CBP60b is a central transcriptional activator of immunity. *Plant Physiol*. 2021;186(3):1645-59. <https://doi.org/10.1093/plphys/kiab164>
 59. Lee TA, Bailey-Serres J. Integrative analysis from the epigenome to transcriptome uncovers patterns of dominant nuclear regulation during transient stress. *Plant Cell*. 2019;31(11):2573-95. <https://doi.org/10.1105/tpc.19.00463>
 60. Tian L, Musetti V, Kim J, Magallanes-Lundback M, DellaPenna D. The *Arabidopsis* LUT1 locus encodes a member of the cytochrome p450 family that is required for carotenoid epsilon-ring hydroxylation activity. *Proc Natl Acad Sci U S A*. 2004;101(1):402-27. <https://doi.org/10.1073/pnas.2237237100>
 61. Kim JE, Cheng KM, Craft NE, Hamberger B, Douglas CJ. Overexpression of *Arabidopsis thaliana* carotenoid hydroxylases individually and in combination with a beta-carotene ketolase provides insight into in vivo functions. *Phytochemistry*. 2010;71(2-3):168-78. <https://doi.org/10.1016/j.phytochem.2009.10.011>
 62. Ko JH, Yang SH, Han KH. Upregulation of an *Arabidopsis* RING-H2 gene, XERICO, confers drought tolerance through increased abscisic acid biosynthesis. *Plant J*. 2006;47(3):343-55. <https://doi.org/10.1111/j.1365-313X.2006.02782.x>
 63. Dal Bosco C, Lezhneva L, Biehl A, Leister D, Strotmann H, Anner G et al. Inactivation of the chloroplast ATP synthase gamma subunit results in high non-photochemical fluorescence quenching and altered nuclear gene expression in *Arabidopsis thaliana*. *J Biol Chem*. 2004;279(2):1060-69. <https://doi.org/10.1074/jbc.M308435200>
 64. Zybailov B, Rutschow H, Friso G, Rudella A, Emanuelsson O, Sun Q et al. Sorting signals, N-terminal modifications and abundance of the chloroplast proteome. *PLoS One*. 2008;3(4):e1994. <https://doi.org/10.1371/journal.pone.0001994>
 65. Wang Y, Zeng L, Xing D. ROS-mediated enhanced transcription of CYP38 promotes the plant tolerance to high light stress by suppressing GTPase activation of PsbO2. *Frontiers in Plant Science*. 2015;6:777. <https://doi.org/10.3389/fpls.2015.00777>
 66. Zheng X, Chen L, Li X. *Arabidopsis* and rice showed a distinct pattern in ZIPs genes expression profile in response to Cd stress. *Botanical Studies*. 2018;59(1):22. <https://doi.org/10.1186/s40529-018-0238-6>
 67. Huang K-C, Lin W-C, Cheng W-H. Salt hypersensitive mutant 9, a nucleolar APUM23 protein, is essential for salt sensitivity in association with the ABA signaling pathway in *Arabidopsis*. *BMC Plant Biology*. 2018;18(1):40. <https://doi.org/10.1186/s12870-018-1255-z>
 68. Ulm R, Revenkova E, di Sansebastiano G-P, Bechtold N, Paszkowski J. Mitogen-activated protein kinase phosphatase is required for genotoxic stress relief in *Arabidopsis*. *Genes and Development*. 2001;15(6):699-709. <https://doi.org/10.1101/gad.192601>
 69. Yao X, Cai Y, Yu D, Liang G. bHLH104 confers tolerance to cadmium stress in *Arabidopsis thaliana*. *Journal of Integrative Plant Biology*. 2018;60(8):691-702. <https://doi.org/10.1111/jipb.12658>
 70. Mittal S, Mallikarjuna MG, Rao AR, Jain PA, Dash PK, Thirunavukkarasu N. Comparative analysis of CDPK family in maize, *Arabidopsis*, rice and sorghum revealed potential targets for drought tolerance improvement. *Frontiers in Chemistry*. 2017;5:115. <https://doi.org/10.3389/fchem.2017.00115>
 71. Beemster G, De Jaeger G, editors. A systems approach to understand cell cycle regulation during leaf development. In: 18th International Conference on *Arabidopsis* Research 2007.
 72. Sánchez-Corrienero Á, Sánchez-Vicente I, González-Pérez S, Corrales A, Krieger-Liszczay A, Lorenzo Ó et al. Singlet oxygen triggers chloroplast rupture and cell death in the zeaxanthin epoxidase defective mutant *aba1* of *Arabidopsis thaliana* under high light stress. *Journal of Plant Physiology*. 2017;216:188-96. <https://doi.org/10.1016/j.jplph.2017.06.009>
 73. Ndamukong I, Jones DR, Lapko H, Divecha N, Avramova Z. Phosphatidylinositol 5-phosphate links dehydration stress to the activity of *Arabidopsis* trithorax-like factor ATX1. *PLoS One*. 2010;5(10):e13396. <https://doi.org/10.1371/journal.pone.0013396>
 74. He XJ, Mu RL, Cao WH, Zhang ZG, Zhang JS, Chen SY. AtNAC2, a transcription factor downstream of ethylene and auxin signaling pathways, is involved in salt stress response and lateral root development. *The Plant Journal*. 2005;44(6):903-16. <https://doi.org/10.1111/j.1365-313X.2005.02575.x>
 75. Ma S, Bohnert HJ. Integration of *Arabidopsis thaliana* stress-related transcript profiles, promoter structures and cell-specific expression. *Genome Biology*. 2007;8(4):R49. <https://doi.org/10.1186/gb-2007-8-4-r49>
 76. Kaye Y, Golani Y, Singer Y, Leshem Y, Cohen G, Ercetin M et al. Inositol polyphosphate 5-phosphatase7 regulates the production of reactive oxygen species and salt tolerance in *Arabidopsis*. *Plant Physiology*. 2011;157(1):229-41. <https://doi.org/10.1104/pp.111.176883>
 77. Abeysinghe JK, Lam KM, Ng DWK. Differential regulation and interaction of homoeologous WRKY 18 and WRKY 40 in *Arabidopsis allotetraploids* and biotic stress responses. *The Plant Journal*. 2019;97(2):352-67. <https://doi.org/10.1111/tpj.14124>
 78. Van Aken O, De Clercq I, Ivanova A, Law SR, Van Breusegem F, Millar AH et al. Mitochondrial and chloroplast stress responses are modulated in distinct touch and chemical inhibition phases. *Plant Physiology*. 2016;171(3):2150-65. <https://doi.org/10.1104/pp.16.00273>
 79. Renault H, El Amrani A, Berger A, Mouille G, Soubigou-taconnat L, Bouchereau A et al. γ -Aminobutyric acid transaminase defi-

- ciency impairs central carbon metabolism and leads to cell wall defects during salt stress in *Arabidopsis* roots. *Plant, Cell and Environment*. 2013;36(5):1009-18. <https://doi.org/10.1111/pce.12033>
80. Dong C-H, Hu X, Tang W, Zheng X, Kim YS, Lee B-h et al. A putative *Arabidopsis* nucleoporin, AtNUP160, is critical for RNA export and required for plant tolerance to cold stress. *Molecular and Cellular Biology*. 2006;26(24):9533-43. <https://doi.org/10.1128/MCB.01063-06>
 81. Stanley Kim H, Yu Y, Snesrud EC, Moy LP, Linford LD, Haas BJ et al. Transcriptional divergence of the duplicated oxidative stress-responsive genes in the *Arabidopsis* genome. *The Plant Journal*. 2005;41(2):212-20. <https://doi.org/10.1111/j.1365-313X.2004.02295.x>
 82. Pulido P, Zagari N, Manavski N, Gawronski P, Matthes A, Scharff LB et al. Chloroplast ribosome associated supports translation under stress and interacts with the ribosomal 30S subunit. *Plant Physiology*. 2018;177(4):1539-54. <https://doi.org/10.1104/pp.18.00602>
 83. Ding Y, Liu N, Virilouvet L, Riethoven J-J, Fromm M, Avramova Z. Four distinct types of dehydration stress memory genes in *Arabidopsis thaliana*. *BMC Plant Biology*. 2013;13(1):229. <https://doi.org/10.1186/1471-2229-13-229>
 84. Martínez-Ballesta M, Moreno-Fernández DA, Castejón D, Ochando C, Morandini PA, Carvajal M. The impact of the absence of aliphatic glucosinolates on water transport under salt stress in *Arabidopsis thaliana*. *Frontiers in Plant Science*. 2015;6:524. <https://doi.org/10.3389/fpls.2015.00524>
 85. Taochy C, Gaillard I, Ipotesi E, Oomen R, Leonhardt N, Zimmermann S et al. The *Arabidopsis* root stele transporter NPF2.3 contributes to nitrate translocation to shoots under salt stress. *The Plant Journal*. 2015;83(3):466-79. <https://doi.org/10.1111/tpj.12901>
 86. Moreno JE, Moreno-Piovano G, Chan RL. The antagonistic basic helix-loop-helix partners BEE and IBH1 contribute to control plant tolerance to abiotic stress. *Plant Science*. 2018;271:143-50. <https://doi.org/10.1016/j.plantsci.2018.03.024>
 87. Chan CW, Wohlbach DJ, Rodesch MJ, Sussman MR. Transcriptional changes in response to growth of *Arabidopsis* in high external calcium. *FEBS Letters*. 2008;582(6):967-76. <https://doi.org/10.1016/j.febslet.2008.02.043>

§§§