



RESEARCH ARTICLE

# Growth enhancement of agricultural crops using seaweed liquid fertilizer

Akshada Parab\* & Sunil Shankhadarwar

Department of Botany, Ramnarain Ruia Autonomous College, Mumbai 400 019, India

\*Email: [parabakshada10@gmail.com](mailto:parabakshada10@gmail.com)

 OPEN ACCESS

## ARTICLE HISTORY

Received: 30 September 2021

Accepted: 19 December 2021

Available online

Version 1.0 (Early Access): 26 January 2022



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, etc. See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Parab A, Shankhadarwar S. Growth enhancement of agricultural crops using seaweed liquid fertilizer. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.1439>

## Abstract

Extensive use of chemical fertilizer is leading to infertility of soil and various hazardous effect on the ecosystem. Using biofertilizer reduces these harmful effects. Many are commercially available, being a poor match to chemical fertilizer they are not used on a large scale. Near coastal regions seaweeds are used as biofertilizers. Recent studies show seaweeds have multiple growth regulators, macro and micronutrients, polysaccharides which are necessary for plant growth. The study is carried out to check the bioactivity of Seaweed liquid fertilizer (SLF) as a growth stimulant. *Sargassum cinereum*, *Ulva rigida* and *Ahnfeltia plicata* found on the Indian coastline were used. SLFs were prepared from each of three species in varying concentration (0.02%, 0.04%, 0.06%, 0.08% and 0.1% v/v) with water and Urea as control. The test plant selected was *Vigna radiata*. Agronomic characters like germination rate, shoot length, flowering and fruiting period of test plants were studied. Phyto-hormones from SLF were detected by thin-layer chromatography. Elemental analysis of micronutrients in SLF was carried by ICP-AES. Along with Chlorophyll and Protein estimation of grown plants were checked. SLF shelf life was studied using chemical and biological preservatives. The effect of 0.06% concentration of all SLFs showed enhancement in growth and phytochemicals in plants. Plants treated with SLF showed flowering earlier than control plants. Thus, using SLF bio-fertilizer can become an alternative method to reduce the use of harmful chemical fertilizers.

## Keywords

Seaweeds, Bio-fertilizer, Phytohormones, ICP-AES, Shelf-life

## Introduction

Agriculture is the backbone of India; various schemes, modifications, hybrid varieties etc. are developed to enhance the yield of crops. The use of fertilizers, insecticides, herbicides etc. has increased after the green revolution. Overuse of chemical fertilizers has increased the crop yield and production but also there is excessive chemical leaching into the soil, causing bioaccumulation and other harmful effects. The health impact is increased in several cancers and other diseases. As a result, farmers are opting to reuse traditional methods. The use of organic fertilizers like manures, bio-stimulants is increasing. Production of organic fertilizers is increasing since 2012 according to reports from the Department of Fertilizers Ministry of Chemicals and Fertilizers Government of India (1). Organic fertilizers include the ones made from living organisms like plants, animals, and microbes (2).

Organic fertilizers are also prepared from algae, both micro and macro. Algae biofertilizers are effective as they are rich in proteins, lipids, polysaccharides, nutrients. Cyanobacteria like *Nostoc* and *Anabaena* are already naturally enhancing agriculture crops, there are emerging studies in the use of microalgae fertilizers but the time taken to grow algae is more (3). Talking about seaweeds, which are grown naturally in coastal areas, are equally rich in all the contents. Seaweeds are marine macroscopic algae that are used as biofertilizers from ancient times i.e., Romans in the 1st century AD as well as by the Chinese (4). Commercial use of seaweeds in agriculture is carried out for the last seven decades (5). Seaweed provides nutrition as it contains macro and micronutrients, amino acids, vitamins, phytohormones, polysaccharides etc (6). These help in increase in growth and development, tolerate environmental stress (7), enhance nutrient uptake from soil (8) and also increase nutrients in the plants itself (9).

Seaweeds are used as biofertilizers, manures as well as biostimulants. The application of foliar spray of the liquid extracts has shown a tremendous effect on the growth and flowering of tomato plants (10). Seaweed fertilizers are more beneficial than chemical and other organic fertilizers as they are biodegradable, non-polluting, non-toxic and as mentioned earlier have various growth-promoting organic compounds (11). SLF was prepared from *Codium decorticatum*, with varying concentrations and tested on capsicum plants (12). SLF was also tested on cucumber plants (13), for growth enhancement. There are studies on using a combination of seaweeds in preparing SLF (14) and all these studies have a common concentration of SLF used.

In India, seaweeds are commercially grown for pharmaceuticals and cosmetics. Agricultural applications of seaweeds are less explored. Seaweed fertilizer available in markets is made from *Ascophyllum nodosum* a brown alga imported from European countries which makes it expensive. As a result, very few people know the use and buy it. Hence, the present work follows the use of indigenous seaweeds i.e. the ones found in our country, like *Sargassum cinereum*, *Ulva rigida* and *Ahnfeltia plicata* as biofertilizers and biostimulants for growth enhancement crops. Concentration of each SLF prepared is 0.02%, 0.04%, 0.06%, 0.08% and 0.1%, which is unlike other studies published as more concentrated SLFs are used (3, 12, 13). The selected crop in this study, Moong - *Vigna radiata* has been selected and grown using SLF and with water and urea as control. Agronomic and biochemical tests were conducted to compare the effects of SLF and chemical fertilizer which is further discussed in the study.

Thus, the study states the preparation of seaweed extracts which are economically cheap and easily available in coastal regions rather than imported ones. The concentrations used are low and have an impact on crops. Also, preparation of SLF is discussed which opens up start-up opportunities for farmers and fishermen, also reducing the use of chemical fertilizer.

## Materials and Methods

### Sample collection

Seaweed samples were collected from the west coast of Maharashtra, India. Three species from each of the classes i.e., Chlorophyceae, Rhodophyceae and Phaeophyceae were selected. *Ulva rigida* C. Agardh was collected from Malvan, *Sargassum cinereum* J. Agardh was collected from Devgad while, *Ahnfeltia plicata* (Hudson) E. M. Fries was collected from Alibaug. The seaweed samples were authenticated from the Botanical Survey of India, Southern Regional Circle T.N.A.U. campus, Coimbatore in November 2019. These seaweeds were identified up to species level.

Collected seaweeds were washed thoroughly to remove the salt and sand attached, along with living impurities like marine organisms along with seashells were separated. These washed seaweeds were then shade dried at room temperature. Further, packed airtight and stored at 37 °C in an incubator to keep the biochemical properties intact. Samples were used in dried form for the study.

### Preparation of seaweed liquid fertilizer

Seaweed liquid fertilizer of each sample was prepared by modifying the method (15). One gm of dried seaweed was taken in 100 ml of distilled water which was further autoclaved at 15 Psi for 30 mins. It was then cooled and filtered through a muslin cloth. The solution was then centrifuged for 10 mins at 3000 rpm. The supernatant was collected and stored in the refrigerator for further use. The extract from was 0.1% extract which was then diluted into different concentrations using distilled water and made into 0.02%, 0.04%, 0.06%, 0.08% and 0.1%. The same concentration was used for germination and further the plants were given SLF every week as a foliar spray and direct application in coco peat.

### Seed selection and treatment

The crop selected for testing the SLF was Moong beans (*Vigna radiata*). The seeds were washed with distilled water two times than with 70% alcohol again with D/W. Further, they were washed with 0.1% HgCl<sub>2</sub> and then with sterile distilled water two to three times to avoid infection. At last, they were dried on filter paper and were further maintained.

Seeds are further soaked in each of the three SLF of different concentrations (0.02%, 0.04%, 0.06%, 0.08% and 0.1%) which were kept overnight. Distilled water was used as control and chemical control as urea (0.05 g). Ten seeds are sown in pots containing coco peat to avoid interactions with soil minerals. Data was noted for 10 days. There was a total of 7 pots with each type of concentration of SLF along with 2 control. Initially, pots were kept in a greenhouse and maintained for temperature and humidity. Further, the study was carried out in direct sunlight. For the rate of germination, the plants were noted for 10 day and agronomic characters were under observation for 10 weeks. As mentioned earlier, SLF was applied at week intervals – 10 ml/week of which 5 ml was applied as a foliar spray and the remaining 5 ml was directly applied in co-peat.

### Estimation of the plant hormone

Endogenously present plant hormones i.e., Auxin - Indole acetic acid (IAA) and Cytokinin - kinetin were detected using thin-layer chromatography. Merk Silica gel 60 F254 TLC plates were used the aqueous liquid extracts were used for both the detection. Initially, for IAA 15 ml of the sample was taken and evaporated to 5 ml. which was then adjusted to pH 2.8 using 7N HCl. Further to this, 15 ml ethyl acetate was added and the aqueous fraction was separated which was again adjusted to pH 7.0 using 7N NaOH. To this water-saturated butanol was added and this aqueous layer was again taken and evaporated. Methanol was used for reconstitution and the sample preparation was stored at 4 °C in dark. The Mobile phase used was n-Butanol: acetic acid: Ethanol (12:3:5) (V: V: V) and Salkowski's reagent was used for spraying and treated at 80 °C. Plates were visualized under visible light at 554 nm. For cytokinin, the aqueous 15 ml extract was taken centrifuged at 40 °C for 20 mins. It was mixed with 15 ml of ethyl acetate of which an aqueous layer was separated and evaporated which was reconstituted with methanol. For this, the mobile phase was n-Butanol: Acetic acid: water (12:3:5) (V: V: V). Plates were visualized at 366 nm (16).

### Elemental analysis

Seaweeds already consist of various minerals. Plants need minerals in major and minor quantities. In the following work, plant minor elements were detected from the aqueous extracts using the Induced Coupled Plasma-Atomic Emission Spectroscopy (ICP-AES) method. The following was carried out at SAIF-IIT Bombay in November 2019. The

phyll a and b and total chlorophylls were estimated. All the plants including control and urea treated were also tested (17).

### Protein estimation

Proteins were also calculated from the leaves of Moong to check enhancement. Control and urea-treated leaves were tested. Lowry's method of estimation was carried out (18). Leaves were extracted into aqueous extracts.

### The shelf life of SLF

It was seen at SLF if stored outside gets contaminated by fungus and bacteria. Hence to avoid it, they were stored in spray bottles in the refrigerator. For storing outside at room temperature was also tested. Sodium benzoate was used in varying concentrations along with natural preservation using clove oil. Clove oil was also used in varying concentrations. The test was carried out for one month.

## Results

### Rate of germination

The Rate of germination was observed in the three SLF compared with control and Urea. It was observed after 48 hrs of treatment that, *U. rigida* treated plants showed maximum germination followed by *A. plicata* and *S. cinereum* (Table 1-3). Maximum growth that was 80% which is 8 out of 10 seeds germinated was in *U. rigida* treated seeds with 0.06% concentrations. While in *S. cinereum* it was 10% of the total in 0.06% concentration. Water treated seeds showed a 40% rate while urea treated ones showed 0% germination (Fig. 6).



Fig. 1. Plants grown in *U. rigida* SLF along with water as control

elements selected were Copper-Cu, Nickel-Ni, Molybdenum-Mb, Zinc-Zn and Manganese-Mn. The following elements are useful in various metabolic pathways in plants. For control, distilled water was tested. 5 ml of each SLF of three seaweeds i.e., *U. rigida*, *S. cinereum* and *A. plicata* were tested.

### Biochemical parameters of plants treated with SLFs

#### Chlorophyll estimation

Chlorophyll content of the Moong plants, when started flowering at 5th week, were tested. Only leaves were taken. Chlorophyll was estimated by Aron's method using 80% acetone extracts at 645 nm, 652 nm and 663 nm. Chloro-

### Agronomic characters

#### Height of plants

The height of shoots was recorded in the initial stages of growth after the seeds were sown in cocopeat. The average height of all plants in each concentration of SLF was recorded. In the first week, 0.2% *S. cinereum* treated plants were tallest followed by 0.8% *S. cinereum* SLF with a height of 8.5 cm and 7.8 cm respectively. It was recorded that plants treated with 0.6% *U. rigida* SLF and 0.6% *A. plicata* SLF showed 6.3 cm plantlets. Urea treated plants were the least of 2.4 cm and water treated was 4.5 cm (Table 1-3).



**Table 1.** (Rate of germination, agronomic characters -the height of plants, number of leaves and number of flowers along with biochemical parameters like chlorophyll content and protein content for *U. rigida* treated plants along with control as water and urea)

Sl. No.	SLF concentration (%)	Rate of germination (48hours)	Shoot height (cm)	Number of leaves	Number of flowers	Chlorophyll A (mg/g) ± SE	Chlorophyll B (mg/g) ± SE	Total chlorophyll (mg/g) ± SE	Protein (mg/g) ± SE
1	0.02%	40%	3.6	6	0	3.14 ± 0.03	3.58 ± 0.007	3.94 ± 0.06	196.3 ± 0.25
2	0.04%	70%	4.8	12	1	6.63 ± 0.01	7.748 ± 0.02	10.26 ± 0.02	237.1 ± 0.48
3	0.06%	80%	6.3	10	0	7.17 ± 0.04	7.35 ± 0.01	9.08 ± 0.004	235.8 ± 0.74
4	0.08%	70%	4.2	8	0	3.76 ± 0.01	4.51 ± 0.003	5.93 ± 0.007	190.7 ± 0.56
5	0.1%	70%	5.8	12	0	6.75 ± 0.05	6.85 ± 0.004	8.41 ± 0.004	200.0 ± 0.28
6	Urea	0%	2.4	6	0	5.82 ± 0.03	3.68 ± 0.02	5.33 ± 0.009	231.5 ± 0.87
7	water	40%	4.5	6	0	4.06 ± 0.01	4.26 ± 0.03	6.72 ± 0.006	263.6 ± 0.98

**Fig. 2.** Plants grown in *S. cinereum* SLF along with water as control.**Table 2.** Rate of germination, agronomic characters -the height of plants, number of leaves and number of flowers along with biochemical parameters like chlorophyll content and protein content for *S. cinereum* treated plants along with control as water and urea

Sr.no	SLF concentration (%)	Rate of germination (48hours)	Shoot height (cm)	Number of leaves	Number of flowers	Chlorophyll A (mg/g) ± SE	Chlorophyll B (mg/g) ± SE	Total chlorophyll (mg/g) ± SE	Protein (mg/g) ± SE
1	0.02%	60%	5.2	8	1	4.40 ± 0.004	5.05 ± 0.04	5.10 ± 0.03	191.1 ± 0.23
2	0.04%	60%	5	11	1	5.25 ± 0.01	7.91 ± 0.003	9.74 ± 0.02	194.4 ± 0.3
3	0.06%	60%	6.3	12	0	4.31 ± 0.08	4.75 ± 0.001	6.41 ± 0.004	190.7 ± 0.81
4	0.08%	50%	4.2	7	0	6.39 ± 0.008	4.84 ± 0.006	7.50 ± 0.064	183.3 ± 0.03
5	0.1%	30%	5.3	8	0	6.55 ± 0.041	4.34 ± 0.002	7.12 ± 0.02	201.9 ± 0.09
6	Urea	0%	2.4	6	0	5.88 ± 0.01	3.68 ± 0.03	5.33 ± 0.04	231.5 ± 0.23
7	water	40%	4.5	6	0	4.08 ± 0.003	4.26 ± 0.02	6.72 ± 0.12	263.6 ± 0.41

**Fig. 3.** Plants grown in *A. plicata* SLF along with water as control

### Number of leaves

As an agronomic character, several leaves were recorded after five weeks of germination. Water and urea treated were used as control. In the fifth week, *U. rigida* and *S. cinereum* were with 12 leaves while *A. plicata* treated leaves were similar to *S. cinereum* with a maximum number of 12


leaves in 0.6% SLF. Water and urea were treated where with 6 leaves each (Table 1-3).

### Number of flowers

Flowering and fruiting were seen in the plants treated with SLF after five weeks after germination. In the fifth week, *S. cinereum* treated plants were with the greatest number of

**Table 3.** Rate of germination, agronomic characters -the height of plants, number of leaves and number of flowers along with biochemical parameters like chlorophyll content and protein content for *A. plicata* treated plants along with control as water and urea

Sr.no	SLF concentration (%)	Rate of germination (48hours)	Shoot height (cm)	Number of leaves	Number of flowers	Chlorophyll A (mg/g) ± SE	Chlorophyll B (mg/g) ± SE	Total chlorophyll (mg/g) ± SE	Protein (mg/g) ± SE
1	0.02%	20%	8.5	12	2	5.92±0.08	7.71±0.06	9.89±0.007	201.2±0.56
2	0.04%	20%	4.6	11	0	8.44±0.005	11.32±0.007	14.11±0.001	184.9±0.03
3	0.06%	10%	5.1	7	1	6.62±0.008	9.69±0.02	12.03±0.003	241.5±0.25
4	0.08%	40%	7.8	8	2	4.85±0.007	6.23±0.009	7.74±0.006	254.7±0.52
5	0.1%	40%	6.2	8	0	4.16±0.004	4.73±0.006	6.275±0.02	281.4±0.49
6	Urea	0%	2.4	6	0	5.8±0.03	3.68±0.004	5.33±0.01	231.5±0.69
7	water	40%	4.5	6	0	4.08±0.06	4.26±0.02	6.724±0.006	263.6±0.30



**Fig. 6.** Rate of germination by SLFs and compared with urea and water.

flowers and buds. It was 2 buds per concentration of 0.2% and 0.6% SLF. While *A. plicata* SLF treated were with 1 flower in 0.2% and 0.4% concentration of SLF. *U. rigida* had only one flower in 0.4% concentration. Urea and water showed no flowering (Table 1-3).

### Biochemical parameters

#### Chlorophyll

The chlorophyll content of plants was checked which were treated with SLF and control as water and urea. The Maximum total chlorophyll content was in plants treated with *S. cinereum* SLF with 14.11 mg/g with maximum chlorophyll a 8.44 mg/g and chlorophyll b as 11.32 mg/g. least was in *A. plicata* with chlorophyll a 4.40 mg/g and chlorophyll b with 5.05 mg/g. Water treated showed the lowest of chlorophyll an of 4.07 mg/g and 3.688 mg/g chlorophyll b. while urea treated showed 5.88 mg/g of chlorophyll a and 4.26 mg/g chlorophyll b (Table 1-3) and (Fig. 7).

#### Proteins

The protein content of plants treated with SLF was estimated along with urea and water as a control. It was observed that *S. cinereum* was with the maximum protein content of 281.4 mg/g while *A. plicata* was lowest with 162.9 mg/g.

water and urea treated were 263.6 mg/g and 231.5 mg/g respectively (Table 1-3).

#### Plant hormones

Auxin TLC was carried out to check the presence, it was seen in all the seaweed extracts. While for cytokinin, it was only detected in *U. rigida* and *S. cinereum* (Fig. 4 & 5).

#### Induced coupled plasma-atomic emission spectroscopy (ICP-AES)

ICP-AES was carried out to check the presence of microelements. It was seen that the maximum content of Copper was in Rhodophyceae, Manganese content was maximum in Phaeophyceae members. Nickel was about 0.053 ppm in Chlorophyceae and Zinc was in the highest amounts in Rhodophyceae. Molybdenum was not detected in any of the seaweed (Table 4).

#### Shelf life

As per the conditions and quantity of sodium benzoate added as well as clove oil, no bacterial or fungal growth was seen in clove oil-treated extracts while there was bacterial growth in all the sodium benzoate treated extracts (Table 5).




Fig. 7. Total chlorophyll content in moong treated with SLFs and compared with urea and water.




Fig. 4. Presence of Auxin by TLC ; from left to right ( Sargassum SLF, Ahnfeltia SLF, Ulva SLF and Standard Auxin IAA)

Table 4. ICP-AES carried out on SLFs along with microelements. ND means less than 0.01ppm

Sample	Cu	Mn	Ni	Zn	Mo
Concentration	ppm	ppm	ppm	ppm	ppm
Distilled water	ND	ND	ND	0.016	ND
<i>A. plicata</i>	0.038	0.099	0.019	0.147	ND
<i>S. cinereum</i>	0.022	0.229	0.019	0.066	ND
<i>U. rigida</i>	0.038	0.25	0.053	0.113	ND

Table 5. Shelf life comparison of different concentration of Sodium benzoate and Clove oil to fungal growth

Sr.no	Concentration	Observations
<u>Sodium benzoate:</u>		
1	0.01g	+
	0.02g	+
	0.03g	-
	0.04g	-
	0.05g	-
<u>Clove oil</u>		
2	0.01ml	-
	0.02ml	-
	0.03ml	-




Fig. 5. Presence of Cytokinin by TLC ; from left to right ( Sargassum SLF, Ulva SLF, Ahnfeltia SLF and Standard kinetin).

### Discussion

Seaweeds are used as biofertilizers for many centuries. Research on Seaweeds as fertilizers is carried out for the last three decades. Seaweed liquid fertilizer (SLF) is applied as a biostimulant. Many seaweeds are worked upon for the same some are also commercially available as of products. (19) *Ascophyllum nodosum* found in Scandinavian countries is commercially available as of SLF. And also, these countries are the main suppliers of these biostimulants. The Effects of these biostimulants are much ranging from enhancement in flowering and fruiting, increase in protein, antioxidant content, increased in yield and as well as reduced in infections and pests (20).



In the present work, three indigenous seaweeds found at the western coast are tested. All three seaweeds are authenticated and used in optimum quantities. Work is carried on *Vigna radiata* as the plant has a short cycle. The three seaweeds are used in five different concentrations each from 0.2% to 1% along with water and urea as control. Seaweed extracts were prepared by autoclaving the seaweed with distilled water. There are many other methods of preparation, but comparatively, this method is easy and cost-effective (21). Method was modified and concentrations were reduced, which also shows the minimum quantity of seaweed can also show enhancement in crops.

The Rate of germination was maximum in *U. rigida* treated seeds i.e. 80% germination at 0.06% concentration. While in *A. plicata* in lower concentrates showed maximum growth. Similar results were seen in pearl millets treated with *Gracilaria dura* as lower concentrates showed a maximum rate of germination, as well as maximum chlorophyll and protein content (22), thus for Red algae lower concentration of SLF is profitable.

Agronomic characters were recorded, the height of shoots was recorded after a week, it was observed the plants treated with SLF were tall compared to plants treated with urea and water. Due to the presence of various endogenous auxins like indole-3-carboxylic acid, indole-3-acetic acid etc (23). Number of leaves was counted as a part of growth parameters, plants were treated with seaweeds had a greater number of leaves compared to control. The number of leaves is proportional to the rate of photosynthesis (24). Several flowers were also counted in the fifth week when it was just started to bloom, the rate was also calculated in the seventh week. Urea and water-treated plants were without flowers. Seaweed extracts contain plant hormones, micro and macronutrients as well as polysaccharides which enhance flowering. Similar results were observed in *Tagetes erecta* with *Sargassum wightii* (25). The effect of seaweed commercial extracts like Kelpak was tested on three varieties of capsicum along with chemical fertilizer as control. It was observed seaweed fertilizer had better results compared to chemical fertilizers (26).

Plant hormones and growth regulators are present in seaweeds (27). Seaweed liquid extracts were tested for the presence of cytokinin and auxins. Auxins were present in all three extracts while cytokinin was present in *U. rigida* and *S. cinereum*. Although there are reports on the presence of cytokinins endogenously present in various red algae species (28). Changes can be due to temperature and storage variations, as mentioned in a work carried out with *Ecklonia maxima* stored at various temperatures thus varying in concentrations of plant hormones were observed (29). Also, extraction of phytohormones from seaweed is carried out earlier, but extraction of auxins and cytokinins from prepared SLF is not reported. Thus, the study becomes unique as actual concentration of growth hormones is been detected.

Trace nutrients were detected using the ICP-AES method. Micronutrients are essential in plant metabolic processes. Copper, Zinc, Nickel and Manganese were pre-

sent in all three extracts while Molybdenum was not detected. There was an earlier report on *Vaucheria sessilis* for the presence of various micro and macronutrients (30). These nutrients play role in various enzymatic activities, thus increasing plants responses to the environment.

Thus, from this study, it was observed that seaweed extracts show enhancement compared to the chemical fertilizer.

## Conclusion

Seaweed liquid fertilizer thus enhances the growth of plants. It was observed from the first treatment to seeds till flowering and fruiting. The Rate of germination was increased in plants treated with SLF. It was observed that in *U. rigida* treated plants, 0.06% was optimum for growth. While in *A. plicata* lower concentration has a maximum germination rate, while for *S. cinereum* treated plants initially, there was less rate followed in 100% germination in the next 24 hrs. Due to the increase in the rate of germination, further, these plants showed enhancement in flowering. Enhancement is observed due to various growth-stimulating hormones, micronutrients and macronutrients (31). There are many reports on the bio stimulating activity of seaweeds on various plants. Moong is a fast-growing crop and an essential legume. It was seen that plants treated only with water and urea had no flowering and fruiting, thus proving that SLF contains bio-stimulating properties. Based on the detection of plant hormones it is clear that there is the presence of auxins in extracts and cytokinins in *S. cinereum* and *U. rigida*. Reports are on the presence of indigenous plant hormones in seaweeds (32). But in aqueous extracts, there are fewer reports and ICP-AES reports confirm the presence of micronutrients. Estimating biochemical parameters help in confirming enhancement due to SLF. From the study, it was concluded that three indigenous present seaweeds can be used for preparing SLF. *S. cinereum* with the best results followed by *U. rigida* and *A. plicata*. Thus, seaweed liquid fertilizer shows significant enhancement in growth and as a source of biostimulant for agriculture crops

## Acknowledgements

The authors are thankful and express gratitude to the institution, Ramnarain Ruia Autonomous college, Matunga, Mumbai for carrying out the study. Also, to Botanical Survey of India, Southern Regional Circle, T.N.A.U. campus, Coimbatore for proper authentication of seaweeds. And to SAIF-IIT Bombay, Mumbai for their ICP-AES facility used in the study.

## Authors contributions

The current study was initiated and planned by SS. AP collected samples, carried out experiments, analysed data and wrote the manuscript. SS guided and edited the manuscript followed by resolving the reviewers' edits.

## Compliance with ethical standards

**Conflict of interest:** The authors do not have any conflict of interests to declare .

**Ethical issues:** None.

## References

- Annual Report 2019-20. Government of India Ministry of Chemicals and Fertilizers Department of Fertilizers <https://fert.nic.in/sites/default/files/2020-09/Annual-Report-2019-20.pdf>
- Dittmar H, Drach M, Vosskamp R, Trenkel ME, Gutser R, Steffens G. Fertilizers, 2. types. Ullmann's Encyclopedia of Industrial Chemistry. 2000 Jun 15. [https://doi.org/10.1002/14356007.a10\\_323](https://doi.org/10.1002/14356007.a10_323)
- Deepika P, Mubarak Ali D. Production and assessment of microalgal liquid fertilizer for the enhanced growth of four crop plants. Biocatalysis and Agricultural Biotechnology. 2020 Sep 1;28:101701 <https://doi.org/10.1016/j.bcab.2020.101701>
- Newton L. Seaweed utilization. Nature. 1951 Jun;167(4260):1004- <https://doi.org/10.1038/1671004a0>
- Booth E. Some properties of seaweed manures. In: Proceedings of the 5<sup>th</sup> International Seaweed Symposium, Halifax, August 25-28, 1965, 1966 Jan 1 (pp. 349-357). Pergamon. <https://doi.org/10.1016/B978-0-08-011841-3.50055-0>
- Thirumarani G, Arumugam M, Arumugam R, Anantharaman P. Effect of seaweed liquid fertilizer on growth and pigment concentration of *Cyamopsis tetragonoloba* (L.) Taub. American-Eurasian Journal of Agronomy. 2009 Jan;2(2):50-56. <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.606.4774&rep=rep1&type=pdf>
- Zhang W, Chapman Dj, Phinney Bo, Spray Cr, Yamane H, Takahashi N. Identification of cytokinins in *Sargassum muticum* (Phaeophyta) and *Porphyra perforata* (Rhodophyta) 1. Journal of Phycology. 1991 Feb;27(1):87-91. <https://doi.org/10.1111/j.0022-3646.1991.00087.x>
- Turan M, Köse C. Seaweed extracts improve copper uptake of grapevine. Acta Agriculturae Scandinavica, Section B-Soil & Plant Science. 2004 Nov 1;54(4):213-20. <https://doi.org/10.1080/09064710410030311>
- Verkleij FN. Seaweed extracts in agriculture and horticulture: a review. Biological Agriculture and Horticulture. 1992 Jan 1;8(4):309-24. <https://doi.org/10.1080/01448765.1992.9754608>
- Crouch IJ, Van Staden J. Effect of seaweed concentrate on the establishment and yield of greenhouse tomato plants. Journal of Applied Phycology. 1992 Dec;4(4):291-96. <https://doi.org/10.1007/BF02185785>
- Dhargalkar VK, Pereira N. Seaweed: promising plant of the millennium. Science and Culture. 2005 Mar;71(3-4):60-66. <http://drs.nio.org/drs/handle/2264/489>
- Vijayakumar S, Durgadevi S, Arulmozhi P, Rajalakshmi S, Gopalakrishnan T, Parameswari N. Effect of seaweed liquid fertilizer on yield and quality of *Capsicum annum* L. Acta Ecologica Sinica. 2019 Oct 1;39(5):406-10. <https://doi.org/10.1016/j.chnaes.2018.10.001>
- Hassan SM, Ashour M, Sakai N, Zhang L, Hassanien HA, Gaber A, Ammar G. Impact of seaweed liquid extract biostimulant on growth, yield and chemical composition of Cucumber (*Cucumis sativus*). Agriculture. 2021 Apr;11(4):320. <https://doi.org/10.3390/agriculture11040320>
- Karthik T, Sarkar G, Babu S, Amalraj LD, Jayasri MA. Preparation and evaluation of liquid fertilizer from *Turbinaria ornata* and *Ulva reticulata*. Biocatalysis and Agricultural Biotechnology. 2020 Sep 1;28:101712. <https://doi.org/10.1016/j.bcab.2020.101712>
- Bhosle NB, Untawale AG, Dhargalkar VK. Effect of seaweed extract on the growth of *Phaseolus vulgaris* L. <http://drs.nio.org/drs/handle/2264/5577>
- Parvin W, Othman R, Jaafar H, Rahman M, Mui W. Chromatographic detection of phytohormones from the bacterial strain UPMP3 of *Pseudomonas aeruginosa* and UPMB3 of *Burkholderia cepacia* and their role in oil palm seedling growth. International Journal of Biotechnology Research. 2015;3(5):73-80. [https://www.researchgate.net/publication/286012705\\_Chromatographic\\_detection\\_of\\_phytohormones\\_from\\_the\\_bacterial\\_strain\\_UPMP3\\_of\\_Pseudomonas\\_aeruginosa\\_and\\_UPMB3\\_of\\_Burkholderia\\_cepacia\\_and\\_their\\_role\\_in\\_oil\\_palm\\_seedling\\_growth](https://www.researchgate.net/publication/286012705_Chromatographic_detection_of_phytohormones_from_the_bacterial_strain_UPMP3_of_Pseudomonas_aeruginosa_and_UPMB3_of_Burkholderia_cepacia_and_their_role_in_oil_palm_seedling_growth)
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the 8 Folin phenol reagent. J Biol Chem. 1951;193:266-75. [https://doi.org/10.1016/S0021-9258\(19\)52451-6](https://doi.org/10.1016/S0021-9258(19)52451-6)
- Arnon DI. Copper enzyme in isolated chloroplasts. Polyphenoloxidase in Beta 8 vulgaris. Plant Physiol. 1949;24:1-15. <https://doi.org/10.1104/pp.24.1.1>
- Craigie JS. Seaweed extract stimuli in plant science and agriculture. Journal of Applied Phycology. 2011 Jun;23(3):371-93. <https://doi.org/10.1007/s10811-010-9560-4>
- Jayaraman J, Norrie J, Punja ZK. Commercial extract from the brown seaweed *Ascophyllum nodosum* reduces fungal diseases in greenhouse cucumber. Journal of Applied Phycology. 2011 Jun;23(3):353-61. <https://doi.org/10.1007/s10811-010-9547-1>
- Mohanty D, Adhikary SP, Chattopadhyay GN, Prakasam VR, Gopamma D, Ramoji G, Srinivas N, Mukherjee J, Ray S, Gantait W, Bhattacharya T. Seaweed liquid fertilizer (SLF) and its role in agriculture productivity. Ecoscan. 2013;3:147-55.
- Devi SD, Paul JJ. Influence of seaweed liquid fertilizer of *Gracilaria dura* (Ag.) J. Ag. (red seaweed) on *Vigna radiata* (L.) R. Wilczek. in Thoothukudi, Tamil Nadu, India. World Journal of Pharmaceutical Research. 2014;3(4):968-78. [http://www.wjpr.net/dashboard/abstract\\_id/995](http://www.wjpr.net/dashboard/abstract_id/995)
- Crouch IJ, Van Staden J. Evidence for the presence of plant growth regulators in commercial seaweed products. Plant Growth Regulation. 1993 May;13(1):21-29. <https://doi.org/10.1007/BF00207588>
- Vasilov SP. The effect of photosynthesis parameters on leaf lifespan. Biology Bulletin Reviews. 2016 Jan;6(1):96-112. <https://doi.org/10.1134/S2079086416010084>
- Sridhar S, Rengasamy R. Studies on the effect of seaweed liquid fertilizer on the flowering plant *Tagetes erecta* in field trial. Advances in Bioresearch. 2010;1(2):29-34. <http://www.soegra.com/abr/vol2/5.pdf>
- Arthur GD, Stirk WA, Van Staden J, Scott P. Effect of a seaweed concentrate on the growth and yield of three varieties of *Capsicum annum*. South African Journal of Botany. 2003 Jul 1;69(2):207-11. [https://doi.org/10.1016/S0254-6299\(15\)30348-3](https://doi.org/10.1016/S0254-6299(15)30348-3)
- Spinelli F, Fiori G, Noferini M, Sproccati M, Costa G. A novel type of seaweed extract as a natural alternative to the use of iron chelates in strawberry production. Scientia Horticulturae. 2010 Jun 28;125(3):263-69. <https://doi.org/10.1016/j.scienta.2010.03.011>
- Yokoya NS, Stirk WA, Van Staden J, Novák O, Turečková V, Pěňčík A, Strnad M. Endogenous cytokinins, auxins and abscisic acid in red algae from Brazil 1. Journal of Phycology. 2010 Dec;46(6):1198-205. <https://doi.org/10.1111/j.1529-8817.2010.00898.x>
- Stirk WA, Arthur GD, Lourens AF, Novak O, Strnad M, Van Staden J. Changes in cytokinin and auxin concentrations in seaweed concentrates when stored at an elevated temperature. Journal of Applied Phycology. 2004 Feb;16(1):31-39. <https://doi.org/10.1023/B:JAPH.0000019057.45363.f5>
- Michalak I, Marycz K, Basińska K, Chojnacka K. Using SEM-EDX and ICP-OES to investigate the elemental composition of green



- macroalga *Vaucheria sessilis*. The Scientific World Journal. 2014 Oct;2014. <https://doi.org/10.1155/2014/891928>
31. Mathur C, Rai S, Sase N, Krish S, Jayasri MA. *Enteromorpha intestinalis* derived seaweed liquid fertilizers as prospective biostimulant for *Glycine max*. Brazilian Archives of Biology and Technology. 2015 Dec;58(6):813-20. <https://doi.org/10.1590/S1516-89132015060304>
32. Stirk WA, Novák O, Hradecká V, Pěňčík A, Rolčík J, Strnad M, Van Staden J. Endogenous cytokinins, auxins and abscisic acid in *Ulva fasciata* (Chlorophyta) and *Dictyota humifusa* (Phaeophyta): towards understanding their biosynthesis and homeostasis. European Journal of Phycology. 2009 May 1;44(2):231-40. <https://doi.org/10.1080/09670260802573717>

§§§