

Research Communication

## Moss flora of Rajasthan and Punjab plains

K. K. Rawat<sup>\*1</sup>, Afroz Alam<sup>2</sup>, P. K. Verma<sup>3</sup>

<sup>1</sup> CSIR-National Botanical Research Institute, Lucknow (India)

<sup>2</sup> Department of Bioscience and Biotechnology, Banasthali University, Rajasthan (India)

<sup>3</sup> Forest Research Institute, Dehra Dun (India)

*Article history*

Received: 16 September 2015

Accepted: 19 September 2015

Published online: 1 October 2015

© Rawat *et al.* (2015)

*Editors*

Afroz Alam

Dipjyoti Chakraborty

*Publisher*

Horizon e-Publishing Group

*Corresponding Author*

K. K. Rawat

✉ [drkkrawat@rediffmail.com](mailto:drkkrawat@rediffmail.com)

**Abstract**

An updated account of 51 taxa of mosses of Rajasthan and Punjab plains is provided along with a note on delimitation of boundaries of this unique bryo-geographical zone. Family Pottiaceae is most dominant and diversified with 17 taxa under 11 genera while genus *Fissidens* (Fissidentaceae) has the maximum diversity with 7 species followed by *Physcomitrium* (Funariaceae) and *Bryum* (Bryaceae) with 6 species each.

**Keywords**

Bryophytes; Flora; Moss; Punjab; Rajasthan

Rawat, K. K. Alam, A., Verma, P. K. 2015. Moss flora of Rajasthan and Punjab plains. *Plant Science Today* 2(4): 154-158. <http://dx.doi.org/10.14719/pst.2015.2.4.151>

### Introduction


Pande (1958) proposed floristic division of India into seven bryo-geographical zones to study the enormous bryo-floristic wealth of country namely (i) West Himalayas, (ii) East Himalayas, (iii) Gangetic plains (iv) Panjab and west Rajasthan plains (v) Central India (vi) Western Ghats and (vii) Eastern Ghats and Deccan plateau (Fig. 1). However, he made the division by thick lines, hence; significant problem arose while discussing the distribution of various taxa due to somewhat ill-defined boundaries of various zones (Alam *et al.*, 2015). In our earlier attempt to solve this problem, boundary of 'Central Indian bryo-geographical zone' was proposed along with a checklist of mosses in the area (Alam *et al.*, 2015). In continuation of the same, the present paper not only proposes the boundaries of 'Rajasthan and Punjab plains bryo-geographic zone' for ease in distributional analysis but also provides updated account of mosses present in this zone.

Instead of west Rajasthan (as named by Pande) it is recommended that entire Rajasthan except south

Rajasthan (Fig. 2) to be included in this zone. South Rajasthan is a part of Central Indian zone (Alam *et al.*, 2015) and therefore excluded. The Rajasthan plain is a unique zone particularly due to presence of 'Thar desert' having its own exclusive desert vegetation. Owing to the dry climate and desert conditions, very low number of bryophyte has been reported from this area.

On the other hand, the Punjab plains broadly consists present Punjab and Haryana states. This part of India is made up of alluvial soil brought by Indus and its tributaries, harbouring its own unique flora. However, some parts of this zone contain highlands (particularly Dhar tehsil in Pathankot district of Punjab and Morni hills in Panchkula district of Haryana) which need to be treated under western Himalayan zone. Kashyap (1929, 1932) though published his monumental work on "Liverworts of Western Himalaya and Punjab plains", however; so far the moss flora of this zone has remain neglected.

Furthermore, the status of Delhi is also questionable. Though treated as a part of Gangetic plains, geo-physical position of Delhi favours its



**Fig. 1.** Bryo-geographical territories of India proposed by Pande (1958) [Figure from "The Journal of the Indian Botanical Society 37(1):10, 1958"]

placement in Punjab plains along with Punjab and Haryana, as it is surrounded from three sides by Haryana. Yamuna river, which marks the western boundary of Gangetic plains as well as Delhi ridges (northern most end of Aravali hills) also separate most of the area of Delhi from Gangetic plains. However, East Delhi or Shahdara zone dissected out by river Yamuna, need to be treated within Gangetic plains.

Hence, in present work we propose Rajasthan (except south Rajasthan including Banswara, Sirohi, Udaipur, Rajsamand and Pali districts), Punjab (except Dhar tehsil of Pathankot district), Haryana (except Morni hills area in Panchkula district) and Delhi (except East Delhi) as parts of 'Rajasthan and Punjab plains' that should be considered as a single bryo-geographic zone.

By and large, this zone has been usually ignored in comparison to other bryo-geographic zones. Rajasthan, the most studied area, too had most of the studies focused in south Rajasthan (Central Indian zone), hence bryological records from other parts of the zone are negligible or meager. The present document provides updated accounts of 57 taxa of mosses from this zone. Distributional details with relevant references are also given, which may be useful for future workers and will encourage bryo-floristic studies in neglected areas.

## Materials and Methods

The present compilation is based on the all available literature till date from above mentioned bryo-geographical zone. The taxa reported without species

epithet, have been excluded. The accepted names are cited in bold. Format of the enumeration of taxa follows Alam et al (2015). The classification scheme follows Goffinet et al. (2008).

### Summary of mosses of Rajasthan and Punjab plains:

**Family Amblystegiaceae:** *Drepanocladus* (1)

**Family Bartramiaceae:** *Philonotis* (1)

**Family Bruchiaceae:** *Trematodon* (3)

**Family Bryaceae:** *Brachymenium* (1) *Bryum* (6)

**Family Ditrichaceae:** *Ceratodon* (1)

**Family Fissidentaceae:** *Fissidens* (7)

**Family Funariaceae:** *Entosthodon* (1) *Funaria* (1)  
*Loiseaubryum* (1) *Physcomitrium* (6)

**Family Hypnaceae:** *Orthothecium* (1)

**Family Leskeaceae:** *Leskea* (1)

**Family Myrinaceae:** *Schwetskeopsis* (1)

**Family Pottiaceae:** *Anoectangium* (1) *Barbula* (2)  
*Gymnostomiella* (1) *Gymnostomum* (1) *Hydrogonium* (3) *Hymenostylium* (1) *Hyophila* (4) *Pseudocrossidium* (1) *Semibarbula* (1) *Timmella* (1) *Tortula* (1)

**Family Splachnobryaceae:** *Splachnobryum* (1)

**Family Timmiaceae:** *Timmia* (1)

### Enumeration of taxa:

***Anoectangium strachyanum*** Mitt., Musc. Ind. Or.: 31, 1859. RAJASTHAN: Ranthambore (Alam et al., 2011)



**Fig. 2. Map of India showing proposed extent of 'Rajasthan and Punjab plains' bryo-geographical zone**

***Barbula indica*** (Hook.) Spreng., Nomenc Bot. 2: 72, 1824. RAJASTHAN (Lal, 2005)

***Barbula javanica*** Doz. & Molk. in Ann. Sci. Nat. Bot. ser. 3,2: 300, 1884. RAJASTHAN: Ranthambore (Alam et al., 2011)

***Brachymenium exile*** (Dozy & Molk.) Bosch & Sande Lac., Bryol. Jav. 1: 139, 1860. RAJASTHAN: Ajmer, Pali-near Harganga, (Chaudhary and Deora, 1993) DELHI (Chopra and Kanta, 1966)

***Bryum bicolor*** Dicks., Pl. Crypt. Brit. Fasc. 4: 16, 1801. RAJASTHAN: Jaipur (Bansal and Nath, 2014)

***Bryum bornholmense*** Wink. & Ruthe, Hedwigia 38: 120, 1899. RAJASTHAN: Bikaner (Bansal and Nath, 2014)

***Bryum bornholmense*** Wink. & Ruthe, Hedwigia 38: 120, 1899. RAJASTHAN: Bikaner (Bansal and Nath, 2014)

***Bryum coronatum*** Schwägr. Schwaegr., Sp. Musc. Frond, suppl. 1(2): 103, 1816. PUNJAB: Gurdaspur, Hoshiarpur (Singh et al., 2006) DELHI (Chopra and Kanta, 1966)

***Bryum dichotomum*** Hedw., Sp. Musc. Frond. 183, 1801. RAJASTHAN: Jaipur (Bansal and Nath, 2014)

***Bryum kliggraeffii*** Schimp., in Klinggr., Höh. Crypt. Preuss. 81, 1858. RAJASTHAN: Alwar, Bikaner, Jaipur. (Bansal and Nath, 2014) DELHI (Chopra and Kanta, 1966)

***Ceratodon purpureus*** (Hedw.) Brid., Bryol. Univ. 1: 480, 1826. DELHI (Chopra and Kanta, 1966)

***Drepanocladus uncinatus*** (Hedw.) Warnst., Beih. Bot. Centralbl. 13: 417, 1903. PUNJAB (Lal, 2005)

***Entosthodon nutans*** Mitt., J. Proc. Linn. Soc. Bot., suppl. 1: 55, 1859. PUNJAB and RAJASTHAN (Lal, 2005)

- Fissidens bryoides** Hedw., Sp. Musc. Frond. 153, 1801. DELHI (Chopra and Kanta, 1966)
- Fissidens ceylonensis** Dozy & Molk., Ann. Sci. Nat., Bot., ser. 3, 2: 304, 1844. PUNJAB: Ropar (Singh et al., 2006)
- Fissidens crispulus** Brid., Muscol. Recent Suppl. 4: 187, 1819. RAJASTHAN: Alwar; Kota (Bapna, 1980 and Chaudhary and Deora, 1993 as *Fissidens sylvaticus*) Bhilwara (Chaudhary and Deora, 1993 as *Fissidens sylvaticus*)
- Fissidens curvato-involutus* Dix., Notes Roy. Bot. Gard. Edinburgh 19: 279, 1938. — **Fissidens involutus** ssp. *curvato-involutus* (Dix.) Gangulee
- Fissidens diversifolius** Mitt., J. Linn. Soc. Bot., suppl. 1:140, 1859. RAJASTHAN: Kumarwara (Bapna, 1980) Sawai Madhopur, Kota, (Chaudhary and Deora, 1993)
- Fissidens gemminiflorus** Doz. et Molk. var. *nagasakiensis* (Besch.) Iwats. Jour. Hattori Bot. Lab. 32: 272, 1969. RAJASTHAN: Gorham ghat, Kota-Dhara hills; (Bapna, 1980) Gorham ghat, Bhilwara, (Chaudhary and Deora, 1993)
- Fissidens involutus** ssp. *curvato-involutus* (Dix.) Gangulee, Mosses E India 2: 548, 1971. RAJASTHAN: Ajmer, Chittorgarh, Jaipur (Bapna, 1980; Chaudhary and Deora, 1993 as *Fissidens curvato-involutus* Dix.)
- Fissidens sylvaticus* Griff., Calcutta J. Nat. Hist. 2: 507, 1842. — **Fissidens crispulus** Brid.
- Fissidens taxifolius** Hedw., Sp. Musc. Frond. 155, 1801. DELHI (Chopra and Kanta, 1966)
- Funaria hygrometrica** Hedw. Spec. Musc. 172, 1801. PUNJAB: Ropar (Singh et al., 2006), RAJASTHAN: common throughout Rajasthan, Bhilwara (Chaudhary and Deora, 1993), Ganganagar (Choyal and Sharma, 2011) DELHI (Chopra and Kanta, 1966)
- Funaria nutans* (Mitt.) Broth. — **Loiseaubryum nutans** (Mitt.) Broth.
- Gymnostomiella vernicosa** (Hook. ex Harv.) M. Fleisch., Musci Buitenzorg 1: 310, 1904. RAJASTHAN: Ajmer (Chaudhary and Deora, 1993), Ranthambore (Alam et al., 2011), DELHI (Chopra and Kanta, 1966)
- Gymnostomum aeruginosum** Sm., Fl. Brit. 3: 1163, 1804. RAJASTHAN (Lal, 2005)
- Hydrogonium consanguineum** (Thwaites & Mitt.) Hilp., Beih. Bot. Centralbl., Abt. 2, 50(2): 626, 1933. RAJASTHAN: Gangangar, Jhunjhunu, (Chaudhary and Deora, 1993)
- Hydrogonium gangeticum** (Müll. Hal.) P.C. Chen, Hedwigia 80: 237, 1941. DELHI (Chopra and Kanta, 1966)
- Hydrogonium javanicum** (Dozy & Molk.) Hilp., Beih. Bot. Centralbl., Abt. 2, 50(2): 632, 1933. PUNJAB and RAJASTHAN (Lal, 2005)
- Hymenostylium recurvirostrum** (Hedw.) Dixon, Rev. Bryol. Lichenol., 6: 96, 1933. RAJASTHAN: Kota (Bapna and Chaudhary, 1989), PUNJAB (Lal, 2005) DELHI (Chopra and Kanta, 1966)
- Hyophila comosa* Dixon et P. de La Varde, arch. Bot. 1: 166, 1927. — **Hyophila rosea** Williams
- Hyophila involuta** (Hoop.) A. Jaegr., Ber. S. Gall. Naturew. Ges. 1871-72: 356, 1873. PUNJAB: Gurdaspur (Singh et al., 2006) RAJASTHAN: Alwar, Jaipur, Kota (Bapna and Chaudhary, 1989), Jhunjhunu (Chaudhary and Deora, 1993), Ranthambore (Alam et al., 2011).
- Hyophila nymaniana** (M. Fleisch.) M. Menzel, 22: 198, 1992. RAJASTHAN: Ranthambore (Alam et al., 2011)
- Hyophila rosea** Williams, Bull. N.Y. Bot. Gard. 8: 341, 1941. RAJASTHAN: Menal near Bhilwara (Chaudhary and Deora, 1993)
- Hyophila spathulata** (Harv.) A. Jaeger., Ber. Thätigk. St. Gallischen Naturwiss. Ges. 1871-72: 353, 1873. DELHI (Chopra and Kanta, 1966) Ranthambore (Alam et al., 2011)
- Leskea nervosa** (Brid.) Myrin., Coroll. Fl. Upsal. 52, 1834. PUNJAB (Lal, 2005)
- Loiseaubryum nutans** (Mitt.) Fife., J. Hattori Bot. Lab. 58: 192, 1985. PUNJAB (Chaudhary and Deora, 1993 as *Funaria nutans*) RAJASTHAN: Jaipur – Sawai Madhopur (Robinson, 1968 as *Funaria nutans*)
- Orthothecium intricatum** (Hartm.) Schimp., Bryol. Eur. 5: 108, 1851. Punjab (Lal, 2005)
- Philonotis angusta** Mitt., J. Proc. Linn. Soc., Bot., suppl. 1: 61, 1859. PUNJAB: Gurdaspur (Singh et al., 2006)
- Physcomitrium coorgens** Broth., Rec. Bot. Surv. India 1(12): 319, 1899. DELHI (Chopra and Kanta, 1966; Lal, 2007)
- Physcomitrium cyathicarpum* Mitt., J. Proc. Linn. Soc., Bot., suppl. 1: 54, 1859 — **Physcomitrium immersum** Sull.
- Physcomitrium eurystomum** Sendtn., Denkschr. Bayer. Bot. Ges. Regensburg 3: 142, 1841. PUNJAB: Hoshiarpur (Singh et al., 2006), RAJASTHAN (Lal, 2005)
- Physcomitrium immersum** Sull., Manual 648, 1848. RAJASTHAN: Jaipur – Sawai Madhopur (Robinson, 1968 as *Physcomitrium cyathicarpum*), PUNJAB (Lal, 2005 as *Physcomitrium cyathicarpum*) DELHI (Chopra and Kanta, 1966 as *Physcomitrium cyathicarpum*)
- Physcomitrium japonicum** (Hedw.) Mitt., Trans. Linn. Soc. London, Bot. 3: 164, 1891. RAJASTHAN: Alwar (Chaudhary and Deora, 1993)
- Physcomitrium delicatum** H.A. Crum and L.E. Anderson, Bryologist 58: 1, 1955. DELHI (Lal and Menon 1971)
- Physcomitrium pulchellum** (Griff.) Mitt., J. Proc. Linn. Soc., Bot. suppl. 1: 54, 1859. DELHI (Vashistha and Chopra, 1983)
- Pseudocrossidium porphyreoneurum** (Müll. Hal.) R.H. Zander, Bull. Buffalo Soc. Nat. Sci. 32: 119, 1993. RAJASTHAN (Lal, 2005)
- Schwetschkeopsis formosana** Nog., J. Hattori Bot. Lab., 5: 41, 1951. PUNJAB: Gurdaspur (Singh et al., 2006)
- Semibarbula orientalis** (F. Weber) Wijk. & Margad., Taxon 8: 75, 1959. RAJASTHAN: Alwar (Bapna

- and Choudhary, 1989; Choudhary and Deora, 1993) DELHI (Chopra and Kanta, 1966)
- Splachnobryum flaccidum* (Harv.) Braithw., Grevillea 1(2): 28, 1872 → *Splachnobryum obtusum* (Brid.) Müll. Hal.
- Splachnobryum indicum* Hampe & Müll. Hal., Linnaea 37: 174, 1972 → *Splachnobryum obtusum* (Brid.) Müll. Hal.
- Splachnobryum obtusum* (Brid.) Müll. Hal., Verh. K.K. Zool.-Bot. Ges. Wien 19: 504, 1869. DELHI (Chopra and Kanta, 1966 as *Splachnobryum indicum* and *Splachnobryum flaccidum*)
- Timmia megapolitana* Hedw., Sp. Musc. Frond. 176, 1801. PUNJAB: Gurdaspur (Singh et al., 2006)
- Timmiella anomala* (Bruch & Schimp.) Limpr., Laubm. Deutschl. 1: 592, 188. RAJASTHAN: Ajmer, Chittorgarh, Jaipur, Jodhpur-Mandore, Sawai Madhopur (Bapna and Choudhary, 1989; Chaudhary and Deora, 1993).
- Tortula muralis* Hedw., Sp. Musc. Frond. 123, 1801. RAJASTHAN: Ajmer, Bikaner, Chittorgarh, Jodhpur, Jaipur, Sawai Madhopur (Bapna and Choudhary, 1989; Chaudhary and Deora, 1993)
- Trematodon brevicalyx* Dixon., J. Indian bot. Soc. 4: 287-97, 1925. DELHI (Vashistha and Chopra, 1983)
- Trematodon longicollis* Michx., Fl. Bor.-Amer 2: 289, 1803. PUNJAB: Gurdaspur (Singh et al., 2006)
- Trematodon subulosus* Griff., Cal. J. Nat. hist. Soc. 2: 493, 1842. RAJASTHAN: Ganganagar (Chaudhary and Deora, 1993) PUNJAB (Lal, 2005)
- Biology*, Goffinet, B. and Shaw, A.J. (eds.) 2<sup>nd</sup> edition, Cambridge University Press. pp. 55-138.
- Kashyap, S. R. 1929. *Liverworts of western Himalayas and Panjab plains*. Vol. 1, University of Panjab, Lahore. 129 Pp.
- Kashyap, S. R., and R. N. Chopra. 1932. *Liverworts of western Himalayas and Panjab plains*. Vol. 2, University of Panjab, Lahore. 137 pp.
- Lal, M. and M. K. Menon. 1971. *Physcomitrium delicatum*, new to the moss flora of India. *Bryologist* 74: 51-54.
- Lal, J. 2005. *A checklist of Indian mosses*. Bishen Singh Mahendra Pal Singh, Dehradun. 164 pp.
- Pande, S. K. 1958. Some aspects of Indian Hepaticology. *J Indian Bot Soc* 37(1): 1-27.
- Singh, D. K., S. K. Singh, and S. Kumar. 2006. Bryophyta. In: *Biodiversity in the Shivalik Ecosystem of Punjab*. Jerath N, Puja and Chadha J (Eds.) Punjab State Council for Science and Technology, Chandigarh. Bishen Singh Mahendra Pal Singh, Dehradun, pp. 234-246.
- Vashistha, B. D. and R. N. Chopra 1983. Addition to the moss flora of North-Western India. *Misc Bryol Lichenol* 9(8): 177-179.



## Acknowledgements

The authors are thankful to Drs. A.K. Asthana and Vinay Sahu, Bryology laboratory, CSIR-National Botanical Research Institute, Lucknow One of the authors (AA) is also grateful to Professor Aditya Shastri, Vice Chancellor and Professor Vinay Sharma, Dean, Faculty of Science and Technology, Banasthali University, Rajasthan, India, and Mr. Sonu Yadav, Bryology Unit, Lucknow University for their kind support for this research work.

## References

- Alam, A., K. K., Rawat, P. K. Verma, V. Sharma, and D. Sengupta. 2015. Moss flora of Central India. *Plant Sci Today* (In Press).
- Alam, A., V. Sharma, and S. C. Sharma. 2011. Bryoflora of Ranthambhore Tiger Reserve, Rajasthan (India). *Arch Bryol* 106: 1-8.
- Bansal, P. and V. Nath. 2014. Present status of genus *Bryum* Hedw., (Bryophyta) in Rajasthan, India. *G-Jour Environ Sci and Technol* 1(6): 118-122.
- Bapna, K. R. 1980. Occurrence of *Fissidens* Hedw. in Rajasthan. *J Indian bot Soc* 59: 20-26.
- Chaudhary, B. L. and G. S. Deora 1993. *Moss flora of Rajasthan (India)*. Himanshu Publication, Delhi, India. 127 pp.
- Chopra, R. N. and V. Kanta. 1966. Some mosses from Delhi, India. *Bryologist* 69(4): 509-510.
- Choyal, R. and S. K. Sharma. 2011. Studies on Bryophytes of Thar desert with particular reference to Ganganagar district (Rajasthan), India. *Ind J Fundl Appl Life Sci* 1(3): 239-245.
- Goffinet, B., W. R. Buck, and A. J. Shaw. 2008. Morphology and classification of the Bryophyta. In: *Bryophyte*