



RESEARCH ARTICLE

Assessment of phthalate esters (PAEs) contamination in *Urtica dioica* L.

Priyanka Rajput* & R. A. Sharma

Department of Botany, University of Rajasthan, Jaipur, PIN- 302 004, Rajasthan, India

*Email: rajputpriyanka82@gmail.com

 OPEN ACCESS

ARTICLE HISTORY

Received: 28 September 2021

Accepted: 01 April 2022

Available online

Version 1.0: 10 May 2022



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, etc. See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Rajput P, Sharma RA. Assessment of phthalate esters (PAEs) contamination in *Urtica dioica* L. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.1521>

Abstract

Phthalate esters are a group of chemical compounds of ubiquitous nature which nowadays have become a colossal threat to the environment, human-animal and plant health, because of its higher potential of accumulation in soil and aquatic habitat leading to environmental contamination due to its widespread industrial and agricultural usage. The present research aims to analyze the phthalate esters accumulation in *Urtica dioica* L. For this study, the *Urtica dioica* is tested for the presence of phthalates by using Gas Chromatography-Mass spectrometry. The Gas Chromatography-Mass spectrometry observations show the presence of 11 phthalate esters, among which diethyl phthalate (DEP) and bis(2-ethylhexyl) phthalate (BEHP), dibutyl phthalate (DBP), and diisobutyl phthalate (DIBP) were found to be in significantly higher amount. The sum concentrations of the phthalate ester in different extracts of plant range from 16.25% to 84.07%. The % composition of diethyl phthalate is found to be comparatively higher than other phthalate esters in methanolic extract of *Urtica dioica* while diisobutyl phthalate and bis (2-ethylhexyl) phthalate accumulation is found relatively higher in the ethyl acetate and diethyl ether fractions. The observations show the contamination of the *Urtica dioica* plant with phthalate esters and also indicate the phthalate accumulating potential of the plant.

Keywords

Phthalic acid esters, *Urtica dioica*, Gas Chromatography-Mass spectrometry, Diethyl phthalate, Bis (2-ethylhexyl) phthalate, Diisobutyl phthalate, Dibutyl phthalate

Introduction

Phthalic acid esters (PAEs) commonly called phthalates are a group of synthetic chemical compounds which are known as 1, 2-benzene dicarboxylate esters or dialkyl or alkyl aryl esters of *o*-phthalic acid (1, 2). PAEs have become a serious concern as they possess potential hazardous effects as an environmental contaminant due to their widespread usage with a total global production of 8M tonnes in 2011 (3). PAEs are used as a common plasticizer that is added to plastic polymers for increasing the strength, flexibility, transparency and durability of plastic (1, 4). The most common usage is found in toys, cosmetics, food and pesticides packaging, medical instruments and many other agricultural and household products (5-7). The PAEs remain free of any covalent bond binding it to the surrounding matrix, so it is readily accessible for leaching into the environment released from the higher molecular mass carbon chains (8-10) and spread itself ubiquitously in water, air and soil (11, 12). PAEs are considered among the priority pollu-

tants as it has been reported to cause oxidative stress, apoptosis and cytotoxicity via DNA damage (13-16). They are often grouped as compounds with endocrine disruptive potential, interfering with the normal process of synthesis, secretion and metabolism of hormones involved in sexual development, thus causing an adverse effect on reproductive health (17, 18). Other health issues attributed to PAEs are hyperactivity disorders, allergy, asthma, thyroid cancer and hypertension (19-22).

The plant chosen is collected from the roadsides, drainage areas (adjacent to the agricultural fields) from the Palampur district of Himachal Pradesh. Major sources of phthalate esters pollutant to be reaching the soil in Himalaya region of Himachal Pradesh is the increased protective cultural practices in the hill slopes, promoting the use of greenhouse/polyhouse farming practices. These polyhouses are made of cheaper polythene or plastic instead of glass on the greenhouse roof (23). Other major sources of PAEs pollutants are wastewater irrigation (24), use of agricultural chemicals and fertilizers (25, 26), and emission of industrial waste (27, 28). From the contaminated soil, the PAEs further get distributed in the environment through biogeochemical cycling (Fig. 1).




Fig. 1. Soil Contamination sources of PAEs.

The past few decades have witnessed a massive surge of inclination in the ideology of people towards the use of medicinal plants due to their easy availability, effectiveness with minimalistic side effects, less toxicity and long-term beneficial potentials (29-33). *Urtica dioica* L., a notorious weed of invasive nature comes under the least concern group of IUCN list, is a plant that has long been used for fiber, food and traditional medicine for its potential property in acting against arthritic conditions, musculoskeletal pain, inflammations and for its anti-diabetic, anti-hypertensive and cardiovascular potential (34-37). Although known for its traditional use in Urtication therapy for the treatment of numb joints in case of paralysis and rheumatoid arthritis (38, 39); the plant could not overcome its reputation and still found growing as a neglected weed species in barren lands, waste places, roadsides, neglected yards, stream banks and ditches. The plant grows best in moist soils that are rich in nutrients (40).

U. dioica is found widespread throughout the cooler, temperate climate of Europe, also occurs in North America, North Africa and parts of Asia (41). The species is although invasive in nature, nowadays growing stinging nettle has been encouraged by conservation groups of UK

wildlife as it supports more than 40 species of insects including tortoise shell, peacock butterflies and various kinds of caterpillars. Nettles serve as a food source for butterflies and moth larvae. The seeds produced from the plant make food source for the native birds. The plant is usually insect-prone and is useful as an insect feed for many types of caterpillars and thus has been deliberately planted (42).

Taxonomic classification of the plant (43, 44) is as follows:

Kingdom	:	Plantae
Sub-Kingdom	:	Tracheophyta
Superdivision	:	Spermatophyta
Division	:	Magnoliophyta
Class	:	Magnoliopsida
Subclass:	:	Hemamelididae
Order	:	Urticales
Family	:	Urticaceae
Genus	:	<i>Urtica</i>
Species	:	<i>dioica</i>

The ever-increasing contamination of agronomically important soil is becoming a serious problem as the presence of contaminants disrupts the normal functioning of biogeochemical balance in the environment. Physicochemical remediation technology has various limitations (45), hence an alternative of phytoremediation has been put forth using plants for removal of pollutants from the environment (46). Various plant species possess an inbuilt mechanism for detoxification of the xenobiotic compound. These plants have high toleration capacity and the potential to hyper accumulate toxic chemicals almost up to 1% of their weight. (47). *U. dioica* is also one such hyper-accumulating plant that has already been evaluated for its phytoremediation potential in eradicating the Cr metal stress (48) and soils contaminated with polychlorinated biphenyls (PCBs) (47). The main objective of the present study is determining the contamination in the phytoconstituent composition of *U. dioica* plant with phthalate esters and evaluation of type PAEs accumulated in the plant through GC-MS analysis.

Materials and Methods

Collection and Preparation of Plant extracts

The whole plant of *U. dioica* was collected from the roadsides and drainage areas (adjacent to the agricultural fields) of Palampur District of Himachal Pradesh, India. These areas are chosen deliberately to test the impact of employed agricultural practices on the phytochemical composition of the plants growing in the vicinity. A dried plant completely pressed in blotting papers was submitted for identification of the plant species and specimen no. PLP-18301 has been given as a herbarium specimen to IHBT Palampur. The collected plant material except the one to be provided for herbarium is at first rinsed in run-

ning tap water to remove the surface impurity and blotted dry to remove any type of excess moisture. The whole plant was then dissected into root, stem and leaves which were then allowed to dry in shade for 7 days, occasionally mixed and cut into smaller pieces. The dried stock samples were then powdered using a grinder and the powdered material was kept stored in an airtight container at 4 °C for future extraction. About 25 g of powdered sample was weighed and extracted with 200 ml methanol using Soxhlet extraction for methanolic (ME) extract until the plant material is completely exhausted and for the diethyl ether (DEM) and ethyl acetate (EAM) fraction of methanol extract, about 25 g of powdered sample is soxhlet extracted for 24 hrs in 80% methanol. Soxhlet extraction is carried out at a temperature of 64 °C for both sets. The solvent of the plant extracts was then evaporated in a hot air oven at 40 °C to obtain a viscous liquid. For the DEM and EAM extract 80% methanol concentrated dried extract was taken and re-extracted with pet ether, diethyl ether and ethyl acetate sequentially. The plant residue collected was discarded and the concentrated solvent extracts were transferred to vials of 15 ml sizes and the remaining solvent is further allowed to evaporate in the laboratory at room temperature. The remaining completely dried extract was weighed to evaluate the yield of the extract and then it is stored in a refrigerator at 4 °C until further analysis.

GC-MS Analysis

Gas Chromatography-Mass Spectrometry analysis of the methanolic plant extract was carried out on a Shimadzu QP-2010 Plus equipped with a Thermal desorption system TD 20, helium carrier gas, Rxi-5Sil MS column (30 m length, 0.25 mm ID, 0.25 µm thickness), Column Oven temperature of 60 °C for ME extract and 100 °C for DEM and EAM extracts, injector temperature was maintained at 260 °C, pressure 81.7 kPa for ME and 90.5 kPa for DEM and EAM extracts, total flow 16.3 ml/min, column flow 1.21 ml/min, linear velocity 40.1 cm/sec, purge flow 3.0 ml/min, the split ratio was 10.0, ion source temperature: 230 °C, Interface temperature: 270 °C, Oven temperature program: 60 °C (hold for 5 min) for ME and 100 °C for DEM and EAM with a hold time of 2 min. raised to 300 °C at the rate of 10 °C/min: ending with an isothermal temperature of 300 °C for 19 minutes, run time: 44.98 min for ME and 40 min for DEM and EAM extracts. The instrument was operated using GC-MS solutions software. Scan range was 40-650 m/z. For GC-MS analysis crude extract was re-dissolved in methanol to make a stock solution. 1 µl stock solution was prepared for GC-MS analysis. The relative abundance of the compound in the extract was calculated by comparing the area of average peak to the total area, software adopted to handle mass spectra and the chromatogram was turbo mass. The relative percentage of every constituent present in the extract was expressed as a percentage with their respective peak area. The phytoconstituents in the extracts were determined by the comparison of their retention time and mass weight with recorded authentic samples obtained by GC as well as from the mass spectra of NIST (National Institute of Standards and Technology, US) a database having 62000 patterns and Wiley pesticide library 3rd edition, for

estimating the presence of probable compounds in the extract, which is further cross-checked using PubChem online database and other research articles for the activities and structure of the detected compound.

Results and Discussion

The main aim of the present study is the extraction and evaluation of methanolic, ethyl acetate, and diethyl ether extract of root, stem and leaves of *U. dioica* for the presence and the extent of accumulation percentage of PAEs. The solvents used for the sample injection in GC-MS were pure with no phthalate contamination and as a precautionary measure, the solvents were put in the dried extract immediately before putting the sample into injection. So, the chance of introduction of any PAEs peak in the GC-MS chromatogram originating from the laboratory contamination is kept minimal.

The completely dried plant methanolic soxhlet extractions of root, leaves and stem of the plant were weighed for the total yielding capacity. The yield of the extract is measured by comparing the weight of the solvent extract with the sample weight of the powdered extract of the plant. The yield is measured in the amount of extract given by 100 g of the sample plant part weight (g/100 g dry weight). The result of the extraction yield obtained from each extract is compiled in Table 1. The methanolic extraction yield is found to be the highest in leaves among the tested plant parts and the root seems to be the one with the lowest yield. To make sure that the phthalate compounds source might not be from any sort of laboratory plastic-ware, all the steps and processes of extraction, storage and analysis were monitored for avoiding the use of any type of plastic container in the laboratory.

Table 1. Extraction Yield of different plant parts of *U. dioica*

Plant Part	Yield (g/100g dry weight)		
	ME	DEE	EAE
Root	6.086±0.0728	0.251±0.0049	3.74±0.001
Stem	7.612±0.0538	0.454±0.107	1.034±0.011
Leaves	10.098±0.109	0.13±0.017	2.34±0.006

ME: Methanol Extract, **DEE:** Diethyl ether Extract, **EAE:** Ethyl Acetate extract
*Each value is expressed as mean±standard deviation (n=3)

GC-MS chromatogram results of the different extracts of *U. dioica* exhibited the presence of about 11 PAEs compounds which are diisobutyl Phthalate, dibutyl phthalate, bis(2-ethylhexyl) phthalate, diethyl phthalate, dimethyl phthalate, diamyl phthalate, butyloctyl phthalate, 5-methylhex-2-yl isobutyl phthalate, 5-methylhex-2-yl butyl phthalate, dioctyl terephthalate, dimethoxyethyl phthalate. The phthalates compounds with major contamination in the *U. dioica* extract are illustrated in Fig. 2 with their molecular structure and their respective mass spectra. The GC-MS chromatograms of the extracts are provided in Fig. 3: Root, Fig. 4: Stem and Fig. 5: Leaves.

Leaves of *U. dioica* show the most considerable amount of contamination both in terms of the accumula-




Fig. 2. GC-MS mass spectra of four major phthalate contaminants in *U. dioica* **A.** Diisobutyl phthalate (DIBP), **B.** Diethyl phthalate (DEP), **C.** Dibutyl phthalate (DBP), **D.** Bis (2-ethylhexyl) phthalate (BEHP).




Fig. 3. Mass Chromatograms of *U. dioica* Root **A.** Methanol Extract, 1. DEP, 2. BEHP; **B.** Ethyl Acetate Extract 1. DEP, 2. DIBP, 3. BOP, 4. DBP, 5. BEHP; **C.** Diethyl ether Extract, 1. DIBP, 2. DBP, 3. BEHP, 4. DOTP.




Fig. 4. Mass Chromatograms of *U. dioica* Stem **A.** Methanol Extract, **1.**DMP, **2.** DEP, **3.** BOP, BEHP; **B.** Ethyl Acetate Extract **1.** DIBP, **2.** DBP, **3.** BOP, **4.** BEHP; **C.** Diethyl ether Extract, **1.** DIBP, **2.** DBP, **3.** BEHP.






Fig. 5. Mass Chromatograms of *U. dioica* Leaves **A.** Methanol Extract, **1.** DMP, **2.** DEP, **3.** DBP, **4.** BEHP; **B.** Ethyl Acetate Extract, **1.** DIBP, **2.** MHIBP, **3.** BOP, **4.** DMEP, **5.** BEHP; **C.** Diethyl ether Extract, **1.** DIBP, **2.** MHIBP, **3.** DBP, **4.** MHBP, **5.** DAP, **6.** BOP, **7.** PPP, **8.** BEHP; **1.** DIBP, **2.** DBP, **3.** BEHP.

tion % as well as in terms of various types of phthalate ester observed from the GC-MS results obtained. The diethyl ether extract of the leaves exhibit the highest accumulation of phthalate compounds taking up about 84.07% peak area of the resultant chromatogram. DEP presence is observed only in the methanolic fraction of all plant parts in significantly higher amounts and only in trace amounts from the ethyl acetate fraction of the root. The methanolic leaves extract exhibits the highest % of peak area covered up by DEP.

The comparative phthalate ester contamination in various extracts of *U. dioica* is illustrated in Fig. 6. A relatively higher concentration of DBP and DIBP is found in the diethyl ether and ethyl acetate extract; while BEHP shows its accumulation in all the extracts in significant quantities.

Previous GC-MS analysis conducted on the hexane and aqueous extract of *U. dioica* leaves reported the presence of dibutyl phthalate (DBP) and diisooctyl phthalate esters and were observed to be responsible for imparting antimicrobial properties to the plant. (49-52). There is no further evidence and not much research has been conducted on testing the phthalate esters being an indigenous plant metabolite in plant *U. dioica*.




Fig. 6. Percentage Contamination of the extracts of *U. dioica* by phthalate esters. [LDE: Leaf diethyl ether extract; LEA: Leaf ethyl acetate; LM: Leaf Methanol; SDE: Stem diethyl ether extract; SEA: Stem ethyl acetate; SM: Stem Methanol; RDE: Root diethyl ether extract; REA: Root ethyl acetate; RM: Root Methanol; Diethyl phthalate (DEP), Dimethyl phthalate (DMP), DBP: Dibutyl phthalate, BEHP: Bis (2-ethylhexyl) phthalate, Diisobutyl phthalate, MHIBP: 5-Methyl hex-2-yl isobutyl phthalate, MHBP: 5-methylhex-2-yl butyl phthalate, BOP: Bis-octyl phthalate, DAP: Diallyl phthalate, PPP: Poly-propylene phthalate, DMPP: Di-(methoxyethyl) phthalate, DOTP: Dioctyl terephthalate].

The detection of mentioned PAE compounds revealed the ability of the plant to absorb these contaminants either from the contaminated soil or the polluted water source, keeping in mind that the plasticizer contamination while handling, storage and extraction is kept minimal. The discussed plant grows as a common weed in the mountain region situated near floating river beds with rich soil. The major source of such contaminations in the soil or the draining water is the absence of proper waste plastic disposal or recycles and the increasing plastic warehouse cultivation practices and use of mulching films in agriculture also increases the chances of leaching of PAEs into the soil and further to the groundwater sources (53, 54). Among the commonly used phthalate esters, DMP shows the highest water solubility followed by DEP (55). BEHP is also reported to be slightly water-soluble in nature making the phthalate ester capable of easily leaching and contaminating the water sources (56-59). The prime cause of phthalate pollution in the environment remains the neglect of the proper waste disposal resulting in a manifold increase in phthalate pollutant percentage in the environment (60).

Phthalates constant release into the environment has raised concern about the deteriorating effect of phthalates on human and animal health (61-65). Toxicological study on PAEs reported DEHP with endocrine disruption effect as well as cytotoxicity in various types of cells while DBP is reported to significantly reduce the cell viability. Phthalates that are reported as reproductive as well as developmental toxicants are DiBP (66-68), DEHP

(69-71), DiNP (66, 69), BBzP (72, 73). Phthalates temper with the normal production and functioning of follicle-stimulating hormone, testicular testosterone (fetal) (74). Major critical ill effects can be seen as the structural and functional disruption of male reproduction (75, 76). Phthalate toxicity also shows effects like retention of nipples/areola in male rodents and a reduced anogenital distance; this marks the first sign of feminization and demasculinization (49, 73). This group of symptoms in animals is known as "phthalate syndrome" (75, 77).

Previous research reported phthalate esters to cause adverse effects on the reproductive health of humans have recently been surveyed which concluded women to have relatively broader exposure risk in day-to-day life compared to men as they are more exposed to use cosmetics and personal care products that uses phthalate in their packaging and storage (78). Although phthalate pollution poses a greater threat upon humans and animals' health, the phyto composition of food crops and medicinal plants are also getting greatly affected due to phthalate pollution in the environment. There have been reports on the higher accumulation of PAEs in three staple food crops *Triticum aestivum*, *Brassica napus*, *Zea mays* (48). In some practical experiments, the researchers used the PAEs accumulation potential of the plant *Benincasa hispida* to eradicate and reduce the risk of absorption of DEHP by the planted vegetable crop, thus adapting a new way for eradicating the PAEs removal from the agriculture fields (79).

Conclusion

The present study reported the phthalate esters accumulation in *U. dioica*. The toxic compound that has been adapted for use as a plasticizer in agriculture, pharmaceutical, cosmetic, personal care product, industrial uses has an easy pathway to get leach from the sources and getting mixed into soil and water sources contaminating them. From here these pollutants get absorbed by the plants through roots. Pollution of medicinal and food crop plants has become a serious problem in the developing countries which requires special attention in standardization and restriction of PAEs plasticizers and increase in the capacity of production as well as promoting and making people aware about using phthalate-free plasticizers. Proper toxicological evaluation and examination of medicinal plants are recommended to determine the extent of phthalate accumulation in the plant before taking into consideration for use as an herbal drug to avoid creating an adverse effect. Although the restriction in usage of PAEs remains the first step, but finding new ways for the eradication of such toxic compounds from the soil and water sources may also have benefits.

Acknowledgements

The presented research is the result of practical experiments and in-lab study and no- grants have been received for the work. The author is thankful for the cooperation of director AIRF, JNU and director of IHBT, CSIR, Palampur.

Authors contributions

All the authors have read and approved the final version of the article. The research conducted and the analysis done was solely done by the corresponding author under the guidance of Dr. RA Sharma (the second author).

Compliance with ethical standards

Conflict of interest: The authors declare no conflict of interest to the presented research.

Ethical issues: None

References

- Wittasek M, Koch HM, Angerer J, Bruning T. Assessing exposure to phthalates- The human biomonitoring approach. *Molecular Nutrition and Food Research*. 2011; Jan;55(1):7-31. <https://doi.org/10.1002/mnfr.201000121>
- Saednia S. Phthalates. *Encyclopedia of Toxicology*. 2014; 3: 928-33. <https://doi.org/10.1016/B978-0-12-386454-3.00963-5>
- Li R, Liang J, Gong Z, Zhang N, Duan H. Occurrence and spatial distribution, historical trend and ecological risk of phthalate esters in the Jiulong River, southeast China. *Science of the Total Environment*. 2017; Feb;580:388-97. <https://doi.org/10.1016/j.scitotenv.2016.11.190>
- Arfaenia H, Fazlzadh M, Taghizadeh F, Saeedi R, Spitz J, Do-baradaran S. Phthalate acid esters (PAEs) accumulation in coastal sediments from regions with different land-use configuration along the Persian Gulf. *Ecotoxicology and Environmental Safety*. 2019; Mar;169:496-506. <https://doi.org/10.1016/j.ecoenv.2018.11.033>
- Wang W, Leung AOW, Chu LH, Wong MH. Phthalates contamination in China: status, trend and human exposure with an emphasis on oral intake. *Environmental Pollution*. 2018; Jul. 1;238:771-82. <https://doi.org/10.1016/j.envpol.2018.02.088>
- Mi L, Xie Z, Zhao Z, Zhong M, Mi W. Occurrence and spatial distribution of phthalate esters in sediments of the Bohai and Yellow seas. *Science of the Total Environment*. 2019; Feb. 25;653:792-800. <https://doi.org/10.1016/j.scitotenv.2018.10.438>
- Zhou B, Zhao L, Sun Y, Li X, Weng L, Li Y. Contamination and human health risks of phthalate esters in vegetable and crop soils from the Huang-Huai-Hai region of China. *Science of the Total Environment*. 2021; Jul. 15;778:146281. <https://doi.org/10.1016/j.scitotenv.2021.146281>
- Clara M, Windhofer G, Hartt W, Braun k, Simon M, Gans O, Scheffknecht C, Chovanek. Occurrence of phthalates in surface runoff, untreated and treated wastewater fate during wastewater treatment, *Chemosphere*. 2010; Feb. 1;78(9):1078-84. <https://doi.org/10.1016/j.chemosphere.2009.12.052>
- Gong M, Weschler CJ, Zhang Y. Impact of clothing on dermal exposure to phthalates: observations and insights from sampling both skin and clothing. *Environment Sci Technol*. 2016; Apr. 19;50:4350-57. <https://doi.org/10.1021/acs.est.6b00113>
- Wu Q, Lam JCW, Kwok KY, Tsui MMP, Lam PKS. Occurrence and fate of endogenous steroid hormones, alkylphenol ethoxylates, bisphenol A and phthalates in sewage treatment systems. *Journal of Environment Sciences*. 2017; Nov. 1;61:49-58. <https://doi.org/10.1016/j.jes.2017.02.021>
- Muller J, Kordel W. Occurrence and fate of phthalates in soil and plants, *Science of The Total Environment*. 1993; Jan. 1;134(1):431-37. [https://doi.org/10.1016/S0048-9697\(05\)80044-0](https://doi.org/10.1016/S0048-9697(05)80044-0)
- Teil MJ, Blanchard M, Chevreuil M. Atmospheric fate of phthalate esters in an urban area (Paris-France). *Science of The Total Environment*. 2006; Feb. 1;354(2-3):212-23. <https://doi.org/10.1016/j.scitotenv.2004.12.083>
- Erkekoğlu P, Rahidi W, Rosa VD, Giray B, Favier A, Hincal F. Protective effect of selenium supplementation on the genotoxicity of di (2-ethylhexyl) phthalate and mono(2-ethylhexyl) phthalate treatment in LNCaP cells. *Free Radical Biology and Medicine*. 2010;49(4):559-66. <https://doi.org/10.1016/j.freeradbiomed.2010.04.038>
- Peropadre A, Freire PF, Martin JMP, Herrero O, Hazen MJ. Endoplasmic reticulum stress as a novel cellular response to di (2-ethylhexyl) phthalate exposure. *Toxicology In Vitro*. 2015; Dec. 25;30(1):281-87. <https://doi.org/10.1016/j.tiv.2015.10.009>
- Molino C, Filippi S, Stopiello GA, Meschini R, Angeletti D. In vitro evaluation of cytotoxic and genotoxic effects of Di (2-ethylhexyl) phthalate (DEHP) on European sea bass (*Dicentrarchus labrax*) embryonic cell line. *Toxicology In Vitro*. 2019; Apr. 1;56:118-25. <https://doi.org/10.1016/j.tiv.2019.01.017>
- Lin W, Jiang R, Xiong Y, Xu J, Zheng J, Zhu P, Ouyang G. Quantification of the combined toxic effect of polychlorinated biphenyls and nano-sized poly styrene on *Daphnia magna*. *Journal of hazardous materials*. 2019; Feb. 15; 364: 531-36. <https://doi.org/10.1016/j.jhazmat.2018.10.056>
- Kim H, Nam K, Oh S, Son S, Jeon D, Gye MC, Shin I. Toxicological assessment of phthalates and their alternatives using human keratinocytes. *Environmental Research*. 2019; Aug. 1;175:316-22. <https://doi.org/10.1016/j.envres.2019.05.007>
- Sedha S, Lee H, Singh S, Kumar S Jain S, Ahmad A et al. Reproductive toxic potential of phthalate compounds state of art review. *Pharmacol Res*. 2021; May. 1;167:105536. <https://doi.org/10.1016/j.phrs.2021.105536>
- Engel SM, Miodovnik A, Canfield RL, Zhu C, Silva MJ, Calafat AM,

- Wolff MS. Prenatal phthalate exposure is associated with childhood behavior and executive functioning. *Environ Health Perspect*. 2010; Apr 118:565-71. <https://doi.org/10.1289/ehp.0901470>
20. Katsikantami I, Sifakis S, Tzatzarakis MN, Vakonaki E, Kalantzi OI, Tsatsakis AM et al. A global assessment of phthalates burden and related links to health effects. *Environ Int*. 2016; Dec; 1;97:212-36. <https://doi.org/10.1016/j.envint.2016.09.013>
 21. Zarean M, Keikha M, Poursafa P, Khalighinejad P, Amin M, Keli-shadi R. A systematic review on the adverse health effects of di (2-ethylhexyl) phthalate. *Environ Sci Pollut Res*. 2016; Dec;23 (24):24642-93. <https://doi.org/10.1007/s11356-016-7648-3>
 22. Liu C, Deng YL, Zheng TZ, Yang P, Jiang XQ, Liu EN et al. Urinary biomarkers of phthalate exposure and risk of thyroid cancer and benign nodule. *J Haz Mat*. 2020; Feb; 5;383:121189. <https://doi.org/10.1016/j.jhazmat.2019.121189>
 23. Vinny. Himachal Pradesh popularising polyhouse farming. New protective cultivation technology. 2009 [Cited 4 Feb 2021]. Wordpress.com weblog. <https://greenhousehimachal.wordpress.com/2009/02/09/himachal-pradesh-popularising-polyhouse-farming/>
 24. Cai QY, Mo CH, Wu QT, Zeng QY, Katsoyiannis A. Occurrence of organic contaminants in sewage sludges from the eleven wastewater treatment plants, China. *Chemosphere*. 2007; Aug; 1;68(9):1751-62. <https://doi.org/10.1016/j.chemosphere.2007.03.041>
 25. Wang XL, Lin QX, Wang J, Lu XG, Wang GP. Effect of wetland reclamation and tillage conversion on accumulation and distribution of phthalate esters residues in soils. *Ecol Eng*. 2013; Feb; 1;51:10-15. <https://doi.org/10.1016/j.ecoleng.2012.12.079>
 26. Gue V, Kannan K. Challenges encountered in the analysis of phthalate esters in foodstuffs and other biological matrices. *Anal Bioanal Chem*. 2013; Nov; 404(9):2539-54. <https://doi.org/10.1007/s00216-012-5999-2>
 27. Weschler CJ, Salthammer T, Fromme H. Partitioning of phthalates among the gas phase, airborne particles and settled dust in indoor environments. *Atmos Environ*. 2008; Mar; 1;42 (7):1449-60. <https://doi.org/10.1016/j.atmosenv.2007.11.014>
 28. Zhu YY, Tian J, Yang HB, Wei EQ, Yu G, Wei FS. Phthalate pollution in atmospheric particles and soils of Tianjin and their contamination and remediation of phthalate acid esters in agricultural soils. *Environ Pollut Control*. 2010;31(10):1535-41.
 29. Bodeker C, Bodeker G, Ong CK, Grundy CK, Burford G, Shein K. WHO Global Atlas of Traditional, Complementary and Alternative Medicine. Geneva, Switzerland: World Health Organization. 2005 [June 2021].
 30. Brandão MGL, Acúrcio FA, Montemor RLM, Marlière LDP. Complementary/alternative medicine in Latin America: Use of herbal remedies among Brazilian metropolitan area population. *Journal of Complementary and Integrative Medicine*. 2006; Jun; 6;3 (1):1-8. <https://doi.org/10.2202/1553-3840.1025>
 31. Bussman RW, Sharon D, Lopez A. Blending traditional and Western medicine medicinal plants use among patients at Clinica Antígona in El Porvenir, Peru. *Ethnobotany Research and Applications*. 2007;5:185-99. <https://doi.org/10.17348/era.5.0.185-199>
 32. Oreagba IA, Oshikoya KA, Amachree M. Herbal medicine use among urban residents in Lagos, Nigeria. *BMC Complementary and Alternative Medicine*. 2011; Dec;11:117-25. <https://doi.org/10.1186/1472-6882-11-117>
 33. Picking D, Younger N, Mitchell S, Delgoda R. The prevalence of herbal medicine home use and concomitant use with pharmaceutical medicines in Jamaica. *Journal of ethnopharmacology*. 2011; Sep; 1;137(1):305-11. <https://doi.org/10.1016/j.jep.2011.05.025>
 34. Warren P. 101 uses for stinging nettles. *Wildeye*, UK. 2006 [Nov, 2020].
 35. Williamson EM. *Potter's Herbal Cyclopaedia*. CW Daniel, Saffron Walden. 2003 [Jul 2021].
 36. El Haouari M, Rosado JA. Phytochemical, anti-diabetic and cardiovascular properties of *Urtica dioica* L. (Urticaceae): A Review. *Mini Reviews in Medicinal Chemistry*. 2019; Jan; 1;1:19(1):63-71. <https://doi.org/10.2174/1389557518666180924121528>
 37. Qayyum R, Qamar HM, Khan S, Salma U, Khan T, Shah AJ. Mechanisms underlying the antihypertensive properties of *Urtica dioica*. *Journal of translational medicine*. 2016; Dec;14(1):1-3. <https://doi.org/10.1186/s12967-016-1017-3>
 38. JR, Weiner MF 2019 Urtication (flogging with stinging nettles) and flagellation (beating with rods) in the treatment of paralysis. *Spinal Cord Series and Cases*. 2019; Sep; 26;5(1):1-5. <https://doi.org/10.1038/s41394-019-0222-8>
 39. Rajput P, Sharma RA. GC Profiling of *Urtica dioica* for the Efficacy of Traditional Urtication Therapy. 2021; Dec;101(4):363-67.
 40. Bayer Crop Science. *Urtica dioica*. 2015; [Sep 2021]. <http://www.cropscience.bayer.com/en/Products-and-Innovation/Crop-Compendium/Pests-Diseases-Weeds/Weeds/Urtica-dioica.aspx>
 41. Kew Seed Information Database. *Urtica dioica* L. Kew, UK: Royal Botanic Gardens. 2015 [Jan 2021] <http://data.kew.org/sid/SidServlet?ID=23779&Num=uLl>
 42. Dennis L. (ed.). *The Ecology of Butterflies in Britain*. Oxford University Press, Oxford. 1992.
 43. USDA Forestry Service. Index of Species Information: *Urtica dioica*. 2009.
 44. USDA-NRC. The PLANTS Database. Baton Rouge, USA: National Plant Data Center. 2015.<http://plants.usda.gov/>
 45. Robinson T, McMullan G, Marchant R, Nigam P. Remediation of dyes in textile effluent: a critical review on current treatment technologies with a proposed alternative. *Bioresource Technology*. 2001; May; 1;77(3):247-55. [https://doi.org/10.1016/S0960-8524\(00\)00080-8](https://doi.org/10.1016/S0960-8524(00)00080-8)
 46. Macek T, Kotrba P, Svatos A, Novakova M, Demnerova K, Mackova M. Novel roles for genetically modified plants in environmental protection. *Trends in Biotechnology*. 2008; Mar; 1;26(3):146-52. <https://doi.org/10.1016/j.tibtech.2007.11.009>
 47. Viktorova J, Jandova Z, Madlenakova M, Prouzova P, Batunek V. Native phytoremediation potential of *Urtica dioica* for removal of PCBs and heavy metals can be improved by genetic manipulations using constitutive CaMV 35S promoter. *PLOS ONE*. 2017, Dec; 8;12(10):e0187053. <https://doi.org/10.1371/journal.pone.0187053>
 48. Shams KM, Tichy G, Fischer A, Sager M, Peer T, Bashar A, Filip K. Aspects of phytoremediation for chromium contaminated sites using common plants *Urtica dioica*, *Brassica napus* and *Zea mays*. *Plant and Soil*. 2010; Mar;328:175-89. <https://doi.org/10.1007/s11104-009-0095-x>
 49. Li RW, Leach DN, Myers P, Leach GJ, Lin GD, Brushett DJ, Waterman PG. Anti-inflammatory activity, cytotoxicity and active compounds of *Tinospora smilacina* Benth. *Phytother Res*. 2004;18:78-83. <https://doi.org/10.1002/ptr.1373>
 50. Dar SA, Yousuf AR, Ganai FA, Sharma P, Kumar N, Singh R. Bioassay-guided isolation and identification of anti-inflammatory and anti-microbial compounds from *Urtica dioica* L. (Urticaceae) leaves. *African Journal of Biotechnology*. 2012;11(65):12910-20. <https://doi.org/10.5897/AJB11.3753>
 51. Mahmoudi R, Amini K, Fakhri O, Alem M. Aroma profile and antimicrobial properties of alcoholic and eques extracts from root, leaves and stock of nettle (*Urtica dioica* L.). *Journal of Microbiology Biotechnology and Food Sciences*. 2014; Jan; 6;4(3):220-24. <https://doi.org/10.15414/jmbfs.2014-15.4.3.220-224>
 52. Zehraw HM, Muotaz Z, Taleb M, Mukhlif B, Al SA, Al JI. Extraction

- and evaluation of the activity of *Urtica dioica* as bleeding stop material. Iraqi Journal of Biotechnology. 2019; Sep. 4;18(2):77-87.
53. Mayer FL, Stalling DLS, Johnson JL. Phthalate esters as environmental contaminants. Nature. 1972; Aug;238:411-13. <https://doi.org/10.1038/238411a0>
 54. Wang J, Chen G, Christie P, Zhang M, Luo Y, Teng Y. Occurrence and risk assessment of phthalate esters (PAEs) in vegetable and soils of suburban plastic film greenhouses. Science of The Total Environment. 2015; Aug. 1; 523: 129-37. <https://doi.org/10.1016/j.scitotenv.2015.02.101>
 55. Mesdaghinia A, Azari A, Nodehi R, Yeghmaeian K, Bharti A, Agarwal S. Investigation on the magnetic absorption separation, catalytic degradation and toxicity bioassay. Journal of Molecular Liquids. 2017;233:378-90. <https://doi.org/10.1016/j.molliq.2017.02.094>
 56. Blount BC, Silva MJ, Caudill SP, Needham LL. Levels of seven urinary phthalate metabolites in a human reference population. Environ Health Perspect. 2000;Oct;108(10):979-82. <https://doi.org/10.1289/ehp.00108979>
 57. Koch HM, Rossbach B, Drexler H, Angerer J. Internal exposure of the general population to DEHP and other phthalates- determination of secondary and primary monoester metabolite in urine. Environ Res. 2003; Oct. 1;93:177-85. [https://doi.org/10.1016/S0013-9351\(03\)00083-5](https://doi.org/10.1016/S0013-9351(03)00083-5)
 58. Silva MJ, Barr DB, Reidy JA, Malek NA. Urinary levels of seven phthalate metabolites in the US population from the National Health and Nutrition Examination Survey (NHANES) 1999-2000. Environ Health Perspect. 2004; Mar;112:331-38. <https://doi.org/10.1289/ehp.6723>
 59. Wittassek M, Wiesmuller GA, Koch HM, Eckard R. Internal phthalate exposure over the last two decades- a retrospective human biomonitoring study. Int J Hyg Environ Health. 2007; May. 22;210:319-33. <https://doi.org/10.1016/j.ijheh.2007.01.037>
 60. Bauer MJ, Herrmann R. Estimation of the environmental contamination by phthalic acid esters leading from household wastes. Science of The Total Environment. 1997; Dec. 3;208:49-57. [https://doi.org/10.1016/S0048-9697\(97\)00272-6](https://doi.org/10.1016/S0048-9697(97)00272-6)
 61. ECB. European Union risk assessment Report for dibutyl phthalate (with addendum 2004), European Chemicals Bureau, Ispra, Italy.2004; [Jun 2021].
 62. ECB. European Union Risk Assessment Report for Bis (2-ethylhexyl) phthalate (Final Report 2008), European Chemicals Bureau, Ispra, Italy. 2008.
 63. Kavlock R, Boekelheide K, Chapin R, Cunningham M. NTP Center for the Evaluation of Risks to Human Reproduction: phthalates expert panel report on the reproductive and developmental toxicity of di-n-octyl phthalate. Reprod Toxicol. 2002;16:721-34. [https://doi.org/10.1016/S0890-6238\(02\)00031-X](https://doi.org/10.1016/S0890-6238(02)00031-X)
 64. Kavlock R, Boekelheide K, Chapin R, Cunningham M. NTP Center for the Evaluation of Risks to Human Reproduction: phthalates expert panel report on the reproductive and developmental toxicity of di(2-ethylhexyl) phthalate. Reprod Toxicol. 2002;16:529-653. [https://doi.org/10.1016/S0890-6238\(02\)00032-1](https://doi.org/10.1016/S0890-6238(02)00032-1)
 65. Kavlock R, Barr D, Boekelheide K, Breslin W. NTP-CERHR expert panel update on the reproductive and developmental toxicity of di (2-ethylhexyl) phthalate. Reprod Toxicol. 2006; Oct. 1;16:291-399. <https://doi.org/10.1016/j.reprotox.2006.04.007>
 66. Borch J, Axelstad M, Vinggaard AM, Dalgaard M. Diisobutyl phthalate has comparable anti-androgenic effects to di-n-butyl phthalate in fetal rat testis. Toxicol Lett. 2006; Jun. 1;163:183-90. <https://doi.org/10.1016/j.toxlet.2005.10.020>
 67. Saillenfait, AM, Sabate JP, Gallissot F. Developmental toxic effects of diisobutyl phthalate, the methyl-branched analogue of di-n-butyl phthalate, administered by gavage to rats. Toxicol Lett. 2006; Aug. 1;165(1):39-46. <https://doi.org/10.1016/j.toxlet.2006.01.013>
 68. Saillenfait AM Sabate JP, Gallissot F. Diisobutyl phthalate impairs the androgen-dependent reproductive development of the male rat. Reprod Toxicol. 2008; Oct. 1; 26:107-15. <https://doi.org/10.1016/j.reprotox.2008.07.006>
 69. Gray LE, Ostby J, Furr J, Price M. Perinatal exposure to the phthalates DEHP, BBP and DINP, but not DEP, DMP or DOTP, alters sexual differentiation of the male rat. Toxicol Sci. 2000; Dec. 1;58:350-65. <https://doi.org/10.1093/toxsci/58.2.350>
 70. Parks LG, Ostby JS, Lambright, CR, Abbott BD. The plasticizer diethylhexyl phthalate induces malformations by decreasing fetal testosterone synthesis during sexual differentiation in the male rat. Toxicol Sci. 2000; Dec. 1;58:339-49. <https://doi.org/10.1093/toxsci/58.2.339>
 71. Andrade AJ, Grande SW, Talsness CE, Gericke C. A dose-response study following in utero and lactational exposure to di - (2-ethylhexyl) phthalate (DEHP): reproductive effects on adult male offspring rats. Toxicol. 2006;228:85-97. <https://doi.org/10.1016/j.tox.2006.08.020>
 72. Nagao T, Ohta R, Marumo H, Shindo T. Effect of butyl benzyl phthalate in Sprague Dawley rats after gavage administration: a two-generation reproductive study. Reprod Toxicol. 2000; Nov. 1;14:513-32. [https://doi.org/10.1016/S0890-6238\(00\)00105-2](https://doi.org/10.1016/S0890-6238(00)00105-2)
 73. Tyl RW, Myers CB, Marr MC, Fail PA. Reproductive toxicity evaluation of dietary butyl benzyl phthalate (BBP) in rats. Reprod Toxicol. 2004; Mar. 1;18:241-64. <https://doi.org/10.1016/j.reprotox.2003.10.006>
 74. Sharpe RM, Irvine DS. How strong is the evidence of a link between environmental chemicals and adverse effects on human reproductive health ? Br Med J. 2004; Feb. 19;328:447-51. <https://doi.org/10.1136/bmj.328.7437.447>
 75. Foster PMD. Disruption of reproductive development in male rat offspring following in utero exposure to phthalate esters. Int J Androl. 2006; Feb;29:140-47. <https://doi.org/10.1111/j.1365-2605.2005.00563.x>
 76. Howdeshell KL, Rider CV, Wilson VS, Gray LE. Mechanisms of action of phthalate esters, individually and in combination, to induce abnormal reproductive development in male laboratory rats. Environ Res. 2008; Oct. 1;108:168-76. <https://doi.org/10.1016/j.envres.2008.08.009>
 77. Wilson VS, Blystone CR, Hotchkiss AK, Rider CV. Diverse mechanisms of anti-androgen action: impact on male rat reproductive tract development. Int J Androl. 2008; Apr;31:178-87. <https://doi.org/10.1111/j.1365-2605.2007.00861.x>
 78. Bao J, Wang M, Ning X, Zhou Y, He Y, Yang J, Gao X, Li S, Ding Z, Chen B. Phthalate concentrations in personal care products and cumulative exposure to female adults and infants in Shanghai. Journal of Toxicology and Environmental Health. 2015; Mar. 4;78(5):325-41. <https://doi.org/10.1080/15287394.2014.968696>
 79. Wu Z, Zhang X, Wu X, Shen G, Du Q, Mo C. Update of di (2-ethylhexyl) phthalate DEHP by the plant *Beninoasa hispida* and its use for lowering DEHP content of intercropped vegetables. Journal of Agricultural and Food Chemistry. 2013; Jun. 5;61 (22):5220-25. <https://doi.org/10.1021/jf401378u>