



REVIEW ARTICLE

Plant growth promoting rhizobacteria (PGPR) and their eco-friendly strategies for plant growth regulation: a review

Debapriya Choudhury, Santanu Tarafdar & Sikha Dutta*

Applied and Molecular Mycology and Plant Pathology Laboratory, CAS Department of Botany, The University of Burdwan, Purba Bardhaman 713 104, West Bengal, India

*Email: sikha.bu.academia@gmail.com

 OPEN ACCESS

ARTICLE HISTORY

Received: 27 November 2021

Accepted: 17 March 2022

Available online

Version 1.0 (Early Access): 17 April 2022



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, etc. See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Choudhury D, Tarafdar S, Dutta S. Plant growth promoting rhizobacteria (PGPR) and their eco-friendly strategies for plant growth regulation: a review. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.1604>

Abstract

As a natural engineer, Plant Growth Promoting Rhizobacteria (PGPR) play an important role in increasing plant growth, yield and enhancing plant tolerance to stressful conditions. These beneficial bacteria take up their position in the rhizosphere, around the plants' root tissues. They may be in, or on their host tissues and help to provide nutrients to their host plants. For sustainable agriculture, PGPR transmit their extensive assistance in ecosystem management, soil structure maintenance, stress management and plant morphology and physiology modulation in an environmentally friendly manner. Plant- PGPR interactions also stimulate nutrient acquisition and accumulation, improve plant performance and enhance plants' tolerance to abiotic and biotic stresses. Beside these, PGPR are good biofertilizers and safe for our environment. Nanotechnological advances with PGPR applications are important today to increase the impact of PGPR in agriculture. Undoubtedly, PGPR concept is intimately involved with agriculture, horticulture, forestry and they are too enough to establish a vibrant environment. In this review we have focused on the versatility of PGPR-their performance and aimed to address some future prospects of PGPR as an eco-friendly tool for plant growth regulation.

Keywords

PGPR, eco-friendly, plant growth regulation, sustainable agriculture, stress management

Introduction

Soil is the core house to millions and millions of microbes, specifically rhizosphere become more precious due to rhizospheric effect. Some of these bacteria provide benefits to plant community and help plants to cope with challenging adversities (1). In the year 1978, Kopper and Schroth applied the term 'Plant growth promoting rhizobacteria' (PGPR) for beneficial rhizospheric free living bacteria (2).

PGPR are involved in improving soil composition through decomposition of crop residues, stabilization of mineral nutrients and mineralization of soil organic matter, acquisition of plant growth promoters including nutrient (biofertilizer), phosphate solvents, nitrogen fixation, nitrification and synthesis of soil organic matters (3).

PGPR maintain their host plants' health through different processes such as (i) enhancement of nutrient acquisition by host plants (ii) improvement of plant growth and crop production by producing phytohormones, siderophore and also by enriching soil nutrients (iii) promotion of nodule

formation in legume and, (iv) enhancement of seedlings' emergence (4-8). Beside these, PGPR facilitate their host plants to manage different types of abiotic stresses like drought (9, 10), salinity (11, 12) and heavy metal absorption is also restricted by some metal resistant PGPR strains (13). These types of bacteria activate antioxidant signalling to reduce temperature stress (14).

PGPR tend to protect the plants against some pests and phyto-pathogens by triggering Induced Systemic Resistance (ISR) through lipopolysaccharides, flagella, homoserine lactones, acetoin and butanediol (5). From the host-PGPR interaction, the plants gain some benefits including increase in seed germination rate, yield, root growth, leaf area, chlorophyll content, protein content, nutrient content, germination percentage, root weight and root length, hydraulic activity, abiotic stress tolerance, delayed aging and biocontrol (15).

Excessive use of artificial fertilizers, herbicides, fungicides and pesticides are the major causes of environmental pollution. The demand for food is increasing rapidly due to the increase in human population. Today civilizations, industrialization, chemical fertilizers in agriculture, excessive use of pesticides are major obstacles to the survival of agriculture. Excessive use of nitrogen fertilizers produces Nitrous oxide (N_2O), a greenhouse gas that leads to global warming. The main drawback of nitrogen fertilizers is the reduction of biological nitrogen fixation. Ammonium nitrate is widely used by farmers as nitrogen fertilizer. Excess ammonium flow into the plant body meets the nitrogen requirements of the plant which reduces the symbiotic association between the plant and the microbe (16).

Reducing the use of chemical fertilizers in agriculture and reducing the burden of pollutants in the environment is possible only through the use of biofertilizer (17).

To date, various reports on PGPR- agricultural sustainability have been documented. Undoubtedly it is a highly promising tool for soil and crop management. The purpose of this review is to provide a concise overview of the PGPR and to highlight how they modify plant growth regulations in various environmental conditions.

Isolation of Plant growth promoting rhizobacteria (PGPR)

In bacteriological research, bacterial isolation is very important and the first step in separating different strains from natural mixtures. PGPR isolation can be done from rhizosphere soil sample or from root nodules. Serial dilution-spread plate is a very common, simplified and well-accepted method for isolating PGPR strains from rhizosphere soil sample. In the first step, it is important to collect soil samples that adhere to the roots. This is followed by sequential steps like, preparation of the stock solution, serial dilution of the stock solution, spread plate and incubation (18) as discussed in Fig. 1.

Endophytic bacteria can be isolated from sterilized root samples (18-21) as discussed in Fig. 2.

Isolation of PGPR strains from crushed root nodules has also been reported (22). Different growth media




Fig. 1. Steps of serial dilution-spread plate method for isolation of PGPR from rhizosphere soil.




Fig. 2. Steps for isolation of endophytic bacteria from root samples.

namely Pikovskayas agar (PKV) media (18, 23), Luria-Bertani (LB) agar media (24), King B agar media (25), Tryptic soy agar media (26), Nutrient agar media (21) are used for bacterial growth.

Culture of Plant growth promoting rhizobacteria (PGPR)

Bacterial culture is basically a method that allows bacteria to multiply under controlled lab conditions. Carbohydrates, proteins and nucleic acids are important ingredients of culture media which are dissolved in water. After isolation, to maintain the purity of bacterial strains it is required to apply pure culture method (27) as presented in Fig. 3.

Sub culturing is a method where bacteria are transferred from stock culture to fresh nutritive medium. Beside solid medium, bacteria can be cultured in liquid medium or broth. Broth is liquid medium without solidifying agent (agar) containing all the important ingredients necessary for the growth of bacteria. PGPR containing broth general-




Fig. 3. Maintenance of PGPR strains (different colours represent different PGPR strains).

ly use for experimental studies including ammonia production, enzymes (Catalase, Oxidase) synthesis, EPS production, IAA production (27-29) etc.

Role of Plant growth promoting rhizobacteria (PGPR) in Plant Growth Regulation

This section is categorized into 3 main parts viz. PGPR and its types, Contributions of PGPR in plant growth regulations, Molecular responses and signalling pathways in response to Plant-PGPR interactions.

PGPR and its types

Plant growth promoting rhizobacteria are important groups of beneficial rhizosphere bacteria that contribute a mutualistic relationship with plants. These provide an eco-friendly safeguarding policy for the plants (30). PGPR maintain the existence of plants by promoting their growth and acting as biocontrol agents (31, 32) and regulate agro-ecological natural equilibrium by increasing productivity with a low chemical assimilation (33). Some common plant growth promoting rhizobacteria and their host plants are reported (Table 1).

Table 1. Plant growth promoting rhizobacteria (PGPR) and their host plants (4, 15, 31, 34-44)

PGPR	Host plants
<i>Rhizobium leguminosum</i>	<i>Phaseolus vulgaris</i> <i>Phaseolus vulgaris</i>
<i>Pseudomonas cepacia</i>	<i>Gossypium hirsutum</i> <i>Cucumis sativus</i>
<i>Pseudomonas sp.</i>	<i>Dianthus caryophyllus</i> <i>Phaseolus vulgaris</i>
<i>Pseudomonas fluorescens</i>	<i>Triticum aestivum</i> <i>Hordeum vulgare</i> <i>Brassica juncea</i>
<i>Azotobacter chroococcum</i>	<i>Triticuma aestivum</i> <i>Saccharum officinarum</i>
<i>Azospirillum brasilense</i>	<i>Zea mays</i>
<i>Bradyrhizobium japonicum</i>	<i>Glycine max</i>
<i>Achromobacter xylosoxidans</i>	<i>Vigna radiata</i> <i>Hordeum vulgare</i>
<i>Bacillus subtilis</i>	<i>Gossypium hirsutum</i> <i>Brassica juncea</i> <i>Eucalyptus globulus</i>
<i>Methylobacterium mesophilicum</i>	<i>Oryza sativum</i>
<i>Serratia sp.</i>	<i>Zea mays</i>
<i>Acinetobacter sp.</i>	<i>Zea mays</i>

Plant growth promoting rhizobacteria colonize in rhizosphere region including root surface, or superficial intercellular space of host root, and frequently they form root nodules. As endophyte they also remain in apoplastic region of host plants (45). Broadly PGPR are categorized into two prime groups as presented in Fig. 4. These are represented as iPGPR: Intracellular plant growth promoting rhizobacteria and ePGPR: Extracellular plant growth promoting rhizobacteria (46).




Fig. 4. Types of PGPR on the basis of their inhabitants (iPGPR and ePGPR indicate intracellular and extracellular Plant Growth Promoting Rhizobacteria respectively).

Mechanisms by which PGPR stimulate plant growth are categorized into biofertilizers, phytostimulators and biopesticides or biocontrol agents (45, 47).

Biofertilizers

Mechanisms like biological nitrogen fixation, ammonia production, phosphate solubilization are covered in this section. PGPR enhance the supply of primary nutrients to the plant body through these processes and stimulate plant growth.

Phytostimulators

This section covers some PGPR mechanisms, such as the production of phytohormones (auxin, cytokinin and gibberellin) and the reduction of ethylene concentration.

Biopesticides or biocontrol agents

Some of the important PGPR mechanisms like competitive exclusion, acquired and induced systemic resistance, production of fungal wall degrading enzymes and production of antibiotics (siderophores, Hydrogen cyanide and anti-fungal metabolites) are included within this category.

Contributions of PGPR in Plant Growth regulations

The starting letters 'PGP' of PGPR directly indicate its promising role in plant growth promotion. Plant growth means plant's overall growth including plant biomass, yield and key regulators (48). PGPR help plants to perform proper physiological functions through photosynthesis, nitrogen metabolism, ammonia production, phytohormone production and regulation, mineral solubilization, metal chelating (49-56) as discussed in Fig. 5.

From ecological point of view PGPR are good biofertilizer (2) and maintain soil fertility by recycling the soil nutrients (57). Agricultural sustainability directly depends upon soil texture quality. A sustainable agricultural system




Fig. 5. Policies raised by PGPR to regulate plants' internal activities (all the solid and dotted lines indicate the management policies by PGPR, where dotted lines are exclusive for indicating stress related management).

can reimpose our daily lives by raising a large financial support (58). At present day it is impossible to imagine a sustainable crop production without involvement of beneficial microbes (59). They colonize in plant root area, and also improve the productivity (60). PGPR can show their performance versatility (45). Some PGPR with their performances are documented (Table 2).

Table 2. Plant growth promoting rhizobacteria (PGPR) and their applications (4, 6, 23, 44, 61-95)

TYPE OF FUNCTIONS		PGPR
Nitrogen fixation		<i>Bacillus tequilensis</i> , <i>Pseudomonas aeruginosa</i> , <i>Klebsiella</i> sp., <i>Azotobacter</i> spp, <i>Azospirillum</i> , <i>Enterobacter cloacae</i> , <i>Bacillus drentensis</i> , <i>Rhizobium</i> , <i>Bacillus pumilus</i> Sol-1, <i>Alcaligenes</i> sp. Mal-4, <i>Providencia vermicola</i> Ama-2, <i>Brevundimonas</i> Kro13, <i>Kluyvera ascorbata</i> SUD165, <i>Pseudomonas putida</i> , <i>Ochrobactrum</i> , <i>Bacillus megaterium</i> , <i>Bacillus mycoides</i>
Phosphate solubilization		<i>Bradyrhizobium</i> , <i>Bacillus tequilensis</i> , <i>Pseudomons</i> , <i>Azotobacter</i> , <i>Azospirillum brasilense</i> , <i>Azospirillum amazonense</i> , <i>Agrobacterium</i>
Siderophore production		<i>Rhizobium ciceri</i> , <i>Bradyrhizobium japonicum</i> , <i>Kluyvera ascorbata</i> , <i>Pseudomonas fluorescens</i> , <i>Pseudomonas putida</i> , <i>Mesorhizobium ciceri</i> , <i>Azotobacter Chroococcum</i> , <i>Serratia marcescens</i> , <i>Proteus vulgaris</i> , <i>Paenibacillus polymyxa</i> , <i>Pseudomonas aeruginosa</i> 4EA, <i>Klebsiella</i> sp., <i>Azotobacter vinelandii</i> , <i>Bacillus megaterium</i> , <i>Bacillus subtilis</i> , <i>Pantoea allii</i> and <i>Rhizobium radiobacter</i>
Phytohormone production	IAA	<i>Bradyrhizobium</i> , <i>Rhizobium</i> , <i>Azotobacter chroococcum</i> , <i>Bacillus</i> , <i>Pseudomons</i> , <i>Azotobacter</i> , <i>Azospirillum</i> , <i>Bradyrhizobium japonicum</i> , <i>Sphingomonas</i> sp., <i>Mycobacterium</i> sp., <i>Bacillus</i> sp., <i>Rhodococcus</i> sp., <i>Cellulomonas</i> sp., <i>Pseudomonas</i> sp., <i>Pseudomonas fluorescens</i> , <i>Brevibacillus</i> sp., <i>Klebsiella oxytoca</i> , <i>Burkholderia</i> , <i>Enterobacter</i> sp., <i>Rahnella aquatilis</i> , <i>Bacillus tequilensis</i> , <i>Bacillus subtilis</i> , <i>Phyllobacterium</i> , <i>Agrobacterium</i>
	Cytokinin	<i>Rhizobium leguminosarum</i> , <i>Azotobacter chroococcum</i>
	ABA	<i>Bacillus subtilis</i>
	GA	<i>Achromobacter xylosoxidans</i> , <i>Gluconobacter diazotrophicus</i> , <i>Acinetobacter calcoaceticus</i> , <i>Rhizobia</i> , <i>Azotobacter</i> sp., <i>Bacillus</i> sp., <i>Azotobacter chroococcum</i> , <i>Serratia nematodiphila</i>

Atmospheric nitrogen fixation

Nitrogen is one of the major constituents of plant cells which present in DNA, RNA, protein, amino acid etc. Eukaryotic plants rely entirely on symbiotic microbes for biological nitrogen fixation because they do not have the ability to break the triple bond between two nitrogen atoms. Some plant growth promoting bacteria namely *Klebsiella* sp, *Acinetobacter* sp, *Bacillus pumilus* (44), *Azotobacter* spp (4), *Burkholderia* (96), *Pseudomonas* (62) have been reported to show atmospheric nitrogen fixing ability. PGPR with atmospheric nitrogen fixing ability can raise the nitrogen

uptake throughout whole plant body (44). In addition to nitrogen fixation, PGPR conduct a 5-step reaction as sketched in Fig. 6 to complete the nitrogen cycle on Earth.




Fig. 6. PGPR involvements in Nitrogen cycle (where Ammonification, Nitrosification, Nitrification, Denitrification and Nitrogen fixation indicate intermediate steps).

Atmospheric nitrogen fixing PGPR show versatility in their performance including disease management, crop growth promotion in addition with the nitrogen maintenance in agricultural land (58). *Bacillus tequilensis* also shows positive response in atmospheric nitrogen fixation in soil though this is a potent phosphate biofertilizer (23).

Ammonia production

Ammonia (NH₃) is the first stable product of nitrogen fixation which can fulfill the nitrogen demand in plant body. As an important nutrient source it is frequently used in agricultural field to improve the crop productivity though

it shows detrimental effect above its optimum level (96, 97). Ammonia producing PGPR are included in the genera *Bacillus*, *Pseudomonas*, *Serratia* (43).

Phosphate solubilization

Plants need phosphate for growth, maturation and function. Usually inorganic phosphate is soluble in nature. Chemical fertilizer contains inorganic phosphate but it becomes immobilized and inaccessible to plants (98). Phosphate solubilizing bacteria are the key convertor of insoluble form of phosphate in soluble form and make it useable for plants (99). Some PGPR can apply this management policy have been mentioned (Table 2). Phosphate solubilizing PGPR also improve plant growth and yield (100). A synergistic efficiency of phosphate solubilisation is found in the combine application of bacteria instead of individual (101).

Phytohormone production

Phytohormones are organic compounds which are key players in plant growth, development and maintenance of plants' cellular activities even under environmental stress conditions (6, 102). Phytohormones such as Auxin, Cytokinin, Gibberellin are involved in the growth and development of the plants (48), are reported to be produced by many PGPR (103-105). Among the various types of natural Auxin produced by plant growth promoting rhizobacteria, IAA is important one (6). Auxin producing PGPR can activate Auxin responsive genes that promote plant growth (106) and induce root length (107) and biomass also (108). It has been documented in previous studies that PGPR which are able to produce Auxin, also involve in hormonal changes, and also regulate some defense genes, wall related genes at transcriptional level (109). Some strains of PGPR have been shown to promote plant shoot growth by eliciting Gibberellins production (110). *Achromobacter xylosoxidans*, *Gluconobacter diazotrophicus*, *Acinetobacter calcoaceticus*, *Rhizobia*, *Azotobacter* spp., *Bacillus* sp., *Herbaspirillum seropedicae* and *Azospirillum* spp. are some Gibberellic acid producing PGPR (94). Increment in Cytokinin concentration with plant growth has been found after *Bacillus subtilis* inoculation in lettuce plant (111). On the other hand, PGPR showed a down regulating effect on ethylene concentration (16) as shown in Fig. 7.




Fig 7. Regulation of Auxin-Ethylene biosynthesis by PGPR in response to stress (where ACC and SAM indicate 1-Aminocyclopropane-1-Carboxylic acid and S-Adenosyl-L-methionine respectively)

Hydrogen cyanide (HCN) production

Hydrolysis of cyanogenic compound and Ethylene biosynthesis are some common pathways of cyanide production for plants. Plants detoxify most of it by a special enzyme β cyanoalanine (99). Depending upon its concentration hydrogen cyanide exerts positive impact on plants. At a low concentration it regulates plant metabolic processes as a signaling molecule whereas it shows its anti herbivore activity at high concentration (112). In the agriculture system HCN is commonly used as a biocontrol agent, and it is a good metal chelator also (113). In a Pea - *Pythium* pathosystem HCN has been reported as biocontrol trait for pseudomonads (114). As potent biofertilizer HCN producing PGPR promote plant vegetative growth, and yield also (113). PGPR which are characterised with of HCN and ammonia production can exert synergistic effect on plant growth and plant metabolite modulation (43). The concentration of HCN produced by PGPR under *in vitro* condition is unable to relate to the geochemical processes instead of biocontrol hypothesis (112).

Siderophore production

These are small, iron chelating low molecular weight secondary metabolites secreted by nearly all bacterial species (115-117). These are categorised on the basis of chemical nature of their iron-chelating group which may be α -hydroxyl-carboxylate, catechol, hydroxamate, or mixed types (116). As a scavenger they scavenge iron from environment and make mineral (115). Siderophores transport Fe^{3+} ions and transport them across cell membranes (118). Siderophore producing microbes are considered as an efficient PGPR which show multifunctional potentiality in plant growth promotion (119), and also put down diseases (120). *Pseudomonas putida* enhances the level of iron content in the natural habitat utilizing heterologous siderophores produced by other microorganisms (34).

Rhizoremediation

At present era it is very urgent to reduce the pollution level to keep away all of the organisms from its cope. Contaminated soil and water is a big problem for agricultural sustainability. Combine effect of phytoremediation and bioaugmentation regulate rhizoremediation (58). Phytoremediation is a promising approach by which plants can extract metals from contaminated soil (121). Biological waste treatment by adding cultured microorganisms is known as bio-augmentation (122). Genetically engineered *Pseudomonas fluorescens*, *Pseudomonas aeruginosa* and certain *Bacillus* sp. are some examples of PGPR having rhizoremediation property (123). PGPR participate in this remediation process by removing organic contaminants, or by applying certain processes such as phytoextraction, rhizofiltration, phytostabilization and phytovolatilization (57) to treat heavy metals (121) and lower the contamination levels.

Stress tolerance

The devastating impact of environmental stress on crop yield is a serious problem at present scenario. Now it is very urgent to adopt some eco-friendly management poli-

cies to secure our agricultural system. Environmental stress may be caused by abiotic factors, or some biotic agents like, fungi, bacteria, nematodes, herbivores, oomycetes etc (124). Worldwide major losses in crop field can be accomplished by abiotic mediators including salinity, floods, drought, radiation, high temperature, extreme cold, heavy metals etc (16, 124). Any kind of stress ultimately converts to oxidative stress and increases the production of some free radicals such as H_2O_2 , O_2^- , $\cdot N=O$, ClO^- , and OH^- which are considered as Reactive Oxygen Species (ROS) having the capacity to damage plants membrane lipids, proteins and nucleic acids (125). PGPR are some key agents which protect plants from multiple stresses (60) through some management policies (Table 3).

Table 3. Plant growth promoting rhizobacteria (PGPR) management policies to protect plant from stress effects (2, 31, 58, 126-129)

Management strategies	PGPR
Protective enzymes production	<i>Pseudomonas fluorescens</i> LPK2, <i>Sinorhizobium fredii</i> KCC5, <i>Trichoderma</i> sp., <i>Bacillus subtilis</i> , <i>Variovorax paradoxus</i> 5C-2, <i>Pseudomonas fluorescens</i> biotype G, <i>Enterobacter</i> sp., <i>Stenotrophomonas maltophilia</i> , <i>Bacillus cepacia</i> , <i>Bacillus licheniformis</i> , <i>Bacillus cereus</i> , <i>Bacillus circulans</i> and <i>Bacillus thuringiensis</i>
Disease resistance antibiosis	<i>Pseudomonas</i> sp., <i>Bacillus</i> sp., <i>Pseudomonas fluorescens</i> and <i>Pseudomonas aeruginosa</i>
Volatile organic compounds secretion	<i>Pseudomonas</i> , <i>Bacillus</i> , <i>Arthrobacter</i> , <i>Stenotrophomonas</i> and <i>Serratia</i> , <i>Bacillus megaterium</i> BOFC15
Exopolysaccharides secretion	<i>Rhizobium leguminosarum</i> , <i>Azotobacter vinelandii</i> , <i>Bacillus drementensis</i> , <i>Enterobacter cloacae</i> , <i>Agrobacterium</i> sp., <i>Xanthomonas</i> sp., <i>Rhizobium</i> sp., <i>Bradyrhizobium</i> sp., <i>Pseudomonas putida</i> , <i>Bacillus subtilis</i> and <i>Azospirillum brasilense</i>

As anti-phytopathogenic agents (130) PGPR produce some cell wall degrading enzyme such as β -1, 3-glucanase, ACC deaminase and chitinase and neutralize pathogens (126). *Burkholderia*, is an ACC deaminase secreting genus has been reported to increase the growth of tomato seedlings by limiting ethylene levels (131). ACC deaminase also reduces the ethylene induced negative changes in the roots (132) as presented in Fig. 7. Various types of plant-microbe interactions are essential for regulating the local and systemic types of plant defence mechanism which help to increase the chances of survivability of the plants living in stress affected area (131). PGPR defeat plants in dehydration by improving plants water use efficiency (WUE) and water absorption through roots (133). Induced systemic resistance (ISR) is a modern approach in agricultural research, inspired by non pathogenic microbes and involved in control of numerous diseases (58). VOCs producing PGPR show their potentiality in plant growth, in plant pathogen control and also in ISR induction (134). Involvement of VOCs produced by *Bacillus subtilis* in hormonal regulation, plant growth development, photosynthesis and phytohormone induction with alleviation of ethylene level have been reported (134). Hormonal regulation plays central role in stress management (135). PGPR change the levels of endogenous phytohormone within wheat seedlings and enhance the ability to overcome the stress like salt and drought (136). *B. subtilis* resists drought stress elevating the levels of Abscisic acid (137). IAA producing bacteria help plants to defend against water deficit condition (138) by increasing root growth and/or enhancing the formation of lateral roots and roots hairs in various plant species (60). In this context EPS secretion is one of the prime characterizations of PGPR which are high molecular weight, biodegradable polymers

and sustain host plants growth, and yield acting as protective barrier against biotic and abiotic type of stress conditions (58). EPS producing PGPR strains enhance plant growth by avoiding water stress (55) and also stimulate the production of enzymatic antioxidant such as Superoxide dismutase (SOD), Catalase (CAT) and Peroxidase (POD) to detoxify ROS (129, 136). Some PGPR showing stress-tolerance capabilities have been listed (Table 4).

Molecular responses and signalling pathways in response to Plant-PGPR interactions

Plant microbe signalling exerts phyto-beneficial roles including crop biomass, nutrient uptake and metabolite up-regulation. Some important traits sustain plants under

Table 4. Stress responsive Plant growth promoting rhizobacteria (PGPR) (127, 129, 139)

Type of stress	PGPR	
ABIOTIC STRESS	Salinity	<i>Azospirillum brasilense</i> , <i>Pseudomonas syringae</i> , <i>Pseudomonas fluorescens</i>
	Drought	<i>Achromobacter pechaudii</i> , <i>Azospirillum brasilense</i> , <i>Acinetobacter</i> sp., <i>Pseudomonas</i> sp., <i>Pseudomonas fluorescens</i> biotype G, <i>Bacillus subtilis</i> and <i>Azospirillum brasilense</i>
	Temperature	<i>Burkholderia phytofirmans</i> , <i>Aeromonas hydrophila</i> , <i>Serratia liquefaciens</i> , <i>Pseudomonas</i> sp. (strain AKM-P6) and <i>Pseudomonas putida</i> (strain AKM-P7)
BIOTIC STRESS	Heavy metal	<i>Pseudomonas putida</i> (Pb and Cd resistance), <i>Brevibacillus</i> sp. (Zn resistance)
		<i>Paenibacillus Polymyxa</i> (strains B2, B3, B4), <i>Paenibacillus favisporus</i> (strain BKB30), <i>Bacillus amyloliquefaciens</i> (strain HYD-B17), <i>Bacillus licheniformis</i> (strain HYTAPB18), <i>Bacillus thuringiensis</i> (strain HYDGRFB19) and <i>Bacillus subtilis</i> (strain RMPB44)

biotic and abiotic types of stress (140). Advanced bioinformatic studies indicate reciprocal gene regulation between bacteria and their host plants during colonization and post colonization also (141). Phyto-beneficial proteins have

been reported to induce in sorghum roots after PGPR inoculation (141). Some phyto-beneficial proteins deal with DNA regulation including Phosphodiesterase, sulfatase, Carbohydrate kinases (FGGY_C) whereas some of these are stress relieving in nature such as HSP70 (141). The important protein family Carbohydrate kinases (FGGY_C) are involved in the production of carbohydrates and control plant biomass (141). PGPR enhance photo pigment contents in Basil leaves under water stressed condition (142), and maintain photosynthesis to manage plant biomass and survivability. Induction of photosynthetic rate, plant biomass, and photosynthetic pigments such as chlorophyll content, carotenoids under salt stress after inoculation of PGPR have been reported (143). Carotenoids are belonging to C₄₀ tetraterpene family, this report also supports the fact that the biosynthesis of terpene is somehow regulated by PGPR. Phosphodiesterase is one of the key players involved in DNA-protein crosslink repair mechanism in plants (141) and cellular homeostasis maintains by HSP70 (141). PGPR regulate phenylalanine concentration in tomato (144). Phenylalanine is an important product of shikimic acid pathway, regulates biosynthesis of phenolic compounds. One of the important phenolic compounds is flavonoids which are non enzymatic antioxidant, synthesized from an extended part of shikimic acid pathway, more specifically from phenyl propanoid acetate pathway. PGPR enhance flavonoid biosynthesis genes (145) and also trigger the core pathway of phenolics biosynthesis by up-regulating the key enzyme PAL in Chickpea (146) as shown in Fig. 8.




Fig. 8. Phenolic compounds-PGPR-ISR interrelationship (where, ISR represents Induced Systemic Resistance. Black arrows are indicators of sequential biochemical steps and green lines indicate the routes for induction of plant defence triggered by PGPR)

In *Solanum*, enhancement of photochemical efficiency after *Bacillus* inoculation has been reported, and this makes plant to cope with abiotic stresses such as salt, drought and heavy metal (146). Flavonoids are 15 C plant phenolics involve in pigmentation, pollination, antimicrobial and UV protection. Side by side this flavonoid enhancement positively correlates with non pathogenic defence management or ISR: Induced Systemic Response. PGPR regulate some biochemical signalling involved in

nitrogen fixation. Nitrogenase is the multi-subunit oxygen liable key enzyme for Nitrogen fixation, present in prokaryotes. Nitrogenase activity has been presented in Fig. 9.




Fig. 9. Schematic presentation of Nitrogenase activity (where S and L represent Small and Large subunit respectively. Fd (R) indicates Reduced Ferredoxin and Fd (OX) indicates Oxidized Ferredoxin)

Nitrogenase reductase, the small component of Nitrogenase enzyme is anchored by nifH. It has been confirmed that *Bacillus* promotes nifH gene expression in sugarcane varieties (65). Signalling system of some defence responsive enzymatic antioxidants such as CAT, GPx and SOD are regulated by plant growth promoting rhizobacteria under environmental adverse situations (147). Phytohormonal regulation is triggered by activation of ACC deaminase, the main enzyme of stress regulation that is shown in Fig. 7.

Conclusion

Agriculture is the only way to fulfill food demand worldwide. But unfortunately it is closely involved in the production of chemical pollutants. The use of chemical fertilizers, pesticides and insecticides by farmers has been increased to meet the growing food demand to feed booming population worldwide. They increase crop yields but reduce soil quality, which ultimately disturbs the ecosystem.

Environmentally PGPR are very good. They provide promising assistance in crop management and maintenance of soil ecosystems in terms of biofertilization, bioremediation, pest management, disease management, biocontrol and stress tolerance. They have also been shown to maintain plant cellular homeostasis in adverse environmental conditions.

In the current situation, it is very important to increase crop production by keeping the ecosystem sustainable. There is no doubt that PGPR are sufficient to increase global agricultural production in an environmentally sound way. Nanotechnology is a potent tool in this regard. Although it is difficult to incorporate nanotechnology into agriculture, it is efficient enough to promote global environmental and agricultural sustainability.

Future Prospects

PGPR are capable to manage plants' internal activities, plant growth and productivity even under adverse environmental conditions and also participate in contaminat-

ed soil management. But the effects of PGPR are gradually diminishing due to some abiotic and biotic factors. PGPR are lost in a large percentage in the conventional method of their application as biofertilizer (16).

Therefore, advanced technology needs to be improved to increase the use of PGPR in the agricultural system. Nanotechnology is a potential tool for improving PGPR services in agriculture (16). Nowadays, in agricultural revolution, nanoagriculture is a potential tool. In nanoagriculture, nanofertilizers improve crop production efficiently by assisting plants in efficient nutrients uptake, monitoring plant growth, enhancing food quality, protecting plants, detecting plant diseases, increasing food production and reducing waste (16, 148). Nanoencapsulation technology is a potential method to extend PGPR service life (16). The application of nanotechnology in agriculture and environmental sustainability is not easy, so more research is needed to make it more convenient. In addition, much effort is required to obtain information on particular steps regulated by PGPR in plant biochemical pathways.

Overall, there has been a lot of work in PGPR over the decades, but the application of nanotechnology in PGPR-agricultural systems requires a lot of work that will play an important role in sustainable agriculture without harming the ecosystems.

Acknowledgements

The Authors acknowledge all the members of Applied and Molecular Mycology and Plant Pathology Laboratory, University of Burdwan, West Bengal, India for their constant support and cooperation. The Authors highly commended UGC, Govt. of India for its financial support and thankful to the Department of Botany Library for providing educational resources for compiling the review article.

Authors contributions

The main concept of the review was provided by SD. Literature study was done by DC. The whole work was discussed by both Authors and finally DC prepared the manuscript. ST helped DC to prepare the figures according to the prescribed format.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflicts of interests to declare.

Ethical issues: None.

References

- Smith DL, Praslickova D, Ilangumaran G. Inter-organismal signaling and management of the phytomicrobiome. *Frontiers In Plant Science*. 2015; 6:722. <https://doi.org/10.3389/fpls.2015.00722>
- Basu A, Prasad P, Das SN, Kalam S, Sayyed RZ, Reddy MS, El Enshasy H. Plant Growth Promoting Rhizobacteria (PGPR) as Green Bioinoculants: Recent Developments, Constraints and Prospects. *Sustainability*. 2021; 13(3):1140. <https://doi.org/10.3390/su13031140>
- Prasad R, Kumar M, Varma A. Role of PGPR in soil fertility and plant health. In: Egamberdieva DE *et al* editors. *Plant-growth-promoting rhizobacteria (PGPR) and medicinal plants*, *Soil Biology*. Cham: Springer; 2015;247-60. https://doi.org/10.1007/978-3-319-13401-7_12
- Bhattacharyya PN, Jha DK. Plant growth-promoting rhizobacteria (PGPR): Emergence in agriculture. *World J Microbiol Biotechnol*. 2012; 28:1327-50. <https://doi.org/10.1007/s12045-016-0421-6>
- Gopalakrishnan S, Sathya A, Vijayabharathi R, Varshney RK, Gowda CLL, Krishnamurthy L. Plant growth promoting rhizobia: Challenges and opportunities. *Biotech*. 2015; 3: 355-77. <https://doi.org/10.1007/s13205-014-0241-x>
- Backer R, Rokem J.S, Ilangumaran G, Lamont J, Praslickova D, Ricci E, Subramanian S, Smith D.L. Plant growth promoting rhizobacteria: Context, mechanisms of action and roadmap to commercialization of biostimulants for sustainable agriculture. *Front Plant Sci*. 2018; 9:1473. <https://doi.org/10.3389/fpls.2018.01473>
- Kalam S, Das SN, Basu A, Podile AR. Population densities of indigenous *Acidobacteria* change in the presence of plant growth promoting rhizobacteria (PGPR) in rhizosphere. *J Basic Microbiol*. 2017; 57:376-85. <https://doi.org/10.1002/jobm.201600588>
- Swarnalakshmi K, Yadav V, Tyagi D, Dhar DW, Kannepalli A, Kumar S. Significance of plant growth promoting rhizobacteria in grain legumes: Growth promotion and crop production. *Plants*. 2020; 9(11):1596. <https://doi.org/10.3390/plants9111596>
- Khan N, Peiman Z, Shahid A, Asif M, Muhammad AS, Jianjun Y. Impact of salicylic acid and PGPR on the drought tolerance and phytoremediation potential of *Helianthus annuus*. *Frontiers in Microbiology*, 2018; 9:2507. <https://doi.org/10.3389/fmicb.2018.02507>
- Pereira SIA, Abreu D, Moreira H, Vega A, Castro PML. Plant growth-promoting rhizobacteria (PGPR) improve the growth and nutrient use efficiency in maize (*Zea mays* L.) under water deficit conditions. *Heliyon*. 2020; 6(10):e05106. <https://doi.org/10.1016/j.heliyon.2020.e05106>
- Nawaz A, Shahbaz M, Asadullah, Imran A, Marghoob MU, Imtiaz M, Mubeen F. Potential of salt tolerant PGPR in growth and yield augmentation of wheat (*Triticum aestivum* L.) under saline conditions. *Frontiers in Microbiology*. 2020; 11. <https://doi.org/10.3389/fmicb.2020.02019>
- Sagar A, Sayyed RZ, Ramteke PW, Sharma S, Marraiki N, Elgorban AM, Syed A. ACC deaminase and antioxidant enzymes producing halophilic *Enterobacter* sp. PR14 promotes the growth of rice and millets under salinity stress. *Physiol Mol Biol Plants*. 2020; 9:1847-54. <https://doi.org/10.1007/s12298-020-00852-9>
- Khanna K, Jamwal VL, Gandhi SG, Ohri P, Bhardwaj R. Metal resistant PGPR lowered Cd uptake and expression of metal transporter genes with improved growth and photosynthetic pigments in *Lycopersicon esculentum* under metal toxicity. *Sci Rep*. 2019; 9: 5855. <https://doi.org/10.1038/s41598-019-41899-3>
- Sarkar J, Chakraborty U, Chakraborty B. High temperature resilience in *Bacillus safensis* primed wheat plants: A study of dynamic response associated with modulation of antioxidant machinery, differential expression of HSPs and osmolyte biosynthesis. *Environmental and Experimental Botany*. 2021; 182:104315. <https://doi.org/10.1016/j.envexpbot.2020.104315>
- Pérez-Montaño F, Alías-Villegas C, Bellogín RA, del Cerro P, Espuny MR, Jiménez-Guerrero I, López-Baena FJ, Ollero FJ, Cubo T. Plant growth promotion in cereal and leguminous agri-

- cultural important plants: From microorganism capacities to crop production. *Microbiological Research*. 2014; 169(5-6): 325-36. <https://doi.org/10.1016/j.micres.2013.09.011>
16. Vejan P, Abdullah R, Khadiran T, Ismail S, Nasrulhaq Boyce A. Role of plant growth promoting rhizobacteria in agricultural sustainability—a review. *Molecules*. 2016 May; 21(5):573. <https://doi.org/10.3390/molecules21050573>
 17. Shahid M. et al. Pesticides Pollution in Agricultural Soils of Pakistan. In: Hakeem K, Akhtar J, Sabir M Editors. *Soil Science: Agricultural and Environmental Perspectives*. Cham: Springer; 2016;199-229. https://doi.org/10.1007/978-3-319-34451-5_9
 18. Lavakush JY, Verma JP. Isolation and characterization of effective plant growth promoting rhizobacteria from rice rhizosphere of Indian soil. *Asian J Biol Sci*. 2012; 5:294-303. <https://dx.doi.org/10.3923/ajbs.2012.294.303>
 19. Hameed S, Yasmin S, Malik K A, Zafar Y, Hafeez FY. *Rhizobium*, *Bradyrhizobium* and *Agrobacterium* strains isolated from cultivated legumes. *Biol Fertil Soils*. 2004; 39: 179–85. <https://doi.org/10.1007/s00374-003-0697-z>
 20. Ramadoss D, Lakkineni VK, Bose P, Ali S, Annapurna K. Mitigation of salt stress in wheat seedlings by halotolerant bacteria isolated from saline habitats. *SpringerPlus*. 2013; 2:6. <https://doi.org/10.1186/2193-1801-2-6>
 21. Albdaawi RN, Khyami-Horani H, Ayad JY, Alananbeh KM, Al-Sayaydeh R. Isolation and characterization of halotolerant plant growth promoting rhizobacteria from durum wheat (*Triticum turgidum* subsp. *durum*) cultivated in saline areas of the dead sea region. *Frontiers in Microbiology*. 2019; 10:1639. <https://doi.org/10.3389/fmicb.2019.01639>
 22. Korir H, Mungai NW, Thuita M, Hamba Y, Masso C. Co-inoculation effect of rhizobia and plant growth promoting rhizobacteria on common bean growth in a low phosphorus soil. *Frontiers in Plant Science*. 2017;8:141. <https://doi.org/10.3389/fpls.2017.00141>
 23. Mukherjee B, Dutta S. Isolation of a phosphate solubilizing bacterial strain *Bacillus tequilensis* mcc 3872 from the rice field of burdwan district and characterization of its plant growth promoting traits. 2018;8(4): 956-62.
 24. Majeed A, Kaleem Abbasi M, Hameed S, Imran A, Rahim N. "Isolation and characterization of plant growth-promoting rhizobacteria from wheat rhizosphere and their effect on plant growth promotion." *Frontiers in Microbiology*. 2015; 6:198. <https://doi.org/10.3389/fmicb.2015.00198>
 25. Rahmoune B, Morsli A, Khelifi-Slaoui M, Khelifi L, Strueh A, Erban A, Kopka J, Prell J, van Dongen JT. Isolation and characterization of three new PGPR and their effects on the growth of *Arabidopsis* and *Datura* plants. *Journal of Plant Interactions*. 2017;12(1):1-6. <https://doi.org/10.1080/17429145.2016.1269215>
 26. Shultana R, Tan Kee Zuan A, Yusop MR, Mohd Saud H, Ayanda AF. Effect of salt-tolerant bacterial inoculations on rice seedlings differing in salt-tolerance under saline soil conditions. *Agronomy*. 2020;10(7):1030. <https://doi.org/10.3390/agronomy10071030>
 27. James C, Natalie S. *Microbiology. A laboratory manual*. Pearson Education; 2014.
 28. Dubois M, Gilles KA, Hamilton JK, Rebers PT, Smith F. Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*. 1956 Mar 1; 28(3):350-56. <https://doi.org/10.1021/ac60111a017>
 29. Bric JM, Bostock RM, Silverstone SE. Rapid in situ assay for indole acetic acid production by bacteria immobilized on a nitrocellulose membrane. *Applied and Environmental Microbiology*. 1991 Feb;57(2):535-38. <https://doi.org/10.1128/aem.57.2.535-538.1991>
 30. Akhtar N, Qureshi MA, Iqbal A, Ahmad MJ, Khan KH. Influence of *Azotobacter* and IAA on symbiotic performance of *Rhizobium* and yield parameters of lentil. *J Agric Res*. 2012; 50:361–72.
 31. Ahemad M, Kibret M. Mechanisms and applications of plant growth promoting rhizobacteria: current perspective. *Journal of King Saud University-Science*. 2014;26(1):1-20. <https://doi.org/10.1016/j.jksus.2013.05.001>
 32. Dutta S, Sarkar A, Dutta S. Characterization of *Pseudomonas aeruginosa* MCC 3198 and its potential for growth promotion of seedlings of the medicinal plant *Celosia cristata* L. *Intern J Curr Microbiol App Sci*. 2019; 8(4):985-97. <https://doi.org/10.20546/ijcmas.2019.804.114>
 33. John C J, Kumar S, Ge M. Probiotic prospects of PGPR for green and sustainable agriculture. *Archives of Phytopathology and Plant Protection*. 2020; 53:1-16. <https://doi.org/10.1080/03235408.2020.1805901>
 34. Rathore P. A review on approaches to develop plant growth promoting rhizobacteria. *International Journal of Recent Scientific Research*. 2014; 5:403-07.
 35. Ramadan RA, Gebriel MG, Kadry HM, Mosallem A. Carbapenem-resistant *Acinetobacter baumannii* and *Pseudomonas aeruginosa*: characterization of carbapenemase genes and E-test evaluation of colistin-based combinations. *Infection and drug resistance*. 2018;11:1261. <https://doi.org/10.2147/IDR.S170233>
 36. Santoro MV, Bogino PC, Nocelli N, Cappellari LD, Giordano WF, Banchio E. Analysis of plant growth-promoting effects of fluorescent *Pseudomonas* strains isolated from *Mentha piperita* rhizosphere and effects of their volatile organic compounds on essential oil composition. *Frontiers in Microbiology*. 2016; 7:1085. <https://doi.org/10.3389/fmicb.2016.01085>
 37. Narożna D, Pudełko K, Królczyk J, Golińska B, Sugawara M, Mądrzak CJ, Sadowsky MJ. Survival and competitiveness of *Bradyrhizobium japonicum* strains 20 years after introduction into field locations in Poland. *Applied and Environmental Microbiology*. 2015;81(16):5552-59. <https://doi.org/10.1128/AEM.01399-15>
 38. Orlandini V, Emiliani G, Fondi M, Maida I, Perrin E, Fani R. Network analysis of plasmidomes: The *Azospirillum brasilense* Sp245 case. *International Journal of Evolutionary Biology*. 2014;2014. <http://dx.doi.org/10.1155/2014/951035>
 39. Ma Y, Rajkumar M, Freitas H. Inoculation of plant growth promoting bacterium *Achromobacter xylosoxidans* strain Ax10 for the improvement of copper phytoextraction by *Brassica juncea*. *Journal of Environmental Management*. 2009;90(2):331-37. <https://doi.org/10.1016/j.jenvman.2008.01.014>
 40. Prathap M, Kumari BR. A critical review on plant growth promoting rhizobacteria. *Journal of Plant Pathology and Microbiology*. 2015;6(4). <https://doi.org/10.4172/2157-7471.1000266>
 41. Oyedele AO, Ogunbanwo TS. Antifungal activities of *Bacillus subtilis* isolated from some condiments and soil. *African Journal of Microbiology Research*. 2014 Apr 30;8(18):1841-49. <https://doi.org/10.5897/AJMR2013.6162>
 42. Sanders JW, Martin JW, Hooke M, Hooke J. *Methylobacterium mesophilicum* infection: case report and literature review of an unusual opportunistic pathogen. *Clinical Infectious Diseases*. 2000 Jun 1;30(6):936-38. <https://doi.org/10.1086/313815>
 43. Agbodjato NA, Noumavo PA, Baba-Moussa F, Salami HA, Sina H, Sèzan A, Bankolé H, Adjanohoun A, Baba-Moussa L. Characterization of potential plant growth promoting rhizobacteria isolated from Maize (*Zea mays* L.) in central and Northern Benin (West Africa). *Applied and Environmental Soil Science*. 2015 Oct;2015. <https://doi.org/10.1155/2015/901656>
 44. Kuan KB, Othman R, Abdul Rahim K, Shamsuddin ZH. Plant growth-promoting rhizobacteria inoculation to enhance vegetative growth, nitrogen fixation and nitrogen remobilisation of

- maize under greenhouse conditions. *PLoS one*. 2016 Mar 24; 11(3):e0152478. <https://doi.org/10.1371/journal.pone.0152478>
45. Vessey JK. Plant growth promoting rhizobacteria as biofertilizers. *Plant and Soil*. 2003 Aug; 255(2):571-86. <https://doi.org/10.1023/A:1026037216893>
 46. Martínez-Viveros O, Jorquera MA, Crowley DE, Gajardo GM, Mora ML. Mechanisms and practical considerations involved in plant growth promotion by rhizobacteria. *Journal of Soil Science and Plant Nutrition*. 2010 Jul; 10(3):293-319. <http://dx.doi.org/10.4067/S0718-95162010000100006>
 47. Somers E, Vanderleyden J, Srinivasan M. Rhizosphere bacterial signalling: a love parade beneath our feet. *Critical reviews in microbiology*. 2004 Jan 1;30(4):205-40. <https://doi.org/10.1080/10408410490468786>
 48. Wu CH, Bernard SM, Andersen GL, Chen W. Developing microbe-plant interactions for applications in plant growth promotion and disease control, production of useful compounds, remediation and carbon sequestration. *Microbial biotechnology*. 2009 Jul; 2(4):428-40. <https://doi.org/10.1111/j.1751-7915.2009.00109.x>
 49. Mehmood U, Inam-ul-Haq M, Saeed M, Altaf A, Azam F, Hayat S. A brief review on plant growth promoting Rhizobacteria (PGPR): a key role in plant growth promotion. *Plant protection*. 2018 Sep 11;2(2):77-82. <http://esciencepress.net/journals/PP>
 50. Liu X, Li Q, Li Y, Guan G, Chen S. *Paenibacillus* strains with nitrogen fixation and multiple beneficial properties for promoting plant growth. *Peer J*. 2019 Sep 23; 7:e7445. <http://dx.doi.org/10.7717/peerj.7445>
 51. Woo OG, Kim H, Kim JS, Keum HL, Lee KC, Sul WJ, Lee JH. *Bacillus subtilis* strain GOT9 confers enhanced tolerance to drought and salt stresses in *Arabidopsis thaliana* and *Brassica campestris*. *Plant Physiology and Biochemistry*. 2020 Mar 1; 148:359-67. <https://doi.org/10.1016/j.plaphy.2020.01.032>
 52. Manasa K, Reddy S, Triveni S. Characterization of potential PGPR and antagonistic activities of *Rhizobium* isolates from different rhizosphere soils. *Journal of Pharmacognosy and Phytochemistry*. 2017; 6(3):51-54. <https://doi.org/10.20546/ijcmas.2017.605.316>
 53. Khan N, Bano A. Exopolysaccharide producing rhizobacteria and their impact on growth and drought tolerance of wheat grown under rainfed conditions. *PLoS One*. 2019 Sep 12;14(9):e0222302. <https://doi.org/10.1371/journal.pone.0222302>
 54. Ghosh D, Gupta A, Mohapatra S. Dynamics of endogenous hormone regulation in plants by phytohormone secreting rhizobacteria under water-stress. *Symbiosis*. 2019 Mar;77(3):265-78. <https://doi.org/10.1007/s13199-018-00589-w>
 55. Naseem H, Ahsan M, Shahid MA, Khan N. Exopolysaccharides producing rhizobacteria and their role in plant growth and drought tolerance. *Journal of Basic Microbiology*. 2018 Dec; 58(12):1009-22. <https://doi.org/10.1002/jobm.201800309>
 56. Lenin G, Jayanthi M. Indole acetic acid, gibberellic acid and siderophore production by PGPR isolates from rhizospheric soils of *Catharanthus roseus*. *Int J Pharmacy Biol Arch*. 2012;3:933-38.
 57. Glick BR. Phytoremediation: synergistic use of plants and bacteria to clean up the environment. *Biotechnology Advances*. 2003 Aug 1;21(5):383-93. [https://doi.org/10.1016/S0734-9750\(03\)00055-7](https://doi.org/10.1016/S0734-9750(03)00055-7)
 58. Gouda S, Kerry RG, Das G, Paramithiotis S, Shin HS, Patra JK. Revitalization of plant growth promoting rhizobacteria for sustainable development in agriculture. *Microbiological Research*. 2018 Jan 1;206:131-40. <https://doi.org/10.1016/j.micres.2017.08.016>
 59. Prasad M, Srinivasan R, Chaudhary M, Choudhary M, Jat LK. Plant growth promoting rhizobacteria (PGPR) for sustainable agriculture: perspectives and challenges. In: Singh AK, Kumar A, Singh PK Editors. *PGPR amelioration in sustainable agriculture*. Woodhead; 2019 Jan 1;129-57. <https://doi.org/10.1016/B978-0-12-815879-1.00007-0>
 60. Dimkpa C, Weinand T, Asch F. Plant-rhizobacteria interactions alleviate abiotic stress conditions. *Plant, Cell and Environment*. 2009 Dec; 32(12):1682-94. <https://doi.org/10.1111/j.1365-3040.2009.02028.x>
 61. Lebrazi S, Niehaus K, Bednarz H, Fadil M, Chraibi M, Fikri-Benbrahim K. Screening and optimization of indole-3-acetic acid production and phosphate solubilization by rhizobacterial strains isolated from *Acacia cyanophylla* root nodules and their effects on its plant growth. *Journal of Genetic Engineering and Biotechnology*. 2020; 18(1):1-2. <https://doi.org/10.1007/s00284-010-9674-6>
 62. Piromy P, Buranabanyat B, Tantasawat P, Tittabutr P, Boonkerd N, Teamroong N. Effect of plant growth promoting rhizobacteria (PGPR) inoculation on microbial community structure in rhizosphere of forage corn cultivated in Thailand. *European Journal of Soil Biology*. 2011 Jan 1; 47(1):44-54. <https://doi.org/10.1016/j.ejsobi.2010.11.004>
 63. Montañez A, Abreu C, Gill PR, Hardarson G, Sicardi M. Biological nitrogen fixation in maize (*Zea mays* L.) by ¹⁵N isotope-dilution and identification of associated culturable diazotrophs. *Biology and Fertility of Soils*. 2009; 45(3):253-63. <https://doi.org/10.1007/s00374-008-0322-2>
 64. Mahmood S, Daur I, Al-Solaimani SG, Ahmad S, Madkour MH, Yasir M, Hirt H, Ali S, Ali Z. Plant growth promoting rhizobacteria and silicon synergistically enhance salinity tolerance of mung bean. *Frontiers in Plant Science*. 2016; 7:876. <https://doi.org/10.3389/fpls.2016.00876>
 65. Singh RK, Singh P, Li HB, Song QQ, Guo DJ, Solanki MK, Verma KK, Malviya MK, Song XP, Lakshmanan P, Yang LT. Diversity of nitrogen-fixing rhizobacteria associated with sugarcane: a comprehensive study of plant-microbe interactions for growth enhancement in *Saccharum* spp. *BMC Plant Biology*. 2020 Dec; 20:1-21. <https://doi.org/10.1186/s12870-020-02400-9>
 66. Deshwal VK, Pandey P, Kang SC, Maheshwari DK. Rhizobia as a biological control agent against soil borne plant pathogenic fungi. *Indian Journal of Experimental Biology*. 2003; 41:1160-64.
 67. Akhtar S, Ali B. Evaluation of rhizobacteria as non-rhizobial inoculants for mung beans. *Australian Journal of Crop Science*. 2011 Jan; 5(13):1723-29.
 68. Thakuria D, Talukdar NC, Goswami C, Hazarika S, Boro RC, Khan MR. Characterization and screening of bacteria from rhizosphere of rice grown in acidic soils of Assam. *Current Science*. 2004 Apr 10; 978-85.
 69. Rajkumar M, Freitas H. Effects of inoculation of plant-growth promoting bacteria on Ni uptake by Indian mustard. *Biore-source Technology*. 2008 Jun 1; 99(9):3491-98. <https://doi.org/10.1016/j.biortech.2007.07.046>
 70. Berraho EL, Lesueur D, Diem HG, Sasson A. Iron requirement and siderophore production in *Rhizobium ciceri* during growth on an iron-deficient medium. *World Journal of Microbiology and Biotechnology*. 1997 Sep; 13(5):501-10. <https://doi.org/10.1023/A:1018553022960>
 71. Wittenberg JB, Wittenberg BA, Day DA, Udvardi MK, Appleby CA. Siderophore-bound iron in the peribacteroid space of soybean root nodules. *Plant and soil*. 1996 Jan; 178(2):161-69. <https://doi.org/10.1007/BF00011579>
 72. Burd GI, Dixon DG, Glick BR. Plant growth-promoting bacteria that decrease heavy metal toxicity in plants. *Canadian Journal of Microbiology*. 2000 Mar 1; 46(3):237-45. <https://doi.org/10.1139/w99-143>

73. Dey RK, Pal KK, Bhatt DM, Chauhan SM. Growth promotion and yield enhancement of peanut (*Arachis hypogaea* L.) by application of plant growth-promoting rhizobacteria. *Microbiological Research*. 2004 Dec 15; 159(4):371-94. <https://doi.org/10.1016/j.micres.2004.08.004>
74. Ahemad M, Khan MS. Effects of insecticides on plant-growth-promoting activities of phosphate solubilizing rhizobacterium *Klebsiella* sp. strain PS19. *Pesticide Biochemistry and Physiology*. 2011 May 1;100(1):51-56. <https://doi.org/10.1016/j.pestbp.2011.02.004>
75. Ahemad M, Khan MS. Biotoxic impact of fungicides on plant growth promoting activities of phosphate-solubilizing *Klebsiella* sp. isolated from mustard (*Brassica campestris*) rhizosphere. *Journal of Pest Science*. 2012 Mar;85(1):29-36. <https://doi.org/10.1007/s10340-011-0402-1>
76. Wani PA, Khan MS, Zaidi A. Synergistic effects of the inoculation with nitrogen fixing and phosphate solubilizing rhizobacteria on the performance of field grown chickpea. *Journal of Plant Nutrition and Soil Science*. 2007 Apr;170(2):283-87. <https://doi.org/10.1002/jpln.200620602>
77. Selvakumar G, Mohan M, Kundu S, Gupta AD, Joshi P, Nazim S, Gupta HS. Cold tolerance and plant growth promotion potential of *Serratia marcescens* strain SRM (MTCC 8708) isolated from flowers of summer squash (*Cucurbita pepo*). *Letters in Applied Microbiology*. 2008 Feb;46(2):171-75. <https://doi.org/10.1111/j.1472-765X.2007.02282.x>
78. Phi QT, Park YM, Seul KJ, Ryu CM, Park SH, Kim JG, Ghim SY. Assessment of root-associated *Paenibacillus polymyxa* groups on growth promotion and induced systemic resistance in pepper. *Journal of Microbiology and Biotechnology*. 2010;20(12):1605-13. <https://doi.org/10.4014/jmb.1007.07014>
79. Naik MM, Dubey SK. Lead-enhanced siderophore production and alteration in cell morphology in a Pb-resistant *Pseudomonas aeruginosa* strain 4EA. *Current Microbiology*. 2011 Feb;62(2):409-14. <https://doi.org/10.1007/s00284-010-9722-2>
80. Ferreira MJ, Silva H, Cunha A. Siderophore-producing rhizobacteria as a promising tool for empowering plants to cope with iron limitation in saline soils: a review. *Pedosphere*. 2019 Aug 1;29(4):409-20. [https://doi.org/10.1016/S1002-0160\(19\)60810-6](https://doi.org/10.1016/S1002-0160(19)60810-6)
81. Rani A, Souche YS, Goel R. Comparative assessment of *in situ* bioremediation potential of cadmium resistant acidophilic *Pseudomonas putida* 62BN and alkalophilic *Pseudomonas monteilii* 97AN strains on soybean. *International Biodeterioration and Biodegradation*. 2009 Jan 1;63(1):62-66. <https://doi.org/10.1016/j.ibiod.2008.07.002>
82. Antoun H, Beauchamp CJ, Goussard N, Chabot R, Lalande R. Potential of *Rhizobium* and *Bradyrhizobium* species as plant growth promoting rhizobacteria on non-legumes: effect on radishes (*Raphanus sativus* L.). In: Hardarson G, Broughton WJ Editors. *Molecular Microbial Ecology of the Soil*. Vol. 83. Dordrecht: Springer; 1998; 57-67. https://doi.org/10.1007/978-94-017-2321-3_5
83. Verma A, Kukreja K, Pathak D, Suneja S, Narula N. *In vitro* production of plant growth regulators (PGRs) by *Azotobacter chroococcum*. *Indian J Microbiol*. 2001;41:305-07.
84. Shaharoon B, Arshad M, Zahir ZA. Effect of plant growth promoting rhizobacteria containing ACC deaminase on maize (*Zea mays* L.) growth under axenic conditions and on nodulation in mung bean (*Vigna radiata* L.). *Letters in Applied Microbiology*. 2006 Feb;42(2):155-59. <https://doi.org/10.1111/j.1472-765X.2005.01827.x>
85. Tsavkelova EA, Cherdyntseva TA, Netrusov AI. Auxin production by bacteria associated with orchid roots. *Microbiology*. 2005;74(1):46-53. <https://doi.org/10.1007/s11021-005-0027-6>
86. Vivas A, Biro B, Ruiz-Lozano JM, Barea JM, Azcon R. Two bacterial strains isolated from a Zn-polluted soil enhance plant growth and mycorrhizal efficiency under Zn-toxicity. *Chemosphere*. 2006 Mar 1;62(9):1523-33. <https://doi.org/10.1016/j.chemosphere.2005.06.053>
87. Jha PN, Kumar A. Endophytic colonization of *Typha australis* by a plant growth promoting bacterium *Klebsiella oxytoca* strain GR 3. *Journal of Applied Microbiology*. 2007 Oct;103(4):1311-20. <https://doi.org/10.1111/j.1365-2672.2007.03383.x>
88. Jiang CY, Sheng XF, Qian M, Wang QY. Isolation and characterization of a heavy metal-resistant *Burkholderia* sp. from heavy metal-contaminated paddy field soil and its potential in promoting plant growth and heavy metal accumulation in metal-polluted soil. *Chemosphere*. 2008 May 1;72(2):157-64. <https://doi.org/10.1016/j.chemosphere.2008.02.006>
89. Kumar KV, Singh N, Behl HM, Srivastava S. Influence of plant growth promoting bacteria and its mutant on heavy metal toxicity in *Brassica juncea* grown in fly ash amended soil. *Chemosphere*. 2008 Jun 1;72(4):678-83. <https://doi.org/10.1016/j.chemosphere.2008.03.025>
90. Mehnaiz S, Bauer JS, Gross H. Complete genome sequence of the sugar cane endophyte *Pseudomonas aurantiaca* PB-St2, a disease-suppressive bacterium with antifungal activity toward the plant pathogen *Colletotrichum falcatum*. *Genome Announcements*. 2014 Feb 27;2(1):e01108-13. <https://doi.org/10.1128/genomeA.01108-13>
91. Yousef NM. Capability of plant growth-promoting rhizobacteria (PGPR) for producing indole acetic acid (IAA) under extreme conditions. *European Journal of Biological Research*. 2018 Sep 10;8(4):174-82. <http://dx.doi.org/10.5281/zenodo.1412796>
92. Noel TC, Sheng C, Yost CK, Pharis RP, Hynes MF. *Rhizobium leguminosarum* as a plant growth-promoting rhizobacterium: direct growth promotion of canola and lettuce. *Canadian Journal of Microbiology*. 1996;42(3):279-83. <https://doi.org/10.1139/m96-040>
93. Liu F, Xing S, Ma H, Du Z, Ma B. Cytokinin-producing, plant growth-promoting rhizobacteria that confer resistance to drought stress in *Platycladus orientalis* container seedlings. *Applied Microbiology and Biotechnology*. 2013; 97(20):9155-64. <https://doi.org/10.1007/s00253-013-5193-2>
94. Deka H, Deka S, Baruah CK. Plant Growth Promoting Rhizobacteria for Value Addition: Mechanism of Action. In: Egamberdieva D, Shrivastava S, Verma A Editors. *Plant-Growth-Promoting Rhizobacteria (PGPR) and Medicinal Plants*. Soil biology. Vol. 42. Cham: Springer; 2015;305-21. https://doi.org/10.1007/978-3-319-13401-7_15
95. Kang SM, Khan AL, Waqas M, You YH, Hamayun M, Joo GJ, Shahzad R, Choi KS, Lee IJ. Gibberellin-producing *Serratia nematodiphila* PEJ1011 ameliorates low temperature stress in *Capsicum annuum* L. *European Journal of Soil Biology*. 2015; 68:85-93. <https://doi.org/10.1016/j.ejsobi.2015.02.005>
96. Elliott GN, Chen WM, Chou JH, Wang HC, Sheu SY, Perin L, Reis VM, Moulin L, Simon MF, Bontemps C, Sutherland JM. *Burkholderia phymatum* is a highly effective nitrogen fixing symbiont of *Mimosa* spp. and fixes nitrogen ex planta. *New Phytologist*. 2007;173(1):168-80. <https://doi.org/10.1111/j.1469-8137.2006.01894.x>
97. Van der Eerden LJ. Toxicity of ammonia to plants. *Agriculture and Environment*. 1982; 7(3-4):223-35. [https://doi.org/10.1016/0304-1131\(82\)90015-7](https://doi.org/10.1016/0304-1131(82)90015-7)
98. Malboobi MA, Owlia P, Behbahani M, Sarokhani E, Moradi S, Yakhchali B, Deljou A, Heravi KM. Solubilization of organic and inorganic phosphates by three highly efficient soil bacterial isolates. *World Journal of Microbiology and Biotechnology*. 2009 Aug;25(8):1471-77. <https://doi.org/10.1007/s11274-009-0037-z>

99. Chen YP, Rekha PD, Arun AB, Shen FT, Lai WA, Young CC. Phosphate solubilizing bacteria from subtropical soil and their tricalcium phosphate solubilizing abilities. *Applied Soil Ecology*. 2006 Nov 1;34(1):33-41. <https://doi.org/10.1016/j.apsoil.2005.12.002>
100. Otieno N, Lally RD, Kiwanuka S, Lloyd A, Ryan D, Germaine KJ, Dowling DN. Plant growth promotion induced by phosphate solubilizing endophytic *Pseudomonas* isolates. *Frontiers in Microbiology*. 2015 Jul 22; 6:745. <https://doi.org/10.3389/fmicb.2015.00745>
101. Baas P, Bell C, Mancini LM, Lee MN, Conant RT, Wallenstein MD. Phosphorus mobilizing consortium Mammoth P™ enhances plant growth. *PeerJ*. 2016; 4:e2121. <https://doi.org/10.7717/peerj.2121>
102. Khan Z, Rho H, Firrincieli A, Hung SH, Luna V, Masciarelli O, Kim SH, Doty SL. Growth enhancement and drought tolerance of hybrid poplar upon inoculation with endophyte consortia. *Current Plant Biology*. 2016;6:38-47. <https://doi.org/10.1016/j.cpb.2016.08.001>
103. Gupta G, Parihar SS, Ahirwar NK, Snehi SK, Singh V. Plant growth promoting rhizobacteria (PGPR): current and future prospects for development of sustainable agriculture. *J Microb Biochem Technol*. 2015 Mar;7(2):96-102. <http://dx.doi.org/10.4172/1948-5948.1000188>
104. Omer ZS, Tombolini R, Broberg A, Gerhardson B. Indole-3-acetic acid production by pink-pigmented facultative methylotrophic bacteria. *Plant Growth Regulation*. 2004 May;43(1):93-96. <https://doi.org/10.1023/B:GROW.0000038360.09079.ad>
105. Kumar A, Bahadur I, Maurya BR, Raghuvanshi R, Meena VS, Singh DK, Dixit J. Does a plant growth-promoting rhizobacteria enhance agricultural sustainability. *J Pure Appl Microbiol*. 2015 Mar; 9(1):715-24.
106. Ruzzi M, Aroca R. Plant growth-promoting rhizobacteria act as biostimulants in horticulture. *Scientia Horticulturae*. 2015 Nov 30; 196:124-34. <https://doi.org/10.1016/j.scienta.2015.08.042>
107. Hong Y, Glick BR, Pasternak JJ. Plant-microbial interaction under gnotobiotic conditions: a scanning electron microscope study. *Current Microbiology*. 1991 Aug; 23(2):111-14. <https://doi.org/10.1007/BF02092259>
108. Llorente BE, Alasia MA, Larraburu EE. Biofertilization with *Azospirillum brasilense* improves *in vitro* culture of *Handroanthus ochraceus*, a forestry, ornamental and medicinal plant. *New biotechnology*. 2016 Jan 25; 33(1):32-40. <https://doi.org/10.1016/j.nbt.2015.07.006>
109. Spaepen S, Bossuyt S, Engelen K, Marchal K, Vanderleyden J. Phenotypical and molecular responses of *Arabidopsis thaliana* roots as a result of inoculation with the auxin producing bacterium *Azospirillum brasilense*. *New Phytologist*. 2014 Feb; 201(3):850-61. <https://doi.org/10.1111/nph.12590>
110. Jha CK, Saraf M. Plant growth promoting rhizobacteria (PGPR). *Journal of agriculture research and development*. 2015;5(2):0108-19.
111. Arkhipova TN, Veselov SU, Melentiev AI, Martynenko EV, Kudoyarova GR. Ability of bacterium *Bacillus subtilis* to produce cytokinins and to influence the growth and endogenous hormone content of lettuce plants. *Plant and Soil*. 2005 May; 272(1):201-19. <https://doi.org/10.1007/s11104-004-5047-x>
112. Siegień I, Bogatek R. Cyanide Action in Plants—from toxic to regulatory. *Acta Physiologiae Plantarum*. 2006 Oct 1;28(5):483-97. <https://doi.org/10.1007/BF02706632>
113. Rijavec T, Lapanje A. Hydrogen cyanide in the rhizosphere: not suppressing plant pathogens, but rather regulating availability of phosphate. *Frontiers in Microbiology*. 2016; 7:1785. <https://doi.org/10.3389/fmicb.2016.01785>
114. Rezzonico F, Zala M, Keel C, Duffy B, Moëne Locozy Y, Défago G. Is the ability of biocontrol fluorescent pseudomonads to produce the antifungal metabolite 2, 4 diacetylphloroglucinol really synonymous with higher plant protection?. *New Phytologist*. 2007; 173(4):861-72. <https://doi.org/10.1111/j.1469-8137.2006.01955.x>
115. Neilands JB. Siderophores: structure and function of microbial iron transport compounds. *Journal of Biological Chemistry*. 1995 Nov 10; 270(45):26723-26. <https://doi.org/10.1074/jbc.270.45.26723>
116. Page MG. The role of iron and siderophores in infection, and the development of siderophore antibiotics. *Clinical Infectious Diseases*. 2019 Nov 13; 69(Supplement_7):S529-37. <https://doi.org/10.1093/cid/ciz825>
117. Albelda-Berenguer M, Monachon M, Joseph E. Siderophores: From natural roles to potential applications. *Advances in Applied Microbiology*. 2019 Jan 1;106:193-225. <https://doi.org/10.1016/bs.aambs.2018.12.001>
118. Sureshababu K, Amaresan N, Kumar K. Amazing multiple function properties of plant growth promoting rhizobacteria in the rhizosphere soil. *Int J Curr Microbiol Appl Sci*. 2016; 5(2):661-83. <http://dx.doi.org/10.20546/ijcmas.2016.501.074>
119. Sayyed RZ, Naphade BS, Chincholkar SB. Ecologically competent rhizobacteria for plant growth promotion and disease management. *Recent Trends in Biotechnology*. 2004;1:1-6.
120. Compant S, Reiter B, Sessitsch A, Nowak J, Clément C, Ait Barka E. Endophytic colonization of *Vitis vinifera* L. by plant growth-promoting bacterium *Burkholderia* sp. strain PsJN. *Applied and Environmental Microbiology*. 2005;71(4):1685-93. <https://doi.org/10.1128/AEM.71.4.1685-1693.2005>
121. Zhuang X, Chen J, Shim H, Bai Z. New advances in plant growth-promoting rhizobacteria for bioremediation. *Environment International*. 2007;33(3):406-13. <https://doi.org/10.1016/j.envint.2006.12.005>
122. Herrero M, Stuckey DC. Bioaugmentation and its application in wastewater treatment: a review. *Chemosphere*. 2015 Dec 1;140:119-28. <https://doi.org/10.1016/j.chemosphere.2014.10.033>
123. Kuiper I, Lagendijk EL, Bloemberg GV, Lugtenberg BJ. Rhizoremediation: a beneficial plant-microbe interaction. *Molecular Plant-Microbe Interactions*. 2004;17(1):6-15. <https://apsjournals.apsnet.org/doi/abs/10.1094/MPMI.2004.17.1.6#>
124. Gull A, Lone AA, Wani NU. Biotic and Abiotic Stresses in Plants. 2019;7:1-19. 10.5772/intechopen.85832 <https://doi.org/10.5772/intechopen.85832>
125. Ramegowda V, Senthil-Kumar M. The interactive effects of simultaneous biotic and abiotic stresses on plants: mechanistic understanding from drought and pathogen combination. *Journal of Plant Physiology*. 2015;176:47-54. <https://doi.org/10.1016/j.jplph.2014.11.008>
126. Goswami D, Thakker JN, Dhandhukia PC. Portraying mechanics of plant growth promoting rhizobacteria (PGPR): A review. *Cogent Food and Agriculture*. 2016;2(1). <https://doi.org/10.1080/23311932.2015.1127500>
127. Vurukonda SS, Vardharajula S, Shrivastava M, SkZ A. Enhancement of drought stress tolerance in crops by plant growth promoting rhizobacteria. *Microbiological Research*. 2016;184:13-24. <https://doi.org/10.1016/j.micres.2015.12.003>
128. Jadhav HP, Sayyed RZ. Hydrolytic enzymes of rhizospheric microbes in crop protection. *MOJ Cell Sci Rep*. 2016;3(5):135-36. <https://doi.org/10.15406/mojcsr.2016.03.00070>
129. Ilyas N, Mumtaz K, Akhtar N, Yasmin H, Sayyed RZ, Khan W, Enshasy HA, Dailin DJ, Elsayed EA, Ali Z. Exopolysaccharides producing bacteria for the amelioration of drought stress in

- wheat. Sustainability. 2020; 12(21):8876. <https://doi.org/10.3390/su12218876>
130. Meena KK, Sorty AM, Bitla UM, Choudhary K, Gupta P, Pareek A, Singh DP, Prabha R, Sahu PK, Gupta VK, Singh HB. Abiotic stress responses and microbe-mediated mitigation in plants: the omics strategies. Frontiers in Plant Science. 2017;8:172. <https://doi.org/10.3389/fpls.2017.00172>
 131. Onofre-Lemus J, Hernández-Lucas I, Girard L, Caballero-Mellado J. ACC (1-aminocyclopropane-1-carboxylate) deaminase activity, a widespread trait in *Burkholderia* species and its growth-promoting effect on tomato plants. Applied and Environmental Microbiology. 2009;75(20):6581-90. <https://doi.org/10.1128/AEM.01240-09>
 132. Saleem M, Arshad M, Hussain S, Bhatti AS. Perspective of plant growth promoting rhizobacteria (PGPR) containing ACC deaminase in stress agriculture. Journal of Industrial Microbiology and Biotechnology. 2007 Oct 1; 34(10):635-48. <https://doi.org/10.1007/s10295-007-0240-6>
 133. Sade N, Gebretsadik M, Seligmann R, Schwartz A, Wallach R, Moshelion M. The role of tobacco Aquaporin1 in improving water use efficiency, hydraulic conductivity and yield production under salt stress. Plant Physiology. 2010; 152(1): 245-54. <https://doi.org/10.1104/pp.109.145854>
 134. Tahir HA, Gu Q, Wu H, Raza W, Safdar A, Huang Z, Rajer FU, Gao X. Effect of volatile compounds produced by *Ralstonia solanacearum* on plant growth promoting and systemic resistance inducing potential of *Bacillus* volatiles. BMC Plant Biology. 2017;17(1):1-6. <https://doi.org/10.1186/s12870-017-1083-6>
 135. Fahad S, Hussain S, Bano A, Saud S, Hassan S, Shan D, Khan FA, Khan F, Chen Y, Wu C, Tabassum MA. Potential role of phytohormones and plant growth-promoting rhizobacteria in abiotic stresses: consequences for changing environment. Environmental Science and Pollution Research. 2015 Apr;22(7):4907-21. <https://doi.org/10.1007/s11356-014-3754-2>
 136. Barnawal D, Bharti N, Pandey SS, Pandey A, Chanotiya CS, Kalra A. Plant growth promoting rhizobacteria enhance wheat salt and drought stress tolerance by altering endogenous phytohormone levels and TaCTR1/TaDREB2 expression. Physiologia Plantarum. 2017 Dec;161(4):502-14. <https://doi.org/10.1111/ppl.12614>
 137. Lau JA, Lennon JT. Evolutionary ecology of plant-microbe interactions: soil microbial structure alters selection on plant traits. New Phytologist. 2011 Oct; 192(1):215-24. <https://doi.org/10.1111/j.1469-8137.2011.03790.x>
 138. Egamberdieva D, Kucharova Z. Selection for root colonising bacteria stimulating wheat growth in saline soils. Biology and Fertility of Soils. 2009 Jul;45(6):563-71. <https://doi.org/10.1007/s00374-009-0366-y>
 139. Khan A, Sayyed RZ, Seifi S. Rhizobacteria: Legendary Soil Guards in Abiotic Stress Management. In: Sayyed R, Arora N, Reddy M Editors. Plant Growth Promoting Rhizobacteria for Sustainable Stress Management. Microorganisms for Sustainability. Vol. 12. Singapore: Springer; 2019;327-34. https://doi.org/10.1007/978-981-13-6536-2_15
 140. Ayaz A, Saqib S, Huang H, Zaman W, Lü S, Zhao H. Genome-wide comparative analysis of long-chain acyl-CoA synthetases (LACSSs) gene family: A focus on identification, evolution and expression profiling related to lipid synthesis. Plant Physiology and Biochemistry. 2021 Apr 1;161:1-11. <https://doi.org/10.1016/j.plaphy.2021.01.042>
 141. Dhawi F. Plant growth promoting Rhizobacteria (PGPR) regulated Phyto and microbial beneficial protein interactions. Open Life Sciences. 2020 Jan 1;15(1):68-78. <https://doi.org/10.1515/biol-2020-0008>
 142. Nedjimi B. Seasonal variation in productivity, water relations and ion contents of *Atriplex halimus* spp. schweinfurthii grown in Chott Zehrez wetland, Algeria. Journal of the Saudi Society of Agricultural Sciences. 2012 Jan 1;11(1):43-49. <https://doi.org/10.1016/j.jssas.2011.08.002>
 143. Pan J, Peng F, Xue X, You Q, Zhang W, Wang T, Huang C. The growth promotion of two salt-tolerant plant groups with PGPR inoculation: a meta-analysis. Sustainability. 2019; 11(2):378. <https://doi.org/10.3390/su11020378>
 144. Mhlongo MI, Piater LA, Steenkamp PA, Labuschagne N, Dubery IA. Metabolic profiling of PGPR-treated tomato plants reveal priming-related adaptations of secondary metabolites and aromatic amino acids. Metabolites. 2020 May; 10(5):210. <https://doi.org/10.3390/metabo10050210>
 145. Garcia-Seco D, Zhang Y, Gutierrez-Manero FJ, Martin C, Ramos-Solano B. Application of *Pseudomonas fluorescens* to blackberry under field conditions improves fruit quality by modifying flavonoid metabolism. PLoS One. 2015 Nov 11; 10(11):e0142639. <https://doi.org/10.1371/journal.pone.0142639>
 146. Basha SA, Chatterjee SC. Effect of PGPR on *Scierotinia scierotiorum* infection through elicitation of phenylalanine ammonia lyase in chickpea. Indian Phytopathology. 2012.
 147. Gururani MA, Upadhyaya CP, Baskar V, Venkatesh J, Nookaraju A, Park SW. Plant growth-promoting rhizobacteria enhance abiotic stress tolerance in *Solanum tuberosum* through inducing changes in the expression of ROS-scavenging enzymes and improved photosynthetic performance. Journal of Plant Growth Regulation. 2013; 32(2):245-58. <https://doi.org/10.1007/s00344-012-9292-6>
 148. Tarafdar A, Raliya R, Wang WN, Biswas P, Tarafdar JC. Green synthesis of TiO₂ nanoparticle using *Aspergillus tubingensis*. Advanced Science, Engineering and Medicine. 2013 Sep 1;5 (9):943-49. <https://doi.org/10.1166/asem.2013.1376>

§§§