



RESEARCH ARTICLE

# Extraction and characterization of oil from seeds of the medicinal plant *Withania coagulans* (Stocks) Dunal (Doda paneer)

Priya Raghav<sup>1</sup>, Priyavart Choudhary<sup>1</sup>, Ajay Singh<sup>2</sup>, Rohit Sharma<sup>3</sup> & Nishesh Sharma<sup>1\*</sup>

<sup>1</sup>Department of Biotechnology, SALS, Uttarakhand University, Dehradun 248 007, India

<sup>2</sup>Department of Chemistry, SALS, Uttarakhand University, Dehradun 248 007, India

<sup>3</sup>Department of Chemistry, SOE, University of Petroleum and Energy Studies, Dehradun 248 007, India

\*Email: [hodbiotechnology@uttarakhanduniversity.ac.in](mailto:hodbiotechnology@uttarakhanduniversity.ac.in)

 OPEN ACCESS

## ARTICLE HISTORY

Received: 29 December 2021

Accepted: 28 April 2022

Available online

Version 1.0: 16 June 2022



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS etc. See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Raghav P, Choudhary P, Singh A, Sharma R, Sharma N. Extraction and characterization of oil from seeds of the medicinal plant *Withania coagulans* (Stocks) Dunal (Doda paneer). Plant Science Today (Early Access). <https://doi.org/10.14719/pst.1649>

## Abstract

*Withania coagulans* (Stocks) Dunal commonly known as Doda paneer/ Indian Rennet is a well-known medicinal plant but remains underutilized mainly due to scarce distribution in nature and extremely poor germination rate. In the present study oil was extracted from seeds of *Withania coagulans* which was subsequently subjected to FTIR and GC-MS analysis to identify phytochemical components of extracted oil. FTIR analysis revealed presence of diverse group of compounds including alcohols, alkane, alkene, aldehyde, ketone and halo compounds. GC-MS analysis depicted presence of 49 phytoconstituents in the oil extracted from seeds of *W. coagulans*. among which hexanoic acid, n-hexadecenoic acid, Vitamin E, gamma-Tocopherol, squalene, fucosterol, 2-Pyrrolidinone, 1-methyl, octadecanoic acid represent major phytoconstituents identified. Several compounds identified to be present in extracted oil have been reported to possess one or more pharmacological activity. Hence, the study suggests validation of plant oil to be utilized as ingredient of different pharmacological, cosmetic and other food products.

## Keywords

FTIR, GC-MS; oil, phytoconstituents, seeds, *Withania coagulans*

## Introduction


The genus *Withania* belonging to the family Solanaceae is a well-recognized genus comprising of several medicinal plants (1). Among 23 reported species of *Withania*, *Withania somnifera* and *Withania coagulans*, are among economically important (2). *W. coagulans* Dunal, is usually referred to as 'Indian cheese maker' or 'vegetable rennet' due to its milk coagulating properties (3, 4). *W. coagulans* has been reported to possess several medicinal properties such as anti-cancer, anti-diabetic, anti-oxidant, anti-fungal, anti-asthmatic, anti-bacterial, and anti-anxiety (5). Withanolides (such as steroidal alkaloids and lactones) being characteristic phytoconstituents of *W. coagulans* besides which flavonoids, tannin (6).

Extraction of essential oil from numerous plant species has been successfully accomplished for medicinal as well as traditional purposes. Medicinal and commercial significance of plant oil is attributed to the presence of aromatic compounds, secondary metabolites with biological activities. Published literature reports antifungal, anti-bacterial, anti-viral, anti-diabetic, anti-cancer, anti-inflammatory, anti-oxidant and repellent activities to be prominent biological activities of essential oil extracted from plant species (7-9). Hydro distillation, steam distillation and Soxhlet extraction

represents most commonly practiced methods for extraction of oil from different parts of plant species. Commercial application of plant essential oil includes their respective utilization in food industry, cosmetics, pharmaceuticals and health care etc. (10, 11). Scientific studies have been conducted to analyze and assess medicinal potential of oil extracted from medicinal plants, aromatic plants and other plant species. Still there are species for which only few studies have been conducted pertaining to extraction of oil along with its phytochemical characterization. *W. coagulans* is one such plant which is not yet explored to its medicinal potential and hence the plant is not commercially utilized in pharmaceutical and other industries whereas related species *Withania somnifera* is a component of several consumable healthcare and cosmetic products due to several validated *in vitro* and clinical Studies. With reference to research conducted on oil of *W. coagu-*

### Extraction of oil

For the extraction 10 g seeds of *W. coagulans* were finely powdered and mixed with 60 ml n-hexane and 60 ml acetone. After 90 cycles of Soxhlet apparatus and the extract was filtered through Whatman filter paper. After evaporation, lipid portion was extracted and collected following which lipid content was poured in separating funnel to which 12 ml diethyl ether was added for separation of lipids. The separating funnel was left undisturbed for 15 min. after which 2 independent layers were obtained in separating funnel. Upper layer represents ether layer and lower layer is water soluble layer. Once the two layers (ether and aqueous) got separated, ether layer was carefully removed and the ether was subsequently evaporated to obtain oil (Fig. 1).



**Fig. 1.** Methodology adopted for extraction of oil from seeds of *W. coagulans* (A) Seeds of *W. coagulans*, (B) Grounded seed powder of *W. coagulans* (C) Extraction of oil through Soxhlet distillation method (D) Collected Solvent with Oil (E) Evaporation of Solvent (F) Layers of Ether and Solvent (G) Evaporation of Ether (H) Extracted oil.

*lans*, a study reported (12) is among the prominent published literature pertaining to extraction and characterization of oil from the plant. Further studies are required to optimize the protocol for extraction of oil (along with its biochemical characterization). In the present study protocol for extraction of oil from seeds of *W. coagulans* was optimized along with its phytochemical characterization.

## Materials and Methods

### Plant material

Commercially available seeds of *W. coagulans* (Stocks Dunal) were utilized as study sample for the present work. The sample was authenticated by Dr. Manjul Dhiman, Head, Department of Botany KLDVA (PG) College Roorkee.

### Gas chromatography – Mass spectrophotometry (GC-MS)

The extracted oil was subjected to GC-MS analysis. Perkin Elmer Auto system was utilized as GC-MS analyzer. Helium gasses acted as carrier with a flowrate (constant) of 1.51 ml/min. An injection volume of 2  $\mu$ l was utilized. Mass spectrum was analyzed through Turbo mass software. Phytoconstituents were identified based upon molecular mass, structure, retention time and mass spectra compared to standard compounds from database NIST98, NIST database.

### Fourier Transmission Infrared Spectroscopy (FTIR)

FTIR Technique has been recognized as an effective bio-analytical tool for identification of different type of compounds or identification of functional group and chemical



**Table 1.** Phytoconstituents identified to be present in oil of seeds of *Withania coagulans* through GC-MS analysis.

Peak	R. Time	Area%	Name of the compound
1	8.809	3.98	Hexanoic acid
2	9.266	1.09	2-Pyrrolidinone, 1-Methyl-
3	10.351	0.23	Phorone
4	10.925	1.29	2,6-Dimethyl-6-nitro-2-hepten-4-one
5	11.816	0.66	2-Pentanol, 2,3-Dimethyl-
6	13.714	0.37	2-Methoxy-4-vinylphenol
7	15.097	0.21	Vanillin
8	16.176	0.12	Phenol, 3,5-bis(1,1-dimethylethyl)-
9	16.364	0.13	Benzene, (1-butylhexyl)-
10	16.735	0.13	Benzene, (1-Ethylloctyl)-
11	16.895	0.29	Dodecanoic Acid
12	17.675	0.27	Benzene, (1-propyloctyl)-
13	17.943	0.37	Benzene, (1-Ethylnonyl)-
14	18.384	0.42	Benzene, (1-Methyldecyl)-
15	18.670	0.18	Benzene, (1-butylloctyl)-
16	18.818	0.17	Benzene, (1-Propylnonyl)-
17	19.122	0.84	Tetradecanoic acid
18	19.513	0.23	Benzene, (1-methylundecyl)-
19	20.589	0.17	Benzene, (1-Methylododecyl)-
20	20.702	0.17	Hexadecanoic Acid, Methyl Ester
21	20.997	0.24	9-Hexadecenoic Acid
22	21.278	18.05	n-Hexadecanoic acid
23	22.346	0.80	9,12-Octadecadienoic acid (Z, Z)-, methyl ester
24	23.025	47.82	9,12-Octadecadienoic acid (Z, Z)-
25	23.193	1.50	Octadecanoic acid
26	24.625	0.33	Cyclohexane, 1,1'-Hexylidenebis-
27	25.629	0.94	Ethanol, 2-(9,12-octadecadienyloxy)-, (Z, Z)-
28	25.806	0.11	3-Heptadecene, (Z)-
29	27.000	0.87	Cyclopropane, 1,1-dichloro-2,2,3,3-tetramethyl-
30	27.184	1.16	(R)-(-)-14-Methyl-8-hexadecyn-1-OL
31	27.331	0.47	7-(3,4-Methylenedioxy)-tetrahydrobenzofuranone
32	27.413	0.35	9,12-Octadecadienoic acid (Z, Z)-2-hydroxy-1-(hydroxym)
33	28.019	0.19	9-Octadecenamamide
34	28.147	0.46	Squalene
35	28.798	0.32	Androst-5-en-3-ol, 4,4-dimethyl-, (3. beta.)-
36	29.171	0.28	delta. -Tocopherol
37	29.959	6.17	gamma. -Tocopherol
38	30.283	0.54	beta. -Tocopherol
39	30.585	1.91	Vitamin E
40	30.963	0.21	Lanostan-7-One
41	31.629	0.68	Stigmasta-5,24(28)-DIEN-3-OL, (3. beta.)-
42	31.704	0.31	Ergost-5-en-3-ol, (3. beta.)-
43	31.962	0.31	Stigmasta-5,22-DIEN-3-OL
44	32.300	0.34	Delta. 24-24-Methylcholester
45	32.662	1.43	gamma. -Sitosterol
46	32.850	0.68	Fucosterol
47	33.642	1.29	Lanost-8-en-3-ol, 24-methylene-, (3. beta.)-
48	34.489	0.69	9,19-Cyclolanostan-3-ol, 24-methylene-, (3. beta.)-
49	36.184	0.21	Octadecanoic Acid, 2,3-Bis [(1-Oxotetradecy

phytoconstituents among which sesquiterpenes and esters to represent major phytoconstituents while acids, alkanes and aldehydes comprised minor proportion of oil. In a specific study conducted antidiabetic potential of methanolic and aqueous extract of *W. coagulans* flower was studied (17). The study reported decrease in blood glucose of STZ induced diabetic rats compared to control rats after 28 days. As evident from the reported studies most of the medicinal and pharmacological properties reported of *W. coagulans* have utilized extracts prepared from either seeds / fruits /flower for their respective study. Findings of the present study reveal presence of several compounds in the oil extracted from seeds which have been reported to possess medicinal value. Hexanoic acid and octadecanoic acid possess anti-oxidant activity and anti-inflammatory activity (18, 19) Fucosterol, 2-Pyrrolidinone, 1-methyl possess anti-cancer activity (20, 21). Stigmasta-5, 22-dien-3-ol and 2-Methoxy-4-vinylphenol, vanillin, octadecanoic acid comprise phytoconstituents present in oil extracted from seeds of *W. coagulans* with reported antimicrobial activity (22, 20-30). Similar to the findings of present earlier studies conducted have also effectively utilized the technique of GC-MS to identify phytoconstituents with reported medicinal properties and biological activities (31, 32).


#### FTIR analysis

FTIR has been recognized as an effective analytical technique to identify different types of chemical bonds as well as functional groups present in organic compounds. The functional groups of different phytoconstituents were identified according to the peak values in region of infrared radiation. The analysis (Fig. 4) revealed extracted oil to possess organic compounds belonging to different classes including alcohol at 3473  $\text{cm}^{-1}$ , alkenes at 3008.9  $\text{cm}^{-1}$  and 2924.64  $\text{cm}^{-1}$ , aldehydes at 2854.12  $\text{cm}^{-1}$ , 2672.75  $\text{cm}^{-1}$  and 1377.49  $\text{cm}^{-1}$ , alkanes at 1465.08  $\text{cm}^{-1}$ , ketones at 1654.08  $\text{cm}^{-1}$ , cyclopentanone at 1744.13  $\text{cm}^{-1}$  and sulfone at 1164.17  $\text{cm}^{-1}$  (33-35). Presence of diverse nature of organic compounds indicates the extracted oil to be highly rich in containing different metabolites with characteristic properties and function. FTIR is a commonly utilized technique to identify different classes of organic compounds or functional groups present in plant species (36, 37). In an earlier study conducted, FTIR analysis of extract of fruits of *W. coagulans* revealed the presence of steroidal lactones (13). FTIR technique (along with NMR and UVvis) was utilized (4) for identification of withanolides in *W. coagulans*. Most of the reported literature pertaining to FTIR of *W. coagulans* have utilized plant extract and comparatively few studies have been conducted related to extraction as well as analytical analysis of oil of *W. coagulans*.

#### Conclusion

*Withania coagulans* (Stocks) Dunal is a well-known medicinal plant which is not yet fully explored to its potential for utilization in pharmaceutical, cosmetics and food industries. The commercial utilization is challenged by slow propagation rate (due to extremely





**Fig. 3.** Mass spectrum of major compounds identified to be present in oil extracted from seeds of *W. coagulans*.

poor germination rate) and present endangered status of the plant. Efforts are also required to accomplish commercial cultivation of the plant to produce sufficient raw material which can serve as industrial feedstock. Along with commercial cultivation, micropropagation can also prove extremely beneficial for mass propagation as well as conservation. Also, at present there exist a research gap to validate the utilization of plant extract, oil for medicinal purposes and as component of other cosmetic and food products. Studies are required for optimization of process and protocols to fully utilize medicinal potential of the plant.

### Acknowledgements

Authors are thankful to Division of Research and Innovation, Uttarakhand University, Dehradun for the support rendered in analytical analysis of studied oil sample.

### Authors contributions

NS and PR identified the research problem and designed the experimental study. PR and PC conducted the experimental study supervised by NS and RS. AS contributed in analysis of GC-MS and FTIR. Initial draft of manuscript was written by RS and AS. Manuscript was revised by PR and NS.




Fig. 4. FTIR analysis of oil extracted from seeds of *W. coagulans*.

### Compliance with ethical standards

**Conflict of interest:** Authors declare that there exists no conflict of interest.

**Ethical issues:** None.

### References

- Jain R, Kachhwaha S. Phytochemistry, pharmacology and biotechnology of *Withania somnifera* and *Withania coagulans*: A review. *J Med Plants Res.* 2015;6(41):5388-99. <https://doi.org/10.5897/JMPR12.704>.
- Panwar J, Tarafdar JC. Distribution of three endangered medicinal plant species and their colonization with arbuscular mycorrhizal fungi. *J Arid Environ.* 2006;65(3):37-350. <http://dx.doi.org/10.1016%2Fj.jaridenv.2005.07.008>.
- Ali N, Ahmad B, Bashir S. Calcium channel blocking activities of *Withania coagulans*. *Afr J Pharmacy Pharmacol.* 2009;3(9):439-42.
- Maher S, Choudhary MI, Saleem F, Rasheed S, Waheed I, Halim SA et al. Isolation of antidiabetic withanolides from *Withania coagulans* Dunal and their *in vitro* and *in silico* validation. *Biology.* 2020;9,197. <http://dx.doi.org/10.3390/biology9080197>
- Khan MI, Maqsood M, Saeed RA, Alam A, Sahar A, Kieliszek M et al. Phytochemistry, Food application and therapeutic potential of the medicinal plant (*Withania coagulans*): A Review. *Molecules.* 2021;26, 6881. <https://doi.org/10.3390/molecules26226881>
- Hemalatha S, Kumar R. *Withania coagulans* Dunal: A review. *Pharmacognosy Reviews.* 2008;2(4):351-58.
- Adorjan B, Buchbauer G. Biological properties of essential oils: an updated review. *Flavor and Fragrance J.* 2010;25(6):407-26. <https://doi.org/10.1002/ffj.2024>.
- Upadhyay BN, Gupta V. A clinical study on the effect of Rishyagandha (*Withania coagulans*) in the management of Prameha (Type II Diabetes Mellitus). *Ayu.* 2011;32(4):507-11. <https://dx.doi.org/10.4103%2F0974-8520.96124>.
- Silva TD, Kher MM. *Withania coagulans* (Stocks) Dunal: Biotechnological achievements and perspectives. *J Horticult Res.* 2015;23(1):5-12. <http://dx.doi.org/10.2478/johr-2015-0001>.
- Ferrentino G, Morozova K, Horn C, Scampicchio M. Extraction of essential oils from medicinal plants and their utilization as food antioxidants. *Curr Pharm Des.* 2020;26(5):519-41. <http://dx.doi.org/10.2174/1381612826666200121092018>. PMID: 31965940.
- Salehi M, Aghamaali MR, Sajedi RH. Purification and characterization of a milk-clotting aspartic protease from *Withania coagulans* fruit. *Int J Biol Macromol.* 2017;98,847-54. <https://doi.org/10.1016/j.ijbiomac.2017.02.034>.
- Ali A, Jameel M, Ali M. Analysis of fatty acid composition of *Withania coagulans* fruits by gas chromatography/mass spectrometry. *Res J Pharmacogn.* 2017;4(4):1-6.
- Peerzade N, Sayed N, Das N. Antimicrobial and phytochemical screening of methanolic fruit extract of *Withania coagulans* L. Dunal for evaluating the antidiabetic activity. *The Pharma Innov J.* 2018;7(1):197-204.
- Pramanick DD, Srivastava SK. Pharmacognostic evaluation of *Withania coagulans* Dunal (Solanaceae)-an important ethnomedicinal plant. *Bioscience Discovery.* 2015;6(1):06-13.
- Azhar MF, Naseer U. Antioxidant and Phytochemical composition of leaves, stem and root extracts of *Withania coagulans* and *Withania somnifera*. *J Medicinal and Spice Plants.* 2020;24(1):27-30.
- Bakhtawar S, Mughal T, Naeem I. Chemical composition of the essential oil of *Withania coagulans*. *Asian J Chem.* 2010;22(1):122-26.
- Kitukale MD, Chandewar AV. Antidiabetic potential of *Withania coagulans* Dunal flower in streptozotacin induced diabetic rats. *Int J Pharm Biol Arch.* 2017;8(4):58-62.
- Ramya B, Malarvili T. and Velavan S. GC-MS analysis of bioactive compounds in *Bryonopsis laciniosa* fruit extract. *Int J Pharm Sci Res.* 2015;6(8):3375-79. [http://dx.doi.org/10.13040/IJPSR.0975-8232.6\(8\).3375-79](http://dx.doi.org/10.13040/IJPSR.0975-8232.6(8).3375-79).
- Abdelhamid MS, Kondratenko EL, Lomteva NA. GC-MS analysis of phytocomponents in the ethanolic extract of *Nelumbo nucifera* seeds from Russia. *J Appl Pharma Sci.* 2015;5(04):115-18. <http://dx.doi.org/10.7324/JAPS.2015.50419>.
- Abdul QA, Choi RJ, Jung HA, Choi JS. Health benefit of fucosterol from marine algae: a review. *J Sci Food Agric.* 2016;96

- (6):1856-66. <https://doi.org/10.1002/jsfa.7489>.
21. Hosseinzadeh Z, Ramazani A. An overview on chemistry and biological importance of pyrrolidinone. *Current Organic Synthesis*. 2017;5(2). <http://dx.doi.org/10.2174/1570179414666170908165445>.
  22. Jebastella J, Reginald AM. Bioactive components of *Cynodon dactylon* using ethanol extract. *World J Pharm Sci*. 2015;3(12):2321-3310.
  23. Rubab M, Chelliah R, Saravanakumar K. Bioactive Potential of 2-Methoxy-4-vinylphenol and benzofuran from *Brassica oleracea* L. var. *capitata f. rubra* (Red Cabbage) on oxidative and microbiological stability of beef meat. *Foods*. 2020; 9(5): 568. <https://doi.org/10.3390/foods9050568>.
  24. Syeda FA, Habib-ur-Rahman. Gas chromatography-mass spectrometry (GC-MS) analysis of petroleum ether extract (oil) and bio-assays of crude extract of *Iris germanica*. *Inter J Genetics Molecular Biology*. 2011;3(7):95-100. Available From: <http://www.academicjournals.org/ijgmb>.
  25. Achi NK, Ohaeri OC. GC-MS Determination of bioactive constituents of the methanolic fractions of *Cnidioscolus aconitifolius*. *British Journal of Pharmaceutical Research*. 2015;5(3):163-72. 10.9734/BJPR/2015/13893. <https://doi.org/10.9734/BJPR/2015/13893>
  26. Chandrasekaran M, Senthilkumar A. Antibacterial and antifungal efficacy of fatty acid methyl esters from the leaves of *Sesuvium portulacastrum* L. *Eur Rev Med Pharmacol Sci*. 2011;15(7):775-80. PMID: 21780546
  27. Hema R, Kumaravel S, Alagusundaram K. GC-MS Determination of Bioactive components of *Murraya koenigii*. *Journal of American Science*. 2011;7(1):80-83. Available From: <http://www.americanscience.org>.
  28. Arya SS, Sharma MM, Das RK. Vanillin mediated green synthesis and application of gold nanoparticles for reversal of antimicrobial resistance in *Pseudomonas aeruginosa* clinical isolates. *Heliyon*. 2019;5(7):1-11.<https://doi.org/10.1016/j.heliyon.2019.e02021>.
  29. Rizvi S, Raza ST, Ahmed F. The role of vitamin E in human health and some diseases. *Sultan Qaboos Univ Med J*.2014;14(2):157-65. PMID: 24790736; PMCID: PMC3997530.
  30. Huang ZR, Lin YK, Fang JY. Biological and pharmacological activities of squalene and related compounds: potential uses in cosmetic dermatology. *Molecules*. 2009;14(1):540-54.<https://doi.org/10.3390/molecules14010540>.
  31. Sharma N, Rawat R, Lalremruati F, Singh A. Complete plant regeneration of *Valeriana wallichii* DC. on auxin enriched medium and phytochemical analysis. *Plant Sci Today*. 2020;7(4):542-50. <https://horizonpublishing.com/journals/index.php/PST/article/view/837> <https://doi.org/10.14719/pst.2020.7.4.837>
  32. Varnika, Sharma R, Singh A, Shalini S, Sharma N. Micropropagation and screening of phytochemicals present among *in vitro* raised and wild plants of *Rauvolfia serpentina*. *WJST*. 2020;17(11):1177-93. <https://doi.org/10.48048/wjst.2020.6492>
  33. Saleem A, Younas U, Ghullam M. Phytochemical screening by FTIR spectroscopy and antimicrobial activity of different solvent fractions from *Murraya koenigii* L. shoots. *Int Res J Pharm*. 2016;7(4):30-37. <http://dx.doi.org/10.7897/2230-8407.07435>.
  34. Janakiraman N, Satshish S, Jhonson M. UV-VIS and FTIR spectroscopic studies on *Peristrophe bicalyculata* (Retz) Nees. *Asian J Pharm Clin Res*. 2011;4(4):125-29. Available From: <https://www.researchgate.net/publication/306287011>.
  35. Subrahmanian H, Suriyamoorthy P. Fourier transform infra-red spectroscopy analysis of *Erythrina variegata* L. *J Pharm Sci and Res*. 2017;9(11):2062-67. Available From: [www.jpsr.pharmainfo.in](http://www.jpsr.pharmainfo.in).
  36. Chaudhari SY, Rajput D. Fourier transform infrared analysis of *Tamra bhasma* at different levels: A preliminary study. *An International Quarterly Journal of Research in Ayurveda*. 2015;36(1):0974-8520. <https://doi.org/10.4103/0974-8520.169013>.
  37. Thenmozhi M, Sangeetha M. Bioactive metabolites and FTIR analysis in *Ziziphus oenoplia* Mill. *Acta Scientific Nutritional Health*. 2021;5(6):2582-1423. Available From: <https://actascientific.com/>

§§§