



RESEARCH ARTICLE

# Characterisation of indigenous plants for herbal formulations preparation based on pharmacognostic and physiochemical data

Banjai Mochahary, Sudem Brahma, Mrinal Kalita & Arvind Kumar Goyal\*

Department of Biotechnology, Bodoland University, Kokrajhar-783 370, Assam, India

\*Email: [arvindgoyal210883@gmail.com](mailto:arvindgoyal210883@gmail.com)



OPEN ACCESS

## ARTICLE HISTORY

Received: 30 January 2022

Accepted: 02 April 2022

Available online

Version 1.0: 9 May 2022



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, etc. See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

**About this article:** This article is part of a special issue meant to publish selected articles presented during National Seminar on “Ethnobotany and Resource Management of the Bodos” held at Bodoland University (India) on 10 December, 2021.

## CITE THIS ARTICLE

Mochahary B, Brahma S, Kalita M, Goyal A K. Characterisation of indigenous plants for herbal formulations preparation based on pharmacognostic and physiochemical data. Plant Science Today 2022;9(sp2):8-17.  
<https://doi.org/10.14719/pst.1709>

## Abstract

The use of plant-based drugs has increased considerably in the modern world for their efficiency in managing diseases with lesser side effects than synthetic drugs. The current study was aimed to confirm the identity, quality and purity of some locally available potential medicinal plants such as *Alstonia scholaris* (bark), *Centella asiatica* (whole plant), *Drymaria cordata* (whole plant), *Hydrocotyle sibthorpioides* (whole plant), *Oroxylum indicum* (bark), *Senna hirsuta* (leaf), *Senna occidentalis* (leaf), *Solanum indicum* (root), *Stephania japonica* (tuber) in powdered form. The powdered plant parts were evaluated for preliminary phytochemicals, pharmacognostical studies, physical characteristics and heavy metals. Preliminary phytochemical screening of the different extracts inferred the existence of carbohydrates, phenolics, alkaloids and amino acids while triterpenoids were absent. Microscopical study of the powder revealed the diagnostic qualities such as stone cells, trichomes, stomata, calcium oxalate crystal, fibres, xylem vessel, pitted spiral vessels, etc. The colour, odour, flavour/taste and texture of the pulverized plant were overall acceptable. The physical characteristics which determine the flow rate of the powder with respect to Carr's index and Hausner's ratio were found to be good to passable except for *Hydrocotyle sibthorpioides* (whole plant), *Oroxylum indicum* (bark), which were not easily passable. The heavy metal test showed the absence of bismuth, cadmium and lead. Thus the present study may serve as a standard reference for the quality control analysis of the herbal drug either singly or in synergy.

## Keywords

Medicinal plants, organoleptic, heavy metal, *Alstonia*, *Centella*, *Drymaria*, *Hydrocotyle*, *Oroxylum*, *Senna*, *Solanum*, *Stephania*

## Introduction

Plant-based chemical moieties have attracted people since ancient times to treat various ailments because of innumerable healing properties (1, 2). This claim is supported by the World Health Organization. As per WHO, 88% of the population among the member countries are dependent on traditional and complementary medicine (3). The use of medicinal herb has been in practice in India since the days of yore due to the pool of secondary metabolites it houses (4). Ayurveda, a unique discipline for disease management using natural products in India, has been accepted globally for its exemplary output and uniqueness (5). Herb-herb combination is the basis of polyherbal formulation which provide collective effects in the treatment of

diseases. A single plant may have therapeutic ability but when compared to polyherbal formulations which have the collection of several plants, it is less potential (6, 7). In traditional Chinese medicine (TCM) herbal combinations are popularly used in order to decrease toxicity and increase efficacy of a formulation. The herb-herb combinations are simpler rather than complex without losing its basic therapeutic value (8, 9). Though Ayurveda medicine is gaining popularity, the main problem that poses is its standardisation and quality control during the manufacturing process (10). Standardisation is considered to be the first and important step of drug development which helps to evaluate the quality of polyherbal drugs and avoid variation from batch to batch (11, 12). This growing interest in naturopathy attracted us to explore the floral diversity in our vicinity for their potential therapeutic uses. A total of 9 plants were selected based on the uses by traditional healers in the locality for hepatoprotection for the current study.

## Materials and Methods

### Plant materials

Based on the traditional use by the Bodo tribe, plants such as *Alstonia scholaris* (bark), *Centella asiatica* (whole plant), *Drymaria cordata* (whole plant), *Hydrocotyle sibthorpioides* (whole plant), *Oroxylum indicum* (bark), *Senna hirsuta* (leaf), *Senna occidentalis* (leaf), *Solanum indicum* (root), *Stephania japonica* (tuber) were collected from various localities of Bodoland Territorial Region and were identified initially by Plant Taxonomist Dr. Sanjib Baruah, Department of Botany, Bodoland University, Kokrajhar, Assam, India. However, the voucher specimens was deposited at Botanical Survey of India, Central National Herbarium, Howrah for final authentication and identification (CNH/Tech.II/2021/42 date 26.11.2021). The plant materials were washed, shade dried, grounded using mechanical grinder, sieved through a sieve of mesh size of 600 µm and stored in airtight glass bottles for further analysis. A brief description of the nine plants selected for the current study are as follows:

*Alstonia scholaris* (L.) R.Br. is locally known as Sitaona in Bodo and blackboard or devil's tree in English (13). It belongs to the family Apocynaceae, an evergreen tree native to southern China, tropical Asia and Australasia (14). Though *A. scholaris* bark is reported to be toxic at a

higher dose (15), yet, it is found to be used in the treatment of myriad diseases (16). It is also reportedly used for medicinal purposes for the treatment of various diseases such as dysentery, diarrhoea, fever and other stomach aches (17), hepatoprotective (18), malaria, jaundice, gastrointestinal troubles and cancer (13).

*Centella asiatica* (L.) Urb. is also known as Indian Pennywort and Asiatic pennywort (English), and Manimuni geder (Bodo). It is an herbaceous plant native to Southeast Asian countries such as India, China, Indonesia, and Malaysia as well as a few African countries and belonging to family Apiaceae (19). *Centella asiatica* has been used as a vegetable and as a medicinal herb (19, 20). Some of the medicinal properties of *C. asiatica* are its use in the management of the liver disorder (21), and having anti-cancerous, anti-bacterial, anti-fungal, anti-inflammatory, neuroprotective, antioxidant, wound healing, anti-depressant activities (21), cognitive function (22), anti-diabetic (23), central nervous system, skin and gastrointestinal disorders (24).

*Drymaria cordata* (L.) Willd. ex Schult, locally named as Jabsri in Bodo and commonly known as chickweed in English, belongs to the family Caryophyllaceae, extensively distributed in Northeast India and originated from tropical America (25). It is mostly considered a weed of gardens but possesses many medicinal properties and find its application in the treatment of snake bites (26), many kinds of diarrhoea (26, 27), skin problems (28), constipation and throat pain (29); moreover, it has been reported to have some properties such as antibacterial property (30), anti-tussive activities and to manage acute cold attacks, coughs, sinusitis (31). Analgesic and antipyretic properties of *D. cordata* are well established (32).

*Hydrocotyle sibthorpioides* Lam. belongs to the family Umbelliferae, a perennial herb widely distributed in parts of Asia (33) and Africa (34). It is known as Lawn pennywort in English and Manimuni pisa in Bodo. The people of India and China traditionally use it for the treatment of various illnesses and disorders (34). *H. sibthorpioides* is also widely used in managing fever, rheumatism, coughing, liver problems, dysentery, sore throat, psoriasis, oedema, herpes zoster infection, hepatitis-B infection, soothing pain, dysmenorrhoea and carbuncles (34). It also possesses anti-inflammatory (35) and anti-chikungunya virus activities (36).

**Table 1.** List of plants with information on parts used, vernacular names and GPS coordinates of the collection site

Botanical Name	Parts Used	Vernacular names		GPS Coordinates	
		Bodo Name	Assamese Name	Latitude	Longitude
<i>Alstonia scholaris</i> (L.) R.Br.	Bark	Sitaona	Chatiyana, Sotiyana, Chatim	26.470425°N	90.296998°E
<i>Centella asiatica</i> (L.) Urb.	Whole plant	Manimuni geder	Bor manimuni	26.4706114°N	90.2969722°E
<i>Drymaria cordata</i> (L.) Willd. ex Schult	Whole plant	Japsri	Thunthuni, Lajjabori, Lai Jabori	26.534360°N	90.015519°E
<i>Hydrocotyle sibthorpioides</i> Lam.	Whole plant	Manimuni pisa	Soru manimuni	26.4697987°N	90.2968682°E
<i>Oroxylum indicum</i> (L.) Kurz	Bark	Karokandai	Dingdinga, Toguna, Bhatghila	26.463155°N	90.297376°E
<i>Senna hirsuta</i> (L.)	Leaf	Sumu bipang	N/A	26.797191°N	90.540234°E
<i>Senna occidentalis</i> (L.) Link	Leaf	Gangrim bipang	Hat-thenga, Jonjoni-goch, Kusum	26.47872°N	90.3053733°E
<i>Solanum indicum</i> (L.) Kurz	Root	Kuntainara	Deuri tita, Tit bhek, Bhot bengena	26.79414°N	90.5303609°E
<i>Stephania japonica</i> (Thunb.) Miers	Tuber	Dibaolu	Galdua, Tubuki Lota, Tubuka-lot	26.790827°N	90.5300655°E

*Oroxylum indicum* (L.) Kurz is a deciduous tree, categorised in the family Bignoniaceae. It is popular as Karokandai in Bodo, and commonly known as Midnight horror and the Indian trumpet tree. It is native to India and countries in the Himalayan foothills like Bhutan, parts of China, Indo-China and the Malesia region. It is used in the management of gastritis, hypertension, liver problem, fever, purgative, headache, epilepsy, muscular pain, general health problems (37, 38) and hepatoprotective (39).

*Senna hirsuta* (L.) H.S.Irwin & Barneby is a type of woody annual shrub belonging to the family Fabaceae and is locally named sumu bipang (Bodo). It is native to Africa and extensively used as a medicinal herb for various ailments because of its therapeutic properties (40). This species is reported to cure bone fractures of humans and live stocks (41). It is efficient in the treatment of rheumatism (42), hypertension (42), indigestion (43), skin disease (43), fevers (42), biliousness (42), ringworm (42), dropsy (42) and eczema (42), anti-diabetic (44), antimicrobial and hepatoprotective properties (45).

*Senna occidentalis* (L.) Link, belonging to the family Fabaceae is distributed all through the tropical and subtropical region of the world (46). It is known as gangrim bipang in Bodo and septic weed, coffee weed and stinking weed in English. *S. occidentalis* is a common additive or substitute to coffee (47). It is reported to possess scavenging properties, larvicidal and mosquitocidal activities (48-50), antioxidant and antimicrobial (51), anti-inflammatory, immunosuppressive, antianxiety, anti-depressant, analgesic, anti-diabetic and antipyretic activities (50, 52).

*Solanum indicum* (L.) Kurz, (Solanaceae) is commonly known as Indian nightshade and Kuntainara in Bodo. This species is commonly found throughout tropical and subtropical India. It is used in the treatment of stomach problems, anorexia, abdominal pain, worm infestation, fever, inflammation, pain, sleeping and urinary disorders (53), lung disorders (bronchitis and asthma), blood, and sexual disorders, heart problems and pruritis (54).

*Stephania japonica* (Thunb.) Miers, belongs to the family Menispermaceae and is popularly known as Snake vine and tape vine (English) and Dibaolu (Bodo). It is a slender twining shrub native to Asia and Australia and found abundantly in Northeast India. It is validated for use in the management of asthma, malaria, cancer, diarrhoea, dysentery, urinary disease, stomach ache, tuberculosis, hepatitis, itches, dyspepsia, fevers (55, 56), antihyperglycemic and antihyperlipidemic property (57), moderate antioxidant and potent analgesic activity (58).

### Preparation of extracts

The powdered plant materials were individually extracted in a Soxhlet apparatus using double distilled water (1:10 w/v ratio of the sample and solvent). The extraction was carried out for 6 hrs at boiling temperature and finally evaporated under pressure at 50 °C. The extract was stored at 4 °C for experimental analysis.

### Organoleptic parameters

In organoleptic evaluation, sensory impressions were used

to examine and characterize the qualities of crude, such as appearance (colour and texture), odour and taste (59). The results of this experiment are based on the response of sensory organs used.

### Microscopic study

For the Microscopical study, the powdered plant materials were mounted in water and safranin on clean slides and observed under the microscope for various characteristics in fragmented form using a binocular microscope (LaboMed vision 2000).

### Determination of physical characteristics of powder

#### Bulk density

Bulk density is interpreted as powder aerated and allowed to settle gently. It is the ratio between the given value of mass and untapped volume. The powdered sample (15 g) was transferred into a 100 ml cylinder. The initial volume was noted, and the ratio of the occupied weight of volume was computed by the following standard formula (60, 61).

$$\text{Bulk density} = W/V_0$$

Where, W = powder mass V<sub>0</sub> = untapped volume.

#### Tapped density

Tapped density was calculated by pouring 15 g of powdered sample into a measuring cylinder and tapped manually for approximately 500 times. The volume before and after tapping was noted (60, 61). The tapped density was calculated by the following standard formula (60).

$$\text{Tapped density} = W/V_f$$

Where, W = mass of the powder V<sub>f</sub> = tapped volume.

#### Carr's index

This is the tendency of the powder to be compressed depending upon the apparent bulk and tapped density (59, 60). The standard formula for calculating Carr's index was adopted as per standard methodology (46) and is expressed in percentage as.

$$\text{Carr's index} =$$

$$(\text{Tapped density} - \text{Bulk density}) / \text{Tapped density} \times 100$$

#### Hausner's ratio

Hausner's ratio determines the flow properties of the powder. It is the ratio of tapped density to the bulk density of powder (60, 61).

$$\text{Hausner's ratio} = \text{Tapped density} / \text{Bulk density}$$

#### Angle of repose

The angle of repose is defined as the maximum angle probable between the powder pile and the horizontal plane. The powder was allowed to pass through a funnel (mouth diameter 100 mm and stem diameter 10 mm) which is fixed on the tripod stand at the height of 2.9 cm. Graph paper was placed below the tripod on the table, and the height and radius of the pile were measured (59, 60). The formula (60) used for the calculation is

$$\text{Angle of repose} = \tan^{-1}h/r$$

Where, h=height of the pile, r = radius of the pile.

### Qualitative phytochemical investigation

The plant samples were phytochemically evaluated for the presence or absence of carbohydrates (Benedict's test, Molisch's test, and Fehling's test), alkaloids (Dragendorff's test), phenolics (Lead acetate test and FeCl<sub>3</sub> test), amino acids (Millon's test), and triterpenoids (Salkowski's test) according to the standard procedure (62, 63).

### Qualitative estimation of heavy metals

The standard procedure (64) was followed to qualitatively determine the occurrence of heavy metals like cadmium, lead and bismuth in different plant parts. It is evaluated to know the safe use of plants. .

## Results and Discussion

### Preliminary phytochemical screening

All the 9 plant extracts showed the presence of alkaloids and carbohydrates (Table 2). However, alkaloid was reported to be absent in the preliminary phytochemical screening of *S. occidentalis* (65). Phenols were present in *D. cordata*, *H. sibthorpioides*, *S. japonica*, *S. occidentalis*. While Amino acids were absent in *S. japonica*, *O. indicum*, *A. scholaris*, *D. cordata*, *C. asiatica* and *H. sibthorpioides*. Triterpenoids were not found in any of the plant extracts. Recently, the presence of secondary metabolites has also been reported in *C. asiatica* and *O. indicum* (66, 67) respectively. It is a known fact that the possible medicinal properties of plants are because of the presence of secondary metabolites such as alkaloids, phenols, amino acids, carbohydrates, triterpenoids and many more (68). Thus, screening of secondary metabolites is the vital step in developing a drug from the potential medicinal plants, though the phytochemical constituents vary qualitatively and quantitatively with seasons and different kind of species and even depending on storage (69).

**Table 2.** Preliminary phytochemical screening of various plant parts under study

Tests performed	<i>S. japonica</i>	<i>O. indicum</i>	<i>A. scholaris</i>	<i>D. cordata</i>	<i>S. occidentalis</i>	<i>S. indicum</i>	<i>C. asiatica</i>	<i>H. sibthorpioides</i>	<i>S. hirsuta</i>
Benedict's test (Carbohydrate)	Green	Brick red	Brick red	Brick red	Green	Green	Orange	Orange	Green
Molisch's test (Carbohydrate)	Violet ring	Violet ring	Violet ring	Violet ring	Violet ring	Violet ring	Violet ring	Violet ring	Violet ring
Fehling's test (Carbohydrate)	+	++	+++	+++	++	+	+++	+++	+
Dragendorff's test (Alkaloids)	++	++	++	++	++	+	++	++	++
Lead acetate test (Phenolic test)	White ppt	White ppt	White ppt	White ppt	White ppt	White ppt	White ppt	White ppt	White ppt
FeCl <sub>3</sub> test (Phenolic test)	Blue	Green	Green	Violet	Blue	Green	Green	Violet	Green
Millon's test (amino acid test)	ND	ND	ND	ND	Red ppt	Red ppt	ND	ND	Red ppt
Salkowski's test (Triterpenoids test)	ND	ND	ND	ND	ND	ND	ND	ND	ND

+ = Low, ++ = Moderate; +++ = High, Brick Red = High, Orange = Moderate, Green (Benedict's test) = Traceable; White ppt = presence of phenols, Red ppt = presence of Amino acids, Green (FeCl<sub>3</sub>) = Catechol, Violet and Blue = Phenols, ND = Not detected

### Organoleptic parameters

The powdered samples were physically evaluated using sensory organs for their colour, taste, texture and odour

and are presented in Table 3. The colour was in the shades of brown for all, except for *O. indicum* powder which was dandelion (shade of yellow). The intensity of taste ranged from strong to slight bitter in *S. japonica*, *A. scholaris*, *S. indicum*, *O. indicum* and *S. hirsuta*. Astringent taste was detected in *D. cordata* and *S. occidentalis* while *C. asiatica* and *H. sibthorpioides* had a pungent taste. The organoleptic analysis resulted in different odours such as characteristics, woody, grassy, barnyard and sweet in nature, as mentioned for various powdered plant parts (Table 3). In terms of texture, all the powders were of moderate to fine quality except for *D. cordata* (whole plant), which was soft spongy. Organoleptic data, when deviating from the standard observation, suggest adulteration or bad quality of powder (70). Organoleptic analysis revealed that the plant was free from foreign materials and had acceptable sensory characteristics.


### Powder microscopy

A microscopic study of powders revealed the occurrence of many features, as illustrated in Fig. 1, 2 and 3. The tuber of *S. japonica* showed the presence of acicular crystal, epidermal cell, fibre and xylem vessel. The powdered stem bark of *O. indicum* displayed acicular crystal, fibres, stone cell, tracheid and trichome. Calcium oxalate crystal, stone cell and fibre were observed in the stem bark powder of *A. scholaris*. The whole plant powder of *D. cordata* showed the presence of fibre, parenchyma cell, sclerenchyma cell and stone cell. The powdered form of leaf of *S. occidentalis* exhibited the presence of stomata, fibre, parenchyma cell and trichome. The roots powder of *S. indicum* revealed the presence of trichome, parenchyma cell, fibre and annular vessel. Fibre, trichome and stomata were observed in the whole plant powder of *C. asiatica*. The whole plant powder of *H. sibthorpioides* showed the presence of fibre, stomata, trichome and hair. *S. hirsuta* powdered leaf revealed the presence of trichome, pitted vessel and stomata. Similar to


organoleptic analysis, microscopical evaluation helps us to ascertain the presence or absence of any foreign matters or adulterants (5). The supporting tissue,

**Table 3.** Organoleptic properties of powdered plants part under study

Plant Species	Characteristics			
	Colour	Taste	Odour	Texture
<i>S. japonica</i>	Brown (584520)	Bitter	Soil type	Fine powder
<i>O. indicum</i>	Dandelion (fbc969)	Slight bitter	Pungent	Moderately fine powder
<i>A. scholaris</i>	Champaign brown (fad6a5)	Strong bitter	Woody	Moderately fine powder
<i>D. cordata</i>	Chamomile brown (dac395)	Astringent	Grassy	Soft spongy
<i>S. occidentalis</i>	Wood brown (Ci9a6b)	Slight astringent	Barnyard	Fine powder
<i>S. indicum</i>	Wheat brown (f5deb3)	Bitter	Woody	Moderately fine powder
<i>C. asiatica</i>	Bird seed brown (e2c28e)	Pungent	Sweet	Moderately fine powder
<i>H. sibthorpioides</i>	Khaki brown (c3b091)	Pungent	Sweet	Moderately fine powder
<i>S. hirsuta</i>	Brown bear (7f6244)	Slight bitter	Grassy	Fine powder



**Fig. 1.** Photomicrographs of microscopic evaluation (400x). 1a. *Stephania japonica* plant, 1b. *S. japonica* powder, 1c. Acicular crystal, 1d. Eplidermal cell, 1e. Fibre, 1f. Xylem vessel, 1g. *Oroxylum indicum* tree, 1h. *O. indicum* bark powder, 1i. Acicular vessel, 1j. Fibre, 1k. Stone cell, 1l. Tracheid, 1m. Trichome, 1n. *Alstonia scholaris* tree, 1o. *A. scholaris* bark powder, 1p. Fibre, 1q. Calcium oxalate crystal, 1r. Fibre, 1s. Stone cell, 1t. Tracheid



**Fig. 2.** Photomicrographs of microscopic evaluation (400x). 2a. *Drymaria cordata* plant, 2b. *D. cordata* whole plant powder, 2c. Fibre, 2d. Parenchyma cell, 2e. Sclerenchyma cell, 2f. Stone cell, 2g. *Senna occidentalis* plant, 2h. *S. occidentalis* leaf powder, 2i. Parenchyma cell, 2j. Fibre, 2k. Trichome, 2l. Stomata, 2m. *Solanum indicum* plant, 2n. *S. indicum* powder, 2o. Annular vessel, 2p. Fibre, 2q. Trichome, 2r. Xylem vessel.



**Fig. 3.** Photomicrographs of microscopic evaluation (400x). 3a. *Centella asiatica* plant, 3b. *C. asiatica* whole plant powder, 3c. Fibre, 3d. Stomata, 3e. Fibre, 3f. Trichome, 3g. *Hydrocotyle sibthorpioides* plant, 3h. *H. sibthorpioides* whole plant powder, 3i. Fibre, 3j. Stomata, 3k. Trichome, 3l. Hair, 3m. *Senna hirsuta* plant, 3n. *S. hirsuta* leaf powder, 3o. Trichome, 3p. Pitted vessel, 3q. Stomata.

i.e. fibre provides mechanical support (71), was seen in all the plant samples except *S. hirsuta*. The trichomes or hairs are mainly responsible for providing protection against pathogens in addition to regulating temperature by reducing evaporation (72) whereas, stomata is associated with gaseous exchange (5).

Reports are on (73) the microscopic evaluation of powdered stem bark of *O. indicum* and found the acicular crystal, fibres, stone cell, along with other characteristics. However, the presence of tracheid and trichome were reported for the first time. Reports are on (74) the presence of trichome and stomata in *C. asiatica* fresh leaves similar to our study in powdered form.

#### Flow ability

The flow property of the prepared powders was evaluated using simple procedures such as bulk density, tapped density, angle of repose, Hausner's ratio and Carr's index. The angle of repose of *A. scholaris* and *D. cordata* was found to

be in the range of  $34.30 \pm 0.50$  and  $34.62 \pm 0.31$  respectively, which signifies the good flow of granules; the value of Carr's index was observed in the range of  $23.68 \pm 0.070$  and  $19.05 \pm 0.025$  indicating poor and fair compressibility of granules. Hausner's ratio was found to be in the range of  $1.31 \pm 0.005$  and  $1.24 \pm 0.010$ , indicating passable and fair flow properties as shown in Table 4. The angle of repose and Hausner's ratio of *S. hirsuta* was found to be fair flow with  $38.09 \pm 0.50$  and  $1.29 \pm 0.010$  respectively and the compressibility was found to be slightly poor. The flow and compressibility properties of other plants were moderate. However, there was no data available for comparison in the case of the plants under study. Previously, there has been no test conducted to compare the flow properties and compressibility of these nine plant species. The insight of flow properties of powder enables pharmaceutical industries to operate smoothly by taking consideration during the processes such as filling, packaging, mixing and transportation (76).

**Table 4.** Physical characteristics of the powder of different plant parts

Plant Species	Angle of repose	Bulk density (g/mL)	Tapped density (g/mL)	Hausner's ratio	Carr's index
<i>Alstonia scholaris</i>	$34.30 \pm 0.50$	$0.39 \pm 0.035$	$0.52 \pm 0.010$	$1.31 \pm 0.005$	$23.68 \pm 0.070$
<i>Centella asiatica</i>	$37.35 \pm 0.15$	$0.25 \pm 0.015$	$0.29 \pm 0.020$	$1.18 \pm 0.005$	$15.00 \pm 0.050$
<i>Drymaria cordata</i>	$34.62 \pm 0.31$	$0.24 \pm 0.005$	$0.29 \pm 0.010$	$1.24 \pm 0.010$	$19.05 \pm 0.025$
<i>Hydrocotyle sibthorpioides</i>	$35.27 \pm 0.29$	$0.28 \pm 0.020$	$0.38 \pm 0.005$	$1.36 \pm 0.010$	$26.42 \pm 0.100$
<i>Oroxylum indicum</i>	$35.94 \pm 0.47$	$0.22 \pm 0.010$	$0.30 \pm 0.005$	$1.34 \pm 0.005$	$25.37 \pm 0.050$
<i>Senna hirsuta</i>	$38.09 \pm 0.50$	$0.15 \pm 0.005$	$0.19 \pm 0.010$	$1.29 \pm 0.010$	$22.33 \pm 0.050$
<i>Senna occidentalis</i>	$36.28 \pm 0.14$	$0.19 \pm 0.020$	$0.25 \pm 0.005$	$1.31 \pm 0.010$	$23.75 \pm 0.025$
<i>Solanum indicum</i>	$35.94 \pm 0.47$	$0.25 \pm 0.005$	$0.30 \pm 0.005$	$1.22 \pm 0.010$	$18.03 \pm 0.100$
<i>Stephania japonica</i>	$35.27 \pm 0.39$	$0.45 \pm 0.010$	$0.60 \pm 0.005$	$1.32 \pm 0.005$	$24.24 \pm 0.100$

Angle of repose= 30–40 Passable (79); Hausner's ratio= 1.12–1.18 (Good), 1.19–1.25 Fair, 1.26–1.34 (Passable) and 1.35–1.45 (Poor) (80)  
Carr's index= 12–16 (Good), 18–31 (Fair to Passable) (79).

## Heavy metal test

Heavy metal contaminations with herbal medicines beyond the safety level affect our health (77). Heavy metal is known for bioaccumulation in humans because of its non-degradable property (78). In the present study, there was no contamination of cadmium, bismuth and lead (Table 5) in any of the plants, which indicates that the prepared powders are safe to use in the formulation of drugs.

of capsule or tablet. Most of the plants included in this study are evaluated for the first time for the above parameters. Though, the evaluation procedures used here are simple and inexpensive but this knowledge of standardisation could help the society whenever traditional medicine is used as a source of cure. It rightly can be state that the plants here are clearly potential candidate for medicinal uses. Thus, the present study may serve as a standard

**Table 5.** Determination of heavy metals in plant parts

Sample solution	Procedure	Heavy metals					
		Cadmium (Cd)	Cadmium (Cd)	Bismuth (Bi)	Bismuth (Bi)	Lead (Pb)	Lead (Pb)
		Sample solution + NH <sub>4</sub> OH	Sample solution + Potassium Ferrocyanide	Sample solution + NH <sub>4</sub> OH	Sample solution + H <sub>2</sub> S	Sample solution + Dil HCl	Sample solution + KI
<i>Stephania japonica</i>	Observation	No white ppt.	No white ppt.	No white ppt.	No dark brown ppt.	No white ppt.	No yellow ppt.
	Inference	Absent	Absent	Absent	Absent	Absent	Absent
<i>Oroxylum indicum</i>	Observation	No white ppt.	No white ppt.	No white ppt.	No dark brown ppt.	No white ppt.	No yellow ppt.
	Inference	Absent	Absent	Absent	Absent	Absent	Absent
<i>Alstonia scholaris</i>	Observation	No white ppt.	No white ppt.	No white ppt.	No dark brown ppt.	No white ppt.	No yellow ppt.
	Inference	Absent	Absent	Absent	Absent	Absent	Absent
<i>Drymaria cordata</i>	Observation	No white ppt.	No white ppt.	No white ppt.	No dark brown ppt.	No white ppt.	No yellow ppt.
	Inference	Absent	Absent	Absent	Absent	Absent	Absent
<i>Senna occidentalis</i>	Observation	No white ppt.	No white ppt.	No white ppt.	No dark brown ppt.	No white ppt.	No yellow ppt.
	Inference	Absent	Absent	Absent	Absent	Absent	Absent
<i>Solanum indicum</i>	Observation	No white ppt.	No white ppt.	No white ppt.	No dark brown ppt.	No white ppt.	No yellow ppt.
	Inference	Absent	Absent	Absent	Absent	Absent	Absent
<i>Centella asiatica</i>	Observation	No white ppt.	No white ppt.	No white ppt.	No dark brown ppt.	No white ppt.	No yellow ppt.
	Inference	Absent	Absent	Absent	Absent	Absent	Absent
<i>Hydrocotyle sibthorpioides</i>	Observation	No white ppt.	No white ppt.	No white ppt.	No dark brown ppt.	No white ppt.	No yellow ppt.
	Inference	Absent	Absent	Absent	Absent	Absent	Absent
<i>Senna hirsuta</i>	Observation	No white ppt.	No white ppt.	No white ppt.	No dark brown ppt.	No white ppt.	No yellow ppt.
	Inference	Absent	Absent	Absent	Absent	Absent	Absent

## Conclusion

Mother Nature has blessed us with medicinal plants that are known fact, and now responsibilities are vested upon researcher to promote healthy life in the society by validation of the traditional knowledge. Pharmacognosy mainly unveils the therapeutic potential of plants in the management of diseases. Standardisation of herbal drugs draws attention of researchers due to its vast sources with high variables. These reasons motivated us to characterize nine indigenous medicinal plants on the basis of phytochemicals (qualitative), organoleptic, microscopy, flow characteristics and heavy metals parameters. The presence of various phytochemical constituents in the different plants parts under study reveals that they have therapeutic potentials. The authentication of the plant species can be achieved by the organoleptic and micro-morphological features, whereas the flow properties is essential in pharmaceutical industry to obtain finished products in the form

reference for the quality control analysis of the herbal drug either singly or in synergy.

## Acknowledgements

All the authors are thankful to the Higher Education Department, Government of Assam for financial assistance vide letter no. AHE.493/2017/110 under the scheme "Tejasvi Navadhitamastu Edu Infra Fund: Astadash Mutukar Unnoyonee Mala" and Department of Biotechnology, Ministry of Science and Technology, Government of India for the project grant vide letter No. BT/IN/Indo-US/Foldscope/39/2015 under the scheme "Proposal for use of Foldscope as a research tool". Banjai Mochahary is grateful to Ministry of Tribal Affairs, Government of India and University Grant Commission, Government of India, for providing the National Fellowship for Higher Education of ST.

## Authors contributions

AKG conceptualized and designed the study. BM, SB and MK carried out the research work and acquired the data. All the authors analyzed the data and wrote the first draft of the manuscript. Finally all the authors edited the manuscript and approved the final version for submission.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** None

## References

- Xiong F, Guan YS. Cautiously using natural medicine to treat liver problems. *World J Gastroenterol.* 2017;23(19):3388-95. <https://doi.org/10.3748/wjg.v23.i19.3388>
- Goyal AK, Middha SK, Usha T, Sen A. Analysis of toxic, anti-diabetic and antioxidant potential of *Bambusa balcooa* Roxb. leaf extracts in alloxan-induced diabetic rats. *3 Biotech.* 2017;7:120. <https://doi.org/10.1007/s13205-017-0776-8>
- World Health Organization. WHO global report on traditional and complementary medicine. World Health Organization; 2019.p. 44-45.
- Usha T, Pradhan S, Goyal AK, Dhivya S, Kumar HP, Singh MK et al. Molecular simulation-based combinatorial modeling and antioxidant activities of Zingiberaceae family rhizomes. *Pharmacogn Mag.* 2017;13(3):S715. [https://doi.org/10.4103/pm.pm\\_82\\_17](https://doi.org/10.4103/pm.pm_82_17)
- Mukhi S, Bose A, Panda P, Rao MM. Pharmacognostic, physico-chemical and chromatographic characterization of Samasharkara churna. *J Ayurveda Integr Med.* 2016;7(2):88-99. <https://doi.org/10.1016/j.jaim.2015.11.004>
- Xie, Z. *Practical Traditional Chinese Medicine*; Foreign Languages Press: Beijing, China, 2000. p. 260.
- Che CT, Wang ZJ, Chow MSS, Lam CWK. Herb-herb combination for therapeutic enhancement and advancement: theory, practice and future perspectives. *Molecules* 2013;18:5125-41. <https://doi.org/10.3390/molecules18055125>
- Wang S, Hu Y, Tan W, Wu X, Chen R, Cao J, Chen M, Wang Y. Compatibility art of traditional Chinese medicine: from the perspective of herb pairs. *J Ethnopharmacol.* 2012; 143(2): 412-23. <https://doi.org/10.1016/j.jep.2012.07.033>
- Shi L, Tang X, Dang X, Wang Q, Wang X, He P, Wang Q, Liu L, Liu X, Zhang Y. Investigating herb-herb interactions: The potential attenuated toxicity mechanism of the combined use of *Glycyrrhizae radix et rhizoma* (Gancao) and *Sophorae flavescens radix* (Kushen). *J Ethnopharmacol.* 2015;165:243-50. <https://doi.org/10.1016/j.jep.2015.02.022>
- Sahoo N, Manchikanti P, Dey S. Herbal drugs: standards and regulation. *Fitoterapia.* 2010;81(6):462-71. <https://doi.org/10.1016/j.fitote.2010.02.001>
- Pattanayak P, Kumar Hardel D, Mohapatra P. Standardization of Vaisvanara churna: A poly-herbal formulation. *Phcog J.* 2010; 2:50-59. <https://doi.org/10.4103/0974-8490.62957>
- Kumari R, Dubey GP. Standardization of Ayurvedic Drug-*Nyctanthes arbor-tristis*, *Hippophae salicifolia*, *Ocimum tenuiflorum* and *Reinwardtia indica* and Combined Herb-Herb. *Nepal J Biotechnol.* 2020;8(1):36-53. <https://doi.org/10.3126/njb.v8i1.30208>
- Khyade MS, Kasote DM, Vaikos NP. *Alstonia scholaris* (L.) R. Br. and *Alstonia macrophylla* Wall. ex G. Don: A comparative review on traditional uses, phytochemistry and pharmacology. *J Ethnopharmacol.* 2014;153(1):1-8. <https://doi.org/10.1016/j.jep.2014.01.025>
- Pandey K, Shevkar C, Bairwa K, Kate AS. Pharmaceutical perspective on bioactives from *Alstonia scholaris*: ethnomedicinal knowledge, phytochemistry, clinical status, patent space and future directions. *Phytochem Rev.* 2020;19(1):191-33. <https://doi.org/10.1007/s11101-020-09662-z>
- Bello I, Bakkouri AS, Tabana YM, Al-Hindi B, Al-Mansoub MA, Mahmud R, Asmawi M. Acute and sub-acute toxicity evaluation of the methanolic extract of *Alstonia scholaris* stem bark. *Med Sci (Basel).* 2016;4(1):4. <https://doi.org/10.3390/medsci4010004>
- Chandra Jagetia G, Shrinath Baliga M. Modulation of antineoplastic activity of cyclophosphamide by *Alstonia scholaris* in the Ehrlich ascites carcinoma-bearing mice. *Journal of Experimental Therapeutics and Oncology.* 2003;3(5):272-82. <https://doi.org/10.1111/j.1533-869X.2003.01098.x>
- Pandey K, Shevkar C, Bairwa K, Kate AS. Pharmaceutical perspective on bioactives from *Alstonia scholaris*: ethnomedicinal knowledge, phytochemistry, clinical status, patent space and future directions. *Phytochem Rev.* 2020;19:191-33. <https://doi.org/10.1007/s11101-020-09662-z>
- Kumar A, Khan MA, Saxena A, Singh RB, Zaman K, Husain A. Hepatoprotective activity of methanolic extract of stem bark of *Alstonia Scholaris* (L.) R. Br. *Am J Pharmtech Res.* 2012;2(2):545-55.
- Orhan IE. *Centella asiatica* (L.) Urban: From traditional medicine to modern medicine with neuroprotective potential. Evidence-based complementary and alternative medicine. 2012;2012: 946259. <https://doi.org/10.1155/2012/946259>
- Joshi K, Chaturvedi P. Therapeutic efficiency of *Centella asiatica* (L.) Urb. An underutilized green leafy vegetable: an overview. *International Journal of Pharma and Bio Sciences.* 2013;4(1):135-49.
- Prakash V, Jaiswal NI, Srivastava MR. A review on medicinal properties of *Centella asiatica*. *Asian J Pharm Clin Res.* 2017;10(10):69-74. <https://doi.org/10.22159/ajpcr.2017.v10i10.20760>
- Gray NE, Magana AA, Lak P, Wright KM, Quinn J, Stevens JF et al. *Centella asiatica*: Phytochemistry and mechanisms of neuroprotection and cognitive enhancement. *Phytochem Rev.* 2018;17(1):161-94. <https://doi.org/10.1007/s11101-017-9528-y>
- Oyenihi AB, Langa SO, Mukaratirwa S, Masola B. Effects of *Centella asiatica* on skeletal muscle structure and key enzymes of glucose and glycogen metabolism in type 2 diabetic rats. *Bio-med Pharmacother.* 2019;112:108715. <https://doi.org/10.1016/j.biopha.2019.108715>
- Torbati FA, Ramezani M, Dehghan R, Amiri MS, Moghadam AT, Shakour N et al. Ethnobotany, Phytochemistry and Pharmacological Features of *Centella asiatica*: A Comprehensive Review. *Adv Exp Med Biol.* 2021;1308:451-99. [https://doi.org/10.1007/978-3-030-64872-5\\_25](https://doi.org/10.1007/978-3-030-64872-5_25)
- Dequan L, Gilbert MG. *Drymaria Willd.* ex J. A. Schultes. In: Beijing, China, Wu Z, Raven PH, editors. *Flora of China* (Caryophyllaceae through Lardizabalaceae), Science Press; and St Louis, Missouri, USA: Missouri Botanical Garden Press. 2001, p. 6.
- Neogi B, Prasad MN, Rao RR. Ethnobotany of some weeds of Khasi and Garo hills, Meghalaya, Northeastern India. *Econ Bot.* 1989;43(4):471-79. <https://doi.org/10.1007/BF02935921>
- Nono NR, Nzowa KL, Barboni L, Tapondjou AL. *Drymaria cordata* (Linn.) Willd. (Caryophyllaceae): Ethnobotany, Pharmacology and Phytochemistry. *Adv Biol Chem.* 2014;4:160-67. <https://doi.org/10.4236/abc.2014.42020>



28. Rao RR. Ethnobotany of Meghalaya: medicinal plants used by Khasi and Garo tribes. *Econ Bot.* 1981;35(1):4-9. <https://doi.org/10.1007/BF02859208>
29. Focho DA, Ndam WT, Fonge BA. Medicinal plants of Aguambu-Bamumbu in the Lebiale highlands, southwest province of Cameroon. *Afr J Pharmacy Pharmacol.* 2009;3(1):1-13.
30. Mukherjee PK, Bhattacharya S, Saha K, Giri SN, Pal M, Saha BP. Antibacterial evaluation of *Centella asiatica* ordata Willd. (Fam. Caryophyllaceae) extract. *Phyto Res.* 1996;11:249-50. [https://doi.org/10.1002/\(SICI\)1099-1573\(199705\)11:3<249::AID-PTR69>3.0.CO;2-W](https://doi.org/10.1002/(SICI)1099-1573(199705)11:3<249::AID-PTR69>3.0.CO;2-W)
31. Mukherjee PK, Saha K, Bhattacharya S, Giri SN, Pal M, Saha BP. Studies on antitussive activity of *Drymaria cordata* Willd. (Caryophyllaceae). *J Ethnopharmacol.* 1997;56(1):77-80. [https://doi.org/10.1016/S0378-8741\(97\)01512-2](https://doi.org/10.1016/S0378-8741(97)01512-2)
32. Akindele AJ, Ibe IF, Adeyemi OO. Analgesic and antipyretic activities of *Drymaria cordata* (Linn.) Willd. (Caryophyllaceae) extract. *Afr J Tradit Complement Altern Med.* 2012;9(1):25-35. <https://doi.org/10.4314/ajtcam.v9i1.4>
33. Huang Q, Zhang S, Huang R, Wei L, Chen Y, Lv S et al. Isolation and identification of an anti-hepatitis B virus compound from *Hydrocotyle sibthorpioides* Lam. *J Ethnopharmacol.* 2013;150(2):568-75. <https://doi.org/10.1016/j.jep.2013.09.009>
34. Hazarika I, Mukundan GK, Sundari PS, Laloo D. Journey of *Hydrocotyle sibthorpioides* Lam.: From traditional utilization to modern therapeutics—A review. *Phytother Res.* 2021;35(4):1847-71. <https://doi.org/10.1002/ptr.6924>
35. Nazmi R, Dillasamola D, Badriyya E, Aldi Y. Anti-inflammatory Activity of Pegagan Embun (*Hydrocotyle sibthorpioides* Lam.) Plant Extract on Topical Application. *Pharmacogn J.* 2021; 13(2):411-17. <https://doi.org/10.5530/pj.2021.13.52>
36. Chan SM, Khoo KS, Sekaran SD, Sit NW. Mode-dependent antiviral activity of medicinal plant extracts against the mosquito-borne Chikungunya virus. *Plants.* 2021;10(8):1658. <https://doi.org/10.3390/plants10081658>
37. Mao AA, Hynniewta TM, Sanjappa M. Plant wealth of Northeast India with reference to Ethnobotany. *Indian J Tradit Knowl.* 2009;8(1):96-03.
38. Samatha TA, Srinivas P, Shyamsundarachary RU, Rajinikanth M, Rama Swamy N. Phytochemical analysis of seeds, stem bark and root of an endangered medicinal forest tree *Oroxylum indicum* (L) Kurz. *Int J Pharma Bio Sci.* 2012;3(3):1063-75.
39. NandaTripathy B, Panda SK, Sahoo S, Mishra SK, Nayak L. Phytochemical examination and hepatoprotective impact of stem bark of *Oroxylum indicum* (L.) Vent. on carbon tetrachloride induced hepatotoxicity in rat. *Technol Innov Pharm Res.* 2021;12(3):113-20. <https://doi.org/10.9734/bpi/tipr/v3/4069D>
40. Essiett UA, Bassey IE. Comparative phytochemical screening and nutritional potentials of the flowers (petals) of *Senna alata* (L.) Roxb., *Senna hirsuta* (L.) Irwin and barneby, and *Senna obtusifolia* (L.) Irwin and Barneby (Fabaceae). *Journal of Applied Pharmaceutical Science.* 2013;3(8):97-101.
41. Rahman MA, Ahmed NU. Phytochemical and biological activities of ethanolic extract of *C. hirsuta* leaves. *Bangladesh J Sci Ind Res.* 2013;48(1):43-50. <https://doi.org/10.3329/bjsir.v48i1.15413>
42. Essien EE, Thomas PS, Ascrizzi R, Setzer WN, Flamini G. *Senna occidentalis* (L.) Link and *Senna hirsuta* (L.) HS Irwin and Barneby: constituents of fruit essential oils and antimicrobial activity. *Nat Prod Res.* 2019;33(11):1637-40. <https://doi.org/10.1080/14786419.2018.1425842>
43. Singh TT, Devi AR, Sharma HM. Ethnopharmacological survey of medicinal plants in Andro Village in Manipur (India). *Res J Life Sci.* 2018;4:606-23. <https://doi.org/10.26479/2018.0405.49>
44. Madankumar D, Pari S. Evaluation of *in-vitro* antidiabetic and antioxidant activity using methanolic extract of medicinal plants (*Senna alata* and *Senna hirsuta*). *Solid State Technol.* 2020;4962-73.
45. Akharaiyi FC, Boboye B, Akpambang VO. Antibacterial and biochemical effects of ethanol leaf extract of *Senna hirsuta* Mill using animal model-mice. *J Microbiol Biotechnol Food Sci.* 2021;4(4):292-96. <https://doi.org/10.15414/jmbfs.2015.4.4.292-296>
46. Barbosa-Ferreira M, Dagli ML, Maiorka PC, Górniak SL. Sub-acute intoxication by *Senna occidentalis* seeds in rats. *Food Chem Toxicol.* 2005;43(4):497-503. <https://doi.org/10.1016/j.fct.2004.11.017>
47. Olapade AA, Ajayi OA. Effect of roasting regime on phytochemical properties of *Senna occidentalis* seeds. *Int. J Food Stud.* 2016;5(2):203-11. <https://doi.org/10.7455/ijfs/5.2.2016.a8>
48. Nuhu A, Aliyu R. Effects of *Cassia occidentalis* aqueous leaf extract on biochemical markers of tissue damage in rats. *Trop J Pharm Res.* 2008;7(4):1137-42. <https://doi.org/10.4314/tjpr.v7i4.14699>
49. Daskum AM, Godly C, Qadeer MA, Ling LY. Effect of *Senna occidentalis* (Fabaceae) leaves extract on the formation of  $\beta$ -hematin and evaluation of *in vitro* antimalarial activity. *Int J Herb Med.* 2019;7(3):46-51. <https://doi.org/10.3390/plants10081658>
50. Issa TO, Mohamed Ahmed AI, Mohamed YS, Yagi S, Makhawi AM, Khider TO. Physiochemical, insecticidal and antidiabetic activities of *Senna occidentalis* Linn. root. *Biochem Res Int.* 2020;2020:8810744. <https://doi.org/10.1155/2020/8810744>
51. Tsado NA, Lawal B, Kontagora GN, Muhammad BM, Yahaya MA, Gboke JA et al. Antioxidants and antimicrobial-activities of methanol leaf extract of *Senna occidentalis*. *J Adv Med Pharm Sci.* 2016;8(2)1-7. <https://doi.org/10.9734/JAMPS/2016/25655>
52. Daskum AM, Godly C, Qadeer MA, Ling LY. Effect of *Senna occidentalis* (Fabaceae) leaves extract on the formation of  $\beta$ -hematin and evaluation of *in vitro* antimalarial activity. *Int J Herb Med.* 2019;7(3):46-51.
53. Deb PK, Ghosh R, Chakraverty R, Debnath R, Das L, Bhakta T. Phytochemical and pharmacological evaluation of fruits of *Solanum indicum* Linn. *Int J Pharm Sci Rev Res.* 2014;25(2):28-32.
54. Saxena HO, Pawar G. Total phenolic and caffeic acid contents in roots of *Solanum indicum* L. from different agroclimatic regions of Madhya Pradesh State of India. *Indian J Pharm Educ Res.* 2019;53(2S):s164-69. <https://doi.org/10.5530/ijper.53.2s.62>
55. Ghani. *Medicinal Plants of Bangladesh: Chemical Constituents and Uses*, 2nd Edition, Asiatic Society of Bangladesh, Dhaka, 2003, p. 1-16.
56. Shankar R, Neyaz S, Anku G, Rungsung LW. Exploration, conservation and cultivation of *Stephania japonica* (Thunb.) Miers. *J Drug Res Ayurvedic Sci.* 2020;5(3):143-44. <https://doi.org/10.5005/jdras-10059-0098>
57. Sultana MC, Alam MB, Asadujjaman M, Akter MA. Antihyperglycemic and antihyperlipidemic effects of *Stephania japonica* (Thunb.) Miers. Tenril in alloxan induced diabetic mice. *Int J Pharm Sci Rev Res.* 2012;3(8):2726-32.
58. Uddin SN, Amin MN, Shahid-Ud-Daula AF, Hossain H, Haque MM, Rahman MS et al. Phytochemical screening and study of antioxidant and analgesic potentials of ethanolic extract of *Stephania japonica* Linn. *Journal of Medicinal Plants Research.* 2014;8(37):1127-33. <https://doi.org/10.5897/JMPR2014.5443>
59. Tripathi R, Verma S, Easwari TS, Shah H. Standardization of some herbal anti-diabetic drugs in polyherbal formulation and their comparative study. *Int J Pharm Sci Res.* 2013;4(8):3256-65. [http://dx.doi.org/10.13040/IJPSR.0975-8232.4\(8\).3256-65](http://dx.doi.org/10.13040/IJPSR.0975-8232.4(8).3256-65)
60. Madhavi N, Kumar D, Naman S, Singh M, Singh PA, Bajwa N,

- Baldi A. Formulation and evaluation of novel herbal formulations incorporated with Amla extract for improved stability. *Journal of Drug Delivery and Therapeutics*. 2019;9(4):212-21.
61. Anilkumar V, Kalyani R, Padmasri, B, Prasanth, D. In-house preparation, development and evaluation of herbal cosmetics face pack using various natural powders. *J Drug Deliv Sci Technol*. 2020;10(5):159-64. <https://doi.org/10.22270/jddt.v10i5.4314>
  62. Trease GE and Evans WC, 1989. *Pharmacognosy*. 11th Edn. Brailliar Tiridel Can. Machmillian publishers.
  63. Goyal AK, Middha SK, Sen A. Evaluation of the DPPH radical scavenging activity, total phenols and antioxidant activities in Indian wild *Bambusa vulgaris* "Vittata" methanolic leaf extract. *Journal of Natural Pharmaceuticals*. 2010;1(1):40-45. <https://doi.org/10.4103/2229-5119.73586>
  64. Ranjith D. Fluorescence analysis and extractive values of herbal formulations used for wound healing activity in animals. *J Med Plants Stud*. 2018;6(2):189-92.
  65. Isah RT, Mohammed MO, Muhammad AT, Sahabi SM, Umar ZU, Mahmud RI et al. Effects of aqueous leaf extracts of *Senna occidentalis* on rat kidney. *Afr J Biomed Res*. 2018;21(2):225-30.
  66. Baby S, Antony AV, John B, Jose D. Phytochemical screening and *in-vitro* anti-inflammatory activity of ethanolic extract of *Centella asiatica*. *Int J Curr Med Pharm Res*. 2020;2(2):181-83. <https://doi.org/10.37022/WJCMR.2020.2218>
  67. Rai B, Bhutia S, Pal P, Kakoti BB. Phytochemical analysis and antibacterial evaluation against selected gram strains by *Oroxylum indicum* (L.) Kurz stem bark extract, a folklore medicine of Sikkim Himalaya. *J Pharmacogn Phytochem*. 2020;9:11-16.
  68. Windyaswari AS, Purba JP, Nurrahmah SS, Ayu IP, Imran Z, Amin AA, Kurniawan F, Pratiwi NT, Iswantari A. Phytochemical profile of sea grass extract (*Enhalus acoroides*): A new marine source from Ekas Bay, East Lombok. *IOP Conf Ser Earth Environ Sci*. 2019;278(1), 012081. IOP Publishing. <https://doi.org/10.1088/1755-1315/278/1/012081>
  69. Tamasi A, Shoge MO, Adegboyega T, Chukwuma E. Phytochemical analysis and *in-vitro* antimicrobial screening of the leaf extract of *Senna occidentalis* (Fabaceae). *Asian J Nat Prod Biochem*. 2021;19(2):57-64. <https://doi.org/10.13057/biofar/f190203>
  70. Kewatkar SM, Paithankar VV, Deshpande SS, Jain SP, Nagore DH, Bhusari DV et al. Comparative pharmacognostical and phytochemical study of *Cassia auriculata* and *Cassia fistula*. *J Nat Remedies*. 2021;21(1):77-85. <https://doi.org/10.18311/jnr/2021/25626>
  71. Aslam I, Iqbal J, Peerzada S, Afridi MS, Ishtiaq S. Microscopic investigations and pharmacognostic techniques for the standardization of *Caralluma edulis* (Edgew.) Benth. ex Hook. f. *Microsc Res Tech*. 2019;82(11):1891-02. <https://doi.org/10.1002/jemt.23357>
  72. Wang X, Shen C, Meng P, Tan G, Lv L. Analysis and review of trichomes in plants. *BMC Plant Biol*. 2021;21(70):1-11. <https://doi.org/10.1186/s12870-021-02840-x>
  73. Bisht A, Zaman K, Singh M, Gupta R, Singh V. Pharmacognostical studies on *Oroxylum indicum* (Linn.) Vent. stem bark. *Indian J Nat Prod Resour*. 2011;2(4):472-78.
  74. Sudhakaran MV. Botanical Pharmacognosy of *Centella asiatica* (Linn.) Urban. *Pharmacognosy Journal*. 2017;9(4):546-58. <https://doi.org/10.5530/pj.2017.4.88>
  75. Dutta S, Das M, Banerji A, Rao MM. Pharmacognostical evaluation of *Alstonia scholaris* (L.) R. Br. (Saptaparṇa), roots. *Journal of Pharmacognosy and Phytochemistry*. 2019;8(2):2096-101.
  76. Kaur P, Kataria SK, Singh B, Arora S. Pharmacognostic Investigation of *Punica granatum* L. Peel. *Int J Pharm Drug Anal*. 2018;6(2):116-21.
  77. Luo L, Wang B, Jiang J, Fitzgerald M, Huang Q, Yu Z, et al. Heavy metal contaminations in herbal medicines: Determination, comprehensive risk assessments and solutions. *Frontiers Pharmacol*. 2020;11: 595335. <https://doi.org/10.3389/fphar.2020.595335>
  78. Alias N, Rosli SA, Sazalli NA, Hamid HA, Arivalakan S, Umar SN, et al. Metal oxide for heavy metal detection and removal. *InMetal Oxide Powder Technologies*. 2020;299-32. <https://doi.org/10.1016/B978-0-12-817505-7.00015-4>
  79. Sharma AK, Kaushik K. Formulation and evaluation of herbal antidiabetic tablet. *J drug Deliv Ther*. 2011;1(1):65-67. <https://doi.org/10.22270/jddt.v1i1.24>
  80. Attia UM, Fones A, Trepleton R, Hamilton H, Davies S, Wimpenny D. HIPing of Pd-doped titanium components: A study of mechanical and corrosion properties. In: *Proceedings of the 11th International Conference on Hot Isostatic Pressing, Stockholm, Sweden*. 2014. p. 9-13.

§§§