



RESEARCH ARTICLE

Impacts of habitat variability on the phenotypic traits of *Aconitum violaceum* Jacq. ex Stapf. at different altitudes and environmental conditions in the Ladakh Himalaya, India

Abdul Hadi^{1, 2*}, Seema Singh¹, Irshad A Nawchoo², Shah Rafiq¹ & Sajjad Ali¹

¹Plant Tissue Culture Research Laboratory, Department of Botany, University of Kashmir, Srinagar 190 006, J & K, India

²Plant Reproductive Biology, Genetic Diversity and Phytochemistry Research Laboratory, Department of Botany, University of Kashmir, Srinagar 190 006, J & K, India

*Email: abhadi.scholar@kashmiruniversity.net



ARTICLE HISTORY

Received: 16 February 2022

Accepted: 24 March 2022

Available online

Version 1.0 (Early Access): 25 April 2022

Version 2.0: 01 July 2022



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, etc. See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Hadi A, Singh S, Nawchoo I A, Rafiq S, Ali S. Impacts of habitat variability on the phenotypic traits of *Aconitum violaceum* Jacq. ex Stapf. at different altitudes and environmental conditions in the Ladakh Himalaya, India. Plant Science Today. 2022; 9(3): 546–554. <https://doi.org/10.14719/pst.1734>

Abstract

Aconitum violaceum Jacq. ex Stapf. is endemic to the Northern Himalayan regions of India, Pakistan and Nepal. To adapt and endure the rigorous stressful situations, phenotypic features of a specific plant species vary across different climatic conditions and elevations. The aim of this study was to reveal the impact of altitude and environmental conditions on the morphological attributes of a threatened species *A. violaceum*. Plants at high elevations (Maan-man, 4010 m asl) were short; whereas the plants grown at ecological conditions of population-I (Khawous, 3220 m asl) showed better growth and development in all morphological attributes. It has been noticed that there is a considerable reduction in the plant height, rhizome, leaf dimension, flower number and leaf number at higher elevations. Regression analysis and Pearson's correlation coefficient demonstrated a positive relationship between plant height and all other morphological attributes. Principal component analysis (PCA) revealed that the habitat, ecological conditions and elevations of population-I (Khawous) and Population-II (Numsuru) were proved to be relatively better for the luxuriant growth and development of an endemic plant species *A. violaceum*. From the observations, it can be inferred that *A. violaceum* grows successfully at an elevation ranges between 3000 to 3500 m asl. This study provides a comprehensive account of phenotypic variability concerning environmental conditions and suggested a suitable environment for sustainable cultivation of this important medicinal plant species in the Ladakh Himalaya.

Keywords

Distribution, habitat variability, morphology, phenotypic variability, regression

Introduction

Phenotypic plasticity is the potential of a genotype to manifest alternative morphological, behavioural and physiological attributes in response to environmental conditions (1, 2). To thrive under adverse environmental conditions, plants can either adjust to the local environmental circumstances genetically or respond to environmental variation through phenotypic plasticity (3, 4). Phenotypic plasticity can play an essential role in evolution by altering or modifying the developmental pathways in plant bodies, thus leading to the creation of phenotypic diversity in nature (1). Variations in general growth forms of resource accumulating organs such as roots, tubers, rhizomes, leaves and flowers of a plant are essential to adjust the available resources (5). Plants acclimatize themselves under acute environ-

mental conditions either by manifesting phenotypic malleability or by adapting genetically (6). Phenotypic flexibility is crucial for plants as they are stationary and must live in a constantly challenging environment (6, 7). Phenotypic variability is believed to exist more regularly at higher elevations, enabling alpine plants to adjust to the temporally fluctuating environment (2, 8). Although; plasticity is beneficial for the plant to survive under extreme climatic conditions; somehow it induces negative impacts on the fitness of plant species (9). With increasing elevations, plants must adapt to altering environmental conditions (10). Plants regulate important physiological processes such as gaseous exchange, photosynthetic rates and vegetative traits in response to climate fluctuations. Reducing plant height with respect to elevation disparity is the most noticeable change in plant morphology (10, 11).

Aconitum is a large genus of family Ranunculaceae with over 300 species distributed in the temperate areas of the Northern Hemisphere; such as Asia, Central Europe and North America (12). *A. violaceum* Jacq. ex Stapf. is an erect biennial herb attaining maximum height of 30 cm tall (13, 14), endemic to the Northern Himalayas (15). In India, *A. violaceum* is randomly distributed in the alpine and sub-alpine regions of Jammu and Kashmir, Himachal Pradesh, Uttarakhand and Ladakh Union Territory. *A. violaceum* is known as Aconite, violet monkshood in English; Atis, Bishmool and Mohand in Kashmir region and Bova-nagpo, Jhimba, Yangtso in Ladakh region (16).

A. violaceum is a highly useful medicinal plant, mostly harvested for its tuberous root, which are rich sources of diterpenoid alkaloids (aconitine, indaconitine etc) (17, 18), quercetin, acylated flavonol glycosides, flavonoids, kaempferol, anthraquinones, saponins, benzoic acid, sparteine, tannins, fatty acids and polysaccharide (19-21). It also has a wide range of biological activities including antioxidant, antimicrobial, anti-inflammatory, antiviral, anti-malarial, anti-proliferative, analgesic and antipyretic properties (19-21). Ethno-medicinally, it is used to cure boils, cough, fever, asthma, gastric trouble, sciatic pain, intestinal worms (22-24). The species are grown luxuriantly along the edges of *Rhododendron*, *Quercus* and *Abies* forests, glacial riverine, moist rocky areas, open grassy alpine slopes, stabilized alpine meadows, moist semi-shaded slopes (15).

Materials and Methods

In the present study, a dynamic technique was used to understand the phenotypic variability under different environmental and altitudinal conditions. The survey was conducted during May 2019 to October 2021.

Exploration, collection and documentation

Various habitats in Ladakh Himalaya (India) from Zojila to Zaskar in the West; Leh to Nubra in the East, were explored extensively and assessed to choose a specific study sites across the region. To investigate the phenotypic variability in *A. violaceum* along the altitudinal gradient and ecological conditions, 3 sites (populations) have been se-

lected based on their accessibility, habitat structure and specific environmental conditions, along with 1 control site at Kashmir University Botanical garden (KUBG). Given due cognizance to the threat status, vigor mature flowering plants of *A. violaceum* was collected quite judiciously from different study sites. The specimens were identified and deposited in the Kashmir University Herbarium (KASH) under voucher number 3739-KASH. 10-15 individuals of the species were randomly selected from each study site and were marked with tags. The tagged individuals were monitored on regular basis from May 2019 to October 2021. Vegetation type, habitat, geo-coordinates, altitudes along with associated species of the selected sites were recorded. The vegetative propagules (rhizome and seeds) of the plant were judiciously collected and transplanted to KUBG (control site) to study the morphological variations.

Species morphology and phenotypic variability

The randomly selected individuals in the populations were observed continuously at regular intervals of 5-10 days. Variations in morphological traits like plant height, rhizome, petiole, leaf, pedicel dimensions and floral characters such as the dimension of flower whorls, follicles and seeds were recorded.

Statistical analysis

LSD and Tukey's test (one-way ANOVA) were done using IBM-SPSS software (version 23) to compare the significant level and the average \pm standard error were computed using Microsoft Excel (office-10). Linear regression and Pearson's correlation coefficient analysis were performed to examine the correlation between various morphological attributes across different environmental conditions and elevations to find an ideal habitat for the better growth and development of this species. Principal component analysis (PCA) was carried out to assess the phenotypic traits in relation to different habitat conditions. Origin-Pro 2019b 64 software was used to analyze the PCA and R- software (version 4.1.2; 64 bit) was used to determine the correlation. To assess the variability in phenotypic features, data was collected for 10-15 plants from each population. The photographs were taken using a Canon DSLR (1300D), an Olympus (EPI-5) camera and a 1-plus 9-pro mobile camera.

Results and Discussion

Distribution

Aconitum violaceum is locally known as Nagpo- boma/ Bona-nagpo, Jhimba, Yangtso. In the Ladakh Union Territory, the species are randomly distributed in the alpine and sub-alpine regions of Drass (Matayeen, Prandrass, Maan-Man, Laabar), Suru Valley (Khawous, Numsuru, Pursa, Kochik, Tangole, Achampore, Choskore, Panikhar) Sapi, Nubra (Chalunka, Turtuk) on the way to Pangong lake, Changthang and Zaskar region at an altitude of 3100 to 4080 m asl. The distribution pattern and selected area are depicted in the map (Fig. 1). In Ladakh it is strictly distributed along the banks of continuous water flow irrigation channels, moist damp slopes under the shade of *Salix* species




Fig. 1. Distribution pattern of *Aconitum violaceum* Jacq. ex Stapf in Ladakh Himalaya, India.

and *Hippophae rhamnoides*, alpine wet meadows, banks of alpine stabilized sloped and bank of alpine streams, strongly associated with species like *Pedicularis rhinanthoides*, *Delphinium cashmerianum*, *Aconitum heterophyllum*, *Rumex acetosa*, *Gentianopsis vvedenskyi*, *Gentianopsis paludosa*, *Parnassia cabulica*, *Bistorta vivipara*, *Caryum carvi*, *Trifolium repens*, *Ranunculus brotherusii*, *Geranium pratense*, *Gentiana carniata*, *Taraxacum officinale*, *Anaphalis* spp. etc (Table 1).

Species morphology and phenotypic variability

The morphological characteristics of *A. violaceum* are enumerated in (Table 2), (Fig. 2. A-F). The phenotypic characters of *A. violaceum* showed significant variation at different altitudes and climatic conditions. This phenotypic variation was observed in 4 selected sites. Viz, Khawous (3220 m asl), Numsuru (3350 m asl), Maan-man, Drass (4020 m asl) and Kashmir University Botanical Garden (KUBG) Srinagar (1595 m asl) as a control site. The morphological

Table 1. Habitat characteristics, threat factor and geo-coordinates of the selected study sites.

Sl. No	Locality	Geo- coordinate		Elevation (m ASL)	Vegetation/Habitat	Threat factor
		Latitude	Longitude			
01	Khawous	N34°15.905	E75°96.040	3220	Oasitic; Moist damp places, along the bank of irrigation canals.	Pre-harvesting pressure, habitat fragmentation, over exploitation.
02	Numsuru	N34°12.705	E75°96.720	3340	Damp places, along the bank of irrigation canals, under the shade of <i>Salix</i> trees.	Pre-harvesting pressure, habitat fragmentation, over exploitation, prolonged seed dormancy.
03	Maan-man, Drass)	N34°38.592	E75°84.305	4010	Alpine grasslands, moist slopes, bank of streams	Snow slides, prolonged seed dormancy, pre-seed maturation frozen temperature, landslides, overgrazing.
04	KUBG	N34°75.717	E74°50.151	1595	Open field	Nil

Table 2. Morphological attributes of *Aconitum violaceum*

Habit	Biennial erect herb
Root	Paired tuberous root, slightly sweetish in taste.
Stem	Simple, erect up to 50 cm high, glabrous below, hairy above.
Leaves	Orbicular-cordate or reinform, deeply bipinnatisect and 5-partite, upper leaves much reduced or withered at the time of flowering. Lower leaves have long slender petioles; upper leaves has reduced petiole.
Bracts	Similar to upper leaves; but much reduced.
Bracteoles	linear to filiform.
Inflorescence	Short, lax or dense raceme or corymb or reduced to a solitary flower.
Sepals	Sepals violet, rarely blue or yellowish-green with blue veins, variegated, pubescent.
Petals	Nectaries, hairy with truncate recurved lips and hood is gibbous dorsally.
Androecium	Stamens numerous, 4-8 mm long, hairy in the upper part, winged in the lower part and these wings end in tiny teeth.
Gynoecium	Carpel 5, oblong, densely covered with soft, spreading hairs,
Flowering	20 th of July to ending August
Fruit	Bunch of 5 follicle, densely hairy
Seed	3-sided, obpyramidal
Senescence	Started from 10 th of September

**Fig. 2.** Morphological attributes of *A. violaceum*. A. Habit; B. Habitat. C. Inflorescence; D. Stamens; E. Ripened follicle; F. Seeds. Scale bar represents 0.5 cm.

traits that were assessed for phenotypic variability are; mean plant height, mean rhizome length, mean rhizome width, mean petiole length, mean leaf length, mean leaf width, mean pedicel length, flower length, flower width,

dimension of individual floral whorls (Sepals, petals, stamens and carpels), mean follicle length, mean seed length and mean seed width (Table 3 and 4). The populations of selected sites showed significant variations for studied

phenotypic traits along with elevations and environmental conditions. The mean plant height ranges from 29.69±2.91 to 66.4±2.41 cm, old rhizome length 2.74±0.16 to 3.66±0.21 4.56±0.29 to 9.06±0.25 cm, apical leaf length 1.15±0.08 to 1.65±0.19 cm, apical leaf width 0.75±0.07 to 1.14±0.16 cm, pedicel length 3.27±0.34 to 3.93±0.33 cm. Besides this; the

Table 3. Phenotypic variability in morphological attributes of *A. violaceum* at different altitudes and environmental conditions

Morphological Characters			Population (m asl)			
			Khawous-I (3220) N = 15	Numsuru-II (3340) 15	Man-Maan-Drass-III (4010) 15	KUBG-IV (1595) Control site 10
Unit (cm)	Code					
Plant Height (cm)	PH	*56.75±3.82 ^c	40.55±2.91 ^b	29.69±2.91 ^a	66.4±2.41 ^c	
Old rhizome length (cm)	ORL	3.44±0.21 ^b	2.95±0.15 ^{a,b}	2.74±0.16 ^a	3.66±0.21 ^c	
New rhizome Length (cm)	NRL	4.52±0.29 ^c	3.79±0.17 ^b	2.99±0.16 ^a	4.41±0.30 ^c	
Old rhizome width(cm)	ORW	0.70±0.05 ^a	0.58±0.04 ^a	0.53±0.04 ^a	0.61±0.03 ^a	
Rhizome New rhizome width (cm)	NRW	1.17±0.05 ^b	0.96±0.05 ^a	0.88±0.06 ^a	1.18±0.11 ^b	
Old rhizome girth (mm)	ORG	2.88±0.14 ^b	2.21±0.16 ^a	1.86±0.15 ^a	2.98±0.17 ^b	
New rhizome girth (mm)	NRG	4.34±0.16 ^c	3.62±0.15 ^b	2.97±0.09 ^a	4.11±0.23 ^c	
Fresh rhizome yield/ plant (g)	FRY	3.94±0.33 ^b	2.76±0.26 ^a	2.39±0.14 ^a	3.41±0.36 ^b	
Petiole Basal petiole length (cm)	BPL	12.23±0.94 ^b	10.18±1.11 ^{a,b}	6.90±0.64 ^a	13.2±1.04 ^b	
Upper petiole length (cm)	UPL	1.42±0.22 ^b	1.74±0.28 ^{a,b}	0.87±0.10 ^a	2.63±0.29 ^c	
Basal leaf length (cm)	BLL	5.53±0.16 ^c	3.26±0.17 ^a	2.81±0.20 ^a	4.52±0.46 ^b	
Basal leaf width (cm)	BLW	9.06±0.25 ^b	5.36±0.38 ^b	4.56±0.29 ^a	8.08±0.91 ^b	
Leaves Middle leaf length (cm)	MLL	3.25±0.15 ^b	2.72±0.12 ^a	2.51±0.09 ^a	3.38±0.27 ^b	
Middle leaf width (cm)	MLW	5.79±0.27 ^b	4.15±0.22 ^a	4.07±0.22 ^a	5.54±0.59 ^b	
Apical leaf length (cm)	ALL	1.37±0.05 ^b	1.28±0.07 ^a	1.15±0.08 ^a	1.65±0.19 ^b	
Apical leaf width (cm)	ALW	0.90±0.08 ^b	0.83±0.09 ^a	0.75±0.07 ^a	1.14±0.16 ^b	
Pedicel pedicel length (cm)	PL	3.51±0.37 ^a	3.35±0.36 ^a	3.27±0.34 ^a	3.93±0.33 ^a	

Different superscript letters (a, b and c) indicates that means are significantly different among different populations. (Tukey test: P ≤ 0.05)

*Mean± Standard Error; N=Sample size

Table 4. Phenotypic variability in floral attributes of *A. violaceum* at different altitudes and environmental condition

Floral Characters			Population (m asl)		
			Khawous-I (3220) N=15	Goma-Numsuru-II (3340) 15	Man-Maan-Drass (4010) 15
Unit (cm)	Code				
Flower	Flower length	FL	2.31±0.10 ^b	2.28±0.08 ^a	1.99±0.08 ^a
	Flower width	FW	2.91±0.08 ^b	2.84±0.06 ^b	2.53±0.10 ^a
	Posterior sepal length	PSL	1.65±0.08 ^b	1.50±0.08 ^{a,b}	1.29±0.05 ^a
	Posterior sepal width	PSW	1.07±0.10 ^a	1.09±0.09 ^a	0.9±0.03 ^a
Sepal	Lateral sepal length	LSL	1.5±0.06 ^a	1.38±0.07 ^a	1.36±0.10 ^a
	Lateral sepal width	LSW	1.0±0.04 ^a	0.978±0.06 ^a	0.952±0.77 ^a
	Inner sepal length	ISL	1.03±0.03 ^a	0.982±0.03 ^a	0.971±0.44 ^a
	Inner sepal width	ISW	0.51±0.05 ^a	0.426±0.02 ^a	0.383±0.38 ^a
Petal with nectaries	Petal length	PtL	1.70±0.02 ^a	1.37±0.16 ^a	1.35±0.07 ^a
Stamen	Filament length	FltL	0.61±0.03 ^a	0.59±0.02 ^a	0.54±0.21 ^a
Carpel	Carpel length	CL	0.6±0.02 ^b	0.51±0.01 ^a	0.46±0.01 ^a
Fruit	Follicle length	FoL	1.50±0.07 ^b	1.27±0.05 ^a	1.18±0.06 ^a
Seed	Seed length (mm)	SL	1.90±0.04 ^b	1.78±0.05 ^b	1.41±0.91 ^a
	Seed width (mm)	SW	0.95±0.05 ^b	0.89±0.02 ^{a,b}	0.79±0.03 ^a

Different superscript letters (a, b and c) indicates that means are significantly different among different populations. (Tukey test: P ≤ 0.05)

Mean± Standard Error; N=Sample size

cm, new rhizome length 2.99±0.16 to 4.52±0.29 cm, old rhizome width 0.53±0.04 to 0.70±0.05 cm, new rhizome width 0.88±0.06 to 1.22±0.11 cm, old rhizome girth 1.86±0.15 to 2.98±0.17 cm, new rhizome girth 2.97±0.09 to 4.34±0.16 cm, basal petiole length 6.90±0.64 to 13.2±1.04 cm, upper petiole length 0.87±0.10 to 2.63±0.29 cm, basal leaf length 2.81±0.20 to 5.53±0.16 cm, basal leaf width 1.50±0.07 cm and seed length varies from 1.41±0.91 to floral trait also showed variability with respect to altitude. The average floral length varies from 1.99±0.08 to 2.31±0.10 cm, floral width 2.53±0.10 to 2.91±0.08 cm; petal length along with nectaries 1.35±0.07 to 1.70±0.02 cm; stamens length 0.54±0.21 to 0.61±0.03 cm; carpel length 0.46±0.01 to 0.6±0.02 cm, follicle length 1.18±0.06 to 1.50±0.07 cm and seed length varies from 1.41±0.91 to

1.90±0.04 cm. The floral traits of control site (KUBG) are not mentioned here because it provided the same result as population-II. Most of the plants grown at the control site (KUBG) dried before reaching to flowering stage, as the environment was not suitable for its establishment. Most of the quantitative morphological traits showed a negative correlation with respect to altitude (i.e., they displayed a steady decline as altitude increased).

The morphological characteristics analyzed vary significantly across the selected populations. The different populations showed considerable morphological variation along with the altitude. Compared to all the populations; Population-I (Khawous 3220 m) exhibits the highest value for nearly all the morphological traits. The population established at the control site (KUBG) showed the highest value in terms of plant height and leaf dimension; however, they lost vigor as the summer temperature gradually increased. The remaining established individual plantlets developed flowers and fruits similar to the population-II, whereas the population-III (Maan-man, Drass) exhibited the lowest values for all the phenotypic characters. There was a positive correlation between plant height and other quantitative characters (i.e., individual plants with large height have larger growth forms for other phenotypic traits such as rhizome, leaf dimension, leaf number and flower number). There was a positive correlation between plant height and new rhizome length ($r^2 = 0.92635$), plant height and basal leaf length ($r^2 = 0.71765$), plant height and basal leaf width ($r^2 = 0.90183$), plant height and number of leaves ($r^2 = 0.41227$), plant height and number of flowers ($r^2 = 0.02309$) as shown in (Fig. 3. A-F). Pearson's correlation coefficient also showed a positive correlation for almost all quantitative phenotypic traits (rhizome dimension, leaf dimension, leaf number, inflorescence length, flower num-




Fig. 3. Graph (A-F). Regression analyses between several morphological characters of *A. violaceum*

ber etc) (Fig. 4). Principle Component Analysis (PCA) revealed that there is a gradual decrease in the size of morphological traits with the increase in altitude and vice versa (Fig. 5). The population growing at high elevation (Maan-man, Drass) showed a decrease in size of all phenotypic traits as compared to other populations, thus sepa-




Fig. 4. Scatter plot demonstrating Pearson's correlation matrix for quantitative phenotypic traits among different populations of *A. violaceum*. (***, $P < 0.000$, **, $P < 0.001$, *, $P < 0.01$).

rating them from the rest of the other populations of the study site, whereas the population established at Kashmir University Botanical Garden (KUBG) grew well at the onset of spring; but lost vigorousness as summer temperature rose. This implies that the environmental conditions are not favorable for the luxuriant growth of *A. violaceum* at

the control site. The environment and altitude of population-I (Khawous) are highly favorable for the plant species. Our results were in accordance with the former observations (25). It has been stated that reducing plant height and leaf dimension along with altitude is beneficial for the plant species as it prevents them from strong winds (26).




Fig. 5. Principal Component Analysis (PCA) of morphological characteristics of *A. violaceum* across the different study sites. ALL. Apical leaf length; PH. plant height; BPL. Basal petiole length; NRL. new rhizome length; NRG. New rhizome girth; BLL. Basal leaf length; LN. Leaf number.

It has been noticed that there is a tangled relationship between the environment and plants. With the change in altitudinal gradient, a range of environmental factors also vary which imposes a challenge to the plants, to adapt and survive under varying environmental and altitudinal gradients. Plants bring variations in their biochemical and morphological features to acclimate and overcome these stress circumstances (27). The change in morphological traits with varying elevation indicates plasticity and the fixed evolutionary changes (genetic) in these traits may lead to speciation (28).

Conclusion

The present study reveals that variations in the environmental conditions proved to have an enormous impact on the growth pattern and development of *A. violaceum*. The plants growing at altitudinal ranges between 3000-3220 m

asl (Population-I) were comparatively much more vigorous and diverse in all morphological characteristics. In addition, flower colours in population-I (khawous) were diverse (violet, blue, light white with blue veins, variegated); whereas the plant specimens grown at (KUBG) grew well in the beginning, however, as the summer temperatures rose, it gradually lost its vigorousness and dried out before reaching the flowering stage. The individuals of population -III (Maan-man, Drass) showed a decreased growth rate in all aspects of morphological traits such as plant height, leaf length, leaf width, number of flowers, rhizome length etc as compared to other population. This study aimed to find a suitable habitat for the establishment, development, cultivation, sustainable economic utilization and conservation of *A. violaceum*, a threatened endemic medicinal plant of the western Himalaya. It can be concluded that the ecology and elevation of population-I (khawous) are suitable for the growth and development of this threatened medicinal plant. Other areas which are also suitable for mass cultivation of this important medicinal plant species in Ladakh are, Nubra valley (Turtuk and Chalunka), Barsoo valley (Tiket, Khandi and Itchoo), Suru valley (Pursa, Thangbo) Drass valley (Matayeen, Prandrass, Mushkoo, Goshan etc) on large scale for commercial purposes as these regions have the same ecological conditions with population-I.

Acknowledgements

The first author is thankful to Council of Scientific and Industrial Research (CSIR) Pusa, New Delhi for providing financial assistance as SRF during this study. The authors are highly thankful of reviewers of journal 'Plant science Today' for valuable suggestions. All authors are grateful of the Head, Department of Botany, University of Kashmir for providing necessary research facilities to conduct this study.

Authors contributions

This is a collaborative research work of all authors. AH collected the primary data wrote the first draft of the manuscript and prepared graphs. SS edited and designed the manuscript and supervised AH. IAN edited, designed and supervised AH. SR performed the statistical analysis. SA helped in collecting the primary data. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors declared that there is no competing of interest.

Ethical issues: None

References

- Garland T, Kelly SA. Phenotypic plasticity and experimental evolution. *Journal of Experimental Biology*. 2006;209:2344-61. <https://doi.org/10.1242/jeb.02244>
- Magray JA, Zargar SA, Islam T, Nawchoo IA. Impact of habitat variability on growth dynamics of *Bergenia ciliata* (Haw.) Sternb. along an altitudinal gradient in Kashmir. *Plant Science Today*. 2021;9:144-49. <https://doi.org/10.14719/pst.1367>
- Becker U, Colling G, Dostal P, Jakobsson A, Matthies D. Local adaptation in the monocarpic perennial *Carlina vulgaris* at different spatial scales across Europe. *Oecologia*. 2006;150:506-18. <https://doi.org/10.1007/s00442-006-0534-9>
- Yaqoob U, Nawchoo IA. Impact of habitat variability and altitude on growth dynamics and reproductive allocation in *Ferula jaeschkeana* Vatke. *Journal of King Saud University – Science*. 2015. <http://dx.doi.org/10.1016/j.jksus.2015.10.002>
- Sattarian A, Akbarian M, Zarafshar M, Bruschi P, Fayyaz P. Phenotypic variation and leaf fluctuating asymmetry in natural populations of *Parrotia persica* (Hamamelidaceae), an endemic species from the Hyrcanian forest (Iran). *Acta Botanica Mexicana*. 2011;97:65-81. <https://doi.org/10.21829/abm97.2011.251>
- Sultan SE. Phenotypic plasticity for plant development, function and life history. *Trends in Plant Science*. 2000;5:537-42. [https://doi.org/10.1016/S1360-1385\(00\)01797-0](https://doi.org/10.1016/S1360-1385(00)01797-0)
- Hoffmann AA, Sgrò CM. Climate change and evolutionary adaptation. *Nature*. 2011;470:479-85. <https://doi.org/10.1038/nature09670>
- Stöcklin J, Kuss P, Pluess AR. Genetic diversity, phenotypic variation and local adaptation in the alpine landscape: case studies with alpine plant species. *Bot Helv*. 2009;119(2):125-33. <https://doi.org/10.1007/s00035-009-0065-1>
- Ghalambor CK, McKay JK, Carroll SP, Reznick DN. Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology*. 2007;21:394-407. <https://doi.org/10.1111/j.1365-2435.2007.01283.x>
- Körner Christian. The use of 'altitude' in ecological research. *Trends in Ecology and Evolution*, 2007;22:569-74. <https://doi.org/10.1016/j.tree.2007.09.006>
- Ma M, Zhou X, Wang G, Ma Z, Du G. Seasonal dynamics in alpine meadow seed banks along an altitudinal gradient on the Tibetan Plateau. *Plant and Soil*. 2010;336:291-302. <https://doi.org/10.1007/s11104-010-0480-5>
- Xiao PG, Wang FP, Gao F, Yan LP, Chen DL, Liu Y. A pharmacophylogenetic study of *Aconitum* L. (Ranunculaceae) from China. *Acta Phytotaxonomica Sinica*. 2006;44:1-46. <https://doi.org/10.1360/aps050046>
- Rledl H. The genus *Aconitum* in Pakistan and adjacent territories. *Linzer Biol Beitr*. 1993;25:67-79.
- Jabeen N, Kozgar MI, Dar GH, Shawl AS, Khan S. Distribution and taxonomy of genus *Aconitum* in Kashmir: Potent medicinal resource of Himalayan valley. *Chiang Mai Journal of Science*. 2013;40:173-86. <http://it.science.cmu.ac.th/ejournal/>
- Ved, D K, Kinhal GA, Ravikumar K, Prabhakaran V, Ghate U, Vijaya SR, Indresha JH. Conservation assessment and management prioritization for the medicinal plants of Jammu and Kashmir, Himachal Pradesh and Uttarakhand. *Foundation for Revitalisation of Local Health Traditions*, Bangalore, India, January. 2003.
- Hadi A, Singh S. Ethno-botanical studies of some threatened medicinal plants and local perception of its population decline in Kargil, Ladakh UT. *J Med Plants Stud*. 2021;9:44-50 <https://doi.org/10.22271/plants.2021.v9.i4a.1311>
- Miana GA, Ikram M, Khan MI, Sultana F. Alkaloids of *Aconitum violaceum*. *Phytochemistry*. 1971;10:3320-22. [https://doi.org/10.1016/S0031-9422\(00\)97417-X](https://doi.org/10.1016/S0031-9422(00)97417-X).
- Braca A, Fico G, Morelli I, De Simone F, Tomè F, De Tommasi N. Antioxidant and free radical scavenging activity of flavonol glycosides from different *Aconitum* species. *Journal of Ethnophar-*

- macology. 2003;86:63-67. [https://doi.org/10.1016/s0378-8741\(03\)00043-6](https://doi.org/10.1016/s0378-8741(03)00043-6)
19. Rawat JM, Rawat B, Mishra S, Negi R. Altitudinal and seasonal variation in bioactive compound aconitine in *Aconitum violaceum*, a threatened medicinal plant of Indian Himalayan Region. International Journal of Advanced Research. 2014;2:981-88.
 20. Sabir S, Arshad M, Hussain M. et al. A probe into biochemical potential of *Aconitum violaceum*: A medicinal plant from Himalaya. Asian Pacific Journal of Tropical Disease. 2016;6:502-04. [https://doi.org/10.1016/S2222-1808\(16\)61076-9](https://doi.org/10.1016/S2222-1808(16)61076-9)
 21. Khan FA, Khan S, Khan NM, Khan H, Khan S. Antimicrobial and antioxidant role of the aerial parts of *Aconitum violaceum*. J Mex Chem Soc. 2021;65:84-93. <https://doi.org/10.29356/jmcs.v65i1.1310>
 22. Batachariya A. Ethnobotanical observations in the Ladakh region of northern Jammu and Kashmir State, India. Economic Botany. 1991;45:305-08. DOI:10.1007/BF02887067
 23. Gairola, S, Sharma J, Bedi YS. A cross-cultural analysis of Jammu, Kashmir and Ladakh (India) medicinal plant use. Journal of Ethnopharmacology. 2014;155:925-86. <https://doi.org/10.1016/j.jep.2014.06.029>
 24. Hadi A, Singh S, Ali S, Mehdi M. Traditional uses of medicinal plants by indigenous tribes of Ladakh Union Territory. Res Jr of Agril Sci. 2022;13:68-77.
 25. Abrahamson WG, Gadgil M. Growth form and reproductive effort in goldenrods (*Solidago*, Compositae). Am Nat. 1973;107(957):651-61. <https://doi.org/10.1086/282864>
 26. Körner CH, Bannister P, Mark AF. Altitudinal variation in stomatal conductance, nitrogen content and leaf anatomy in different plant life forms in New Zealand. Oecologia. 1986;69:577-88. <https://doi.org/10.1007/BF00410366>
 27. Ncube. B, Finnie JF, Van SJ. Quality from the field: The impact of environmental factors as quality determinants in medicinal plants. South African Journal of Botany. 2012;82:11-20. <https://doi.org/10.1016/j.sajb.2012.05.009>
 28. Kim E, Donohue K. Demographic, developmental and life-history variation across altitude in *Erysimum capitatum*. Journal of Ecology. 2011;99:1237-49. <https://doi.org/10.1111/j.1365-2745.2011.01831.x>

§§§