



RESEARCH ARTICLE

# Acute toxicity and anticancer potential of knobweed (*Hyptis capitata*) ethanolic leaf extract and fraction

Nelsiani To'bungan\*, Wibowo Nugroho Jati, & Felicia Zahida

Faculty of Biotechnology, Universitas Atma Jaya Yogyakarta, Babarsari Street, Sleman 55281, Yogyakarta, Indonesia

\*Email: nelsiani.tobungan@uajy.ac.id

 OPEN ACCESS

## ARTICLE HISTORY

Received: 23 April 2022  
Accepted: 19 June 2022

Available online  
Version 1.0 : 28 August 2022  
Version 2.0 : 01 October 2022



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS etc. See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

To'bungan N, Jati WN, Zahida F. Acute toxicity and anticancer potential of knobweed (*Hyptis capitata*) ethanolic leaf extract and fraction. Plant Science Today. 2022; 9(4): 955–962. <https://doi.org/10.14719/pst.1847>

## Abstract

Ethanolic leaf extract of *Hyptis capitata* Jacq. showed a toxic effect on *Artemia salina* larvae. The extract was simplified through fractionation using vacuum liquid chromatography to trace compounds with biological activity. The F1, F2, F3 and F4 fractions were used in the acute toxicity test on *A. salina* larvae. F1 showed the highest toxicity value of  $196.77 \pm 2.7$   $\mu\text{g/ml}$ . As an initial screening for its anticancer potential, ethanolic leaf extract and F1 fraction were subjected to cytotoxicity test on T47D cells to obtain an  $\text{IC}_{50}$  value of  $45.32 \pm 2.0$  and  $>1000$   $\mu\text{g/mL}$ . The  $\text{LC}_{50}$  of the F1 fraction and its cytotoxicity level decreased compared to the ethanolic leaf extract. Based on GC-MS analysis, the ethanolic leaf extract and the F1 fraction contained neophytadiene, hexadecanoic acid, methyl ester, hexadecanoic acid, ethyl ester and heptadecanoic acid, 16-methyl-, methyl ester. Meanwhile,  $\alpha$ -sitosterol, docosanoic acid, ethyl ester and squalene are only found in the ethanolic leaf extract. The compounds contained in ethanolic leaf extract work synergistically. This implies the ethanolic leaf extract has the potential to be developed as an anticancer candidate.

## Keywords

Anticancer, cytotoxicity, *Hyptis capitata*, GC-MS, toxicity

## Introduction

Plants contain secondary metabolites that can be utilized for human benefit. In Indonesia, the cultural heritage of using many plants as medicinal ingredients is the basis for further exploration of the efficacy and safety of certain plants. For example, Knobweed or *Hyptis capitata* is one of the plants widely used by the community as a medicinal ingredient (1, 2). This plant is used to treat fever, open and internal wounds (1, 2) and also has opportunity to act as an antioxidant, antiviral, antibacterial and anticancer (3). However, the safety in its utilization and scientific potential still needs to be studied in depth to underlie its development in the future.

A previous research conducted an acute toxicity test of *H. capitata* ethanolic leaf extract using the Brine Shrimp Lethality Test Method on *Artemia salina* larvae and obtained  $\text{LC}_{50}$  183.91  $\mu\text{g/ml}$  (4) for the toxic category (4, 5). An early acute toxicity test can be the basis for the next stage (6). The initial screening methods determine the potential of a test material to be developed as an anticancer candidate (5). This research develops previous results comparing ethanolic leaf extract's toxicity level and fraction. It also associated the results of acute toxicity tests with their potential as anticancer based on cytotoxicity activity. The fractionation procedures simplify

and trace the secondary metabolite components in the ethanolic leaf extract and the toxicity activity using the Vacuum Liquid Chromatography method. The results were reused for acute toxicity tests using the Brine Shrimp Lethality Test method. The fraction that showed the highest activity was subjected to a cytotoxicity test on T47D breast cancer cells. A cytotoxicity test was carried out as an initial screening of the potential of the ethanolic leaf extract and its fraction as an anticancer candidate. Screening of metabolite compounds was also conducted using the Gas Chromatography-Mass Spectrometry Method.

## Materials and Methods

The material used was *H. capitata* Jacq. (Fig. 1.) leaves taken from Bungapati village, Tana Lili Regency, North Luwu District, South Sulawesi. This plant has been identified in the plant systematics laboratory, Faculty of Biology,



Fig. 1. *Hyptis capitata*. Habit.

Universitas Gadjah Mada, with a certificate number of 014535/S.Tb/III/2019 (7).

### Extraction

The *H. capitata* leaves were taken in the morning, washed thoroughly with running water, covered with a black cloth and dried in the sun (until the dry weight was stable). The leaves were mashed with a grinder, filtered and then weighed. Subsequently, *H. capitata* leaf dry powder was extracted using absolute ethanol solvent in a ratio of 1:5.

### Vacuum chromatography-mass spectrometry

A total of 2.5 g of ethanolic leaf extract was mixed with 5 g of silica gel 60 GF254 powder (Merck, Germany) to obtain a homogeneous mixture. The stationary phase used was 12 g of silica gel 60 GF254. Furthermore the eluent composition was n-hexane:chloroform 3:1(v/v), 2:2 (v/v), 1:3 (v/v), 100% chloroform, chloroform : ethanol 3:1 (v/v), 2:2 (v/v), 1:3 (v/v) and ethanol 100%. The obtained fraction was collected in an Erlenmeyer, poured into a porcelain dish and air-dried. VLC was followed by thin-layer chromatography to monitor the separation of the compound (8).

### Thin-layer chromatography

The mobile and stationary phase used was ethyl acetate: methanol (7:3 v/v) and silica gel F254 (Merck, Germany). The chromatogram pattern was observed under UV light at 254 and 366 nm wavelengths and combined into one fraction.

### A. salina hatching

A total of 0.5 g of artemia eggs were put in a plastic bottle filled with 1 liter of seawater and the hatchery was placed at room temperature. It was given an aerator and left for 36 hrs until the eggs hatched into larvae (4).

### Brine shrimp lethality test

*H. capitata* leaf fraction was dissolved with Dimethyl-sulfoxide (DMSO) in a ratio of 1:5. Each fraction was made into a concentration series of 300; 150; 75; 37.5; 18.75 µg/ml. Meanwhile, the tube was filled with a volume of 20 ml and 10 larvae of 36 hrs old were put in the solution. The test was accompanied by a negative and solvent control (DMSO) and each concentration was made in 3 replications. There was also a negative control tube filled with seawater and 10 larvae. The tube was stored for 24 hrs without a cover and lamp irradiation. Furthermore, the number of live larvae was counted to determine those that died due to the treatment (4).

### Cytotoxicity assay

A cytotoxicity test was performed on T47D cells cultured using 96 well-plates with a density of  $10^4$  cells/well in a complete RPMI culture medium (Gibco, Canada). In addition, 100 µl of cell suspension filled the well, and the cells were incubated for 24 hrs in a CO<sub>2</sub> incubator at 36 °C and 5% CO<sub>2</sub>. These cells were treated when they reached 70-80% confluency. Ethanolic leaf extract and fraction 1 of 10 mg were weighed and dissolved in 100 µl DMSO (Merck, Germany). The series concentration was made in 200 µg/ml, 100 µg/ml, 50 µg/ml, 25 µg/ml, and 12.5 µg/ml with 3 replications. The cultured cells were incubated for 24 hrs

in a CO<sub>2</sub> incubator. The conditions were observed and documented before adding MTT (Bio Basic, Canada) (5 mg/ml PBS) in a ratio of 1:9. Cells were incubated for 4 hrs in a CO<sub>2</sub> incubator and 100 µl Reagent Stop (SDS) (Merck, Germany) was added. The well-plate was wrapped in aluminum foil and incubated at room temperature overnight. The absorbance was measured with a microplate reader machine at 595 nm (9).

#### Gas chromatography-mass spectrometry (GC-MS)

Secondary metabolites were analyzed by the GC-MS method at the Integrated Research and Testing Laboratory (LPPT), Universitas Gadjah Mada, Yogyakarta, using an HP-5MS UI capillary column with a column length of 30 m, a diameter of 0.25 mm and thickness of 0.25 m. Furthermore, the sample was dissolved in pure ethanol and injected using a split technique. The initial conditions with the splitless model were as follows: split-flow 50 ml/min, with a front inflow of 1.00 ml/min. Helium was used as the carrier gas, and the injector temperature was programmed at 260 °C. The ion source temperature was 200 °C, while the mass spectrometry transfer line was 250 °C, with a speed of 5 °C/min. The gas saver time was 5 minutes, purge flow was 3 ml/min, and flow of gas saver was 5 ml/min. Meanwhile, the Mass spectrometry transfer line temperature was (MS) 250 °C and the detected peaks were analyzed descriptively (7).


#### Data analysis

The data obtained were analyzed quantitatively or qualitatively, and the LC<sub>50</sub> value was determined using probit analysis. The IC<sub>50</sub> value was determined by log concentration regression analysis. Furthermore, the data obtained were processed using SPSS 26 and analyzed with a one-way analysis of variance at a 95% confidence level and a significance value of P <0.05.

## Results

#### Fractionation with vacuum liquid chromatography

The results of the fractionation of the ethanolic leaf extract



**Fig. 2.** Thin-Layer Chromatography Fraction leaves of *H. capitata*. **A.** Visualization with UV 365 nm; **B.** Visualization with UV 254 nm. 1 = n-hexane:chloroform fraction 3:1 (v/v); 2 = n-hexane:chloroform fraction 2:2 (v/v); 3 = n-hexane:chloroform fraction 1:3 v/v; 4 = 100% chloroform fraction; 5 = chloroform: ethanol fraction 3:1 v/v; 6 = chloroform:ethanol fraction 2:2 v/v; 7 = chloroform: ethanol fraction 1:3 v/v; 8 = 100% ethanol fraction. Based on the visualization, the fractions were grouped into four, namely fraction 1 separately, then referred to as the first fraction (F1); fractions 2, 3, 4 were combined and were referred to as second fractions (F2); fraction 5 separately then referred to as the third fraction (F3) and fractions 6, 7 and 8 were combined and were referred to as the fourth fraction (F4).

can be seen in Fig. 2. The obtained fractions were visualized by thin-layer chromatography, and the separation profile was observed under ultraviolet light. Based on the figure, the 8 fractions obtained were simplified in number by combining those with a similar pattern. The 8 fractions were simplified to 4 and used in an acute toxicity test on *A. salina*. The most toxic LC<sub>50</sub> value was continued for the cytotoxicity test on T47D cells.



#### Acute toxicity test and cytotoxicity


Based on the LC<sub>50</sub> value (Table 1), the F1-F4 treatment was toxic to *A. salina*. The obtained LC<sub>50</sub> value was <1000 µg/ml,

**Table 1.** Acute toxicity and cytotoxicity of ethanolic leaf extract *H. capitata* and its fraction.

Treatment	LC <sub>50</sub> Average±SD (µg/ml) on <i>A. salina</i>	IC <sub>50</sub> Average±SD (µg/ml) on T47D cells
Ethanolic leaf extract	N.D	45.32±2.0
F1	196.77±2.7 <sup>a</sup>	>1000
F2	246.15±5.8 <sup>b</sup>	N.D
F3	286.93±10.23 <sup>b</sup>	N.D
F4	390.275±32 <sup>c</sup>	N.D

which can be categorized as toxic (5). The treatment of the F1 fraction showed the highest acute toxicity value against *A. salina*. Meanwhile, the F1 cytotoxicity test on T47D cells showed a value of >1000 µg/ml (Tabel 1), hence, it is not cytotoxic. This is different from the IC<sub>50</sub> value of ethanolic leaf extract treatment on T47D cells (Table 1), which was 45.32±2.0 µg/ml, classified as moderate. The morphology of cells exposed to ethanolic leaf extract and F1 fraction





**Fig. 3.** Effect of Ethanolic leaves extracts and Fraction F1 treatment on T47D Cells using an inverted microscope. **A.** T47D cell control, **B.** ethanolic leaf extract treatment on T47D cells, **C.** Fraction F1 treatment on T47D cells.

can be seen in Fig. 3. Cells treated with the extract showed variations in morphology and the cytotoxicity effect was indicated by the change in the shape of the cells into small rounds. Different variables were shown in T47D cells exposed to the F1 fraction having the same morphology as control cells.

#### Identification of bioactive components


The peak chromatogram of the content of metabolite compounds in the ethanolic leaf extract and the F1 frac-

tion can be seen in Fig. 4. It shows that 96 peaks were detected in the ethanolic leaf extract and F1 fraction 44 was detected. Meanwhile, the list of compounds with a similarity index value above 750 is presented in Table 2. In the ethanolic leaf extract, there were 27 components, while the F1 fraction had 12 compounds. The compounds contained in the extract and fraction F1 were neophytadiene, hexadecanoic acid, methyl ester, hexadecanoic ethyl ester, heptadecanoic acid and phytol.

#### Discussion

The acute toxicity and cytotoxicity test of the *H. capitata* Jacq. leaf fraction are presented in Table 1. It shows that the  $LC_{50}$  value of the F1 fraction treatment on *A. salina* larvae was  $196.77 \pm 2.7 \mu\text{g/ml}$ , classified as toxic (5). The toxic effects of compounds can also be useful for treating several diseases (12). These are caused by the compounds contained in the extract. Based on statistical tests, these values are known to be significantly different from the  $LC_{50}$  of other fractions. Therefore, acute toxicity tests are conducted to evaluate the side effects after the organism is exposed to the test material within 24 hrs (6).

The  $LC_{50}$  value of the F1 fraction decreased when compared with ethanolic leaf extract in previous research ( $183.91 \mu\text{g/ml}$ ). Meanwhile, the treatment of the extract on



**Fig. 4.** Chromatogram of metabolite compounds from the leaves ethanolic extract and fraction F1. (A) Chromatogram of metabolite compounds of the *H. capitata* leaves ethanolic extract; 96 peaks were detected. (B) Chromatogram of the metabolite compounds fraction F1; 44 peaks were detected.

**Table 2.** Metabolite compounds of ethanolic leaf extract and fraction F1

Sl. No.	RT	%Area	Formula	Compound
<b>Ethanolic leaf extract</b>				
1	22.83	0.06	C <sub>15</sub> H <sub>24</sub>	β-copaene
2	23.04	0.11	C <sub>15</sub> H <sub>24</sub>	1-Isopropyl-4,7-dimethyl-1,2,3,5,6,8a-hexahydronaphthalene
3	29.06	0.07	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	Tetradecanoic acid, ethyl ester
4	29.87	0.08	C <sub>20</sub> H <sub>40</sub>	2-Hexadecene, 3,7,11,15-tetramethyl-, [R-[R*,R*-(E)]]-
5	29.97	0.65	C <sub>20</sub> H <sub>38</sub>	Neophytadiene
6	30.85	0.22	C <sub>20</sub> H <sub>40</sub> O	3,7,11,15-Tetramethyl-2-hexadecen-1-ol
7	31.62	0.07	C <sub>37</sub> H <sub>76</sub> O	1-Heptatriacotanol
8	31.73	0.61	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	Hexadecanoic acid, methyl ester
9	32.64	0.10	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	E-11-Hexadecenoic acid, ethyl ester
10	32.94	0.50	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	Ethyl 9-hexadecenoate
11	33.06	17.12	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	Hexadecanoic acid, ethyl ester
12	33.75	0.12	C <sub>28</sub> H <sub>48</sub> O	Cholestan-3-ol, 2-methylene-, (3β,5α)-
13	34.92	0.66	C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	Ethyl 14-methyl-hexadecanoate
14	35.04	1.47	C <sub>19</sub> H <sub>32</sub> O <sub>2</sub>	9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)-
15	35.30	11.17	C <sub>20</sub> H <sub>40</sub> O	Phytol
16	35.51	0.52	C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	Heptadecanoic acid, 16-methyl-, methyl ester
17	36.13	2.62	C <sub>20</sub> H <sub>36</sub> O <sub>2</sub>	9,12-Octadecadienoic acid, ethyl ester
18	36.26	12.34	C <sub>20</sub> H <sub>34</sub> O <sub>2</sub>	Ethyl 9,12,15-octadecatrienoate
19	36.34	0.80	C <sub>21</sub> H <sub>36</sub> O <sub>4</sub>	9,12,15-Octadecatrienoic acid, 2,3-dihydroxypropyl ester, (Z,Z,Z)
20	36.17	8.18	C <sub>20</sub> H <sub>40</sub> O <sub>2</sub>	Heptadecanoic acid, 15-methyl-, ethyl ester
21	37.14	0.17	C <sub>20</sub> H <sub>40</sub> O <sub>2</sub>	Ethanol, 2-(9-octadecenyloxy)-, (Z)-
22	37.95	0.02	C <sub>26</sub> H <sub>44</sub> O <sub>5</sub>	Ethyl iso-allocholate
22	38.72	0.55	C <sub>29</sub> H <sub>50</sub> O	γ-sitosterol
23	40.08	3.89	C <sub>22</sub> H <sub>44</sub> O <sub>2</sub>	Eicosanoic acid, ethyl ester
24	44.07	1.70	C <sub>24</sub> H <sub>48</sub> O <sub>2</sub>	Docosanoic acid, ethyl ester
25	46.72	1.99	C <sub>29</sub> H <sub>50</sub> O <sub>2</sub>	Vitamin E
26	51.72	9.89	C <sub>30</sub> H <sub>50</sub>	Squalene
27	58.30	0.17	C <sub>30</sub> H <sub>48</sub> O <sub>2</sub>	Betulinaldehyde
<b>F1</b>				
1	29.97	0.55	C <sub>20</sub> H <sub>38</sub>	Neophytadiene
2	30.11	0.49	C <sub>17</sub> H <sub>36</sub> O	1-Hexadecanol, 2-methyl-
3	31.31	0.35	C <sub>17</sub> H <sub>32</sub> O <sub>2</sub>	9-Hexadecenoic acid, methyl
4	31.73	7.98	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	Hexadecanoic acid, methyl ester
5	33.05	5.23	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	Hexadecanoic acid, ethyl ester
6	34.92	3.49	C <sub>19</sub> H <sub>34</sub> O <sub>2</sub>	Methyl 9-cis,11-trans-octadecadienoate
7	35.03	15.30	C <sub>19</sub> H <sub>36</sub> O <sub>2</sub>	trans-13-Octadecenoic acid, methyl ester
8	35.51	4.98	C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	Heptadecanoic acid, 16- methyl-, methyl ester
9	36.15	2.13	C <sub>20</sub> H <sub>40</sub> O	Phytol
10	41.88	6.17	C <sub>30</sub> H <sub>50</sub> O	A-amyrin
11	43.14	2.90	C <sub>30</sub> H <sub>48</sub> O	Lup-20(29)-en-3-one
12	2.11	51.67	C <sub>30</sub> H <sub>52</sub> O	2,2,4-Trimethyl-3-(3,8,12,16-tetramethyl-heptadeca-3,7,11,15-

T47D cells showed a moderate level of cytotoxicity. Different variables were seen in the IC<sub>50</sub> value of F1 fraction treatment on T47D cells. The level of cytotoxicity of the ethanolic leaf extract was moderate, indicating an opportunity to be developed as an anticancer candidate. On the other hand, the LC<sub>50</sub> and IC<sub>50</sub> values of the F1 fraction in T47D cells decreased compared to the crude extract. This

is influenced by the composition of metabolite compounds in toxicity and cytotoxicity (13).

Based on the results of the GC-MS analysis, the ethanolic leaf extract contained several compounds which played a role in toxicity and cytotoxicity. The compounds are neophytadiene; hexadecanoic acid, methyl ester; hexa-

decanoic acid, ethyl ester; heptadecanoic acid, 16- methyl-, methyl ester; phytol;  $\alpha$ -sitosterol; docosanoic acid, ethyl ester; and squalene. They were also detected in the F1 fraction, except  $\alpha$ -sitosterol; docosanoic acid, ethyl ester; and squalene.

Neophytadiene and phytol are terpenoids with antimicrobial and anti-inflammatory properties (14). Neophytadiene is also one of the main components in *Acalypha segetalis* leaf essential oil, which play a role in the toxicity activity of *A. salina* larvae (15). Furthermore, the toxicity of this compound isolated from *Selenicereus hamatus* showed a high value for *A. salina* larvae, with an LC<sub>50</sub> of 16.52  $\mu$ g/ml (16). Meanwhile, phytol shows pharmacological activity and can be used as an anticancer (17, 18). Phytol enhances pro-carcinogens and induces genotoxicity as well as apoptosis in breast cancer cells (19). The content of phytol and, hexadecanoic acid, methyl esters in *Calotropis gigantea* leaves can also influence toxicity activity against *A. salina* larvae (20). In the ethanolic leaf extract and the fraction F1, these compounds had a toxic effect on *A. salina* larvae.

Hexadecanoic acid, methyl ester (palmitic acid, methyl ester) (21), hexadecanoic acid ethyl ester (palmitic acid, ethyl ester) (22), Heptadecanoic acid, 16- methyl-, methyl ester (methyl isostearat) (23) and docosanoic acid ethyl esters (Ethyl docosanoate) is a group of fatty acid ester compounds (24) with anticancer activity. Hexadecanoic acid can affect the growth of colon cancer cells (25), acts as a growth inhibitor, triggers apoptosis and inhibits the cell cycle in the G0/G1 phase (26). In addition, the ethyl ester derivative can inhibit DNA topoisomerase I and is an inducer of apoptosis in leukemia and neuroblastoma cells (27-29). Ethyl docosanoate has also been reported as a secondary metabolite in the cytotoxic activity of *Xetospongia testudinaria* isolate (30).

$\alpha$ -sitosterol has anticancer activity and can inhibit the growth of MCF-7 and A549 cells. In addition, this compound triggers arrests in the G2/M phase of the cell cycle and reduces the expression of the transcription factor c-Myc in cell division (31).

In line with the previously described metabolites compounds, squalene also has cytotoxicity and antitumor potential against HeLa cells. In combination with the anticancer agent Adriamycin, it shows a synergistic effect (32). Furthermore, squalene has the potential as an anti-inflammatory and can inhibit inflammation by increasing the production of prostaglandin E2 during chemotherapy in tumor-bearing mice (33). The decreased level of cytotoxicity in the F1 was thought to be caused by the absence of  $\alpha$ -sitosterol, docosanoic acid ethyl esters and squalene.

Under certain circumstances, some compounds may exhibit a synergistic or positive effect. This is shown when the combination of many secondary metabolites is greater than the single effect (34). Therefore, the compounds detected in the ethanolic leaf extract work synergistically to provide a strengthening effect on toxicity and cytotoxicity. The reduction in the composition of compounds in the F1 fraction indicated a decrease in the acute

toxicity and cytotoxicity effects. The absence of  $\alpha$ -sitosterol, docosanoic acid, ethyl ester and squalene in the F1 fraction greatly influenced the cytotoxicity level of the F1 fraction in T47D cells.

This study shows the opportunity of ethanolic leaf extract of *H. capitata* to be developed as an anticancer candidate. Several *Hyptis* species also have cytotoxic activity on cancer cell lines. These species include *H. suaveolens* (35, 36), *H. brevipes* (37), *H. martiusii* (38), *H. incana* (39) and *H. pectinata* (40).

## Conclusion

The acute toxicity and cytotoxicity of ethanolic leaf extract are higher than the F1 fraction. The extract contained neophytadiene; hexadecanoic acid, methyl ester; hexadecanoic acid, ethyl ester; heptadecanoic acid, 16- methyl-, methyl ester; phytol;  $\alpha$ -sitosterol; docosanoic acid, ethyl ester; and squalene. These compounds can positively impact toxicity and cytotoxicity, working synergistically. Therefore, the ethanolic leaf extract of *H. capitata* can be developed as an anticancer candidate.

## Acknowledgements

We extend our gratitude to Universitas Atma Jaya Yogyakarta for providing support for this research through the UAJY Internal Grant.

## Authors contributions

NT conducts research, processes data and manuscript writing. FZ contributed to provide important suggestions related to research and manuscript writing. WNJ conceived of the study and coordination. All authors read and approved the final manuscript.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** None.

## References

- To'bungan N. Utilization and phytochemicals screening of knobweed (*Hyptis capitata* Jacq.). *Biota J Ilm Ilmu-Ilmu Hayati*. 2020;5:149-54. <https://doi.org/10.24002/biota.v6i1.3577>
- Rupa D, Sulistyarningsih CY, Dorly, Ratnadewi D. Identification of secretory structure, histochemistry and phytochemical compound of medical plant *Hyptis capitata* Jacq. *Biotropia*. 2019;24:94-103. <https://doi.org/10.11598/btb.2017.24.2499>
- To'bungan N, Widyarini S, Nugroho L H, Pratiwi R. Ethnopharmacology of *Hyptis capitata*. *Plant Science Today*. 2022;9(3):593-600. <https://doi.org/10.14719/pst.1602>
- To'bungan N, Jati WN, Zahida F. Acute toxicity of knob weed leaves ethanolic extract (*Hyptis capitata* Jacq.) using brine shrimp lethality test (BSLT). *Sciscitatio*. 2020;1:64-69. <https://doi.org/10.21460/sciscitatio.2020.12.30>
- Meyer BN, Ferrigni NR, Putnam JE. Brine shrimp: A convenient

- general bioassay for active plant constituents. *Planta Med.* 1982;45:31-34. <https://doi.org/10.1055/s-2007-971236>
6. Harmita, Radji M. Buku Ajar Analisis Hayati. Jakarta: Erlangga; 2008. p. 42-77.
  7. To'bungan N, Pratiwi R, Widyarini S, Nugroho LH. Cytotoxicity extract and fraction of knobweed (*Hyptis capitata*) and its effect on migration and apoptosis of T47D cells. *Biodiversitas.* 2022;23:572-80. <https://doi.org/10.13057/biodiv/d230162>
  8. Dewatisari WF, Retnaningrum E, Purwestri YA, Nugroho LH. The potency of *Sansevieria trifasciata* and *S. cylindrica* leaves extracts as an antibacterial against *Pseudomonas aeruginosa*. *Biodiversitas.* 2021;22:408-15. <https://doi.org/10.13057/biodiv/d220150>
  9. Anonymous. 2019. *In vitro* test protocol, cancer chemoprevention researches center (CCRC) Universitas Gadjah Mada [Internet]. [cited 2019 September 13]. Available from: [http://ccrc.farmasi.ugm.ac.id/en/?page\\_id=240](http://ccrc.farmasi.ugm.ac.id/en/?page_id=240)
  10. Hameed E. Phytochemical studies and evaluation of antioxidant, anticancer and antimicrobial properties of *Conocarpus erectus* L. growing in Taif, Saudi Arabia. *Eur J Med Plants.* 2012;2(2): 93-112. <https://doi.org/10.9734/ejmp/2012/1040>
  11. Srisawat T, Chumkaew P, Heed-Chim W, Sukpondma Y, Kanokwiroon K. Phytochemical screening and cytotoxicity of crude extracts of *Vatica diospyroides* Symington type LS. *Trop J Pharm Res.* 2013;12(1):71-76. <https://doi.org/10.4314/tjpr.v12i1.12>
  12. Madariaga-Mazón A, Hernández-Alvarado RB, Noriega-Colima KO, Osnaya-Hernández A, Martínez-Mayorga K. Toxicity of secondary metabolites. *Phys Sci Rev.* 2019;4:1-11. <https://doi.org/10.1515/psr-2018-0116>
  13. Lopes AP *et al.* Antioxidant and cytotoxic effects of crude extract, fractions and 4-nerolidylcatechol from aerial parts of *Pothomorphe umbellata* L. (Piperaceae). *Biomed Res Int.* 2013;1-5. <http://dx.doi.org/10.1155/2013/206581>
  14. Venkata RB, Samuel LA, Pardha SM, Narashima RB, Naga VKA, Sudhakar M, Radhakrishnan TM. Antibacterial, antioxidant activity and GC-MS analysis of *Eupatorium odoratum*. *Asian J Pharm Clin Res.* 2012;5(2):99-106.
  15. Aboaba SA, Aiyelaagbe OO, Ekundayo O. Chemical composition, toxicity and larvicidal activity of the essential oil from the whole plant of *Acalypha segetalis* from south-west Nigeria. *Nat Prod Commun.* 2010;5:481-83. <https://doi.org/10.1177/1934578x1000500328>
  16. Umaru JI, Ahmed UM, Mohammed, A. Characterization of chemical constituent and evaluation of antioxidant, cytotoxicity, potential of *Selenicereus hamatus* crude extract. *Asian J Biochem.* 2020;15:50-59. <https://doi.org/10.3923/ajb.2020.50.59>
  17. Abdel-Aal El, Amany MH, Jalan M. 2015. Successive solvent extraction and GC-MS analysis for the evaluation of the phytochemical constituent of the filamentous green algae *Spirogyra longata*. *Egypt J Aquat Res.* 2014;(41):233-46. <https://doi.org/10.1016/j.ejar.2015.06.001>
  18. Pratama OA, Tunjung WAS, Sutikno, Daryono BS. Bioactive compound profile of melon leaf extract (*Cucumis melo* L. 'Hikapel') infected by downy mildew. *Biodiversitas.* 2019;20:3448-53. <https://doi.org/10.13057/biodiv/d201143>
  19. de Alencar MVOB *et al.* Phytol as an anticarcinogenic and anti-tumoral agent: An *in vivo* study in swiss mice with DMBA-Induced breast cancer. *IUBMB Life.* 2019;71:200-12. <https://doi.org/10.1002/iub.1952>
  20. Istikasari F. Uji toksisitas ekstrak daun rovinga (*Calotropis gigantea*) dengan metode BSLT (Brine Shrimp Lethality Test) pada larva *Artemia salina* [Thesis]. Indonesia: Fakultas kedokteran dan ilmu kesehatan. Universitas Tadulako. 2016.
  21. Anonymous. Methyl palmitate, National Center for Biotechnology Information [Internet]. [Cited 2022 April 7]. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/Methyl-palmitate>
  22. Anonymous. Ethyl palmitate, National Center for Biotechnology Information [Internet]. [cited 2022 April 7]. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/Ethyl-palmitate>
  23. Anonymous. Methyl isostearate, National Center for Biotechnology Information [Internet]. [Cited 2022 April 7]. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/Methyl-isostearate>
  24. Anonymous. Ethyl docosanoate, National Center for Biotechnology Information [Internet]. [Cited 2022 April 7]. Available from <https://pubchem.ncbi.nlm.nih.gov/compound/Ethyl-docosanoate>
  25. Ravi L, Krishnan K. Cytotoxic potential of N-hexadecanoic acid extracted from *Kigelia pinnata* leaves. *Asian J Cell Biol.* 2016;12:20-27. <https://doi.org/10.3923/ajcb.2017.20.27>
  26. Bharath B, Perinbam K, Devanesan S, AlSalhi MS, Saravanan M. Evaluation of the anticancer potential of Hexadecanoic acid from brown algae *Turbinaria ornata* on HT-29 colon cancer cells. *J Mol Struct.* 2021;1235. <https://doi.org/10.1016/j.molstruc.2021.130229>
  27. Harada H, Yamashita U, Kurihara H, Fukushi E, Kawabata J, Kamei Y. Antitumor activity of palmitic acid found as a selective cytotoxic substance in a marine red alga. *Anticancer Res.* 2002;22(5):2587-90.
  28. Pereira DM, Correia-da-Silva G, Valentao P, Teixeira N, Andrade PB. Palmitic acid and ergosta-7,22-dien-3-ol contribute to the apoptotic effect and cell cycle arrest of an extract from *Marthasterias glacialis* L. in neuroblastoma cells. *Marine drugs.* 2013;12(1):54-68. <https://doi.org/10.3390/md12010054>
  29. Mellado M *et al.* *In vitro* antioxidant and antiproliferative effect of the extracts of *Ephedra chilensis* K Presl aerial parts. *BMC Complement. Altern Med.* 2019;19:1-10. <https://doi.org/10.1186/s12906-019-2462-3>
  30. Swantara MD, Rita WS, Suartha N, Agustina KK. Anticancer activities of toxic isolate of *Xestospongia testudinaria* sponge. *Vet. World.* 2019;12:1434-40. <https://doi.org/10.14202/vetworld.2019.1434-1440>
  31. Sundarraj S. *et al.*  $\gamma$ -Sitosterol from *Acacia nilotica* L. induces G2/M cell cycle arrest and apoptosis through c-Myc suppression in MCF-7 and A549 cells. *J Ethnopharmacol.* 2012;141:803-09. <https://doi.org/10.1016/j.jep.2012.03.014>
  32. Nakagawa M, Komiyama S, Fukawa H, Kuwano M, Akiyama SI. Potentiation by squalene of the cytotoxicity of anticancer agents against cultured mammalian cells and murine tumor. *Japanese J Cancer Res.* 1985;76:15-320.
  33. Bhilwade NH *et al.* The Adjuvant effect of squalene, an active ingredient of functional foods on doxorubicin-treated allograft mice. *Nutr Cancer.* 2019;71:1153-64. <https://doi.org/10.1080/01635581.2019.1597900>
  34. Caesar LK, Cech NB. Synergy and antagonism in natural product extracts: when 1 + 1 does not equal 2. *Nat Prod Rep.* 2019;76(4):139-48. <https://doi.org/10.1039/c9np00011a>
  35. Poonkodi K, Karthika J, Tamilselvi V, Anitha R, Vasanthamani S. Chemical Composition of essential oils of *Hyptis suaveolens* (L.) Poit and its *in vitro* anticancer activity. *Journal of Pharmacy Research.* 2017;1(5):410-13.
  36. Bayala B, Nadembega C, Guenné S, Buñay J, Zohoncon TM, Djigma FW *et al.* Chemical composition, antioxidant and cytotoxic activities of *Hyptis suaveolens* (L.) Poit. essential oil on prostate and cervical cancers cells. *Pakistan Journal of Biological Sciences.* 2020; 23(9):1184-92. <https://doi.org/10.3923/pjbs.2020.1184.1192>

37. Deng Y. Bioactive constituents of two medicinal plants from Indonesia. Pharmacy Program [Dissertation]. Ohio: Ohio State University; 2010.
38. Costa-Lotufo, Araújo L, Lima E, Moraes M, Pessoa E, Silveira C *et al.* 2004. Antiproliferative effects of abietane diterpenoids isolated from *Hyptis martiusii* Benth. (Labiatae). *Die Pharmazie*. 2004;59:78-79.
39. Tabata K, Kim M, Makino M, Satoh M, Satoh Y, Suzuki T. Phenolic diterpen derived from *Hyptis incana* induce apoptosis G2/M arrest of neuroblastoma cells. *Anticancer Research*. 2012;32:4781-90.
40. Suzery M, Cahyono B. Evaluation of cytotoxicity effect of *Hyptis pectinata* Poit (Lamiaceae) extracts using BSLT and MTT methods. *J Sains dan Mat*. 2014;22:84-88.

§§§