



RESEARCH ARTICLE

Physiological responses of seven varieties of soybean [*Glycine max* (L.) Merr.] to salt stress

Naila Khalid* & Ahsan Saeed

Institute of Botany, Bahauddin Zakariya University, Multan 60000, Pakistan

*Email: nailakhalid673@yahoo.com

 OPEN ACCESS

ARTICLE HISTORY

Received: 09 July 2022
Accepted: 03 November 2022
Available online
Version 1.0 : 30 December 2022
Version 2.0 : 01 January 2023



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS etc. See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Khalid N, Saeed A. Physiological responses of seven varieties of soybean [*Glycine max* (L.) Merr.] to salt stress. Plant Science Today. 2023; 10(1): 199–209.
<https://doi.org/10.14719/pst.1986>

Abstract

In agriculture, salinity is one of the most significant abiotic stress that plants confront and harms agricultural productivity, physiological growth and development processes. In the present study, there were 7 different varieties of soybean (Ajmeri, William-82, D.A, PSC-60, Rawal-1, NARC-1 and NARC-2) were tested under NaCl concentration level (0 mM and 150 mM) to determine their physiological performance under control and experimental conditions. The present investigation aimed to select salt tolerant varieties. Under salt stress, different varieties have differ significantly in the biological yield, chlorophyll contents, antioxidant activity and ionic concentrations. The results showed that among the seven varieties evaluated NARC-1 and NARC-2 were producing higher biological yield and antioxidant activity than others under 150 mM NaCl. NARC-1 and NARC-2 under 150 mM NaCl concentration produced significantly higher biomass in comparison with other varieties and similarly enhance the antioxidant activity by decreasing the catalase activity. The relative water content (RWC) of plants was measured 15, 30, 45 and 60 days after the treatment was applied, as well as at harvest time, along with the grain yield and characters related to yield. The 7 different soybean varieties tested showed significant differences in grain yield and yield-associated characters when exposed to salinity. The salinity had a greater impact on Ajmeri and William than on NARC-1 and NARC-2. Under salt stress, the grain yield of the NARC-1 and NARC-2 varieties was 70% and 65% respectively, while the yields of the Ajmeri and william-82 varieties were 41% and 38% respectively. The salinity-induced decrease in grain yield was traced to fewer pods per plant, fewer seeds per pod and a lighter weight per 100 grains. However, the number of pods per plant was most affected compared to the other characters. It was also observed that Na⁺ ion concentrations were elevated in the shoot under salt stress in all varieties. However, NARC-1 and NARC-2 showed low salt concentration in shoot as compared to other varieties. SDS-PAGE revealed significant variations in the protein profile of seedling soybean varieties. NARC-1 and NARC-2 have shown a unique banding pattern under salt stress with a molecular weight of 60 and 130 kDa. The results indicate that salinity (NaCl) triggered an antioxidant response in tolerant varieties (NARC-1 and NARC-2) of *Glycine max* (L.). This study suggested that both varieties have more capability and appropriate survival under salt stress as compared to other varieties.

Keywords

Antioxidant activity, biological yield, chlorophyll constituents, *Glycine max*, salinity

Introduction

Legume *Glycine max* (L.) Merr. is widely regarded as one of the most significant crop on the earth (1). As a source of food, fuel and energy, soybean is one of the world's most important crop (2) and is recognized as the fourth most important yield producing crop in the world (3). Protein, hormones, phospholipids and antioxidants are all found in abundance in this plant (4). Tocopherols, sugars, fatty acids and organic acids, as well as sterols and volatile chemicals, are abundant in soybeans (5). Dicot crops have been classified on the basis of variable salt tolerance level. However, soybean is considered as "salt sensitive crop" because of their ability to thrive under adverse conditions (6).

Biotic and abiotic stresses exert a negative impact on yield potentials of the field-grown crops such stresses cause poor growth and development in crops. Salinity, as one of the challenge for irrigation farming, affects soil fertility and limit production (7). Salinity causes adverse effects on the growth and development of plants by interfering with vital metabolic processes (8). It is a limiting factor for plant biomass accumulation, antioxidant enzyme degradation, malondialdehyde (MDA) and hydrogen peroxide imbalance, proteins and peptides shortage and stress-related protein accumulation (9). The chlorophyll contents of plants also become reduced as salinity stress deteriorates photosynthetic pigments in a plant which directly affects the plant's production and growth (10). Additionally, salt stress damages proteins, down-regulation of photosynthesis-related proteins and reduces photosynthesis that leading to a reduction in plant growth (11). Different physio-biochemical changes occur in plants grown under salinity stress (12). Salinity is involved in the overproduction of reactive oxygen species (ROS) in plants which cause oxidative damage to proteins and plant cells (13). To minimize the effects of all these problems it is necessary to screen such varieties which are salinity tolerant and can grow on land affected by high salts concentrations (14). Salt tolerance in plants depends upon growth stages, type of plant, the nutrients provided to the plant and its environment. Various biochemical and physiological markers have been used for salinity tolerance along with changes in genetic makeup identified in soybean for its growth in saline conditions (15).

This work has been undertaken to relate the effect of salt on seven soybean varieties that are commonly grown in Pakistan. The study aims to screen out the varieties that performed well even under salinity and indicate a preference for breeding soybean varieties to develop salt tolerance.

Materials and Methods

There were 7 soybean varieties seeds obtained from the Ayub Agriculture Research Institute in Faisalabad in Pakistan: Ajmeri, William-82, D.A., PSC-60, Rawal-1 NARC-1 and NARC-2. Under natural day/night conditions (14h light/10h dark) at 28 °C in a greenhouse at Bahauddin Zakariya University Multan, Pakistan, the experiment was conducted.

Before conducting experiments, soybean seeds were treated with 5% sodium hypochlorite for 5 min to sterile the surface. Seeds were sowed in 21 cm plastic pots with a diameter of 15.5 cm and a depth of 20 cm, filled with thoroughly washed 4 kg sand. The muslin fabric was used to cover the drainage holes in the bottom of each pot. Five replicates were used for the experiment's arrangement, which was completely randomized. A complete solution of Hoagland's solution was used to irrigate all pots. In just one week, the seeds sprouted and were thinned to a maximum of eight plants per pot. The number of plants in each pot was reduced to five after the second week of growth. The two weeks old seedlings were subjected to salt treatment 0 mM and 150 mM NaCl up until they matured. Two plants for each sample were taken 15, 30, 45, 60 and 80 days for chemical and molecular study, some samples were maintained at -80 °C, while others were dried for additional phytochemical investigation.

Fresh and dry weight

Fresh biomass of plants was subjected to standard method (16) to separating root and shoot with sterilized blade after 2 weeks of salinity treatment and dry biomass was to be recorded by placing the plants in oven at 75 °C for 72 hrs (17) and measured by electric balance and the mean of 3 plants was recorded for accuracy of data.

Relative water content (RWC)

It was determined that the leaflets of the second fully expanded leaf from the top of the main stem had a relative water content (RWC) that could be measured at 15, 30, 45 and 60 days after imposition of salt stress. After the leaves had been harvested and brought to the laboratory, a measurement of the leaf's fresh weight was performed. Following an 8 hr soaking in distilled water, the leaf samples were blotted for surface drying and then the water saturated leaf weight was determined. After being dried in an oven at a temperature of 80 °C until they reached a weight that was constant, the leaf dry weight was determined. The following is the formula that was used to determine RWC (18):

$$\text{RWC (\%)} = (\text{fresh weight} - \text{dry weight} / \text{turgid weight} - \text{dry weight}) \times 100$$

Total proteins and free amino acids

The water-soluble protein quantification was done by described method (19) and the protein samples were prepared by blending 1 g leaf in sodium phosphate buffer (50 mM; pH 7.5). Protein contents were quantified as mg/g of seedling using standard Bovine Serum Albumin (BSA). The absorbance at 595 nm wavelength of protein samples was obtained through UV visible spectrophotometer (Hitachi U-2900, Tokyo Japan). The free amino acids were estimated using a described standard procedure (20). Total free amino acids were estimated by pyridine-ninhydrin as standard and absorbance were taken at 570 nm.

Chlorophyll estimation

Leaf photosynthetic pigments were evaluated by using described acetone method (21). Plant leaves were blended and mixed in 80% acetone and optical densities (OD) were

taken at 750, 663, 652, 645 and 470 nm using a spectrophotometer (Hitachi U-2900, Tokyo Japan).

Malondialdehyde Content (MDA) Estimation

Malondialdehyde contents were determined by described thiobarbituric acid method (22). The sample (0.5 g) was grinded in pre-chilled pestle and mortar on precooled phosphate buffer (pH 7.8). Centrifuge the sample up to 8000-13000 rotation per minute for 15 min at 4 °C and finally mixed with 0.5% thiobarbituric acid along with 5% trichloroacetate. The final mixtures were heated at 95 °C for 30 min placed water bath and cooled to room temperature. Finally, mixtures were centrifuged at 6000 × g for 15 mins and obtained supernatants were subjected to detection at a wavelength of 450, 532 and 600 nm. The MDA concentrations were calculated by following described formula:

$$\text{MDA } (\mu\text{mol/ml}) = 6.45 \times (D_{532} - D_{600}) - 0.56 \times D_{450}$$

H₂O₂ determination

H₂O₂ was observed by using a described protocol (23) 0.5 g of a leaf was blended in 5 ml of 0.1% (w/v) trichloroacetate. The paste was centrifuged at 10000 × g for 30 min. The obtaining supernatant was diluted with an equal volume of sodium phosphate buffer and finally dissolved in 1 ml of potassium iodide buffer. It was shaken well and recorded absorbance at 390 nm with a spectrophotometer.

Sodium and potassium ions Analysis

Ions analysis was performed by oven-dried roots and shoots at 72 °C for 24 hrs. Oven dry samples were ground and 0.2 g powder was digested with 0.1% Conc. HNO₃ at 115 °C for 4 hrs. The digested samples were subjected to a flame photometer to determine the sodium and potassium ions (24).

Dynamic study of catalase activity (CAT)

The catalase enzyme activity was evaluated through H₂O₂ decomposition at 240 nm (25). The samples were placed in

a precooled container filled with liquid nitrogen and 0.5 g of sample was blended in 8 ml sodium phosphate buffer. The samples were mixed with 0.1 ml of H₂O₂ and readings were recorded at 240 nm with a period of 0-30 seconds.

SDS-PAGE

The protein banding profile of seedlings was analyzed using SDS-PAGE as the standard described protocol (26). After electrophoresis, gels were placed in a solution of methanol/acetic acid/water solution (50:10:40) with 0.25% Coomassie brilliant blue R-250 for 30 minutes (27) and destained with the same solvent without Coomassie dye. The protein banding pattern of plant samples and protein marker (Bio-basic BG00363) were compared on SDS-PAGE to determine the molecular weight of protein.

Statistical analysis

Data were statistically examined through One Way Completely Randomized Analysis of Variance (1WCR ANOVA) by using COSTAT software (Cohort Software, Berkeley, California) (28).

Results and Discussion

Morphological and physiological processes were adversely affected by salt stress. Salinity reduces growth in soybean by interfering with physiological pathways, as well as identifying salt-sensitive, moderate and tolerant soybean types in the current study. The morphological and development characteristics of seven soybean varieties were compared under well water condition (Fig.1). The results revealed that some varieties (Ajmeri, William-82, D.A., PSC-60, Rawal-1) were found to be salt sensitive but NARC-1 and NARC-2 were showed salt-tolerant varieties. These results demonstrated that the wild plants retained a higher water status under salinity conditions than the more sensitive cultivars.

Seven varieties were found to have substantial differences in fresh and dry weight reduction (Table 1). Salt stress showed a significant drop in the fresh and dry plant




Fig. 1. Salt stress-related phenotypes of soybean varieties (Ajmeri, William-82, D.A., PSC-60, Rawal-1, NARC-1, and NARC-2) under salt stress (0 mM and 150 mM) for 7 to 15 days.

Table 1. Analysis of variance for biomass (g/plant) of Soybean varieties after salinity treatment.

Source of Variance	Df	Shoot f. wt. (g/plant)	Shoot d. wt. (g/plant)	Root f. wt. (g/plant)	Root d. wt. (g/plant)
Variety	6	5.029*	0.679***	0.776***	1.936***
Salinity	1	500.585***	8.447***	11.160***	7.541***
Variety×Salinity	6	2.832*	0.091*	0.068*	0.040*
Error	42	2.122	0.042	0.152	0.031

*, *** shows Significant level of 0.05 and 0.001 respectively

biomass of the salt-sensitive varieties Ajmeri and William-82. Two varieties (NARC-1 and NARC-2) revealed higher biomass as compared to the control and other varieties. Thus, 7 varieties were classified in response to salinity in the 3 sets, i.e., salt tolerant (NARC-1 and NARC-2), moderate salt-tolerant varieties (Rawal-1, D.A and PSC-60) and the salt-sensitive Ajmeri and William-82 (Fig. 2). Changes

Total free amino acid contents were reduced in all tested 7 varieties as compared to the control. The variation in the content of the amino acid under salinity of all soybean varieties is illustrated in Fig. 4. Mean of data was calculated by two-way completely randomized ANOVA showed a significant increase in the NARC-1 and NARC-2 ($p < 0.001$) as compared to the other varieties (Table. 2). The




Fig. 2. Fresh and dry biomass (g/plant) of seven soybean varieties after under salt stress.

plant dry weight have been reported like that of the plant fresh weight. Significant reduction in both shoot and root growth have been observed under saline conditions (29). Morphologically, the symptoms of salt stress injury were detained plant growth due to hindrance in the cell expansion (30).

Therefore, limited water absorption causes serious cellular growth and developmental consequences which lead to a reduction in stem and root growth (31). On the first day of salt treatment, the RWC differences between the control varieties of soybean and the salt-treated varieties were not statistically significant (Fig. 3). After 30 days of salt treatment and again at harvest (60 days), there was a significant difference in RWC between non-stressed and salt-treated varieties of soybean. According to the findings of our research, out of a total of 7 varieties, NARC-1 and NARC-2 have maintained better water relations when subjected to salt stress.

previous studies supported the above results that free amino acids contents were increased under salt stress (32). Plants that accumulate free amino acid content in the cell during salt stress have been suggested as a chief contribution to the osmotic adjustment (33), inhibit ROS production and stabilized the protein as well as cellular membranes integrity (34).

The total soluble proteins were estimated with Bradford's assay. It is indicated that NARC-1 and NARC-2 showed significantly higher protein concentrations as compared to other varieties (Table 2). The control had a greater protein quantity as compared to all tested varieties under salinity (Fig. 4). It is commonly known that the physiological status of plants was determined through the presence of the total soluble proteins. It was reported that salt stress stimulated signal transduction pathways that modulated the gene expression




Fig. 3. Leaf relative water content of soybean varieties under salt stress.




Fig. 4. Soluble proteins, free amino acids, MDA and H₂O₂ content (mg/g) of soybean varieties.

Table 2. Analysis of variance for biochemical attributes of Soybean varieties.

Source of Variance	Df	proteins (mg/g)	Amino acids (mg/g)	MDA (mg/g)	H ₂ O ₂ (mg/g)
Variety	6	366.479 ***	131209.65***	605.654***	0.805***
Salinity	1	2638.079***	1027320.5***	3344.785***	12.620***
Variety×Salinity	6	5.435**	5033.534***	15.136*	0.161***
Error	42	2.811	415.703	5.261	0.003

*, **, *** shows significant level at 0.05, 0.01 and 0.001 respectively

and relative proportion of protein (35). Under salt stress reduction in the protein, concentration was observed by

many authors (36). Another report stated that protein concentration was decreased in the 3 soybean cultivars

in the saline environment (37). The decline in the protein concentration due to the presence of NaCl causes a toxic effect on protein synthesis (38). The high salt concentration affects many physiological processes in plants like lower water potential and induces osmotic stress that leads to developing secondary oxidative stress which interferes with CO₂ assimilation and protein synthesis (8).

MDA and H₂O₂ contents were elevated in soybean grown under salt stress (Table 2). H₂O₂ and MDA contents have been significantly increased under salt stress in all tested varieties of soybean as compared to their control. William-82 and Ajmeri showed relatively higher levels of H₂O₂ and MDA contents than other varieties. The NARC-1 and NARC-2 showed significantly lower H₂O₂ and MDA compared to other varieties (Fig. 4). H₂O₂ has been accumulated in plants at the arrival of various biotic and abiotic stresses (39). Osmotic stresses accelerate the production of ROS such as H₂O₂, which develops harmful effects on subcellular compartments and metabolic processes by causing oxidative cellular destruction (40). ROS production is also involved in membrane destruction and increases MDA content in the cells (41, 42). A higher level of MDA is involved in the peroxidation of lipids in cell membranes and is considered an indicator during oxidative stress (43). MDA is considered a biomarker of biotic and abiotic stresses that are strongly correlated with the increasing level of MDA with electrolyte leakage and loss of membrane integrity.

The grain yield and characteristics associated with yield were adversely affected by NaCl salinity across all seven soybean varieties (Table 3). However, the salinity had a greater impact on the characters of PSC-60, D.A.,

was increased to 150 mM NaCl, NARC-1 and NARC-2 produced 72% pods per plant, whereas PSC-60, D.A., Rawal-1, Ajmeri and William-82 produced only 32%, 33%, 27%, 25%. When compared to other varieties, Ajmeri and William-82 were the ones whose number of seeds per pod was significantly impacted the most. NARC-1 and NARC-2 had a relative number of 72% and 66% of seeds per pod respectively, when compared to the control; all of the other varieties had a relative number of seeds per pod that was less than 50%. The relative weight of 100 seeds for NARC-1 and NARC-2 was 80% and 83% respectively, whereas for the other varieties, it was less than 50% with the exception of PSC-60, which was 53%. The salinity also had a significant impact on the decrease in the grain yield. When compared to the grain yield of NARC-1 and NARC-2 (12.65 and 12.32 g/plant), the grain yield of Ajmeri and William-82 control was noticeably higher (14.98 and 15.09 g/plant respectively). However, the decrease in grain yield caused by NaCl salinity was more noticeable in Ajmeri and William-82 than it was in NARC-1 and NARC-2. This was the case in both of these locations. The cumulative reduction in all of the yield associated characters was what was thought to be responsible for the lower grain yield that salinity caused. However, among the characteristics that were associated with yield, salinity had a significant impact on the number of pods produced by each plant in soybean varieties. When compared to seed setting and seed development, the findings suggested that salinity had the greatest impact on pod setting. When compared to those of Ajmeri and William-82, the characters of NARC-1 and NARC-2 have a lower level of affected yield and yield associated traits. Salinity stress limited the number of pods produced per plant, the number of seeds produced within

Table 3. Effect of salinity on grain yield and yield associated characters of seven soybean varieties

Varieties	No. of Pods/Plant		No. of seeds/Pod		100-Seeds Weight (g)		Grain Yield (g) / Plant	
	Control	Saline	Control	Saline	Control	Saline	Control	Saline
PSC-60	86.54	27.34 (32%)	1.91	1.37 (47%)	6.34	3.36 (53%)	13.34	6.65 (50%)
D.A	91.21	30.1 (33%)	2.84	1.35 (47%)	5.89	2.77 (47%)	11.72	5.38 (45%)
Rawal-1	70.3	19.21 (27%)	2.81	1.32 (46%)	6.64	3.24 (49%)	10.98	5.58 (51%)
Ajmeri	71.56	18.23 (25%)	2.64	1.19 (45%)	9.67	4.63 (48%)	14.98	6.15 (41%)
William-82	68.6	16.12 (23%)	2.7	1.21 (44%)	9.53	4.16 (44%)	15.09	5.69 (38%)
NARC-1	36.23	26.12 (72%)	2.98	2.16 (72%)	7.87	6.26 (80%)	12.65	8.84 (70%)
NARC-2	32.24	23.37 (72%)	3.21	2.12 (66%)	7.12	5.93 (83%)	12.32	8.12 (65%)

Rawal-1, Ajmeri and William-82 than it did on the characters of NARC-1 and NARC-2. PSC-60, D.A., Rawal-1, Ajmeri and William-82 produced a significantly higher number of pods per plant than NARC-1 and NARC-2 did when the conditions were controlled. However, when the salinity

each pod, the individual grain weight and the grain yield produced by each plant in soybean (44). Soil salinity caused significant damage to plants at any stage of their development, primarily as a result of the osmotic and ionic stress that it produced (45, 46).

Salt stress in soybean significantly influenced the K^+/Na^+ ratio (Table 4). However, under salinity more severe

salt stress, as consequence, many metabolic processes are inhibited that depend on the K^+ ions. It was concluded that

Table 4. Analysis of variance for Na^+ and K^+ ion (mg/g) of seven Soybean varieties.

Source of Variance	Df	Root K^+ ion	Shoot K^+ ion	Root Na^+ ion	Shoot Na^+ ion
Variety	6	807.792***	998.967***	1267.909***	14631.485***
Salinity	1	27542.237***	23776.11***	119200.85***	1083953.5***
Variety×Salinity	6	480.879***	377.706***	975.772***	32076.643***
Error	42	46.323	45.337	187.471	185.340

*** shows Significant level of 0.001.

significant rise in Na^+ content and a reduction in the K^+/Na^+ ratio. The NARC-1 and NARC-2 showed higher K^+ ion concentration in shoots rather than Na^+ ions and similar results have been observed as compared to all other varieties (Fig. 5). Plants are affected under salt stress in 3

K^+ ion observed as a higher concentration in the tolerant varieties than in the susceptible led to a reduction in the toxicity of sodium ions.

Several reports had been revealed that the biochemical action of green pigment (chlorophyll) is extraor-




Fig. 5. Ion analysis for seven varieties of soybean under salt stress.

Table 5. Analysis of variance in chlorophyll contents for seven varieties of Soybean.

Source of Variance	Df	Chlorophyll a (g/ml)/ plant	Chlorophyll b (g/ml)/ plant	Chlorophyll a/b (g/ml)/ plant	Total chlorophyll (g/ml)/ plant
Variety	6	1.260***	0.304***	3.382***	0.403*
Salinity	1	39.009***	30.386***	0.016 ns	24.698***
Variety×Salinity	6	0.929**	0.313***	1.887***	0.332*
Error	42	0.212	0.022	0.131	0.129

*, **, *** showing significant level of 0.05, 0.01 and 0.001 respectively.

different possible ways: It minimizes the water potential, disturb ions balance and also deteriorate ions homeostasis (47). Significant uptake of sodium ions causes severe growth retardation or leads to death of salt-sensitive of glycophytes species (48) i.e. barley (49) and soybean (50). For plants, both Na^+ and K^+ are competitive ions. Potassium ion plays a vital role in the various cellular processes proceeded in plants. If K^+ ions become outcompeted under

dinarily vulnerable to drastic effects of ions and cellular dehydration that altered chemical change by the destruction of chlorophyll directly or dissembling chemical action instrumentality (51). Our findings showed that salinity caused a significant loss of green pigment (Table 5) that's supported by the finding of (52).

The reduction in chlorophyll content; however, was less in soybean varieties NARC-1 and NARC-2 (Fig. 6), indi-




Fig. 6. Chlorophyll estimation for seven varieties of soybean under salt stress.

cating that these varieties resisted chlorophyll destruction and thus had the flexibility to stay up chemical action performance under saline conditions. The decline in chlorophyll and carotenoid under salt stress reported in this study agrees with several studies reported about glycohytes (53). The membrane deterioration is the major cause of the decline in chlorophyll contents under salt stress (54). It was reported that the generation of H_2O_2 and OH^- were considered the major destruction agents of chlorophyll contents and chloroplast ultra-structure (55). Salt stress also interferes with chlorophyll synthesis (56) and also developed instability in chloroplastic membrane and pigment complex protein. It was reported that salinity reduced the activity of 5-aminolevulinic acid dehydratase (ALA-D) which involves in porphyrin ring formation during chlorophyll synthesis (57).

The variation in antioxidant enzyme activity such as catalase (CAT) in soybean varieties during salt stress is presented in Table 6. The CAT activity was observed to be

Table 6. Analysis of variance for catalase activity of Soybean varieties

Source of Variance	Df	CAT (mM/gFW)
Variety	6	0.198***
Salinity	1	3.083***
Variety×Salinity	6	0.035***
Error	42	4.562

*** shows significant at 0.001 level.

significantly decreased in control as compared to salt-treated plants. Our data revealed that increasing level of CAT activities in NARC-1 and NARC-2 and moderately tolerant D.A and Rawal-1, PSC-60 than in the salt-sensitive William-82 and Ajmeri (Fig. 7) under salt stress. This result provides information about the antioxidant defense sys-

tem and suggested that NARC-1 and NARC-2 detoxify H_2O_2 and MDA cellular destruction.




Fig. 7. Catalase activity of seven soybean varieties under salt stress.

The SDS-PAGE showed different protein banding patterns of seven soybean varieties grown under salt stress. In control assays, all types displayed a comparable banding pattern between 10 kDa and 50 kDa (Fig. 8A, B). Salinity, on the other hand, resulted in the disappearance of the protein band (MW 10 kDa) in five kinds of Ajmeri. However, the salt-grown seedling has shown some variance. The 60 kDa and 130 kDa bands found on NARC-1 and NARC-2 were the only ones of their kind (Fig. 8B; Lane 3 and 4 respectively).

Gene expressions were influenced by salinity exposure via the synthesis of novel polypeptides and the loss and over-expression of proteins, respectively (58). The polypeptides with a molecular weight of 40, 34, 32, 29 and 14 kDa were expressed in the salt treat callus of *Mesembryanthemum crystallinum* (59). The proteins have a key role in salinity tolerance with molecular weights of 68, 52, 46, 43, 35, 33, 18 and 11 kDa (60). These proteins have been suggested that these peptides have a vital role in osmoprotectant or protecting cellular structures (61).




Fig. 8. SDS-PAGE of Soybean seedling under control and salt stress. **(A)** is showing the banding profile of Ajmeri, D.A, PSC-60, Rawal-1 and William-82 under control (Lane 1 to 5) and salt stress (Lane 6 to 10). **(B)** is protein banding pattern of NARC-1 and NARC-2, Lane 1 and 2 loaded with control and lane 3, 4 loaded with salt treated plant samples of NARC-1 and NARC-2 respectively. The headed arrows showing the disappearance and newly formed protein under salt stress. Lane M containing protein marker to compare the electrophoretic mobility of protein samples in the gel.

Conclusion

In conclusion, we confirmed the high salt tolerance associated with the relative grain production of the soybean varieties NARC-1 and NARC-2 in comparison to other soybean varieties evaluated. The high relative grain output of NARC-1 and NARC-2 was correlated with the least impacted number of pods per plant, number of seeds per pod and 100-seed weight by NaCl salinity compared to other cultivars examined in this study. This salt tolerance was achieved by NARC-1 and NARC-2 through increased antioxidant activity and decreased peroxidation activity. On the basis of our findings, we suggest that this inherited characteristic of the NARC-1 and NARC-2 soybean will be valuable in research initiatives aimed at creating more salt-tolerant germplasm as well as in genetic and physiological studies aimed at figuring out the mechanisms underlying increased crop yields on high-salinity soils. Further investigations have to be undertaken to recognize molecular markers or salinity-responsive genes to realize a stronger understanding of mechanisms underlying resistance in soybean varieties.

Authors contributions

NK carried out the experimental work and write-up improvement. AS made the drafting of original manuscript. The authors have read and agreed to the published version of the manuscript.

Compliance with ethical standards

Conflict of interest: No conflict of interest in this study.

Ethical issues: No animals/humans were used for studies that are the basis of this research.

References

1. Arshad M, Ali N, Ghafoor A. Character correlation and path coefficient in soybean *Glycine max* (L.) Merrill. *Pakistan Journal of Botany*. 2006;38(1):121.
2. Katerji N, Van Hoorn J, Hamdy A, Mastrorilli M. Salt tolerance of crops according to three classification methods and examination of some hypothesis about salt tolerance. *Agricultural Water Management*. 2001;47(1):1-8. [https://doi.org/10.1016/S0378-3774\(00\)00099-8](https://doi.org/10.1016/S0378-3774(00)00099-8)
3. Escobar JC, Lora ES, Venturini OJ, Yáñez EE, Castillo EF, Almanzan O. Biofuels: environment, technology and food security. *Renewable and sustainable energy reviews*. 2009;13(6-7):1275-87. <https://doi.org/10.1016/j.rser.2008.08.014>
4. Choi M-S, Rhee KC. Production and processing of soybeans and nutrition and safety of isoflavone and other soy products for human health. *Journal of medicinal food*. 2006;9(1):1-10. <https://doi.org/10.1089/jmf.2006.9.1>
5. Kumar V, Rani A, Dixit AK, Bhatnagar D, Chauhan G. Relative changes in tocopherols, isoflavones, total phenolic content and antioxidative activity in soybean seeds at different reproductive stages. *Journal of Agricultural and Food Chemistry*. 2009;57(7):2705-10. <https://doi.org/10.1021/jf803122a>
6. Waluyo SH, Lie TA, Mannetje Lt. Effect of phosphate on nodule primordia of soybean (*Glycine max* Merrill) in acid soils in rhizotron experiments. 2004. 5(2):37-44. <https://doi.org/10.21082/ijas.v5n2.2004.p37-44>
7. Turan S, Cornish K, Kumar S. Salinity tolerance in plants: breeding and genetic engineering. *Australian Journal of Crop Science*. 2012;6(9):1337-48.
8. Khan MN, Siddiqui MH, Mohammad F, Khan M, Naeem M. Salinity induced changes in growth, enzyme activities, photosynthesis, proline accumulation and yield in linseed genotypes. *World J Agric Sci*. 2007;3(5):685-95.
9. Nawaz K, Hussain K, Majeed A, Khan F, Afghan S, Ali K. Fatality of salt stress to plants: Morphological, physiological and biochemical aspects. *African Journal of Biotechnology*. 2010;9(34): 5475-80.

10. Netondo GW, Onyango JC, Beck E. Sorghum and salinity: II. Gas exchange and chlorophyll fluorescence of sorghum under salt stress. *Crop Science*. 2004;44(3):806-11. <https://doi.org/10.2135/cropsci2004.0806>
11. Jamil M, Lee KJ, Kim JM, Kim H-S, Rha ES. Salinity reduced growth PS2 photochemistry and chlorophyll content in radish. *Scientia Agricola*. 2007;64:111-18. <https://doi.org/10.1590/S0103-90162007000200002>
12. Chaves M, Flexas J, Pinheiro C. Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. *Annals of Botany*. 2009;103(4):551-60. <https://doi.org/10.1093/aob/mcn125>
13. Gill SS, Tuteja N. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology and Biochemistry*. 2010;48(12):909-30. <https://doi.org/10.1016/j.plaphy.2010.08.016>
14. Yamaguchi T, Blumwald E. Developing salt-tolerant crop plants: challenges and opportunities. *Trends in Plant Science*. 2005;10(12):615-20. <https://doi.org/10.1016/j.tplants.2005.10.002>
15. Ashraf M, Harris P. Potential biochemical indicators of salinity tolerance in plants. *Plant Science*. 2004;166(1):3-16. <https://doi.org/10.1016/j.plantsci.2003.10.024>
16. Liu H, Song J, Dong L, Wang D, Zhang S, Liu J. Physiological responses of three soybean species (*Glycine soja*, *G. gracilis* and *G. max* cv. Melrose) to salinity stress. *Journal of Plant Research*. 2017;130(4):723-33. <https://doi.org/10.1007/s10265-017-0929-1>
17. Bohm W. Methods of studying root systems. Berlin: Springer-Verlag. 1979.
18. González L, González-Vilar M. Determination of relative water content. *Handbook of Plant Ecophysiology Techniques*: Springer. 2001; p. 207-12. https://doi.org/10.1007/0-306-48057-3_14
19. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*. 1976;72(1-2):248-54. [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
20. Lee YP, Takahashi T. An improved colorimetric determination of amino acids with the use of ninhydrin. *Analytical Biochemistry*. 1966;14(1):71-77. [https://doi.org/10.1016/0003-2697\(66\)90057-1](https://doi.org/10.1016/0003-2697(66)90057-1)
21. Lichtenthaler HK, Wellburn AR. Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. Portland Press Ltd.; 1983. <https://doi.org/10.1042/bst0110591>
22. Hodges DM, DeLong JM, Forney CF, Prange RK. Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. *Planta*. 1999;207(4):604-11. <https://doi.org/10.1007/s004250050524>
23. Velikova V, Yordanov I, Edreva A. Oxidative stress and some antioxidant systems in acid rain-treated bean plants: protective role of exogenous polyamines. *Plant Science*. 2000;151(1):59-66. [https://doi.org/10.1016/S0168-9452\(99\)00197-1](https://doi.org/10.1016/S0168-9452(99)00197-1)
24. Wang B. Comparison of extractive methods of Na⁺, K⁺ in wheat leaves. *Plant Physiol Commun*. 1995;31:50-52.
25. Aebi H. [13] Catalase in vitro. *Methods in Enzymology*. 105: Elsevier. 1984; p. 121-26. [https://doi.org/10.1016/S0076-6879\(84\)05016-3](https://doi.org/10.1016/S0076-6879(84)05016-3)
26. Laemmli UK. Cleavage of structural proteins during the as of the head of bacteriophage T4. *Nature*. 1970;227(5259):680-85. <https://doi.org/10.1038/227680a0>
27. Sambrook J, Fritsch EF, Maniatis T. *Molecular cloning: a laboratory manual*: Cold Spring Harbor Laboratory Press. 1989.
28. Wallenstein S, Zucker CL, Fleiss JL. Some statistical methods in circulation research. *Circulation Research*. 1980;47(1):1-9. <https://doi.org/10.1161/01.RES.47.1.1>
29. Siddiqti S, Kumar S. Effect of salinisation and desalinisation and development of pea (*Pisum sativum* L.). *Indian J Plant Physiol*. 1985;28(2):151-56.
30. Bandoğlu E, Eyidoğan F, Yücel M, Avni Öktem H. Antioxidant ses of shoots and roots of lentil to NaCl-salinity stress. *Plant Growth Regulation*. 2004;42(1):69-77. <https://doi.org/10.1023/B:GROW.0000014891.35427.7b>
31. Cavalcanti FR, Lima JPMS, Ferreira-Silva SL, Viégas RA, Silveira JAG. Roots and leaves display contrasting oxidative response during salt stress and recovery in cowpea. *Journal of Plant Physiology*. 2007;164(5):591-600. <https://doi.org/10.1016/j.jplph.2006.03.004>
32. Panneerselvam R, Muthukumarasamy M, Rajan S. Amelioration of NaCl stress by triadimefon in soybean seedlings. *Biologia Plantarum*. 1998;41(1):133-37. <https://doi.org/10.1023/A:1001737221832>
33. Hare PD, Cress WA, Van Staden J. Dissecting the roles of osmolyte accumulation during stress. *Plant, Cell & Environment*. 1998;21(6):535-53. <https://doi.org/10.1046/j.1365-3040.1998.00309.x>
34. Keutgen AJ, Pawelzik E. Contribution of amino acids to strawberry fruit quality and their relevance as stress indicators under NaCl salinity. *Food Chemistry*. 2008;111(3):642-47. <https://doi.org/10.1016/j.foodchem.2008.04.032>
35. Kosova K, Prasil IT, Vitamvas P. Protein contribution to plant salinity response and tolerance acquisition. *International Journal of Molecular Sciences*. 2013;14(4):6757-89. <https://doi.org/10.3390/ijms14046757>
36. Mahboobeh R, Akbar EA. Effect of salinity on growth, chlorophyll, carbohydrate and protein contents of transgenic *Nicotiana glauca* over expressing P5CS gene. *Journal of Environmental Research and Management*. 2013;4:0163-70.
37. Ghassemi-Golezani K, Taifeh-Noori M, Oustan S, Moghaddam M, Rahmani SS. Physiological performance of soybean cultivars under salinity stress. *Journal of Plant Physiology and Breeding*. 2011;1(1):1-7. <https://doi.org/10.5772/14741>
38. Omar MS, Yousif DP, Al-Jibouri A-JM, Al-Rawi MS, Hameed MK. Effects of gamma rays and sodium chloride on growth and cellular constituents of sunflower (*Helianthus annuus* L.) callus cultures. *Journal of Islamic Academic of Science*. 1993;6(1):69-72.
39. Kaur H, Bhardwaj RD, Grewal SK. Mitigation of salinity-induced oxidative damage in wheat (*Triticum aestivum* L.) seedlings by exogenous application of phenolic acids. *Acta Physiologiae Plantarum*. 2017;39(10):1-15. <https://doi.org/10.1007/s11738-017-2521-7>
40. Siboza XI. Methyl jasmonate and salicylic acid enhance chilling tolerance in lemon (*Citrus limon*) fruit 2013. <https://doi.org/10.17660/ActaHortic.2012.928.53>
41. Yadav V, Singh H, Singh A, Hussain I, Singh N. Salicylic acid induced changes on some physiological parameters symptomatic for oxidative stress in maize (*Zea mays* L.) grown under cinnamic acid stress. *Russian Agricultural Sciences*. 2018;44(1):9-17. <https://doi.org/10.3103/S1068367418010202>
42. Gunes A, Inal A, Alpaslan M, Eraslan F, Bagci EG, Cicek N. Salicylic acid induced changes on some physiological parameters symptomatic for oxidative stress and mineral nutrition in maize (*Zea mays* L.) grown under salinity. *Journal of Plant Physiology*. 2007;164(6):728-36. <https://doi.org/10.1016/j.jplph.2005.12.009>
43. Jain M, Mathur G, Koul S, Sarin N. Ameliorative effects of proline on salt stress-induced lipid peroxidation in cell lines of groundnut (*Arachis hypogaea* L.). *Plant Cell Reports*. 2001;20(5):463-68. <https://doi.org/10.1007/s002990100353>
44. Ghassemi-Golezani K, Taifeh-Noori M, Oustan S, Moghaddam M. Response of soybean cultivars to salinity stress. *J Food Agric Environ*. 2009;7(2):401-04. <https://doi.org/10.15835/nsb224590>

45. Munns R. Comparative physiology of salt and water stress. *Plant, Cell & Environment*. 2002;25(2):239-50. <https://doi.org/10.1046/j.0016-8025.2001.00808.x>
46. Benlloch-González M, Fournier JM, Ramos J, Benlloch M. Strategies underlying salt tolerance in halophytes are present in *Cynara cardunculus*. *Plant Science*. 2005;168(3):653-59. <https://doi.org/10.1016/j.plantsci.2004.09.035>
47. Parida AK, Das AB. Salt tolerance and salinity effects on plants: a review. *Ecotoxicology and Environmental Safety*. 2005;60(3):324-49. <https://doi.org/10.1016/j.ecoenv.2004.06.010>
48. Maathuis FJ, Amtmann A. K⁺ nutrition and Na⁺ toxicity: the basis of cellular K⁺/Na⁺ ratios. *Annals of Botany*. 1999;84(2):123-33. <https://doi.org/10.1006/anbo.1999.0912>
49. Flowers T, Hajibagheri M. Salinity tolerance in *Hordeum vulgare*: ion concentrations in root cells of cultivars differing in salt tolerance. *Plant and Soil*. 2001;231(1):1-9. <https://doi.org/10.1023/A:1010372213938>
50. Umezawa T, Shimizu K, Kato M, Ueda T. Enhancement of salt tolerance in soybean with NaCl pretreatment. *Physiologia Plantarum*. 2000;110(1):59-63. <https://doi.org/10.1034/j.1399-3054.2000.110108.x>
51. Zhang H, Ye Y-K, Wang S-H, Luo J-P, Tang J, Ma D-F. Hydrogen sulfide counteracts chlorophyll loss in sweetpotato seedling leaves and alleviates oxidative damage against osmotic stress. *Plant Growth Regulation*. 2009;58(3):243-50. <https://doi.org/10.1007/s10725-009-9372-1>
52. Ayala-Astorga GI, Alcaraz-Meléndez L. Salinity effects on protein content, lipid peroxidation, pigments and proline in *Paulownia imperialis* (Siebold & Zuccarini) and *Paulownia fortunei* (Seemann & Hemsley) grown *in vitro*. *Electronic Journal of Biotechnology*. 2010;13(5):13-14. <https://doi.org/10.2225/vol13-issue5-fulltext-13>
53. Mitra A, Banerjee K. Pigments of *Heritiera fomes* seedlings under different salinity conditions: perspective sea level rise. *Mesopotamian Journal of Marine Sciences*. 2010;25(1):1-10.
54. Djanaguiraman M, Ramadass R. Effect of salinity on chlorophyll content of rice genotypes. *Agric Sci Digest*. 2004;24(3):178-81.
55. Yamane K, Rahman MS, Kawasaki M, Taniguchi M, Miyake H. Pretreatment with antioxidants decreases the effects of salt stress on chloroplast ultrastructure in rice leaf segments (*Oryza sativa* L.). *Plant production science*. 2004;7(3):292-300. <https://doi.org/10.1626/pp.s.7.292>
56. Garcia-Sanchez F, Jifon JL, Carvajal M, Syvertsen JP. Gas exchange, chlorophyll and nutrient contents in relation to Na⁺ and Cl⁻ accumulation in 'Sunburst' mandarin grafted on different rootstocks. *Plant Science*. 2002;162(5):705-12. [https://doi.org/10.1016/S0168-9452\(02\)00010-9](https://doi.org/10.1016/S0168-9452(02)00010-9)
57. Aldesuquy H, Baka Z, Mickky B. Kinetin and spermine mediated induction of salt tolerance in wheat plants: Leaf area, photosynthesis and chloroplast ultrastructure of flag leaf at ear emergence. *Egyptian Journal of Basic and Applied Sciences*. 2014;1(2):77-87. <https://doi.org/10.1016/j.ejbas.2014.03.002>
58. Patharkar OR, Cushman JC. A stress-induced calcium-dependent protein kinase from *Mesembryanthemum crystallinum* phosphorylates a two-component pseudo-response regulator. *The Plant Journal*. 2000;24(5):679-91. <https://doi.org/10.1046/j.1365-313x.2000.00912.x>
59. Yen HE, Zhang D, Lin JH, Edwards GE, Ku MS. Salt-induced changes in protein composition in light-grown callus of *Mesembryanthemum crystallinum*. *Physiologia Plantarum*. 1997;101(3):526-32. <https://doi.org/10.1034/j.1399-3054.1997.1010311.x>
60. Barakat H. Interactive effects of salinity and certain vitamins on gene expression and cell division. *International Journal of Agriculture Biology*. 2003;3:219-25.
61. Close TJ, Lammers PJ. An osmotic stress protein of cyanobacteria is immunologically related to plant dehydrins. *Plant Physiology*. 1993;101(3):773-79. <https://doi.org/10.1104/pp.101.3.773>

§§§