



RESEARCH COMMUNICATION

# *Phlogacanthus cornutus*: chemical profiles and antioxidant effects

Duy Hoang Truong<sup>1\*</sup>, Thao Nguyen Luu<sup>1</sup>, Nguyen Tuong An Huynh<sup>2</sup>, Ngoc Hung Ngo<sup>3</sup>, Van Son Le<sup>4</sup>, Quoc Hung Nguyen<sup>5</sup> & Thanh Tho Le<sup>5</sup>

<sup>1</sup>Institute of Biotechnology and Food Technology, Industrial University of Ho Chi Minh City, Ho Chi Minh City 700000, Vietnam

<sup>2</sup>Institute of International and Postgraduate Education, Industrial University of Ho Chi Minh City, Ho Chi Minh City 700000, Vietnam

<sup>3</sup>Faculty of Fundamental Science, Industrial University of Ho Chi Minh City, Ho Chi Minh City 700000, Vietnam

<sup>4</sup>Binh Chau-Phuoc Buu Nature Reserve, Ba Ria-Vung Tau Province 78000, Vietnam

<sup>5</sup>Center of Analytical Services and Experimentation HCMC, Ho Chi Minh City 700000, Vietnam

\*Email: [truonghoangduy@iuh.edu.vn](mailto:truonghoangduy@iuh.edu.vn)



## ARTICLE HISTORY

Received: 15 August 2022

Accepted: 29 October 2022

Available online

Version 1.0 : 30 December 2022



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS etc. See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Truong D H, Luu T N, Huynh N T A, Ngo N H, Le V S, Nguyen Q H, Le T T. *Phlogacanthus cornutus*: chemical profiles and antioxidant effects. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.2063>

## Abstract

*Phlogacanthus cornutus* is a rare species and the chemical profiles and the bioactivities of this plant are unknown. In present study, the chemical components of the acetone extract as well as the antioxidant activity of acetone extract and its fractions such as n-hexane, chloroform and ethyl acetate of *P. cornutus* were firstly reported. A total of 33 constituents were identify in the acetone extract of this plant using Gas Chromatography/Mass Spectrometry assay, in which trans-cinnamic acid (21.26%), neophytadiene (6.36%), linolenic acid (5.86%), dihydroagathic acid (5.71%), n-hexadecanoic acid (5.53%), phytol (4.14%) and cis-cinnamic acid (3.23%) were the major compounds. The acetone extract and its fractions such as n-hexane, chloroform and ethyl acetate of *P. cornutus* showed DPPH radical scavenging activity with IC<sub>50</sub> value of 234.31, 185.95, 758.65 and 458.52 µg/mL respectively.

## Keywords

*Phlogacanthus cornutus*, GC/MS, chemical components, antioxidant activities

## Introduction


The genus *Phlogacanthus* is belonging to the family Acanthaceae with around 49 species. This genus is widely found in tropical regions of Asian countries, including Bhutan, Bangladesh, India, China, Myanmar, Indonesia and Vietnam (1-3). Previous studies provided that the various solvent extracts of the different *Phlogacanthus* species possessed many biological effects, including anti-inflammatory, antibacterial, antioxidative and cytotoxic activities (4, 5). About 7 *Phlogacanthus* species have been recorded for the flora of Vietnam such as *P. annamensis*, *P. geoffrayi*, *P. colaniae*, *P. pubiflorus*, *P. pyramygdalis*, *P. turgidus* and *P. cornutus* (1-3, 6).

*Phlogacanthus cornutus* Benoist is an endemic species of Vietnam. It was described for the first time by Benoist in 1927 of which its specimens were collected from Dinh mountain, Ba Ria-Vung Tau province, Vietnam (7). To date, *P. cornutus* is a rare species and the chemical compositions and bioactivities of this species are still unknown. Herein, we firstly provide the chemical profiles of the acetone extract, as well as the antioxidant effects of the acetone extract and its fractions, including-hexane, chloroform and ethyl acetate of *P. cornutus* grown in Binh Chau-Phuoc Buu Nature Reserve, Ba Ria-Vung Tau province, Vietnam.

## Materials and Methods

### Plant material

The leaves of *P. cornutus* were collected from Binh Chau-Phuoc Buu Nature Reserve, Ba Ria-Vung Tau province, Vietnam, location about 10°37'01.5"N 107°31'19.2"E (Fig. 1).



**Fig. 1.** *Phlogacanthus cornutus*. A. The species in habitat, B. Flower. Photos: Van Son Le.

The scientific name of this species was identified by Van Son Le, a botanist from Binh Chau-Phuoc Buu Nature Reserve. The vouchered specimen (BC 396) are deposited at the Herbarium of Binh Chau-Phuoc Buu Nature Reserve.

### Extraction

The plant sample was dried in an oven at 50 °C until constant weight and ground into fine powder using an electric blender. An amount of 100 g of powder was macerated in 500 ml of 99% acetone (Thermo Fisher Scientific, USA) at room temperature for 3 days. The extract was collected and filtered through Whatman filter paper No. 4. The solid material was further soaked in 300 ml and 200 ml sequentially of the same solvent and filtered to collect the extracts. All the three acetone extracts were then combined and condensed on an a rotary evaporator at 40 °C to remove the solvent (8).

### Fractionation of acetone extract

30 ml of distilled water was used to dissolve 3 g of the acetone extract. The suspension was subsequently mixed with 30 ml of n-Hexane (Thermo Fisher Scientific, USA), shaken and allowed to stand until layers were formed. The n-hexane layer (upper) was collected. This procedure was repeated 2 more times to obtain 90 ml of hexane extract. The extract was subjected to a rotary evaporator at 40 °C to remove the solvent. The lower layer was then used to collect the ethyl acetate and chloroform (Thermo Fisher Scientific, USA) fractions using the same procedure (8).

### GC/MS based chemical profiling of the acetone extract

The acetone extract of the studied species was subjected to chemical composition analysis on TRACETM 1310 Gas Chromatograph (Thermo Fisher Scientific Inc., Waltham, MA, USA) equipped with ISQ 7000 single quadrupole mass

spectrometer and DB-5MS column (30 m × 0.25 mm × 0.25 μm). Samples were injected into the GC system thermostated at 250 °C and at a split ratio of 30:1 at the flow rate 36 mL/min with splitless time of 1 min. Helium was used as the carrier gas at a flow rate of 1.2 mL/min. The transfer line temperature was set at 250 °C. The oven temperature was initially set to be hold at 80 °C for 5 min and was then increased to 280 °C at a rate of 20 °C/min and held at this temperature for 10 min. The electron impact ionization was set as 70 eV and the filament source temperature was set at 250 °C. The acquisitions scan mass range of MS was 29-650 m/z with the scanning frequency of 2 scans/sec. The chemical components of the acetone extracts obtained from *P. cornutus* were identified based on the comparison between their mass spectra with NIST 2017 library.

### DPPH Free Radical Scavenging Assay

The antioxidant effects of the studied extracts were investigated using DPPH (2,2-Diphenyl-1-picrylhydrazyl) radical scavenging assay (8). First, 2700 μl of DPPH (Thermo Fisher Scientific, USA) solution (0.1 mM) was added to 100 μl of sample. The mixture was then gently shaken and left in the darkness for 30 min at 37 °C. The mixture was measured for absorbance at a wavelength of 517 nm. DPPH Free Radical Scavenging effect of the studied samples was determined using the following formula (I). The radical scavenging activity of samples was expressed as the inhibition concentration (IC<sub>50</sub>), which is defined as the concentration of sample being able to reduce the initial DPPH concentration by 50%.

$$\% \text{ Antioxidant activity} = \frac{[(\text{OD}_{\text{control}} - \text{OD}_{\text{sample}})]}{\text{OD}_{\text{control}}} \times 100 (\text{I})$$

In which OD<sub>control</sub> = The optical density of DPPH in acetone and OD<sub>sample</sub> = The optical density of DPPH mixed with the extract.

## Results and Discussion

### Chemical compositions of acetone extract of *P. cornutus*

The chemical profiles of the acetone extract isolated from *P. cornutus* leaves were shown in the Table 1 and Fig. 2. A total of 33 components were identified from the studied extract, of which trans-cinnamic acid (21.26%), neophytadiene (6.36%), linolenic acid (5.86%), dihydrogathic acid (5.71%), n-hexadecanoic acid (5.53%), phytol (4.14%)

**Table 1.** Chemical profiles of acetone extract obtained from *Phlogacanthus cornutus*.

Retention time (min)	Compounds	% Composition
4.61	Hexanoic acid	0.58
4.49	Glycerin	0.36
5.6	trans-2-Hexenoic acid	0.38
5.91	Benzyl alcohol	0.11
7.26	Phenylethyl Alcohol	0.10
8.11	Cinnamaldehyde	1.05
8.15	Benzoic acid	0.88
8.73	Coumaran	0.95
9.24	7a-Methyl-1,2,3,6,7,7a-hexahydro-5H-inden-5-one	0.29
9.73	4-Hydroxy-2,6-dimethylbenzotrile	0.79
10.08	cis-Cinnamic acid	3.23
10.68	trans-Cinnamic acid	21.62
10.75	Bicyclo[4.2.0]octa-1,3,5-triene-7-carboxylic acid	0.57
11.24	2(4H)-Benzofuranone, 5,6,7,7a-tetrahydro-4,4,7a-trimethyl-, (R)-	0.22
11.94	2,6-Dimethoxyhydroquinone	0.57
12.26	3-Oxo-7,8-dihydro- $\alpha$ -ionol	0.33
12.71	6-Hydroxy-4,4,7a-trimethyl-5,6,7,7a-tetrahydrobenzofuran-2(4H)-one	0.34
12.92	Neophytadiene	6.36
12.95	Hexahydrofarnesyl acetone	0.41
13.04	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	1.02
13.59	n-Hexadecanoic acid	5.53
14.31	Phytol	4.14
14.44	cis-9,cis-12-Octadecadienoic acid	1.98

14.48	Linolenic acid	5.86
14.56	Stearic acid	0.47
18.38	Dihydrogathic acid	5.71
18.68	Unknown compound	23.86
22.69	dl- $\alpha$ -Tocopherol	0.12
24.66	Campesterol	0.13
25.2	Stigmasterol	1.00
26.2	Sitosterol	1.33
27.07	Lup-20(29)-en-3-one	0.31
27.55	Lup-20(29)-en-3-ol, (3 $\beta$ )-	3.74
<b>Total</b>		94.32%

and cis-cinnamic acid (3.23%) were the major compounds. Notably, there was an unknown compound (at retention time 18.68 min) found in the acetone extract of *P. cornutus* leaves. This component possessed the highest percentage (23.86%) in the studied extract. Several fragments of this compound have been recorded of which the parent mass possessed m/z as 320 (Fig. 3). Furthermore, the mass spectra of this compound did not match with any compound included in NIST 2017 library in spite of the fact that NIST 2017 library, the most innovative library, was used in this study. This compound, thus, should be investigated further to determine whether they are new compounds or not.

The chemical components of other *Phlogacanthus* plants have been reported by previous studies. For instance, turgidol, a new compound, together with 5 known triterpenoids such as lupenone, lupeol, betulin, taraxer and betulinic acid were obtained from the chloroform extract of *Phlogacanthus turgidus* (9). Similarly, 6 chemical components of *P. turgidus* was identified in the ethyl acetate extract, including one derivative of phenylethanoid glycosides (martynoside), one lignan ((+)-syringaresinol), 2 steroids (bsitosterol and daucosterol) and two nor-isoprenoids ((+)-dehydrovomifoliol and (3S,5R,6R,7E,9S)-megastiman-7-ene-3,5,6,9-tetrol) (10). Studies also demonstrated that *Phlogacanthus pulcherrimus* had the content of nutrition (protein, fat, fiber and moisture) and mineral (calcium, sodium, potassium, magnesium, chromium and iron) (11). Furthermore, *P. pubinervius* and *P. Jenkinsii* have been reported to possess some phytochemical contents, including macro nutrients (nuphosphorus, sodium and potassium), micro nutrients (iron, copper, manganese), total free amino acid, total sugar, total free phenol, bound phenol and tannin (12). In addition, the chemical constituents of methanol extracts obtained from 3 *Phlogacanthus* species such as *P. pubinervius*, *P. thyriflorus* and *P. curviflorus* were investigated by using GC/MS assay. Accordingly, the *P. pubinervius* and *P. curviflorus* extracts were found to be rich in squalene, n-hexadecanoic acid and phytol while 2-ethyhexane, n-hexadecanoic acid




Fig. 2. Gas chromatogram of acetone extract from *Phlogacanthus cornutus*.




Fig. 3. Chromatogram of unknown compound at retention time 18.68 min.

and octadeca-9,12,15-trienoic acid were the major compounds in the *P. thrysiflorus* extract (13).

#### **Antioxidant effects of acetone extract and its fractions of *P. cornutus***

The antioxidant effects of the acetone extract and its fractions of *P. cornutus*, including ethyl acetate, hexane and

chloroform were determined by using 1,1-diphenyl-2-picrylhydrazol (Fig. 4). Accordingly, the hexane fraction showed the highest antioxidant activity with  $IC_{50}$  value of 185.95  $\mu\text{g/mL}$ , followed by acetone extract (234.31  $\mu\text{g/mL}$ ), ethyl acetate fraction (458.52  $\mu\text{g/mL}$ ), and chloroform fraction (758.65  $\mu\text{g/mL}$ ). Meanwhile, the positive control




Fig. 4. Radical scavenging activity of acetone extract and its fractions of *P. cornutus*.

(ascorbic acid) displayed activity against DPPH radical-scavenging with  $IC_{50}$  value of 1.84 µg/mL.

The antioxidant activities of other *Phlogacanthus* species were also reported by previous studies. For instance, the methanol extract of *P. pubinervius* and *P. jenkinsii* showed the DPPH radical scavenging activity with 93.96% and 357.72% respectively while the percentages of radical scavenging activity of chloroform extract were 20.14% and 32.89% respectively (12). In another report, Kripasana and Xavier provided the DPPH radical scavenging activities of the methanol extracts of three *Phlogacanthus* species such as *P. pubinervius*, *P. thyriflorus*, *P. curviflorus*. Accordingly, the *P. pubinervius* extract had an inhibitory effect on 57.93% radical DPPH-scavenging at concentration of 150 µg/mL whereas the extracts of *P. thyriflorus* and *P. curviflorus* were found to be effective against 92.94% and 94.20% radical DPPH at concentration of 100 µg/mL (13). Previous study also showed the DPPH radical scavenging activities of 4 extracts of *Phlogacanthus pulcherrimus*, including aqueous, dichloromethane, n-butanol, ethyl acetate extracts. As a result, the ethyl acetate extract possessed the highest antioxidant effect with  $IC_{50}$  values of 730.28 µg/mL, followed by n-butanol extract (1422.93 µg/mL), aqueous extract (>2000 µg/mL) and dichloromethane extract (>2000 µg/mL) (14).

## Conclusion

The present study identified 33 compounds in the acetone extract of *P. cornutus* leaves in which trans-cinnamic acid, neophytadiene, linolenic acid, dihydrogathic acid, n-

hexadecanoic acid, phytol and cis-cinnamic acid were the major compounds. The acetone extract and its fractions such as n-hexane, chloroform and ethyl acetate of *P. cornutus* showed DPPH radical scavenging activity with  $IC_{50}$  value of 234.31, 185.95, 758.65 and 458.52 µg/mL respectively. The results of this study will provide the evidence for future applications of *P. cornutus* in medicine and other related fields.

## Acknowledgements

This research is supported by Industrial University of Ho Chi Minh City, Vietnam. The authors would like to thank MSc. Nguyen Thi Hong Van, officials of Institute of Biotechnology and Food Technology, Industrial University of Ho Chi Minh City for her collaboration.

## Authors contributions

DHT designed this study. The samples were collected by VSL, a staff of Binh Chau-Phuoc Buu Nature Reserve. All authors performed experiments, handled the research data and data analysis. Duy Hoang Truong drafted the manuscript and resolved all the queries of reviewers.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** None

## References

1. Pham H. Araceae, in: Pham-hoang, H. (Editor.) *Cây cỏ Việt Nam: An Illustrated Flora of Vietnam*, vol. 3. Youth Publishing House, Ho Chi Minh City. 2000.
2. Scotland RW. Pollen morphology of Andrographideae (Acanthaceae). *Rev Palaeobot Palynol.* 1992 Jul;72(3-4):229-43. [https://doi.org/10.1016/0034-6667\(92\)90028-F](https://doi.org/10.1016/0034-6667(92)90028-F)
3. Scotland RW, Vollesen K. Classification of Acanthaceae. *Kew Bull.* 2000;55(3):513. <https://doi.org/10.2307/4118776>
4. Chakravarty S, Kalita JC. Antihyperglycaemic effect of flower of *Phlogacanthus thyrsoiflorus* Nees on streptozotocin induced diabetic mice. *Asian Pac J Trop Biomed.* 2012 Jan;2(3):S1357-61. [https://doi.org/10.1016/S2221-1691\(12\)60416-X](https://doi.org/10.1016/S2221-1691(12)60416-X)
5. Mukherjee A, Chaliha M, Das S. Study of analgesic activity of ethanol extract of *Phlogacanthus thyrsoiflorus* on experimental animal models. *Bangladesh J Pharmacol.* 2009 Apr;4(2). <https://doi.org/10.3329/bjpv.v4i2.3808>
6. Hai D Van, Hoan DT, Choudhary RK. *Phlogacanthus geoffrayi* Benoist (Acanthaceae): A New Record to the Flora of Vietnam. *VNU J Sci Nat Sci Technol.* 2020 Jun;36(2). <https://doi.org/10.25073/2588-1140/vnunst.4962>
7. Benoist R. Acanthacées nouvelles asiatiques. *Bull la Soc Bot Fr.* 1927;74(5):907-12. <https://doi.org/10.1080/00378941.1927.10837070>
8. Le HT, Luu TN, Nguyen HMT, Nguyen DH, Le PTQ, Trinh NN et al. Antibacterial, antioxidant and cytotoxic activities of different fractions of acetone extract from flowers of *Dipterocarpus intricatus* Dyer (Dipterocarpaceae). *Plant Sci Today.* 2021 Apr;8(2):273-77. <https://doi.org/10.14719/pst.2021.8.2.1086>
9. Tri MD, Phat NT, Trung NT, Phan CTD, Minh PN, Chi MT et al. A new 26-norlanostane from *Phlogacanthus turgidus* growing in Vietnam. *J Asian Nat Prod Res.* 2022;24(2):196-202. <https://doi.org/10.1080/10286020.2021.1913125>
10. Phat NT, Chi MT, Minh PN, Hien DC, Dinh Tri M. Chemical constituents of the whole plant of *Phlogacanthus turgidus* (Fua ex Hook. f.) Lindau. *Sci Technol Dev J - Nat Sci.* 2021 Jun;5(3):first. <https://doi.org/10.32508/stdjns.v5i3.1071>
11. Jongrungruangchok S, Wongwiwatthananutit S. Evaluation of nutrient and mineral content of the leaves of *Phlogacanthus pulcherrimus* cultivated in Thailand. 2014.
12. Laitonjam W, Yumnam R, Asem S, Wangkheirakpam S. Evaluative and comparative study of biochemical, trace elements and antioxidant activity of *Phlogacanthus pubinervius* T. Anderson and *Phlogacanthus jenkinsii* C. B. Clarke leaves. *Indian J Nat Prod Resour.* 2013;4(1):67-72.
13. Kripasana K, Xavier J. Phytochemical analysis and antioxidant activity of leaf extracts of some selected plants of the family Acanthaceae. *Plant Sci Today.* 2020 May;7(2):264-74. <https://doi.org/10.14719/PST.2020.7.2.717>
14. Poeaim S, Lordkhem P, Charoenying P, Laipasu P. Evaluation of antioxidant, cytotoxic activities and total phenolic content from leaf extracts of *Phlogacanthus pulcherrimus*. *Int J Agric Technol.* 2016;12(1):1657-67.

§§§