



RESEARCH ARTICLE

Planting date and genotype effect on morpho-agronomic traits of Burkina Faso sweet grain sorghum

Wendmanegda Hermann Tondé¹, Nerbéwendé Sawadogo^{1*}, Josiane Tiendrébéogo², Oumar Boro³, Pingawindé Sawadogo⁴, Mariam Kiébré¹, Kouka Fidèle Tiendrébéogo², Iriisa Yaméogo⁵ & Mahamadou Sawadogo¹

¹Equipe Génétique et Amélioration des Plantes (EGAP), Université Joseph KI-ZERBO, Ouagadougou, 03 BP 7021, Burkina Faso

²Institut Supérieur du Développement Durable, Université de Fada N'Gourma, Fada N'Gourma, BP 54, Burkina Faso

³West Africa Centre for Crop Improvement (WACCI), University of Ghana, Accra, PMB 30 Legon, Ghana

⁴Centre Universitaire de Tenkodogo, Université Thomas SANKARA, Saaba, 04 BP 8938, Burkina Faso

⁵Département d'Anglais, Université Joseph KI-ZERBO, Ouagadougou, 03 BP 7021, Burkina Faso

*Email: nerbewende.sawadogo@ujkz.bf



ARTICLE HISTORY

Received: 29 October 2022

Accepted: 05 February 2023

Available online

Version 1.0 : 13 May 2023

Version 2.0 : 01 July 2023



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Tondé W H, Sawadogo N, Tiendrébéogo J, Boro O, Sawadogo P, Kiébré M, Tiendrébéogo K F, Yaméogo I, Sawadogo M, Planting date and genotype effect on morpho-agronomic traits of Burkina Faso sweet grain sorghum. Plant Science Today. 2023;10(3): 149–159. <https://doi.org/10.14719/pst.2208>

Abstract

Sweet grain sorghum is an under-exploited crop mainly grown around dwelling houses. Its production faces harsh environmental conditions. This study aims to assess sowing date effect on morpho-agronomical traits of sweet grain sorghum. Thus, 30 genotypes of sweet grain sorghum were assessed under 2 planting dates (June 26 and July 20) 24 days apart in a Randomized Complete Block Designs with 3 replications using 10 traits. The results showed a significant effect of sowing date on most of the traits, except internode length. All genotypes were sensitive to photoperiod variation by reducing their sowing-flowering cycle from 08 to 20 days, size and yield at the second planting date. Delayed sowing also resulted in a decrease in plant height (66.4 cm), 100 grain weight (8.3%), panicle weight (16.84%) and grain yield per plant (18.93%). The genotypes expressed a differential sensitivity to photoperiod variation with a mean coefficient of 0.59. Finally, a clustered flowering of all genotypes between September 11 and 27 was observed for both sowing dates. These results could be exploited by sweet sorghum breeding programme in the definition of the cropping calendar.

Keywords

Burkina Faso, climate, genetic diversity, photoperiodism, sweet sorghum

Introduction

Sorghum [*Sorghum bicolor* (L.) Moench] is integral to the survival of the population and livestock in Burkina Faso (1-3). It is cultivated in almost the entire territory over more than 1907651 ha, for a production around 1849595 tons during the 2019-2020 agricultural season (4).

Several types of sorghum, with varied but little-known potential, are grown and maintained by farmers. The management of this diversity by farmers allows for an evolutionary adjustment to a heterogeneous environment, but also to meet population needs (5). Among the types of sorghum, sweet grain sorghum, is sparsely valued by the country's breeding programme (3). One of the main characteristics of sweet grain sorghum is its high amount of carbohydrates such as sucrose, glucose and fructose content in the grains (3, 6). Despite its cultivation in the 4 agro-climatic zones in the country, sweet grain sorghum remains a marginal crop. It is not much used by the population and its cultivation is still practiced in small farming

land, especially in cage fields (7, 8). The grains of this sorghum are consumed directly in the doughy state and the marketing of panicles provides substantial income to farmers and traders (6, 7). Previous studies on this local food crop have focused mainly on farmers' knowledge of its cultivation (7, 8), its agro-morphological and genetic diversity (9-11), its response to mineral fertilization, the biochemical composition of its grains (6) and its genetic relationships with other types of sorghum (12-14). However, (7, 8) reported a gradual abandonment of sweet grain sorghum cultivation due to the intensification of economically profitable crops (cotton, maize), the development of formal seed programs supported by national policies (15, 16) and the contrasting Sahelian agro-climatic conditions such as erratic rainfall, gradual reduction of arable land and low fertilizer inputs (17-19).

The recurrence of early and late cycle droughts in West Africa has led to the improvement of very early varieties by agricultural extension programme as they seem to be better adapted to short wintering periods (20). However, most farmers remain attached to their traditional varieties with longer cycles than modern varieties (5, 21). Traditional varieties, despite their limited productivity, have high yield stability (18). Indeed, several previous studies have shown a very high sensitivity of most West African sorghums to photoperiod variation (22-24). This trait gives

present study aims to evaluate the effect of delayed sowing on the expression of morpho-agronomical traits and to determine the level of the photoperiod sensitivity of sweet grain sorghum genotypes.

Materials and Methods

Experimental site

The experimental study was conducted in the experimental station of the Rural Development Institute (IDR) in "Gampèla" during the 2019-2020 cropping season. "Gampèla" is a locality located about 18 km east of Ouagadougou with geographical coordinates 12°25'N and 1°12'W (31). The experimental site is characterized by a unimodal rainy season from June to October with an annual rainfall of between 600 and 900 mm (32). In 2019, the first rains were recorded in March and the cumulative rainfall recorded at the experimental station was 762.3 mm on 47 rainy days. Rainfall peaked in July with 321.3 mm in 13 rainy days and then decreased from September onwards (138.5 mm) and stopped in mid-October (112.5 mm) (33). The average monthly temperature fell from March (34.7 °C) to August (27.4 °C) and then increased to a stable value of 29 °C from September onwards (Fig. 1) (34).

During the trial conducted between June and October, the average monthly insolation decreased from June




Fig. 1. Decadal precipitation diagram (33) and monthly average temperature curve of the experimental area (34).

them an important evolutionary advantage due to the continental area, characterized by a high variation in rainfall (24-26). The photoperiod sensitivity of these cultivars ensures that flowering is synchronized with the end of the rainy season, irrespective of the sowing (25, 27, 28). Photosensitive species generally readjust their reproductive cycle to natural variations in photoperiod to ensure production before the end of the rainy season (29, 30).

In view of the wide variation in rainy season establishment periods affecting planting dates, it becomes important to better understand the response of sweet grain sorghum genotypes to photoperiod variation. Thus, the

onwards, with an average daily duration of 8.2 h/day, reaching a threshold of 5.7 h/day in August, before increasing by 08 h/day from September onwards. Atmospheric humidity increased from the beginning of the crop year to reach a peak in August (77%) and then decreased from September onwards to 72%. The 2 points of intersection of the combined curves of variations in monthly average insolation and atmospheric humidity (Fig. 2) marked the beginning and end of the crop year respectively. Thus, the first substantial rains of the 2019 crop year were recorded in the last 10 days of June and the last rains were recorded in the first 20 days of October (34).




Fig. 2. Monthly average distribution of insolation and atmospheric humidity in 2019 (34).

Plant material

The plant material consists of 30 sweet grain sorghum genotypes from the gene bank of the “Laboratoire Biosciences” of the “Université Joseph KI-ZERBO” (Table 1). The collection was carried out between 2008 and 2014 as part of the preservation of the agro-biodiversity of sweet sorghum in Burkina Faso. The genotypes studied were selected by drawing lots and taking care to include genotypes from the different agro-climatic areas of collection. Previous work by (8) reported that sweet grain sorghum is mainly grown in the sub-Saharan and northern Sudanian zones, hence their predominance in the study sample.

Experimental design

A randomized complete block design was used in this

Table 1. List of genotypes and agro-climatic origins of the collection

Climatic zones	Number of genotypes	Genotype Codes
Sub-Saharan	16	SKA3; YOU4; KBA1; PBO5; BKO3; YOU1; YOH4; BZ11; GB14; YOH3; SPI2; PLA1; PBO4; BKO1; YOU5; PGO3.
North Sudanian	9	BIP4; BKB1; BKB2; BKB4; KBZ1; KBZ4; MB07; MDE5; MTC2.
Southern Sudanian	5	SBR1; SBR5; SBR7; STO5; STO2.

study and was replicated 2 different planting dates. The 2 trials separated from each other by 4 m. Each trial was set up with 3 replicates 2 m apart. Each replication included the 30 genotypes, each planted in a 5.2 m long row. The row spacing was 80 cm and the distance between the patches was 40 cm, i.e., 14 patches per row. The area of each trial was approximately 455 m² (23.2 m × 19.6 m). Two factors were evaluated through this setup. The genotype factor and the sowing date factor which has 2 levels, namely first and second staggered sowing dates (D1 and D2), to measure the photoperiod sensitivity of the genotypes (23).

Cultivations techniques

Sowing trials took place on 26 June and 20 July, 2019 respectively, i.e., a difference of 24 days between the sowing dates, on plots previously ploughed with a tractor and levelled. These 2 dates correspond to the usual sowing periods in case of early and late rains. A first manual weeding followed by a thinning of 1 plant per hole was undertaken 2 weeks after sowing. NPK mineral fertilizer 14-23-14 and urea were applied at the same dose of 100 kg/ha (12) respectively 7 days after thinning and simultaneously during the ridging of the field at the morphogenesis stage.

Data collection

Data were collected on 10 agro-morphological traits. The morphological traits measured were the number of productive tillers (NPT), main stem diameter (DIS), plant height (PHT), length (INL) and number of internodes (NIN). The number of days from sowing to flag leaf appearance (NDS), number of days from sowing to flowering (NDF), panicle weight (PAW), panicle grain weight (WGP) and 100-grain weight (HGW) were the agronomic traits measured. The duration of daily photoperiod variations (alternating days and nights) has been obtained also (34).

Statistical analysis

The Excel spreadsheet was used to enter the collected data, calculate the averages, produce the graphs and calculate the photoperiodism coefficient (Kp) according to the formula (23, 35).

$$Kp = \frac{(NDS1 - NDS2)}{ID1 - D2I}$$

It corresponds, in fact, to the ratio of the difference in the sowing-appearance times of the ligule of the flag leaf of the main stem between the first and second sowing dates (NDS1-NDS2) to the difference between the two sowing dates, expressed in Julian calendar (23, 35). The photoperiod coefficient Kp thus varies from 0 (for genotypes

insensitive to photoperiod variation) to 1 for very photosensitive genotypes because the shortening of the vegetative period compensates for the sowing offset by grouping flowering at the same date.

Analysis of daily photoperiod variation data using R software version 1.5-10 allowed the determination of annual variations in photoperiodicity (long days, short days). Analysis of variance (ANOVA) was performed using Xlstat 2016 software to assess the effect of sowing dates and differences between genotypes for the different traits studied.

Results

Variation in morphological traits of genotypes

The results of the analysis of variance reported in Table 2 revealed variability in morphological traits of the 30 sweet grain sorghum genotypes studied and a significant effect of sowing date on their expression. Indeed, most of the morphological traits except the number of productive tillers ($p=0.88$) and main stem diameter ($p=0.086$) discriminated significantly between genotypes at 5%.

The sowing date factor strongly influenced the expression of the performance of morphological traits ($p<0.0001$) except for internode length trait ($p=0.522$). This

sowing date.

The sowing date x genotype interaction was significant for the traits number of productive tillers, plant height and the number of internodes and not statistically significant for the main stem diameter and internode length (Table 2).

Variation in agronomic traits of genotypes

The results of the analysis of variance reported in Table 3 revealed a variation in mean values of the agronomic traits of the measured genotypes between the 2 sowing dates. All genotypes showed highly significant differences for all agronomic traits assessed between the first and second sowing dates.

Phenological traits such as days to flowering and days to flag leaf appearance showed a significant reduction of the cycle by about 14 days at the second sowing date. Days to flag leaf appearance and days to flowering ranged from 64.3 to 74.7 days and 75.3 to 87.7 days respectively at the first sowing date compared to 48 to 62.3 days and 58 to 69 days at the second sowing date. In addition, a significant decrease in the average values of panicle weight per cluster, panicle kernel weight and hundred kernel weight were recorded. The weight of panicles per genotype varied from 206.5 to 342 g and from 138.4 to 321.1 g between the first and second planting dates respectively.

Table 2. Morphological traits variation of the genotypes according to the sowing dates

Traits		NPT	DIS (cm)	PHT (cm)	NIN	INL (cm)
Minimum	D1	0.7	2.2	290.3	14.4	17.7
	D2	0.4	1.9	230	10.9	16
Maximum	D1	2.2	2.5	392	16.5	23.3
	D2	1.6	2.3	314.7	12.5	22.9
Mean	D1	1.46	2.4	348.91	15.11	20.5
	D2	0.97	2.1	282.51	11.56	21.23
ΔD	D2-D1	-0.5	-0.26	-66.4	-3.55	0.72
Variation	$(\Delta D/D1) *100$	-34.17	-10.86	-19.03	-23.5	3.5
Genotype	F	0.69	1.45	5.95	3.18	4.06
	Pr.	0.88	0.086	<0.0001	<0.0001	0.019
Date	F	27.57	259.93	294.69	34.99	0.56
	Pr.	<0.0001	<0.0001	<0.0001	<0.0001	0.522
Date x Genotype	F	1.34	1.21	1.84	3.53	0.88
	Pr.	0.013	0.24	0.012	<0.0001	0.640

D1: first sowing date; **D2:** second sowing date; **ΔD :** difference in mean values between sowing dates; **F:** value file; **Pr:** probability of the factor; **NPT:** number of productive tillers; **PHT:** plant height; **DIS:** stem diameter; **NIN:** number of internodes; **INL:** internode length.

influence resulted in a significant decrease in the mean values of morphological traits ($D2-D1 < 0$) in the range of -10.86% to -34.17% for delayed sowing. Plant height ranged from 3.3 to 3.92 m at the first sowing date compared to 2.3 m to 3.15 m at the delayed sowing date with mean stem diameters ranging from 2.2 to 2.5 cm and 1.9 to 2.3 cm between the first and second sowing dates respectively. The plants produced an average of 1.46 productive tillers and 15 internodes at the first sowing date compared to 0.97 productive tillers and 11 internodes at the delayed

Panicle grain weight/genotype ranged from 177.1 to 299.1 g on the first and 100.2 to 250 g on the second planting date. Thus, between the first and second sowing dates, the mean values of the panicles weight/genotype varied from 274.16 g to 227.99 g, of the panicle grain weight from 232.36 g to 188.37 g and the 100-grain weight from 4.6 g to 3.3 g. The performance of the three traits decreased by 17%, 19% and 8.3% respectively, when sowing was delayed. The sowing date x genotype interaction was signifi-

Table 3. Performance of agronomic traits of genotypes at both sowing dates

Traits		NDS (days)	NDF (days)	HGW (g)	PAW (g)	WGP (g)
Minimum	D1	64.3	75.3	3.8	206.5	177.1
	D2	48	58	3.4	138.4	100.2
Maximum	D1	74.7	87.7	5.2	342.0	299.1
	D2	62.3	69	4.7	321.1	250.9
Mean	D1	71.3	80.1	4.6	274.16	232.36
	D2	57.48	66.07	3.3	227.99	188.37
ΔD	D2-D1	-13.81	-14.05	-0.35	-46.17	-43.99
Variation	$(\Delta D/D1)*100$	-19.36	-17.53	-8.3	-16.84	-18.93
Genotype	F	6.52	3.96	16.35	1.13	1.66
	Pr.	<0.0001	<0.0001	<0.0001	0.31	0.03
Date	F	2533.20	1498.682	44.83	22.56	38.82
	Pr.	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
Date x Genotype	F	2.19	3.82	5.97	1.43	1.82
	Pr.	0.002	<0.0001	<0.0001	0.095	0.013

D1: first sowing date; **D2:** second sowing date; **ΔD :** difference in mean values between sowing dates; **F:** value file; **Pr:** probability of the factor; **NDS:** number of days from sowing to appearance of the flag leaf; **NDF:** number of sowing-flowering days; **HGW:** 100-grain weight; **PAW:** panicle weight; **WGP:** panicle grain weight.

cant for all agronomic traits measured except panicle weight.

Evolution of the vegetative period of the genotypes

The results of the evolution of the sowing-flag leaf cycle (NDS) of the genotypes in the 2 trials are shown in Fig. 3. Indeed, in both trials, the sowing-flag leaf cycle showed a quasi-similar variation for the same genotypes with how-

ever, different amplitudes. At the second sowing date (curve B), all genotypes expressed a lower cycle compared to the first sowing date (curve A).

The cycle difference of the genotypes for each sowing date is however smaller compared to that of the 24-days sowing date shift. KBZ1, KBZ4, GB14, ST05, YOU1 and YOH3, reduced their cycle slightly at the second sowing date, which is reflected in the weak closeness of the 2




Fig. 3. Variation in the duration of the vegetative phase between the two sowing dates. **A:** evolution curve of the number of days sowing-appearance of flag leaves at the first sowing date; **B:** evolution curve of the number of days sowing-appearance of flag leaves on the second sowing date.

curves. The genotypes KBZ1 and BKB1 occupied extreme positions with gaps of 8.4 and 19.6 days respectively between the 2 sowing dates.

Evolution of flowering during the two sowing dates

The evolution of flowering for the 2 sowing dates over time was reported in Fig. 4. For the first sowing date (curve A), flowering started 76 days after sowing (DAS), on 11 September 2019 for the earliest genotype (PBO4). The number of flowering genotypes increased until 15 September, then

flowering (from 11 to 17 September) and a gap of 08 days between the end of flowering (from 19 to 27 September) of the 2 sowing dates was recorded.

The monthly average day lengths in a 24 h cycle corresponding to the variation in photoperiod recorded in Fig. 5 show that days are indeed longer than nights from May to August (>12 h 30 mins). Flowering, thus began when the average day length fell below an inductive photoperiod of less than 12 h 15 mins.




Fig. 4. Curves of the development of flowering for the two sowing dates. **A:** Curve of the development of flowering at the first sowing date; **B:** flowering evolution curve at the second sowing date.

decreased in the following days and recovered intensively until 19 September (84 days). For the second sowing date (Curve B), flowering started at 58 days (September 17th) for the earliest genotype (PBO4) and reached its optimum on 23 September. It stabilized in 2 days (September 23 and 25) and then dropped very quickly to end on September 27th (68 days). A gap of 06 days between the beginning of

Variation of the photoperiod coefficient

The results in Table 4 showed that all genotypes reacted to the difference in sowing date by reducing the cycle. Genotype KBZ1 with a cycle reduction of 08 days was the least photoperiodic ($K_p = 0.35$) and genotype BKB2 was the most photoperiodic ($K_p = 0.82$) with a cycle reduction of 20




Fig. 5. Curve of the evolution of the daily average monthly photoperiod duration.

Table 4. Variation in photoperiodism coefficient of studied sweet grain sorghum genotypes

Genotypes	NDS1	NDS2	NDS1 - NDS2	Kp	DSP
KBZ1	70.7	62.3	8.4	0.35	LoS
KBZ4	68.7	59.7	9	0.38	LoS
GBI4	70	58.7	11.3	0.47	LoS
STO5	70.7	59.3	11.4	0.48	LoS
YOU1	71	59.6	11.4	0.48	LoS
YOH3	70	58.3	11.7	0.49	LoS
PLA1	71.3	58.7	12.6	0.53	MoS
YOU4	69.3	56.7	12.6	0.53	MoS
STO2	72	59.3	12.7	0.53	MoS
BKB2	70.7	57.7	13	0.54	MoS
BZI1	72.3	59.3	13	0.54	MoS
KBA1	72.3	58.7	13.6	0.57	MoS
MTC2	67.3	53.7	13.6	0.57	MoS
MB07	70	56	14.33	0.58	MoS
BIP4	72	57.7	14.3	0.60	HoS
SPI2	73.3	59	14.3	0.60	HoS
SBR5	73	58.7	14.3	0.60	HoS
SBR1	73	58.3	14.7	0.61	HoS
PGO3	72.7	58	14.7	0.61	HoS
YOU5	70.7	56	14.7	0.61	HoS
MDE5	73	58.3	14.7	0.61	HoS
BKO1	73.3	58.3	15	0.63	HoS
BKB4	72.7	57.7	15	0.63	HoS
SBR7	72.3	56.7	15.6	0.65	HoS
PBO4	64	48	16	0.7	HoS
YOH4	72.3	56.7	17	0.71	HoS
PBO5	70.3	53.3	17	0.71	HoS
SKA3	74.7	57.7	17	0.71	HoS
BKB1	73.3	53.7	19.6	0.82	VHS

NDS1: Number of sowing days – the appearance of the flag leaf on the main stem from the first sowing date; **NDS2:** Number of sowing days – the appearance of the flag leaf on the main stem of the second sowing date; **Kp:** photoperiod coefficient; **DSP:** degree of photoperiod sensitivity; **LaS:** low photoperiod sensitivity; **MoS:** moderate photoperiod sensitivity; **HoS:** high photoperiod sensitivity; **VHS:** very high photoperiod sensitivity

days. The cycle was reduced by an average of 14 days overall and the average photoperiodic coefficient was 0.59. Four groups of sensitivity to photoperiod variation were identified. Thus, 20% of the genotypes showed low sensitivity ($0.3 \leq Kp < 0.5$), and 26.67% moderate sensitivity ($0.5 \leq Kp < 0.6$). A large number of genotypes, i.e. 46.67% showed high sensitivity ($0.6 \leq Kp \leq 0.8$) and a very small proportion (6.67%) showed very high sensitivity ($0.7 < Kp < 1$) to photoperiod variation.

Discussion

Variation in morphological traits of genotypes

The variability of most morphological traits except for the characters, number of productive tillers (NPT) and stem diameter (DIS) observed within the germplasm is similar to the results reported earlier (8). Indeed, these genotypes come from different agro-ecological zones of collection (8).

The variability of the genotypes was also reflected in the response to sowing date shift. Indeed, most morphological traits, except the length of internode (INL), were strongly influenced by the sowing date. This, variation was reflected in a significant decrease in the performance of the traits at the second sowing date. Delayed sowing resulted in a reduction in the size of vegetative organs of the genotypes varying by 10.86%, 34.1%, 19.03% and 25.5% respectively for the diameter and height of the main stem (DIS and PHT), the number of internodes (NIN) and the number of productive tillers (NPT). This reduction in the size of vegetative organs of the genotypes at the second sowing date observed in this study would be much more related to the number of internodes and stem thickness than to the internode length, which was not significantly affected by the sowing date ($P = 0.522$). This result corroborates the statements of (36) that the reduction in size is a direct result of the reduction in the number of phytomeres produced but rarely in their size. The reduction in the average number of

useful tillers at the second date would be related to the fact that the tillers could not accumulate enough photosynthates to ensure grain formation following the delay in sowing.

Variation in agronomic traits of genotypes

Moreover, the variability of most agronomic traits, with the exception of the panicle weight/bunch (PAW) trait, observed within the plant material studied, indicates the presence of significant variability within the sweet grain sorghum genotypes of Burkina Faso. This variability is believed to be due to the farmer's production system, which consists in producing seeds by mass selection, resulting in population varieties with a broad genetic base (37). The variability of genotypes was strongly revealed in the response to the shift in sowing date. Indeed, all agronomic parameters were strongly influenced by sowing date ($P < 0.0001$). The variation of all agronomic traits could reflect the importance of the effect of environmental factors such as rainfall regime, temperature, humidity, insolation and photoperiod on the aptitude of annual crop species. These climatic factors are characterized by high spatio-temporal variability and unpredictability in Sahelian regions (38, 39). The cycle would be the main parameter between sowing dates that is strongly influenced, affecting the expression of other morphological and agronomic traits. Phenology, in fact, is strongly affected by sowing date, which influences the variation in the rate of development of vegetative organs and grain production (27, 40, 41). However, several authors have reported that the phenology of photoperiod-sensitive sorghum is also influenced by the variation in photoperiod duration (42, 43). Indeed, sensitivity to photoperiod variation is a characteristic of local sorghum varieties allowing it to naturally adjust its cycle length to the likely duration of the rainy season (25, 36). Shifting the sowing date resulted in a variation of the cycle from medium (70-80 days) to relatively long (> 80 days) when sowing is early and from very short (50-60 days) to short (60-70 days) when sowing is delayed. The variation in doughy 100-grain weight (HGW) is thought to be due to a difference in the concentration of water in the grain related to the staggered harvesting time of the different genotypes. The decrease in panicle and grain weight per cluster (PAW and WGP) at the second sowing date could be explained by the shortened cycle of the genotypes at the delayed sowing (24).

Evolution of the vegetative period of the genotypes

The reduction of the vegetative period by about 8 to 20 days of all genotypes at the second sowing date shows a reduction of the cycle when sowing is delayed. This could reflect a sensitivity of the genotypes studied to photoperiod variation, because the reduction of the sowing-flowering cycle when sowing is delayed is one of the characteristics of photoperiodic plants (25, 38). Previous studies have shown a reduction in growth during late sowing regardless of water availability or other resources (36, 44, 45) and little variation in photoperiod near the equator (23). However, the results of this study show that sweet grain sorghum is very sensitive to photoperiod variation and its cycle can be modified by very small photoperiod

variations of the order of a few minutes (20 mins) or other climatic factors. Indeed, the decrease in rainfall (130 mm) and atmospheric humidity (72%) and the rapid increase in temperature to 29 °C and insolation duration to about 08 h recorded in September coincided with the floral initiation period. These variations in the external environment would have acted as a signal to lift inhibition that caused the genotypes to stop vegetative development in favor of reproductive development. Earlier studies by (46, 47) reported that the triggering of panicle initiation is mainly under the dual dependence of photoperiod and temperature. Furthermore, (45, 46) have pointed out that the issue of photoperiod cannot be treated independently of temperature and even humidity. There is successively a transition from apical to floral meristem triggered by the variation of photoperiod and then initiation of flowering provided by temperature. Thus, this increase in temperature, which coincided with the end of the rains, created a warm climate that slowed down the vegetative development of the plants. This would have triggered an action on the inhibitory genes of the vegetative phase, as at least 2 major and several minor genes control the vegetative cycle according to (23, 48). The present study shows that, in fact, photosensitive sorghum plants are maintained in the vegetative state when days are long (> 12 h 30 mins), until the day length reaches a critical threshold below 12 h 15 mins. Moreover, this interruption is made without taking into account the full development of the plant but acts as a distress signal that intervenes to allow the transition from the vegetative to the reproductive phase in order to ensure the sustainability of the species (40).

Evolution of flowering during the two sowing dates

The difference of six (06) days for the beginning and eight (08) days for the end of flowering between the 2 sowing dates shows that all the genotypes studied are photoperiodic. Indeed, in non-photosensitive species, the difference in sowing date is compensated for at flowering date (23, 49). A sowing date shift of 24 days will result in a 24-day shift in the flowering date of the second sowing date compared to the first sowing date. This is not the case in the present study. Similar results on photoperiodism have been reported for West African sorghum (23), grain sorghum (1, 5, 24) and sweet stalk sorghum from Burkina Faso (50). The shortening of the cycle at the second sowing date immediately resulted in a clustered flowering in 17 days of the genotypes from both sowing dates combined towards the end of the rainy season. This specificity of photoperiodic plants to interrupt the development of vegetative growth in favor of grain production (20, 24, 40), would give them an important evolutionary advantage due to the continental climate of the area, characterized by a strong interannual variation in rainfall regime. A variety is considered adapted to a zone if it flowers 2-3 weeks before the rainy season ends (45). In reality, grain yield and quality are closely linked to the flowering date, as grain of varieties that flower too early is attacked by birds and damaged by molds and insects. Varieties that flower too late deplete soil water reserves before grain filling is complete.

Variation of the photoperiod coefficient

Analysis of the results of the photoperiodism coefficients (Kp) shows a high sensitivity of sweet grain sorghum to photoperiod variation ($0.35 \leq Kp < 0.9$). Earlier studies by (24) found an average photoperiodism coefficient of 0.40 for ordinary grain sorghum which is relatively low compared to that of sweet grain sorghum ($Kp = 0.59$). This difference could be explained by the greater time lag of 30 days between sowing dates compared to 24 days in the present study, but it also reflects the great capacity of sweet grain sorghum to adapt to environmental conditions. Indeed, (50) found an average coefficient of variation of 0.54 for a 24 day sowing interval in sweet stalk sorghum. The results of the present study also show that the earliness of the genotypes is not related to the degree of sensitivity to photoperiod variation but rather to a more intense meristematic activity and rapid panicle development. This characteristic provides sweet grain sorghum with a possibility of continuous production. Earlier studies (23) reported that when highly photosensitive sorghums are sown under low photoperiods, the duration of the vegetative period is minimal and its value represents the intrinsic earliness of the variety. In addition to rainy season production, sweet grain sorghum could be sown in early January to be harvested before the end of March in Burkina Faso.

Conclusion

The study showed that all agro-morphological traits were influenced by variation in sowing date except for internode length. Delayed sowing resulted in a significant reduction in phenological, morphological and agronomic traits. Phenology was most affected by the sowing date. All sweet grain sorghum genotypes studied were sensitive to photoperiod variation and significantly reduced their cycle when sowing was delayed by 24 days. The coefficient of photoperiodism varied from genotype to another. KBZ1 was the least photoperiodic ($Kp = 0.35$) while BKB1 was the most photoperiodic ($Kp = 0.82$). The results of this study could contribute to the definition of the cropping calendar of sweet grain sorghum.

Acknowledgements

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Authors contributions

This work was carried out in collaboration with all authors. All authors read and approved the final manuscript. IY also provided the English translation.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None.

References

- Zongo JD. Ressources génétiques des sorghos (*Sorghum bicolor* (L.) Moench) du Burkina Faso : Evaluation agro-morphologique et génétique. Sciences naturelles [Thèse d'état]. Abidjan, Côte d'Ivoire : Université d'Abidjan;1991.
- Nebié B. Diversité génétique des sorghos à tige sucrée (*Sorghum bicolor* (L.) Moench) du Burkina Faso. Génétique et Amélioration des Plantes [Thèse de doctorat]. Ouagadougou, Burkina Faso : Université de Ouagadougou;2014.
- Sawadogo N. Diversité génétique des sorghos à grains sucrés (*Sorghum bicolor* (L.) Moench) du Burkina Faso. Génétique et Amélioration des Plantes [Thèse de doctorat]. Ouagadougou, Burkina Faso : Université de Ouagadougou;2015. <https://doi.org/10.4314/jab.v84i1.3>
- DGESS/MAAH. Résultats Définitifs de la Campagne Agricole et de la Situation Alimentaire et Nutritionnelle 2019/2020. Burkina Faso : Ministère de l'Agriculture et des Aménagements Hydrauliques. 2020 ;88p.
- Barro-Kondombo CP. Diversité agromorphologique et génétique de variétés locales de sorgho (*Sorghum bicolor* (L.) Moench) au Burkina Faso. Eléments pour la valorisation des ressources génétiques locales. Génétique et Amélioration des Plantes [Thèse de doctorat]. Ouagadougou, Burkina Faso : Université de Ouagadougou;2010.
- Sawadogo N, Ouédraogo MH, Traoré RE, Nanéma KR, Kiébré Z, Bationo-Kando P *et al.* Effect of agromorphological diversity and botanical race on biochemical composition in sweet grains Sorghum (*Sorghum bicolor* (L.) Moench) of Burkina Faso. J Bio Sci Biotech. 2017;6(1):263-69.
- Nebié B, Gapili N, Traore RE, Nanema KR, Bationo-Kando P, Sawadogo M *et al.* Diversité phénotypique de sorgho à grains sucrés du centre-nord du Burkina Faso. Sciences et techniques, sciences naturelles et agronomie. 2012;32(1&2):73-84.
- Sawadogo N, Nebié B, Kiébré M, Bationo -Kando P, Nanéma KR, Traoré RE *et al.* Caractérisation agro-morphologique des sorghos à grains sucrés (*Sorghum bicolor* (L.) Moench) du Burkina Faso. Int J Biol Chem Sci. 2014;8(5):2183-97. <https://doi.org/10.4314/ijbcs.v8i5.22>
- Sawadogo N, Nanema KR, Bationo-Kando P, Traore RE, Nebie B, Tiama D *et al.* Évaluation de la diversité génétique des sorghos à grains sucrés (*Sorghum bicolor* (L.) Moench) du Nord du Burkina Faso. J Appl Biosci. 2014;84(1):7654-64. <https://doi.org/10.4314/jab.v84i1.3>
- Sawadogo N, Batiéno TBJ, Kiébré Z, Ouédraogo MH, Zida WPMF, Nanéma KR *et al.* Assessment of genetic diversity of Burkina Faso sweet grain sorghum using microsatellite markers. Afr J Biotechnol. 2018;17(12):389-95. <https://doi.org/10.5897/AJB2017.16344>.
- Tiendrebeogo J, Sawadogo N, Kiebre M, Kabore B, Bationo-Kando P, Kiendrebeogo T *et al.* Evaluation comparative de la production de grains et du fourrage de sorgho à grains sucrés du Burkina Faso. Science et Technique, Sciences Naturelles et Appliquées. 2018;Spécial hors-série n° 4:261-71.
- Tiendrebéogo J. Performances agronomiques, potentialités fourragères du sorgho grains sucrés (*Sorghum bicolor* (L.) Moench) et relation génétique avec les autres types de sorgho cultivés au Burkina Faso. Génétique et Amélioration des Plantes [Thèse de Doctorat]. Ouagadougou, Burkina Faso : Université de Joseph KI-ZERBO;2020.
- Sawadogo N, Naoura G, Ouoba A, Yaméogo N, Tiendrebeogo J, Ouédraogo MH. Phenotypical characteristics and genetic diversity of three types of sorghum [*Sorghum bicolor* (L.) Moench] cultivated in Burkina Faso based on qualitative traits. Moroccan Journal of Agricultural Sciences. 2022a;3(2):109-16. <https://doi.org/10.5772/intechopen.105984>
- Sawadogo N, Ouédraogo MH, Bougma LA, Yaméogo N, Tondé WH, Tiendrebéogo J *et al.* Assessment of genetic variability of three types of sorghum cultivated in Burkina Faso using mor-

- phoagronomic quantitative traits and brix. In: Çalışkan M, Aydin S, editors. Genetic Diversity - Recent Advances and Applications. London: IntechOpen. 2022b;p. 1-16. <https://www.intechopen.com/online-first/82862> <https://doi.org/10.5772/intechopen.105984> (Last accessed on 22 Aug 02). <https://doi.org/10.5772/intechopen.105984>
15. MAHRH. Diagnostique du secteur semencier du Burkina Faso. Etat des lieux et perspectives de développement. Burkina Faso : Ministère de l'Agriculture, de l'Hydraulique et de Ressources Halieutiques. 2010;38p.
 16. OCDE/FAO. L'agriculture en Afrique subsaharienne : Perspectives et enjeux de la décennie à venir, dans Perspectives agricoles de l'OCDE et de la FAO. 2016-2025. Paris : Éditions OCDE ;2016. http://dx.doi.org/10.1787/agr_outlook-2016-5-fr (Last accessed on 2020 Jan 10).
 17. Somé L. Diagnostic agro-climatique du risque de sécheresse au Burkina Faso. Etude de quelques techniques agronomiques améliorant la résistance pour les cultures de Sorgho, Mil, Maïs. Agro-climatologie [Thèse de doctorat]. Montpellier, France : Université de Montpellier II;1989.
 18. Bambara D, Bilgo A, Lompo F, Hien V. Influence du changement climatique sur la diversité inter et intra-spécifique des plantes cultivées à Tougou au nord du Burkina Faso. *Int J Biol Chem Sci.* 2011;5(6):2415-33. <https://doi.org/10.4314/ijbcs.v5i6.21>
 19. Gnoumou XN, Yaméogo JT, Traoré M, Bazongo G, Bazongo P. Adaptation aux changements climatiques en Afrique subsaharienne : impact du zaï et des semences améliorées sur le rendement du sorgho dans les villages de Loaga et Sika (province du Bam), Burkina Faso. *IJIAS.* 2017;19(1):166-74.
 20. Trouche G, Chantereau J, Zongo JD. Variétés traditionnelles et variétés améliorées de sorgho dans les régions sahéliennes. In : Bacci L, Reyniers FN, eds. Le futur des céréales photopériodiques pour une production durable en Afrique tropicale semi-aride. Actes du séminaire international, 27-30 avril 1998, Florence, Italie. Florence (Italie) ; Montpellier (France) : Ed. CeSIA, Cirad éditions.1998; p. 197-208.
 21. Trouche G, Fliedel G, Chantereau J, Kondombo-Barro CP. Productivité et qualité des grains de sorgho pour le tô en Afrique de l'Ouest : les nouvelles voies d'amélioration. *Agriculture et développement.* 2000;23:94-107.
 22. Clerget B. Le rôle du photopériodisme dans l'élaboration du rendement de trois variétés de sorgho cultivés en Afrique de l'Ouest. Agronomie [Thèse de Doctorat]. France: Institut National Agronomique Paris-Rignon; 2004.
 23. Vaksman M, Traore SB, Niangado O. Le photopériodisme des sorghos africains. *Agriculture et développement.* 1996;9:13-18.
 24. Gapili N, Sawadogo M, Nanéma KR, Nebié B, Sawadogo N, Zongo JD. Etude du photopériodisme des écotypes de sorgho du Burkina Faso. *IJIAS.* 2015;13(4):901-09.
 25. Kassam AH, Andrews DJ. Effects of sowing date on growth, development and yield of photosensitive sorghum at Samaru, Northern Nigeria. *Exp Agric.* 1975;11(03):227-40. <https://doi.org/10.1017/S0014479700006761>.
 26. Lacy SM, Cleveland DA, Soleri D. Farmer choice of sorghum varieties in southern Mali. *Human Ecology.* 2006;34(3):331-53. <https://doi.org/10.1007/s10745-006-9021-5>
 27. Andrews DJ. Effects of date of sowing on photosensitive Nigerian sorghums. *Exp Agric.* 1973;9(4):337-46. <https://doi.org/10.1017/S0014479700010139>
 28. Cocheme J, Franquin P. A study of the agroclimatology of semi-arid area South of the Sahara in West Africa. *FAO/UNESCO.*1967;325p.
 29. Vigouroux Y, Mariac C, De Mita S, Pham JL, Gérard B, Kapran I, Bezançon G. Selection for earlier flowering crop associated with climatic variations in the Sahel. *PLoS ONE.* 2011;6(5):e19563. <https://doi.org/10.1371/journal.pone.0019563>.
 30. Cockram J, Jones H, Leigh FJ, O'Sullivan D, Powell W, Laurie DA, Greenland AJ. Control of flowering time in temperate cereals: genes, domestication and sustainable productivity. *J Exp Bot.* 2007;58:1231-44. <https://doi.org/10.1093/jxb/erm042>
 31. IGB/BNDT. Données topographiques. Burkina Faso : Institut Géographique du Burkina, Base Nationale des Données Topographiques; 2012. www.igb.bf (Last accessed on 2022 jan 24)
 32. Thiombiano A, Kampman D. Atlas de la biodiversité de l'Afrique de l'Ouest. Tome II. Ouagadougou (Burkina Faso) et Frankfurt/Main. 2010;625p.
 33. IDR. Données pluviométriques. Ouagadougou : Institut de Développement Rural; 2019.
 34. ANAM. Données climatiques. Burkina Faso : Agence Nationale de la Météorologie; 2019. <https://www.meteoburkina.bf> (Last accessed on 2022 Jan 24)
 35. Clerget B, Rattunde HFW, Dagnoko S, Chantereau J. An easy way to assess photoperiod sensitivity in sorghum: Relationships of the vegetative-phase duration and photoperiod sensitivity. *Journal of SAT Agricultural Research.* 2007;3(1):1-4. <http://www.icrisat.org/journal/>.
 36. Clerget B, Dingkuhn M, Goze E, Rattunde HFW. Variability of phyllochron, plastochron and rate of increase in height in photoperiod sensitive sorghum varieties. *Annals of Botany.* 2008;101:579-94. <https://doi.org/10.1093/aob/mcm327>
 37. Gapili N. Etude de la diversité génétique et du photopériodisme d'accessions de sorgho grains [*Sorghum bicolor* (L.) Moench] du Burkina Faso. Thèse de doctorat Unique. Thèse de doctorat. Génétique et amélioration des plantes. Burkina Faso: Université de Ouagadougou; 2016.
 38. Muchow RC, Carberry PS. Phenology and leaf area development in a tropical grain sorghum. *Field Crops Res.* 1990;23(3&4):221-37. [https://doi.org/10.1016/0378-4290\(90\)90056-H](https://doi.org/10.1016/0378-4290(90)90056-H)
 39. Agossou DSM, Tossou CR, Vissoh VP, Agbossou KE. Perception des perturbations climatiques, savoirs locaux et stratégies d'adaptation des producteurs agricoles béninois. *Afr Crop Sci J.* 2012;20(s2):565-88.
 40. Traoré SB, Reyniers F, Vaksman M, Kone B, Sidibe A, Yorote A et al. Adaptation à la sécheresse des écotypes locaux de sorghos du Mali. *Sécheresse.* 2000;11:227-37.
 41. Kouressy M, Traore S, Vaksman M, Grum M, Maikano I, Soumare M et al. Adaptation des sorghos du Mali à la variabilité climatique. *Cah Agric.* 2008;17(2):95-100. <https://doi.org/10.1684/agr.2008.0189>
 42. Clerget B, Dingkuhn M, Chantereau J, Hemberger J, Louarn G, Vaksman M. "Does panicle initiation in tropical sorghum depend on day-to-day change in photoperiod?" *Field Crops Res.* 2004;88(1):11-27. <https://doi.org/10.1016/j.fcr.2003.11.008>
 43. Dingkuhn M, Kouressy M, Vaksman M, Clerget B, Chantereau J. Applying to sorghum photoperiodism the concept of threshold-lowering during prolonged appetite. *Eur J Agron.* 2008;28(2):74-89. <https://doi.org/10.1016/j.eja.2007.05.005>
 44. Miller FR, Barnes DK, Cruzado HJ. Effect of tropical photoperiods on the growth of sorghum when grown in 12 monthly plantings. *Crop Science.* 1968;8:449-502. <https://doi.org/10.2135/cropsci1968.0011183X000800040032x>
 45. Kouressy M, Dingkuhn M, Chantereau J, Vaksman M. Leaf longevity and grain yield of sorghum plant types differing in photoperiod sensitivity and plant height. *Eur J Agron.* 2008;28(3):195-209. <https://doi.org/10.1016/j.eja.2007.07.008>
 46. Bangbol SH. Genetic characterization of flowering time in sorghum. *Plant Biology Genetics* [PhD Thesis]. Queensland, Australian: University of Queensland;2013.
 47. Crossa J. From genotype x environment interaction to gene x

- environment interaction. *Current genomics*. 2012;13(3):225-44. <https://doi.org/10.2174/138920212800543066>
48. Amasino RM, Michaels SD. The timing of flowering. *Plant Physiology*. 2010;154:516-20. <https://doi.org/10.1104/pp.110.161653>
49. Franquin P. Formulation des phénomènes apparents de photoperiodisme en conditions naturelles. *Principes de base. Cahiers ORSTOM, série Biologie*. 1974;23:31-43.
50. Sawadogo N, Tiendrebéogo J, Naoura G, Béré TLK, Tondé WH, Sawadogo B, Bationo-Kando P. Photoperiod sensitivity and variability of agromorphological traits and brix content of sweet sorghum cultivated in Burkina Faso under two sowing dates. *Adv Agric*. 2022;2022:1-10. <https://doi.org/10.1155/2022/9504150>.