Almaraz-Abarca N, Ayala-Chaidez AI, Wallander-Compean L, Ávila-Reyes JA, Delgado-Alvarado EA, Naranjo-Jiménez N, Rosas-Medina I, Colmenero-Robles A. Variation of the phenolic composition and  $\alpha$ -glucosidase inhibition potential of seeds, soaked seeds, and sprouts of four wild forms and four varieties of common bean (*Phaseolus vulgaris*). Plant Science Today (Early Access). https://doi.org/10.14719/pst.2250

## **Supplementary Figure**




Fig. S1. UV spectra of the phenolic compounds found in the seeds, soaked seeds and sprouts of eight wild forms and eight varieties of common bean (*Phaseolus vulgaris*).

S: Accumulated in seeds, I: Accumulated in soaked seeds, G: Accumulated in sprouts.




Fig. S1 *cont*. UV spectra of the phenolic compounds found in the seeds, soaked seeds and sprouts of eight wild forms and eight varieties of common bean (*Phaseolus vulgaris*). S: Accumulated in seeds, I: Accumulated in soaked seeds, G: Accumulated in sprouts.




Fig. S1 *cont*. UV spectra of the phenolic compounds found in the seeds, soaked seedsand sprouts of eight wild forms and eight varieties of common bean (*Phaseolus vulgaris*). S: Accumulated in seeds, I: Accumulated in soaked seeds, G: Accumulated in sprouts.




Fig. S1 *cont*. UV spectra of the phenolic compounds found in the seeds, soaked seeds and sprouts of eight wild forms and eight varieties of common bean (*Phaseolus vulgaris*). S: Accumulated in seeds, I: Accumulated in soaked seeds, G: Accumulated in sprouts.




Fig. S1 *cont*. UV spectra of the phenolic compounds found in the seeds, soaked seeds and sprouts of eight wild forms and eight varieties of common bean (*Phaseolus vulgaris*). S: Accumulated in seeds, I: Accumulated in soaked seeds, G: Accumulated in sprouts.




Fig. S1 *cont*. UV spectra of the phenolic compounds found in the seeds, soaked seeds and sprouts of eight wild forms and eight varieties of common bean (*Phaseolus vulgaris*). S: Accumulated in seeds, I: Accumulated in soaked seeds, G: Accumulated in sprouts.




Fig. S1 *cont*. UV spectra of the phenolic compounds found in the seeds, soaked seeds, and sprouts of eight wild forms and eight varieties of common bean (*Phaseolus vulgaris*). S: Accumulated in seeds, I: Accumulated in sprouts.




Fig. S1 *cont*. UV spectra of the phenolic compounds found in the seeds, soaked seeds and sprouts of eight wild forms and eight varieties of common bean (*Phaseolus vulgaris*). S: Accumulated in seeds, I: Accumulated in soaked seeds, G: Accumulated in sprouts.