



RESEARCH COMMUNICATION

A new variety of *Gymnosporia emarginata* (Celastraceae) from the Coromandel Coast of Peninsular India

Umamaheswari P^{1,2}, Balachandran N^{1*}, Kumaresan V² & Yuvarani K²

¹Ecology Department, French Institute of Pondicherry, Puducherry – 605001, India

²Department of Botany, Kanchi Mamunivar Government Institute for Postgraduate Studies and Research, Puducherry – 605008, India

*Email: balachandran.n@ifpindia.org



ARTICLE HISTORY

Received: 29 November 2022

Accepted: 19 March 2023

Available online

Version 1.0 : 20 April 2023



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care etc. See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Umamaheswari P, Balachandran N, Kumaresan V, Yuvarani K. A new variety of *Gymnosporia emarginata* (Celastraceae) from the Coromandel Coast of Peninsular India. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.2255>

Abstract

A new variety, *Gymnosporia emarginata* (Willd.) Thwaites var. *coromandelica* N.Balach. & P. Umamaheswari (Celastraceae) has been described from Tamil Nadu, India. The diagnostic characters of this variety are: long stamens, ovary immersed in the disc, style sessile and stigma lobes converged. Detailed descriptions, differences in characters between the 2 varieties, ITS based phylogenetic analysis and related images are provided for easy identification.

Keywords

Dry evergreen forest, *Maytenus*; Pondicherry, scrub vegetation, Tamil Nadu

Introduction

Gymnosporia (Wight & Arn.) Hook.f. is one of the largest genus of the family Celastraceae, globally represented by about 115 species (1, 2). Loesener delimited *Gymnosporia* to Old World species which are characterised by the presence of thorns and/or brachyblasts and fasciculate leaves, while similar plants that are unarmed with spiral or distichous (but not fasciculate) leaves are assigned to *Maytenus* Molina (4). Later Marais (5) and Robson (6-7) included all spine-bearing members under *Maytenus*. However, *Gymnosporia* has been reinstated by Jordaan, Jordaan & Van Wyk and Archer & Jordaan (8-11) and as per the circumscription established by them, the genus is characterised by 3-locular capsules with orange seeds. The other diagnostic characters to differentiate the taxa include growth form, pubescence of leaves, branches and peduncles, orientation of spines, presence and shape of stipules, colour of the flowers (white or red) and the shape of capsules.

On the basis of fruit morphology, seed characters and in addition to leaf anatomical evidences, the genus *Gymnosporia* is subdivided into eight sections. Jordaan and Wyk (16) included *G. emarginata* (Willd.) Thwaites under the section *Tenuispinae* Jordaan, comprising 49 species, six subspecies and two varieties. There are 16 species recorded in India, Sri Lanka and Myanmar (12). However, Ding Hou (13) merged the Indian *Gymnosporia montana* (Roth) Benth. with *Maytenus emarginata* (Willd.) Ding Hou against the study of Roxburgh (14) and Lawson (15). Meanwhile Jordaan and Wyk (16) revised the genus and treated the *G. emarginata* as a distinct species; according to them it is confined to the plains of SE India and Sri Lanka and it does not occur further eastwards. Ramamurthy recorded 18 species and two varieties in India (17), but recently Mao and Dash recognized 16 species and two varieties (18). Narasimhan included only seven species to

the state Tamil Nadu (19).

During the last four years of field exploration in and around the Puducherry region, as part of PhD program of the first author we came across few interesting deviations in the characters of the flowers of some samples of *Gymnosporia emarginata*. To know the evolutionary changes of the species Neighbor-Joining method (20) was inferred, evolutionary analyses were conducted using MEGA11 (21) software, followed by construction of

phylogenetic tree based on ITS marker (OP680785.1 *Gymnosporia* sp.) and the optimal tree is presented (Fig. 1). Critical scrutiny of specimens of *G. emarginata* housed at AURO, HIFP, MH, RHT and K and study of literature (16, 17, 22-25) showed that the present specimens clearly differ from the typical form by certain constant characters (Table 1) which merit recognition as a distinct variety. Hence, the same is described here as a new variety with comparative field images (Fig. 2). A key to the species of *Gymnosporia* occurring in Tamil Nadu is appended to facilitate easy




Fig. 1. Phylogenetic analysis among the species of *Gymnosporia* and *Maytenus*, showed the differences in the nucleotide sequences.

Table 1. Character differences between the two varieties of *Gymnosporia emarginata*.

No	Plant parts	var. <i>coromandelica</i>	var. <i>emarginata</i>
1	Stamen filament	5–6 mm long, longer than stigma	1–2 mm, shorter than stigma
2	Style	sessile	1–2 mm long
3	Stigma	lobes converged	distinct, lobes diverged
4	Disc	covered more than half of the ovary	Covered at base of the ovary





Fig. 2. *Gymnosporia emarginata* A-D: var. *emarginata*; E-H: var. *Coromandelica*.



TYPE

27883



INSTITUT FRANÇAIS PONDICHÉRY

Famille : CELASTRACEAE HIFP. Wpp 176

Espèce : *Gymnosporia emarginata* (Willd.) Thunberg
 = (*Maytenus emarginata* (Willd.) Ding Hou)

Date : 26.2.22 Par : Uma Maheswari P. Barathan,
 Balachandran N

Lieu :
 Menveille, Lake estate,
 Osuti.

Alt :
 Lat : 11.9634
 Long : 79.7637.

Observations : Straggler, leaf
 simple, alternate, leaf apex emarginate,
 flower white colour.

Fig. 3. Holotype of *Gymnosporia emarginata* var. *coromandelica* N.Balach. & P.Umamaheswari var. nov.

Taxonomic Treatment

Gymnosporia emarginata (Willd.) Thwaites var. ***coromandelica*** N.Balach. & P.Umamaheswari, **var. nov.**, differing from var. *emarginata* by the lax inflorescence with less number of flowers, 5-6 mm long staminal filaments, style sessile and the disc embracing more than half of the ovary (Table 1).

Type: INDIA, Tamil Nadu, Ossudu, Merveille, 11° 57' 48.24" N, 79° 45' 48.24" E, 26.02.2022, P.Umamaheswari & N. Balachandran 027883 (holotype HIFP!; isotype MH!). (Fig. 3)

Armed shrub, c. 2 m high; spines 2–3 cm long. Leaves obovate to oblanceolate, glabrous, coriaceous, 2.5–8.0 × 1.5–4.0 cm, lamina dark green above pale below, apex obtuse-emarginate, margin usually entire, rarely crenate, lateral nerves 5–6 pairs, obscure above, prominent below; petioles 2–12 mm long. Inflorescence solitary-corymbose (rarely dichasial) cymes, lax in the axils of the leaves, in a short condensed shoot, or on the spines; peduncle subsessile, to 5 mm long. Flowers 4–6 mm across, white, bracteate; pedicels slender, 5–12 mm long. Calyx lobes broadly ovate-orbicular, c. 1 mm long. Petals oblong, 3–4 × 1.0–1.5 mm, reflexed, margin revolute. Disc fleshy, sinuate. Stamens 5, filament 5–6 mm long, equal to petal, inserted at the base of disk; anthers broadly ovoid. Ovary immersed in the disk, style sessile; stigmas 3-lobed, convergence. Capsules broadly obovoid, 8–10 × 7–8 mm, 3-angled and celled, seeds 2 per cell. Seeds red, ovoid or ellipsoid; aril fleshy, white, reduced to a rim at the base of the seed.

Phenology: Flowering: December – January (in very short period, 3–7 days, rarely in August–September); Fruiting: March–April.

Etymology: The specific epithet is derived from the study area Coromandel Coast, the region to which the species is largely confined.

Additional Specimen examined (paratypes): India, Tamil Nadu, South Arcot, Ulunderpet RF, 31.01.1975, K.Balasubramaniam, 004252 (HIFP!); Villupuram, Puthupet, 12° 03' N, 79° 53' E, 19.12.1995, W.F.Gastmans & C. De Riddar, 5475 (AURO!); Chengalpattu, Vallam RF, 12° 41' N, 80° 02' E, 40 m ASL, 07.01.1998, J.D.Hollander, P.Blanchflower & W.F.Gastmans, F 6281 (AURO!).

Distribution: India - endemic.

Habitat: Plains from the coast, sacred groves, reserve forest, scrub forests, and in unclassified vegetation to hillocks (0-750 m). Locally abundant, densely branched and profusely armed with spreading and interlocked branched and thus forming impenetrable thickets. This new taxon is associated with *Benkara malabarica* (Lam.) Tirveng., *Carissa spinarum* L., *Jatropha gossypifolia* L., *Gmelina asiatica* L., *Pterospermum suberifolium* (L.) Raeusch., *Memecylon umbellatum* Burm.f., *Hugonia mystax* L.

Conservation Status: Found growing side by side with the var. *emarginata*, fairly common and widely distributed along the Coromandel Coast of Southern

India. Following the IUCN Red list criteria, Area of Occupancy and Extent of Occurrence (27) this taxon treated as Least Concern (LC).

Key to the species of *Gymnosporia* in Tamil Nadu

- | | |
|---|------------------------|
| 1a. Spines bearing leaves and flowers | 2 |
| 1b. Spines not bearing leaves and flowers | 4 |
| 2a. Fruits globose; ovary 2-loculed; branchlets straight | <i>G. senegalensis</i> |
| 2b. Fruits turbinate; ovary 3-loculed; branchlets zig-zag | |
| 3a. Inflorescences c. 2.5cm long; aril attached to the base of the seed | <i>G. wallichiana</i> |
| 3b. Inflorescences c. 1cm long; aril embracing the seed base | <i>G. emarginata</i> |
| 4a. Flowers in cymes only; fruits obcordate or turbinate | 5 |
| 4b. Flowers in cymes or in fascicles; fruits obovoid | <i>G. ovata</i> |
| 5a. Aril embracing about half the seed; leaves ovate-elliptic | <i>G. heyneana</i> |
| 5b. Aril embracing the seed base only; leaves oblong or lanceolate | 6 |
| 6a. Inflorescences reddish; cymes simple | <i>G. rufa</i> |
| 6b. Inflorescences yellowish green; cymes dichotomous | <i>S.G. acuminata</i> |

Acknowledgements

The authors are thankful to Dr. Blandine Ripert, Directress and Dr. Dorris Barboni, Head of Ecology, French Institute of Pondicherry and the Director and Head of the Department of Botany, Kanchi Mamunivar Government Institute for Postgraduate Studies and Research, Puducherry for their constant support and encouragement; Sri Aurobindo Trust, UK for the financial support to the botanical survey; the Officer In-charge, Botanical Survey of India, Coimbatore, Tamil Nadu granted permission to consult the herbarium specimens and to refer the library; the Curator, AURO Herbarium, Auroville permitted to refer and cite the specimen and the two anonymous reviewer(s) who improved this manuscript in better shape.

Authors contributions

The first two authors conducted field survey, confirmed the identity and drafted the manuscript; KV and YK carried out the DNA extraction, participated in the sequence alignment and performed phylogenetic analysis. All authors read and approved the final version of the manuscript.

Compliance with ethical standards

Conflict of interest: The authors declare that there are no conflict of interest

Ethical issues : None.

References

- Jordaan M, van Wyk AE. *Gymnosporia swazica* (Celastraceae), a new species from southern Africa. *Phytotaxa*. 2015;192(4):296-300.
- Mabberley DJ. *Mabberley's Plant-Book: A portable dictionary of plants, their classification and uses*. First South Asia edition, Rajkamal electric press, Kundli, Haryana. 2018; p. 4133.
- Loesener LET. Celastraceae. *Celastroideae-Eucelastreae*. Die natürlichen Pflanzenfamilien 3, 5. Engelmann, Leipzig. 1942;203-11. <https://doi.org/10.5962/bhl.title.4635>
- Marais W. An enumeration of the *Maytenus* species of southern Africa. *Bothalia*. 1960;381 – 86.
- Robson NKL. Celastraceae. In: *Flora Zambesiaca*. editors, A W Exell, A Fcmandes, H Wild. Cmwn Agents for Oversea Government and Administrations. London . 1966; Vol. 2. p. 355- 418.
- Robson NKL. Celastraceae. *Maytenus*. In: *Flora of tropical East Africa Celastraceae*. RM Polhill. 1994; p. 1-21.
- Jordaan M. A taxonomic revision of the spiny members of subfamily Celastroideae (Celastraceae) in southern Africa. MSc [thesis], University of Pretoria, Pretoria. 2013.
- Jordaan M, Van Wyk AE. Systematic studies in subfamily Celastroideae (Celastraceae) in southern Africa: reinstatement of the genus *Gymnosporia*. *South African Journal of Botany*.1999a;177-81. [https://doi.org/10.1016/S0254-6299\(15\)30958-3](https://doi.org/10.1016/S0254-6299(15)30958-3)
- Jordaan M, Van Wyk AE. Systematic studies in subfamily Celastroideae (Celastraceae) in southern Africa: two new species of *Gymnosporia* from the Maputaland Centre of Endemism. *South African Journal of Botany*. 1999 b;315-20. [https://doi.org/10.1016/S0254-6299\(15\)31018-8](https://doi.org/10.1016/S0254-6299(15)31018-8)
- Archer RH, Jordaan M. Celastraceae. In: Leistner OA, editor. *Seed Plants of Southern Africa-Families and Genera*. *Strelitzia* 10. National Botanical Institute, Pretoria. 2000; p. 214-20.
- Jordaan M, Van Wyk AE. Reinstatement of *Gymnosporia* (Celastraceae): implications for the Flora Malesiana region, *Telopea*. 2003a;155-67.
- Jordaan M, Van Wyk AE. Sectional classification of *Gymnosporia* (Celastraceae), with notes on the nomenclatural and taxonomic history of the genus. *Taxon*. 2006; 515-25. <https://doi.org/10.2307/25065602>
- Ding Hou. Celastraceae I. In: van Steenis, CGGJ, editor. *Flora Malesiana*, series 1, 2. Dijkstra Printers, Groningen. 1962; Vol. 6. P. 227-91.
- Roxburgh W. *Celastrus*. In: Wallich N, editor. *Flora Indica or Descriptions of Indian Plants*. Mission Press, Serampore. 1824;Vol. 2. p. 386-401.
- Lawson MA. Celastrineae. In: Hooker J D. editor. *The Flora of British India*. Reeve & Co. London. 1875; vol 1(3). p. 606-29.
- Ramamurthy K. Celastraceae In: Singh NP, Vohra JN, Hajra PK, Singh DK, editors. *Flora of India*, Botanical Survey of India, Calcutta, India. 2000; Vol. 5. p. 124-25.
- Mao AA, Dash SS. *Flowering Plants of India – As Annotated Checklist (Dicotyledons)*, Botanical Survey of India, Kolkata. 2020; vol 1. p. 258-64.
- Narasimhan D, Sheeba JI. *Flowering plants of Tamil Nadu: A compendium*; 2021.
- Saitou N, Nei M. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*. 1987; 406-25.
- Tamura K, Stecher G, Kumar S. MEGA 11: Molecular Evolutionary Genetics Analysis version 11. *Molecular Biology and Evolution*. 2021. <https://doi.org/10.1093/molbev/msab120>.
- Matthew KM. *The Flora of Tamil Nadu Carnatic, The Rapinath Herbarium, Tiruchirappalli*. 1983; Vol 3 (1). p. 258.
- Wadhwa BM. Celastraceae. In: *A Revised Handbook to the Flora of Ceylon*, editors. MD Dassanayake, WD Clayton. 1996; Vol 10. p. 75-106.
- Liu Quanru, Michele Funston. *Gymnosporia*. In: *Flora of China* [e-book]; 2008 cited 4 Aug 2022]. Vol 11: 474-77. Available from: <http://flora.huh.harvard.edu/china/PDF/PDF11/Gymnosporia.pdf> .
- Britto SJ. *The Flora of Central and North Tamil Nadu, Part – 1 Cabombaceae –Zygophyllaceae (APG – IV)*, The Rapinath Herbarium, Tiruchirappalli. 2019; p 624.
- IUCN. *The IUCN Red List of Threatened Species*. Version 2022-1 [Internet]. 2022 [cited 17 Oct 2022]. Available from: <https://www.iucnredlist.org>