



RESEARCH ARTICLE

Studies on *in vitro* evaluation of antibacterial and antioxidant activity of *Hydrilla verticillata* (L.f.) Royle

Bhagyeswari Behera & Kunja Bihari Satapathy*

Department of Botany, School of Applied Sciences, Centurion University of Technology and Management, Odisha, India


*Email: kbs_bot@rediffmail.com



ARTICLE HISTORY

Received: 13 December 2022 Accepted: 08 May 2023 Available online

Version 1.0 : 03 August 2023 Version 2.0 : 01 October 2023



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an openaccess article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (https://creativecommons.org/licenses/by/4.0/)

CITE THIS ARTICLE

Behera B, Satapathy K B. Studies on *in vitro* evaluation of antibacterial and antioxidant activity of *Hydrilla verticillata* (L.f.) Royle. Plant Science Today. 2023; 10(4): 66–71. https://doi.org/10.14719/pst.2312

Abstract

The purpose of the present investigation was to evaluate the therapeutic effect of Hydrilla verticillata (L.f.) Royle by examining its antibacterial and antioxidant properties in in vitro condition. For this study the methanolic and n-hexane extracts of Hydrilla verticillata (L.f.) Royle was tested against 5 pathogenic bacteria such as "Enterococcus faecalis, Escherichia coli, Klebsiella pneumoniae, Shigella flexneri and Vibrio choleriae" by using agar well diffusion method and for antioxidant activity DPPH scavenging assay was carried out. An antibiotic Ciprofloxacin was taken as reference. The result of the study revealed that, the n-hexane extract of H. verticillata showed highest zone of inhibition against Enterococcus faecalis (20±1.41 mm) and least against Vibrio choleriae (12±1.63 mm) at concentration of 15 mg/mL. While, methanolic extract showed highest zone of inhibition against Shigella flexneri (26±0.75 mm) and the least zone of inhibition against Klebsiella pneumoniae (20±1.24 mm). The significant result for antioxidant activity was evaluated. The IC₅₀ value of n-hexane and methanolic extracts were 36.26 and 16.84 respectively. The methanolic extract was found to be having better result than n-hexane extract. The n-hexane extract showed the presence of alkaloids, triterpenoids and polyphenolic compounds, tannin, glycoside while phytoconstituents like alkaloids, saponins, tannins, steroids, flavonoid, glycosides, triterpenoids and polyphenolic compounds were found to be present in methanolic extract of *H. verticillata*. Further study can be done on the evaluation of secondary metabolites compound analysis for therapeutic effect.

Keywords

Hydrilla verticillata; antibiotic resistant; antibacterial; antioxidant activities

Introduction

Since ancient time, it was believed that all the plants retain some pharma-cological properties due to the presence of secondary metabolites such as "alkaloids, flavonoids, glycosides, saponins, steroids, tannins and triterpenoids" etc. which should be utilized to fight against the disease-causing pathogens (1-3). Now a day's significant progress has been noticed in the field of drug discovery by naturally and synthetically with the innovation of science and technology (4). Antibiotic are considered as most promising solution for bacterial causing diseases. However, only one third of the infectious diseases known have been treated from these synthetic products (5). New therapeutics are needed to control the disease resistance bacterial borne diseases which posing an increasing therapeutic problem. Several

human pathogenic bacteria are found to exhibit antibiotic resistance due to misuse or over dose of drug (6-8). The use of herbal plants as traditional health remedies is the most common in Asia, Latin America and Africa, majority world's population has reported to have minimal side effects (9). One such example is Shigella flexneri, a grampositive bacterium which causes common diseases like diarrhoea, abdominal cramp, fever and also causes serious disease like ulcer in rectum and colon. It can cause acute condition in immune compromised body and HIV infected individual. Shigella flexneri was found to be resistant to Azithromycin. Another example is Methicillin-resistant Staphylococcus aureus (MRSA), a gram-positive bacterium, has been documented as one of the most disease-causing bacteria. It causes infection in skin and soft tissues of body and also infects the lungs causing pneumonia. If it left untreated, it may transmit to blood stream called bacteremia and also to bone. Micrococcus luteus a gram-positive bacterium causes meningitis, septic, arthritis, cutaneous infection and bacteremia and also found in HIV positive patient. This bacterium is a rare pathogen of endocarditis (10). This bacterium is resistant to both tetracycline and erythromycin (11, 12). Escherichia coli, a multi-drug resistant gram-negative bacterium causes many acute diseases including bloody diarrhoea, stomach cramp and urinary tract infection (13). Now-a-days multi-drug resistant (MDR) pathogens are observed to play major role to make failure in organ transplantation. New natural therapeutics and alternative approaches are needed to control the multidrug resistant bacteria. One way to control these MDR bacteria by using antibiotic resistant inhibitors from plant extract (14, 15). Plants are known to produce a variety of compounds to protect themselves against a large number of pathogens. It is expected that plant extracts showing target sites other than those used by antibiotics might be active against drug resistant pathogens (16). The crude extracts of plants have been shown to provide better therapeutic agent than the isolated compound without any side effect (17, 18).

Hydrilla, a submerged aquatic plant, attracts the attention of many researchers due to its aggressive competitive growing nature (invasive) and contains many nutritive components. It is loaded with high level of Vitamins including rare Vitamin-B12, antioxidant and minerals (19). Although this plant is having lots of nutritional properties, it is not commonly utilized as a pharmaceutical product. Additive worth can be given to this plant if it is proven to have medicinal properties like anti-bacterial and antioxidant activity. In the present study, methanolic and n-hexane extracts of Hydrilla verticillata have been evaluated for their antimicrobial and antioxidant behaviour.

Materials and Methods

Collection and preparation of plant extract

The experimental plant sample *Hydrilla verticillata* (L.f.) Royle was collected from the water body located in the CUTM (Centurion University of Technology and Management) campus, Bhubaneswar, Odisha in the month of Au-

gust, 2019. The plant specimen (*Hydrilla verticillata* (L.f.) Royle; Voucher No. BB-341) was identified by following the available literature (Saxena & Brahmam, 1994-96) and matching with the preserved specimen at Herbarium unit, IMMT-CSIR, Bhubaneswar (20). Collected plant sample washed thoroughly in running tap water then distilled water; kept them in room temperature to get them dry. Dried samples were then grinded properly to make a powder form. 15 g of powdered plant materials were successively extracted using 2 different solvents i.e. 250 mL each of n-Hexane and methanol by Soxhlet apparatus (Borosil, India) separately (21). All the extracts were evaporated at room temperature and kept in woven at 40 °C over night to evaporate all the solvent and stored in sterile glass vials at 4 °C for further use.

Preliminary phytochemical analysis

For preliminary phytochemical analysis, standard protocol was used to detect the presence of different secondary metabolites in experimental plant *Hydrilla verticillata*. 2 mL of plant extracts were taken and treated with various chemical reagents as per the test methods (22-24).

Antioxidant Assay

The *in vitro* antioxidant study was done by free radical scavenging assay (DPPH assay). In this technique, stock solution for the experiment was prepared by dissolving 50 mg of extract in 50 mL of methanol and n-hexane. Different concentration ranges from (300, 250, 200, 150, 100 and 50μg/mL) of methanolic and n-hexane extract of *H. verticil*lata was prepared by diluting the stock solution. Then, from above solution 1.5 mL sample was taken and 1.5 mL of DPPH was added to it and kept them in dark condition at room temperature for 30 min for allowing them to react properly. Then, the absorbance was taken using UV spectrophotometer at 517nm. Standard stock solution was prepared by dissolving dry ascorbic acid in 10 mL of methanol and n-hexane. And 5 different aliquots were prepared such as 50, 100, 150, 200, 250 and 300 μg/mL by diluting it. The percentage of free radical scavenging was calculated based on the absorbance on extent of reduction in the colour. The IC₅₀ value was calculated for the expression of antioxidant activities of the extracts (25).

The ability to scavenge the DPPH radical was calculated using the following equation:

% of radical scavenging activity = [(Ac-As)/Ac] × 100

Where Ac = Absorbance of control, As = Absorbance of sample.

Collection of Bacteria

The microbial strains employed in the study included "Staphylococcus aureus (MTCC-1430), Escherichia coli (MTCC-0614), Klebsiella pneumoniae (MTCC-109), Shigella flexneri (MTCC-1457), Micrococcus luteus (MTCC-1809)", procured from MTCC (Microbial Type Culture Collection Centre) and Gene Bank, Chandigarh, India. All the microbial strains were cultured and sub-cultured in nutrient agar slants. Then, the agar slant stored in refrigerator at 4°C for further experimental use.

Microbial culture media

The culture media for the antibacterial susceptibility analysis was done by using solid nutrient agar media applying on petridishes and the test was performed by using agar well diffusion method. To execute the antibacterial investigation, nutrients agar (28 g/L) "(HIMEDIA), REF- M001-500G, LOT-0000145979" was used for developing surface colony growth. The suspension culture for bacterial cells growth was done by preparing nutrient broth "(HIMEDIA), (REF-M088-500G, LOT- 0000154058)". The sterilization process was then followed for both nutrient solid agar media and broth culture media at 121 °C for 20 min. Sterilized cultured tube was used to make fresh media for slant preparation which was used to maintain bacterial strains.

Preparation of test plates

Sterile petridishes were kept in UV light for 20 min to avoid contamination. About 30 mL of nutrient solid media was poured in to each sterile petridishes in an aseptic condition making a depth of approximately 4 mm. The plate was kept in undisturbed condition to make them dry; then swabbing process was carried out using sterile cotton swab with 8 hrs old broth culture of each test microorganism. Then, plates were kept preserved to apply drug sample.

Anti-Bacterial Activity Assay Using Agar Well Diffusion Method

To determine the antibacterial assay of *H. verticillata* agar well diffusion method was followed (26). The wells in each of these swabbed plates were made by using sterile cork borer with 10 mm diameter and about 2 cm apart. About 50 µl of methanolic and n-hexane extract was poured with micropipette in agar well and allowed them to diffuse at room temperature for 2 hrs. without disturbed. Ciprofloxacin drug was taken as standard at concentration of 0.5 mg/mL. The inoculated plates were transferred to incubator at 37 °C for the growth of bacteria with triplicate for 18-20 hrs. The experiment was repeated thrice; the result was recorded in form of zone of inhibition in diameter in 3 different fixed directions.

Preparation of drug for antibacterial analysis

The stock solution (plant sample) for the drug assay was prepared as 500 mg/mL using dimethylsulphoxide (DMSO). Working solution of the drug was prepared by 1% of DMSO diluted with culture media and the fraction is taken as 15 mg/mL followed by 12 mg/mL, 9 mg/mL, 6 mg/mL and 3 mg/mL. "Ciprofloxacin" was taken as standard drug (Reference Antibiotic). The stock solutions of RA were prepared in 10 % DMSO to give a concentration of 0.5 mg/mL for RA. The diameter of zone of inhibition (in mm) was measured by using transparent scale.

Results and discussion

Preliminary Phytochemical analysis

The preliminary phytochemical study of the plant extracts revealed the presence of alkaloids, glycosides, flavonoids, saponins, steroids, triterpenoids, tannins and polyphenolic compounds (Table 1). The n-hexane extract showed the presence of alkaloids, triterpenoids and polyphenolic compounds, tannin, glycoside while phytoconstituents like alkaloids, saponins, tannins, steroids, flavonoid, glycosides, triterpenoids, tannins and polyphenolic compounds were found to be present in methanolic extract of *H. verticillata*. The result, thus indicated that the methanolic extract contains a greater number of phytoconstituent than the n-hexane solvent extract.

Table 1. Preliminary phytochemical analysis of Hydrilla vericillata

Sl.No	Phytochemicals	Hydrilla verticillata Ex- tracts in solvents		
		Methanol	n-Hexane	
1	Alkaloids	+ve	+ve	
2	Flavonoids	+++ve	-ve	
3	Saponins	++ve	-ve	
4	Steroids	++ve	-ve	
5	Tannins and polyphenolic compound	+++ve	++ve	
6	Triterpenoids +++ve		+ve	
7	Glycosides	+ve	+ve	

(+ve) indicating presence of chemical compounds , (-ve) indicating absence of chemical compounds $% \left(\frac{1}{2}\right) =0$

Natural remedies are now being center of attraction for curing any type of disease. Phenolic compounds present in medicinal plants can prevent from oxidative damage (27, 28). Earlier it was reported that the mutagenesis and carcinogenic effect could be reduced by the act of phenolic compounds, when consumed through vegetables and fruits (29). Tannin content has been considering another important phytoconstituent which can scavenge free radicals. Thus, they reduce hazardous free radicals from our body (29).

Antioxidant Assay

The result of antioxidant assay is summarized in Fig. 1. The free radical scavenging activity of both methanolic and n-hexane extract of *H. verticillata* with different concentra-




Fig. 1. Antioxidant assay of *H. verticillata* with standard Ascorbic Acid.

tion ranges from 50 µg/mL-300µg/mL was evaluated in dose dependent manner. The effect of antioxidant on free radicals is thought to be because of their hydrogen donating capacity. In DPPH assay, the reduction capability of "1, 1-diphenyl-2-picrylhydrazyl" was determined by decrease absorbance at 517 nm. Absorbance was decreased significantly by increasing the concentration of the crude extract. Whole experiment was conducted by taking ascorbic acid as standard. From the above experiment it was found that the methanolic extract showed comparatively higher antioxidant activity than n-hexane extract. The half maximal inhibitory concentration (IC50) value is widely used to calculate the antioxidant activity. The IC50 value of n-hexane extract, methanol extract and ascorbic acid was found 36.86, 16.71 and 12.63 respectively.

Present study was conducted to evaluate antioxidant competence of *H. verticillata* which is beneficial at both nutritional levels as well pharmaceutical sectors. Phytoconstituent like phenolic compounds, flavonoid, tannin, alkaloid and glycosides are believed to be responsible to control disease related to stress and are considered as good antioxidant substances (30). Generally, salt tolerant macrophytes are reported to have a greater number of phytochemicals and with higher capacity to scavenge oxidative damage than the normal land plants (31). *Hydrilla verticillata* was also previously proven to enhance the activity of superoxide dismutase and peroxidase after expose to saline conditions (32).

Antibacterial Analysis

The extracts of this plant subjected to antibacterial screening against 5 bacterial pathogens causing different human diseases. The results indicated that methanolic extract of *Hydrilla verticillata* exhibited highest zone of inhibition against *Shigella flexneri* (26±0.75 mm) and least against *Klebsiella pneumoniae* (20±1.24 mm) in concentration of 15 mg/mL. This extract showed moderate activity against *Micrococcus luteus* (22±0.53 mm) and *Escherichia coli* (21±1.21 mm). While n-hexane extract exhibited highest zone of inhibition on *Staphylococcus aureus* (20±1.41 mm)

at concentration of 15 mg/mL of drug, least against *Micrococcus luteus* (12±1.63 mm) and moderate against *Escherichia coli* (18±0.47 mm) and *Klebsiella pneumoniae* (16±1.23 mm) (Table 2; Fig. 2 and 3). The GC-MS study revealed that the presence of "phytol and hexadecanoic acid ethyl ester" as major component in ethanolic extract of *Hydrilla* (33). This may play a major role for antimicrobial and antioxidant properties.




Fig. 2. Antibacterial assay of H. verticillata with methanolic extract.




Fig. 3. Antibacterial assay of *H. verticillata* with n-Hexane extract.

 $\textbf{Table 2.} \textit{In vitro} \ \textbf{Antibacterial activity (zone of inhibition) of different extracts of } \textit{Hydrilla verticillata}$

Test substance	Concentration	Diameter of zone of inhibition (mm)				
		Staphylococcus aureus	Shigella flexneri	Escherichia coli	Micrococcus luteus	Klebsiella pneumoniae
Ciprofloxacin 0.5mg/mL (RA)	0.5 mg/mL	37±0.21	33±0.43	36±0.63	29±0.51	32±0.76
Methanol extract	15 mg/mL	24±0.46	26±0.75	21±1.21	22±0.53	20±1.24
	12 mg/mL	22±0.81	24±0.61	20±0.47	19±0.47	19±0.21
	9 mg/mL	21±0.94	23±0.81	18±1.41	18±0.4	16±0.48
	6 mg/mL	20±0.84	18±0.82	15±1.24	17±1.69	14±1.31
	3 mg/mL	18±1.24	16±0.24	14±1.62	13±0.23	13±0.38
n-Hexane extract	15 mg/mL	20±1.41	14±0.81	18±0.47	12±1.63	16±1.23
	12 mg/mL	18±o.81	13±0.47	16±0.46	11.66±0.94	15±0.48
	9 mg/mL	14±0.52	13±0.21	15±0.51	11±0.29	14±0.61
	6 mg/mL	12±0.12	12±0.65	14±0.42	11±0.11	12±0.21
	3 mg/mL	12±0.31	11±1.21	13±0.62		11±0.68

Result expressed as mean ±S.D (n=3), (---) denotes no zone of inhibition

Conclusion

Results from the present investigation revealed that methanolic extract of Hydrilla verticillata had a strong antibacterial effect against all the test pathogenic bacteria and also showed good results on antioxidant activity. In antibacterial study; methanolic extract showed highest zone of inhibition against Shigella flexneri whereas n-hexane showed good result against Staphylococcus aureus. In study of DPPH assay, the IC₅₀ value was found to be lowest in methanolic extract than n-hexane. From these inhibitory effects of the extracts of *H. verticillata* was proven that it can be used as potential herbal drugs against all tested pathogenic organisms and also a good antioxidant product. In future, compound isolation and analysis can be examined for its specific antibacterial and anti oxidant activity. As it has great nutritional value as well pharmaceutical value, this plant can be recommended as the super food for human.

Acknowledgements

The authors are thankful to the administration and management of Centurion University of Technology and Management, Odisha, India, for their support during the investigation.

Authors contributions

Kunja Bihari Satapathy and Bhagyeswari Behera conceived the idea. Bhagyeswari Behera performed the experiments. Kunja Bihari Satapathy and Bhagyeswari Behera analyzed the information. Both authors have made significant contributions in drafting the manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None.

References

- Kamali HH, Amir MYEL. Antibacterial activity and phytochemical screening of ethanolic extracts obtained from selected sudanese medicinal plants. Current Research Journal of Biological Sciences. 2010; 2(2):143-46.
- Lalitha P, Arathi KA, Shubasini K, Hemalatha S, Jayanti P. Antimicrobial activity and phytochemical screening of an ornamental foliage plant, *Pothos aurea* (Linden ex Andre). An International Journal of Chemistry. 2010;1(2):63-71.
- 3. Hussain H, Badawy A, Elshazly A, Krohn K, Riaz M, Schulz B. Chemical constituents and antimicrobial activity of *Salix subserrata*. Records of Natural Product. 2011;5(2):133-37.
- Preethi R, Devanathan VV, Loganathan M. Antimicrobial and antioxidant efficacy of some medicinal plants against food borne pathogens. Advance In Biological Research. 2010;4(2):122 -25.
- Sharma H. Antibacterial activity of ethanolic extracts of some arid zone plants. International Journal of Pharma Tech Research. 2011;3(1):283-86.

- Enne VL, Livermore DM, Stephens P. Persistence of sulphonamide resistance in *Escherichia coli* in the UK despite national prescribing restriction. The Lancet. 2001;28:1325-28. https://doi.org/10.1016/S0140-6736(00)04519-0
- Westh H, Zinn CS, Rosdah VT. An international multicenter study of antimicrobial consumption and resistance in *Staphylococcus* aureus isolates from 15 hospitals in 14 countries. Microbial Drug Resistance. 2004;10:169-76. https:// doi.org/10.1089/1076629041310019
- Davies J, Davies D. Origins and evolution of antibiotic resistance. Microbiology and Molecular Biology Reviews. 2010;74
 (3):417-33. https://doi.org/10.1128/MMBR.00016-10
- 9. Khatoon A, Arzoo A, Mohapatra A, Satapathy KB. Studies on *in vitro* evaluation of antibacterial activities of *Eucalyptus globulus* Labill leaf. International Journal of Current Pharmaceutical Research. 2017;9:140-42.
- Khan A, Aung TT, Chaudhuri D. The first case of native mitral valve endocarditis due to *Micrococcus luteus* and review of the literature. Case Report in Cardiology. 2019;5907319. Published Dec 4. https://doi.org/10.1155/2019/5907319
- Anne, Eady E, Patricia C, Jeremy I, Ross, Hamid A, Ratyal, Jonathan H. Antibiotic resistance patterns of aerobic coryneforms and furazolidone-resistant gram-positive *Cocci* from the skin surface of the human axilla and fourth toe cleft. Journal of Antimicrobial Chemotherapy. 2000;46(2):205-13. https://doi.org/10.1093/jac/46.2.205
- Liebl W, Kloos WE, Ludwig W. Plasmid-borne macrolide resistance in *Micrococcus luteus*. Microbiology (Reading). 2002;Aug;148(Pt 8):2479-87. doi: 10.1099/00221287-148-8-2479. PMID: 12177341. https://doi.org/10.1099/00221287-148-8-2479
- Ramírez-Castillo FY, Moreno-Flores AC, Avelar-González FJ, Márquez-Díaz F, Harel, Guerrero-Barrera AL. An evaluation of multidrug-resistant *Escherichia coli* isolates in urinary tract infections from Aguascalientes, Mexico: cross-sectional study. Annals of Clinical Microbiology and Antimicrobials. 2018;17 (1):34. https://doi.org/10.1186/s12941-018-0286-5
- Kim H, Park SW, Park JM, Moon KH, Lee CK. Screening and isolation of antibiotic resistance inhibitors from herb material Resistance Inhibition of 21 Korean plants. Natural Product Science. 1995;1:50-54.
- 15. Alagesaboopathi C. Antimicrobial potential and phytochemical screening of *Andrographis affinis* Nees an endemic medicinal plant from India. International Journal of Pharmacy and Pharmaceutical Sciences. 2011;3(2):157-59.
- 16. Ahmad, Beg AZ. Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multiple drug resistant human pathogens. Journal of Ethnopharmacology. 2011;74:113-23. https://doi.org/10.1016/S0378-8741(00)00335-4
- 17. Lal AA, Murthy PB, Pillai KS. Screening of hepatoprotective effect of a herbal mixture against CCl4 induced hepatotoxicity in Swiss albino mice. Journal of Environmental Biology. 2007;28:201-07.
- Rasoanaivo P, Wright CW, Willcox WL, Gilbert B. Whole plant extracts versus single compounds for the treatment of malaria: synergy and positive interactions. Malaria Journal. 2011;10:1-12. https://doi.org/10.1186/1475-2875-10-S1-S4
- Pal DK, Nimse S. Little known uses of common aquatic plant, *Hydrilla verticillata* (Linn. f.) Royle. Natural Product Radience. 2006;5:108-11.
- Saxena HO, Brahmam MM. The flora of Orissa. Orissa Forest Development Corporation, Bhubaneswar, Odisha. 1-4:1994-96.
- Khatoon, A, Jethi S, Nayak SK, Sahoo S, Mohapatra A, Satapathy KB. Evaluation of *in vitro* antibacterial and antioxidant activities of *Melia azedarach* L. bark. IOSR. Journal of Pharmacy and Biological Sciences. 2014;9(6):14-17. https://doi.org/10.9790/3008-09641417

- Siddiqui S, Verma A, Rather AA, Jabeen F, Meghvansi MK. Preliminary phytochemicals analysis of some important medicinal and aromatic plants. Advance in Biological Research. 2009;3(5-6):188-95.
- 23. Harborne JB. Phytochemical methods-A guide to modern technique of plant analysis. Chapman and Hall, UK, 3rd ed. 1998;1-5.
- 24. Kokate CK. Practical Pharmacognosy. 4th ed., Vallabh Prakashan, New Delhi. 2005;7-9.
- Liyanapathirana C, Shahidi F. Optimization of extraction of phenolic compounds from wheat using response surface methodology. Food Chemistry. 2005;93(1):47-56. https://doi.org/10.1016/j.foodchem. 2004.08.050
- Balouiri M, Sadiki M, Ibnsouda SKN. Methods for *in vitro* evaluating antimicrobial activity: A review. Journal of Pharmaceutical Analysis. 2016;6:71-79. https://doi.org/10.1016/j.jpha.2015.11.005
- Silva BA, Ferreres F, Malva JO, Dias ACP. Phytochemical and antioxidant characterization of *Hypericum perforatum* alcoholic extracts. Food Chemistry. 2005;90:157-67. https:// doi.org/10.1016/j.foodchem.2004.03.049
- Meena H, Pandey HK, Pandey P, Arya MC, Ahmed Z. Evaluation of antioxidant activity of two important memory enhancing medicinal plants *Bacopa monnieri* and *Centella asiatica*. Indian Journal of Pharmacology. 2012;44(1):114-17. https://doi.org/10.4103/0253-7613.91880

- 29. Tanaka M, Kuei CW, Nagashima Y, Taguchi T. Application of antioxidative maillrad reaction products from histidine and glucose to sardine products. Nippon Suisan Gakkaishi. 1998;54:1409-14. https://doi.org/10.2331/suisan.54.1409
- 30. Tyagi T, Agarwal M. Phytochemical screening and GC-MS analysis of bioactive constituents in the ethanolic extract of *Pistia stratiotes* L. and *Eichhornia crassipes* (Mart.) Solms. Journal of Pharmacognosy and Phytochemistry. 2017;6(1):195-206.
- Muthazhagan K, Thirunavukkarasu P, Ramanathan T, Kannan D. Studies on phytochemical screening, antimicrobial and antiradical scavenging effect coastal salt mash plant of a Suaeda monoica. Research Journal of Phytochemistry. 2014;8:102-11. https://doi.org/10.3923/rjphyto.2014.102.111
- Rout NP, Shaw BP. Salt tolerance in aquatic macrophytes: possible involvement of the antioxidative enzymes. Plant Science. 2001;160(3):415-23. https://doi.org/10.1016/S0168-9452(00) 00406-4
- Prabha PS, Rajkumar J. Phytochemical screening and bioactive potential of *Hydrilla verticillata*. Journal of Chemicals and Pharmaceutical Research. 2015;7(3):1809-15.