



RESEARCH ARTICLE

# Molecular diversity in maize inbred lines for turcicum leaf blight resistance

D. Keerthana<sup>1\*</sup>, T. Haritha<sup>1</sup>, I. Sudhir Kumar<sup>1</sup> & D. Ramesh<sup>2</sup>

<sup>1</sup> Department of Genetics and Plant Breeding, Acharya N G Ranga Agricultural University, Andhra Pradesh, India

<sup>2</sup> Department of Statistics and Computer Applications, Acharya N G Ranga Agricultural University, Andhra Pradesh, India

\*Email: [keerthanadhanapati@gmail.com](mailto:keerthanadhanapati@gmail.com)



## ARTICLE HISTORY


Received: 30 December 2022

Accepted: 05 March 2023

Available online

Version 1.0 : 12 May 2023

Version 2.0 : 01 July 2023



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Keerthana D, Haritha T, Sudhir Kumar I, Ramesh D. Molecular diversity in maize inbred lines for turcicum leaf blight resistance. Plant Science Today. 2023;10(3): 385–392. <https://doi.org/10.14719/pst.2368>

## Abstract

Maize (*Zea mays* L.), third most important cereal crop in the world, its productivity can be limited partly by an important foliar disease called turcicum leaf blight caused by *Exserohilum turcicum*. In order to design efficient breeding programmes for resistance to leaf blight, the germplasm must be thoroughly characterized. This study evaluated the diversity of maize inbred lines using 26 simple sequence repeat (SSR) primers. The polymorphism information content (PIC) value of the SSR loci ranged from 0.61 to 0.71, with an overall mean of 0.65. It was highest for the primer bnl1335 (0.71) and lowest for the primer bnl1666 (0.61). The markers produced one to four alleles, with an average of 2.5 alleles per marker. Using the DARwin 6.0 programme, the inbred lines were grouped into different clusters. The cluster A was solitary, with the inbred line VL171488-2 having a resistant reaction against leaf blight. This line can be used in crossing programmes with divergent parents to develop leaf blight resistant hybrids. The results of this study can be used to design efficient breeding programmes for resistance to leaf blight.

## Keywords

Dendrogram, Jaccard's similarity coefficient, maize, SSRs, TLB

## Introduction

Maize is an important cereal crop in the world for feed, food, fodder and raw materials of industries (1). Maize, coupled with rice and wheat, supplies at least 30% of the calories consumed by approximately 4.5 billion people in 94 developing countries (2). In India, it is cultivated over an area of 9.89 m ha with a production and productivity of 31.60 million tonnes and 3199 kg ha<sup>-1</sup> respectively. In Andhra Pradesh, maize is cultivated in an area of 0.3 m ha, with a production and productivity of 1.78 million tonnes and 5917 kg ha<sup>-1</sup> respectively (3).

The turcicum leaf blight, commonly known as northern leaf blight, is a foliar disease of maize caused by the ascomycetes fungus, *Setosphaeria turcica* and its conidial stage *Exserohilum turcicum* (Pass.) Leonard and Suggs. (syn. *Helminthosporium turcicum* Pass) affects the photosynthesis and reduces kernel yield to an extent of 28 to 91% (4, 5). The disease is characterized by long elliptical, greyish green or brown leaf lesions which emerge first on the lower leaves and gradually extend throughout the foliage. If the disease starts at an early stage, it causes premature death of blighted leaves. As a result, the crop loses their nutritive value as fodder, have reduced germination capacity, vigor, grain yield, total sugar content,

restricted starch formation, chaffy kernels and infected plants are liable to infection with stalk rots. Several disease management options have been recommended to

**Table 1.** List of parental lines used in the present study

S. No.	Entry	Source of origin	S. No.	Entry	Source of origin
<b>Lines</b>			<b>Testers</b>		
1	VL171488-2	ARS, Peddapuram	12	BML6	MRC, Rajendranagar
2	VL18828	ARS, Peddapuram	13	BML7	MRC, Rajendranagar
3	VL19978-6	ARS, Peddapuram	14	LM13	PAU, Ludhiana
4	VL19705-8	ARS, Peddapuram	15	LM14	PAU, Ludhiana
5	VL19255	ARS, Peddapuram	<b>Checks</b>		
6	VL18142	ARS, Peddapuram	16	P3396	Pioneer
7	CAL1733-13	ARS, Peddapuram	17	DKC8171	Monsanto Bayer
8	VL175869-14	ARS, Peddapuram	18	P3546	Pioneer
9	SNL19564-20	ARS, Peddapuram	19	DKC9120	Monsanto Bayer
10	SNL19582-22	ARS, Peddapuram	20	PAC751	Adventa
11	SNL19588-23	ARS, Peddapuram			

reduce the impact of maize foliar diseases. Among these practices, planting of resistant cultivars can effectively reduce the rate of disease development and is widely recommended. Breeding for resistance is a practical, cost-effective means to manage the disease (6). The development of resistance against turcicum leaf blight will have large effect on the maize crop improvement programmes.

Genetic diversity using molecular markers provides the potential to examine changes in crop diversity over time and place. Among the several PCR-based markers, microsatellite markers based on Simple Sequence Repeats (SSRs) are widely exploited due to their ease of use, reproducibility, rapid analysis, low cost, simple scoring patterns and greater allelic diversity (1). "In the present era, DNA-based markers (molecular markers) are frequently used for genetic diversity and grouping of the populations. Among the various high throughput DNA-based marker techniques available, single nucleotide polymorphisms (SNPs) and simple sequence repeats (SSRs) are the markers of choice because they are co-dominant in nature, locus-specific, reproducible, highly informative and easy to use" (1). Moreover, SSR markers are more informative than biallelic SNP markers because they can detect multiple alleles per locus. It was reported that SSRs are 7 to 11 times more accurate than SNPs (7). "The SSR markers will allow detection of polymorphisms at the DNA level which will facilitate the separation of inbreds at DNA level by the estimates of genetic distance" (8).

## Materials and Methods

### Field Evaluation

A total of 15 parental lines along with 5 checks were screened against TLB under artificial epiphytotics using sorghum grain inoculation technique at Agricultural college farm, Bapatla, Andhra Pradesh, situated at 15°55' North latitude and 80°30' East longitude and an altitude of 5.49 m above mean sea level. Each entry is planted in one row of 1.6 meters length, adopting a spacing of 60 x 20 cm for screening against *Exserohilum turcicum*. The block was

flanked by 3 border rows of susceptible cultivar, P3396. List of entries used in the present investigation are presented in Table 1.

Turcicum leaf blight severity was recorded on five plants in each entry at the time of tasseling, twenty days after tasseling and at maturity using 1-9 disease rating scale (9, 10) and is presented in Table 2.

### Molecular Evaluation

#### DNA Isolation

DNA was extracted from the leaf tissue collected at 15 days after sowing from all the genotypes using the method described by (11, 12) with certain modifications.

#### Principle

The prerequisite for DNA isolation is to extract DNA from the cells. Addition of the extraction buffer containing CTAB assists breakage of nuclear membrane and cell lysis. This results in the dispersion of all the cell components in to the buffer. Tris HCl assures that the pH of the solution is maintained around 8.0. EDTA forms complexes (chelates) with the several metal ions such as magnesium and calcium which are required as cofactors by the majority of DNases. NaCl being a salt decreases the solubility of DNA in the buffer solution and also increases the osmoticity of the buffer to facilitate the process of cell lysis. Phenol: chloroform: Isoamyl alcohol (25:24:1) removes proteins by denaturing them and aggregate in the intermittent phase along with cell debris. For DNA precipitation isopropanol is used. RNA is removed by the RNase treatment. After precipitation the pellet is washed with 70% ethanol for removing any salts retained after precipitation. The DNA is preferably dissolved in a base solution like Tris EDTA rather than in sterile distilled water since Tris solution provides buffering for the DNA and EDTA in the buffer ensures that the DNA is stable by inactivating DNases.

#### Simple Sequence Repeats (SSRs)

A total of 26 primers was used to access the genetic diversity in 15 parental lines along with 5 checks. The primer information for the selected SSR markers was retrieved from the maize GDB database and is available in public domain (13) and summary of the SSR primers used in the present study are given in Table 3.

**Table 2.** Disease scoring scale (1-9) for turicum leaf blight

Rating scale	Degree of infection (per cent DLA*)	PDI**	Disease reaction
1.0	Nil to very slight infection ( $\leq 10\%$ ).	$\leq 11.11$	Resistant (R) Score: $\leq 3.0$ DLA: $\leq 30\%$ PDI: $\leq 33.33$
2.0	Slight infection, a few lesions scattered on two lower leaves (10.1-20%).	22.22	
3.0	Light infection, moderate number of lesions scattered on four lower leaves (20.1-30%).	33.33	
4.0	Light infection, moderate number of lesions scattered on lower leaves, a few lesions scattered on middle leaves below the ear (30.1-40%).	44.44	Moderately resistant (MR) Score: 3.1-5.0 DLA: 30-50% PDI: 33.34-55.55
5.0	Moderate infection, abundant number of lesions scattered on lower leaves, moderate number of lesions scattered on middle leaves below the ear (40.1-50%).	55.55	
6.0	Heavy infection, abundant number of lesions scattered on lower leaves, moderate infection on middle leaves and a few lesions on two leaves above the ear (50.1-60%).	66.66	Moderately susceptible (MS) Score: 5.1-7.0 DLA: 50.1-70% PDI: 55.56-77.77
7.0	Heavy infection, abundant number of lesions scattered on lower and middle leaves and moderate number of lesions on two to four leaves above the ear (60.1-70%).	77.77	
8.0	Very heavy infection, lesions abundant scattered on lower and middle leaves and spreading up to the flag leaf (70.1-80%).	88.88	Susceptible (S) Score: $>7.0$ DLA: $>70\%$ PDI: $>77.77$
9.0	Very heavy infection, lesions abundant scattered on almost all the leaves, plant prematurely dried and killed ( $>80\%$ ).	99.99	

**Table 3.** List of SSR primers used for PCR amplification in the present investigation

The below SSR primers are selected based on the earlier reports of (1) (25)

S. No	Primer	F/R	Sequence	T <sup>m</sup>
1	bnlg198	F	GTTTGGTCTTGCTGAAAAATAAAA	50.9
		R	GCTGGAGGCCTACATTATTATCTC	54.8
2	bnlg1335	F	GAAGGTTGCTCTTCCACTGG	55.8
		R	TGGTTTGTGCAAGTGTCACC	56.1
3	umc2210	F	GATGCTACCATTTCAGTGAGCGAT	57.2
		R	AGCGGGTCGATCTTCTCTTAGTT	57.7
4	umc1665	F	CAATCAGGAGCCAGGGAGATG	57.5
		R	CTTAAACTGTGCGAGACGGTCCTG	57.3
5	umc1029	F	AACACCTGCTGGATATGGATCACT	58.0
		R	GGAAGAAAAATGTCGACCTGCTC	56.2
6	bnlg1666	F	GCTGGTAGCTTTCAGATGGC	56.2
		R	TGTCCCTCCTCAGTTTCAC	56.5
7	bnlg240	F	AAGAACAGAAGGCATTGATACATAA	52.6
		R	TGCAGGTGTATGGGCAGCTA	58.8
8	umc1728	F	AGTACTTTCAGGCAGGGACCTTCT	59.6
		R	AACGCACTTCTGTAGCTGTAGGG	58.9
9	phi056	F	ACTTGCTTGCCTGCCGTTAC	58.5
		R	CGCACACCACTCCAGAA	57.9
10	umc1293	F	GTATCCGTTTCTCATGCAACACAC	56.4
		R	GATCTCGATCTGCTTCATCATCTG	55.1
11	umc1811	F	AGATAGCCGCCGAGACCAAG	55.1
		R	ACTCACTCGACGGACTTCTCGAC	53.5
12	umc1144	F	ATGGCCCACTCATCATATCTCTGT	56.0
		R	TGTGTTGATTAGCAGCGGATAAAA	57.8
13	umc1551	F	CACCGGAACACCTTCTTACAGTTT	63.4
		R	CGAAACCTTCTCGTGATGAGC	65.1
14	umc2077	F	AAACTCACTGAACATGATCCTGGC	54.8
		R	CTGGTTCGGATGCAAGTAGTCAG	56.6

15	umc1086	F	CATGAAAGTTTTCTGTGCAGATT	65.0
		R	GGGCAACTTTAGAGGTCGATTTATT	67.7
16	umc1559	F	CTTGCTAGAGTCGGTGAACAACAA	56.1
		R	AACCAAGCTCCTTAATGAGGTCAC	56.9
17	umc1644	F	CCATAAACTGTTCTTTGGCACAC	58.3
		R	CTTTCACGTGTTAAGGGAGACACC	56.2
18	umc2169	F	ACTACTCCTCGGATAGCCACG	53.7
		R	GACGAGTAGAGGCTCTGGGAC	56.0
19	bnlg2086	F	CGGAACCTGCTGCAGTTAAT	55.9
		R	GAGATGCAGGAATGGGAAAA	53.7
20	phi330507	F	GTAAGTACGATGCGCCTCCC	58.0
		R	CGGGGTAGAGGAGAGTTGTG	56.9
21	umc1568	F	AAGTCCAGCCAAGTTCATCAAGA	56.8
		R	ACTGTAATAAACTGGGTGTGCC	59.0
22	nc003	F	ACCCTTGCCTTTACTGAAACACAACAGG	61.0
		R	GCACACCGTGTGGCTGGTTC	61.8
23	umc2214	F	ACCCCTGATTCTCTTACGTTT	57.6
		R	CTGGATGAGGAGGAAGAATACGAG	56.3
24	phi085	F	AGCAGAACGGCAAGGGCTACT	61.3
		R	TTTGGCACACCACGACGA	57.5
25	bnlg238	F	CTTATTGCTTTCGTCATACACACATTCAT	57.7
		R	GAGCATGAGCTTGCATATTTCTGTGG	58.5
26	phi054	F	AGAAAAGAGAGTGTGCAATTGTGATAGAG	60.8
		R	AATGGGTGCCTCGACCAAG	56.9

F – forward primer, R- Reverse primer, T<sup>m</sup> – Annealing temperature

### Genomic DNA amplification in PCR using SSR markers

The polymerase chain reaction (PCR) was carried out using a programmable thermocycler (prima 96™, HIMEDIA). Amplifications were performed in 10 µl reaction mixture having the following components 3.00 µl of DNA template (50 ng/µl); 1.50 µl of Taq buffer (10X); 0.35 µl of DNTPs (2.5 mM); 0.75 µl of forward and reverse primer each; 0.25 µl of Taq polymerase (5 U/µl); 7.40 µl of Autoclaved double distilled H<sub>2</sub>O. Amplification was performed in a Himedia Thermal Cycler programmed as mentioned below

### Agarose gel electrophoresis for resolution of SSR markers

Amplified products thus obtained were separated on 3% agarose gel using horizontal gel electrophoresis assembly at 80-100 volt for 2 h in 1X TAE buffer. The DNA fragments were then visualized under UV-transilluminator and documented using gel documentation system (SYNGENE Gene flash, U.K.)

### Analysis of SSR data

#### Scoring of data

Data were scored for computer analysis on the basis of the presence or absence of the PCR products. If a product was present in a genotype, it was designated as '1' and if absent it was designated as '0'. The data were entered into

MS-Excel data sheet. The data were maintained in the data sheet format for further analysis. The Microsoft Excel 2013 (Microsoft, USA) sheet was used for further statistical analysis.

### Cluster analysis

The genetic associations between the genotypes were calculated by using the Jaccard's similarity coefficients for pair wise comparisons based on the proportions of shared bands produced by the primers (14). Similarity matrix was generated using DARwin software 6.0 version (15). The similarity coefficients were used for cluster analysis and dendrogram was constructed by Unweighted Pair-Group Method with Arithmetic Average (UPGMA).

### Determination of Polymorphism Information Content (PIC)

The Polymorphism Information Content (PIC) value for each locus was calculated on the basis of allele frequency by the given formula (16).

$$PIC = 1 - \sum_{i=1}^n (P_{ij})^2$$

Where, P<sub>ij</sub> is the frequency of the j<sup>th</sup> band for marker i, and summation extends over n<sup>th</sup> band.

## Results and discussion

### Field Screening

Among the inbreds, six lines viz., VL171488-2, VL18828, VL19705-8, VL18142, VL175869-14, SNL19582-22 and one tester, LM13 were categorized as resistant with disease score  $\leq 3.0$ . Three lines viz., VL19978-6, VL19255, SNL19588-23 and one tester, LM14 were categorized as moderately resistant lines. One line, CAL1733-13 and two male testers, BML6 and BML7 were categorized as moderately susceptible lines. Only one line, SNL19564-20 with disease score 7.5 was categorized as susceptible to disease TLB. Disease reaction of entries were presented in Table 4.


**Table 4.** Disease reaction of 15 parental lines and 5 checks

S. No.	Entries	Disease score	PDI	Disease Reaction	S. No.	Entries	Disease score	PDI	Disease Reaction
<b>Lines</b>					<b>Testers</b>				
1	VL171488-2	3	6.7	R	12	BML6	5	11.1	MR
2	VL18828	3	6.7	R	13	BML7	4	8.9	MR
3	VL19978-6	4	8.9	MR	14	LM13	3	6.7	R
4	VL19705-8	3	6.7	R	15	LM14	4	8.9	MR
5	VL19255	3.5	7.8	MR	<b>Checks</b>				
6	VL18142	2	4.4	R	16	P3396	6	13.3	MS
7	CAL1733-13	6	13.3	MS	17	DKC8171	6	13.3	MR
8	VL175869-14	2	4.4	R	18	P3546	3	6.7	R
9	SNL19564-20	5	11.1	MR	19	DKC9120	4	8.9	MR
10	SNL19582-22	2.5	5.6	R	20	PAC751	4	8.9	MR
11	SNL19588-23	4	8.9	MR					

### SSR Polymorphism

To know the extent of genetic diversity among 20 genotypes, DNA profiling of these genotypes was generated by using SSR markers which were reported to be associated with TLB resistance in maize. Out of 26 SSR primers, only

informative. The similar results were obtained earlier (17) having PIC range from 0.58 to 0.81 with mean of 0.71; (18) with mean PIC of 0.54; (19) having PIC range from 0.32 to 0.85 with mean of 0.68 and (1) having PIC range 0.04 to 0.86.



**Fig. 1.** SSR marker profiles of 15 parental inbreds along with 5 checks generated by various primers.

### Similarity Index

The binary data from the polymorphic primers were used for computing Jaccard's similarity indices. The similarity index values obtained for each pair wise comparison among the 20 genotypes. The overall range of similarity among the 20 genotypes was found to be 0.17 to 0.78. Among the 20 genotypes the highest similarity index (0.78) was observed between BML7 and LM14 followed by (0.77) between LM13 and BML7 indicating that they are genetically less divergent (or) more similar as compared to other genotypes. The lowest similarity index (0.17) was observed between SNL19582-22 and VL171488-2 indicating that they are more divergent (or) less similar.

Similar results were obtained earlier (20) found genetic similarities ranged from 0.27 to 0.99 and (21) found similarity matrix value ranges from 0.17 to 0.84.

### Cluster Analysis


The similarity coefficients during the present investigation were used as input data for UPGMA based clustering method. Twenty inbreds were grouped into 2 major clusters (A and B). Cluster A (VL171488-2) was solitary and cluster B was subdivided into ten subclusters at 0.52 Jaccards similarity coefficient (Fig. 2). Cluster A consists of single inbred.

These results were in accordance with earlier work categorization of genotypes into 3 distinct clusters (22); noted 3 distinct clusters (23); categorization of the genotypes into 4 clusters (16) and categorization of genotypes into 3 clusters based on UPGMA dendrogram (24).

### TLB Resistance and SSR markers

The field screening of these genotypes identified very good variability for resistance indicating the importance of these genotypes in the breeding programmes. The inbred lines, VL171488-2, VL18828, VL19705-8, VL18142, VL175869-14, SNL19582-22 and LM13 recorded resistance reaction to the pathogen. Most of the reports indicated the involvement of more number of genes and the polygenic nature of inheritance of this trait. When these genotypes were tested for polymorphism using SSR markers, they showed very good variation in the banding pattern indicating their usefulness in differentiating the genotypes.

All the resistant genotypes produced a band of 120bp for the marker, bnlg 1335 and 160bp for the marker, bnlg 2086 except in the genotype, VL174488-2 indicating these sequences may not be involved in the resistance to TLB in maize. The primer pair, umc1029, produced a specific band of 80bp in the genotype, VL18828 and a 200bp



**Fig. 2.** UPGMA dendrogram generated based on Jaccards similarity index using SSR markers showing variability among 15 parental inbreds in maize (*Zea mays* L.).

The subcluster I of cluster B consists of 3 inbreds viz., VL18828, VL19978-6, VL19705-8; Subcluster II consists of single genotype VL19255; Subcluster III consists of 2 inbreds viz., VL18142, CAL1733-13; Subcluster IV, V, VI and VII were solitary clusters. VIII subcluster was the largest having 6 genotypes of 4 testers BML6, BML7, LM13, LM14 and 2 checks P3396 and DKC8171; IX subcluster had 2 genotypes P3546 and DKC9120 and the subcluster X had the single genotype *i.e* PAC751.

band in the genotype, P3546. The markers, bnlg238 and bnlg1666, produced 170bp and 60bp bands only in the inbreds, VL18142 and SNL19582-22 and VL19705-8 & VL175869 respectively, indicating these markers are useful for the identification of these genotypes from others and the new sequences present in these genotypes may be associated with the TLB resistance.

Thus, the present study identified lot of variability in the expression of the reported markers associated with

TLB resistance in resistant genotypes indicating the quantitative inheritance and involvement of a greater number of alleles. Further, there is a need to use a greater number of TLB associated markers to get the clear picture of resistance in maize.

## Conclusion

In the present study 26 SSR markers were used for assessment of molecular diversity in 15 parental inbred lines along with 5 checks. Out of these primers seven primers viz., bnlg1335, bnlg1666, bnlg2086, bnlg238, Phi054, umc1029 and umc2077 recorded PIC value more than 0.6; These markers also noted lot of variation in the resistant genotypes indicating their usefulness. Further these markers also produced varied expression of bands in the resistant genotypes indicating the polygenic nature of the trait. Further, these resistant sources can be exploited in developing a novel maize genotypes with desirable yield and TLB resistance.

## Acknowledgements

All authors thank to Acharya N.G Ranga Agricultural University and ARS Peddapuram for providing all the facilities in the form of lab and stipend for successful completion of research and Associate Professor Dr. Lal Ahamed for his guidance in manuscript writing.

## Authors contributions

DK carried out the molecular genetic studies and drafted the manuscript. TH supervised the study and participated in design and coordination. ISK participated in gathering of primers for molecular analysis and sufficient guidance throughout the research work. DR performed the statistical analysis and necessary suggestions. All authors read and approved the final manuscript.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** None.

## References

- Kumar B, Choudhary M, Kumar P, Kumar K, Kumar S, Singh BK et al. Population structure analysis and association mapping for turcicum leaf blight resistance in tropical maize using SSR markers. *Genes*. 2022;13(4):618. <https://doi.org/10.3390/genes13040618>
- Shiferaw B, Prasanna BM, Hellin J, Banziger M. Crops that feed the world 6. Past successes and future challenges to the role played by maize in global food security. *Food Security*. 2011;3(3):307-27. <https://doi.org/10.1007/s12571-011-0140-5>
- Ministry of Agriculture, Government of India. *Indiastat*. 2020-21. <https://www.indiastat.com>.
- Bindhu KG, Pandurangegowda KT, Lohithaswa HC, Madhuri R, Mallikarjuna N. Genetics of resistance to turcicum leaf blight caused by *Exserohilum turcicum* (Pass.) Leonard and Suggs in maize (*Zea mays* L.). *International Journal of Current and Applied Microbiology*. 2017;6(11): 964-69. <http://doi.org/10.18782/2320-7051.6378>
- Robert AL. Some of the leaf blights of corn. Year Book of Agriculture. United States Department of Agriculture, North Carolina. 1953;380-85. <https://naldc.nal.usda.gov/download/IND43894350/pdf>
- Fehr WR. Principles of Cultivar Development. Macmillan, co., New York, USA. 1987.
- Van Inghelandt D, Melchinger AE, Lebreton C, Stich B. Population Structure and genetic diversity in a commercial maize breeding program assessed with SSR and SNP markers. *Theoretical and Applied Genetics*. 2010;120:1289-99. <https://doi.org/10.1007/s00122-009-1256-2>
- Menkir A, Melake BA, Ingelbrecht CI, Adepoju A. Grouping of tropical mid-altitude maize inbred lines on the basis of yield data and molecular marker. *Theoretical Applied Genetics*. 2004;108: 1582-90. <https://doi.org/10.1007/s00122-004-1585-0>
- Chung CL, Longfellow JM, Walsh EK, Kerdieh Z, Van Esbroeck G, Balint-Kurti P, Nelson RJ. Resistance loci affecting distinct stages of fungal pathogenesis: use of introgression lines for QTL mapping and characterization in the maize-*Setosphaeria turcica* pathosystem. *BMC Plant Biology*. 2010;10(1):1-25. <http://www.biomedcentral.com/1471-2229/10/103>
- Mitiku M, Eshte Y, Shiferaw W. Evaluation of maize variety for northern leaf blight (*Trichometasphaeria turcica*) in South Omo zone. *World Journal of Agricultural Research*. 2014;2 (5):237-39. <https://doi.org/10.12691/wjar-2-5-6>
- Irfan M, Ting ZT, Yang W, Chunyu Z, Qing M, Lijun Z, Feng L. Modification of CTAB protocol for maize genomic DNA extraction. *Research Journal of Biotechnology*. 2013;8(1):41-45.
- Vivekananda Y, Thangjam K. A simple modified DNA extraction method of maize. *International Journal of Agricultural Science and Research*. 2018;8(3):67-72. <https://doi.org/10.24247/ijasrjun20186>
- <http://www.maizegdb.org>: maize genome database
- Jaccard P. Nouvellesrechessurla distribution florale. *Bulletin de la Societe Vaudoise des Sciences Naturelles*. 1908;44: 223-70. <https://ci.nii.ac.jp/naid/10020758887/>
- Perrier X, Jacquemoud-Collet JP. 2006. DARwin software <http://darwin.cirad.fr/darwin.html>. Accessed 10 June 2022
- Anderson JA. Optimizing parental selection for genetic linkage maps. *Genome*. 1993;36:181-86. <https://doi.org/10.1139/g93-024>
- Salami HA, Sika KC, Padonou W, Aly D, Yallou C, Adjanohoun A, Kotchoni S, Baba-Moussa L. Genetic diversity of maize accessions (*Zea mays* L.) cultivated from Benin using microsatellites markers. *American Journal of Molecular Biology*. 2016;6: 12-24. <http://dx.doi.org/10.4236/ajmb.2016.61002>
- Wende A, Shimelis H, Gwata ET. Genetic variability for resistance to leaf blight and diversity among selected maize inbred lines. In: M. El-Esawi (ed.), *Maize Germplasm - Characterization and Genetic Approaches for Crop Improvement*. 2018. IntechOpen, London. <https://doi.org/10.5772/intechopen.70553>
- Adu GB, Awuku FJ, Amegbor IK, Haruna A, Manigben KA, Aboyadana PA. Genetic characterization and population structure of maize populations using SSR markers. *Annals of Agricultural Sciences*. 2019;64 (1):47-54. <https://doi.org/10.1016/j.aogas.2019.05.006>
- Nikolic A, Ignjatovic MD, Kovacevic D, Camdzija Z, Filipovic M, Mladenovic, Drinic S. Genetic diversity of maize inbred lines as inferred from SSR markers. *Genetika*. 2015;47(2):489-98. <https://doi.org/10.2298/GENSR1502489N>

21. Mahar KS, Agrawal PK, Kalyana Babu B, Gupta HS. Assessment of genetic diversity among the elite maize (*Zea mays* L) genotypes adapted to North-Western Himalayan region of India using microsatellite markers. *Journal of Plant Biochemistry and Biotechnology* 2009;18(2):217-20. <https://doi.org/10.1007/BF03263322>
22. Zunjare R, Hossain F, Muthusamy V, Vishwakarma AK, Pandey N, Kumar P, Sekhar JC, Jha SK, Nepolean T, Gupta, HS. Analyses of genetic diversity among exotic-and indigenous-maize inbreds differing for responses to stored grain weevil (*Sitophilus oryzae* L.) infestation. *Maydica*. 2015;60:M28. <https://doi.org/10.1080/23311932.2015.1137156>
23. Nyaligwa L, Hussein S, Amelework B, Ghebrehiwot H. Genetic diversity analysis of elite maize inbred lines of diverse sources using SSR markers. *Maydica*. 2015;60:M29. [oai:journals.crea.ugov.it/article/1278](https://oai:journals.crea.ugov.it/article/1278)
24. Mukri G, Patil MS, Motagi, BN, Bhat JS, Singh C, Jeevan Kumar SP, Gadag RN, Gupta NC, Simal-Gandara J. Genetic variability, combining ability and molecular diversity-based parental line selection for heterosis breeding in field corn (*Zea mays* L.). *Molecular Biology Reports*. 2022;1-8. <https://doi.org/10.1007/s11033-022-07295-3>
25. Kiran KK. Development and evaluation of turcicum leaf blight resistant hybrids using newly developed inbred lines of maize (*Zea mays* L.). *Agricultural Science*, Ph.D (thesis). Dharwad, India: University of Agricultural Sciences; 2016.