




ISSN: 2348-1900

Plant Science Today

<http://horizonepublishing.com/journals/index.php/PST>



Research Article

Phytoextraction of cadmium and lead by three vegetable-crop plants

Syed Yakub Ali, Sambhu Nath Banerjee and Shibani Chaudhury*

Department of Environmental Studies, Siksha-Bhavana, Visva-Bharati, Santiniketan 731235, India

Article history

Received: 28 June 2016
Accepted: 6 August 2016
Published: 15 August 2016

© Ali et al. (2016)

Editor

K K Sabu

Publisher

Horizon e-Publishing Group

Corresponding Author

Shibani Chaudhury
✉ shibani.chaudhury@visva-bharati.ac.in

Abstract

Phytoextraction, is an effective and promising means to cure soil contamination with heavy metals. The present study investigates the ability of three vegetables plants for removal of heavy metals from the contaminated soil and metal mobilization to different plant parts. The three plants selected for the study, *Momordica charantia*, *Vigna unguiculata* and *Solanum melongena* were grown for 90 days in soils artificially contaminated with cadmium (Cd) and lead (Pb) (50mg metal/kg of soil). The concentrations of the two metals were observed to be higher in roots of *M. charantia* and *V. unguiculata* than in soil, but root Pb level of *S. melongena* was slightly lower than that of soil after 90 days. Translocation potential of the heavy metals indicated higher accumulation of Cd in roots of *M. charantia* and *S. melongena* than in leaves while the pattern was completely opposite in *V. unguiculata*. Lead accumulation was higher in roots than in leaves for all the three plant species studied. The Translocation Factor (TF) of Cd for the three plants was in the range of 1.16 to 2.29 whereas, TF values of Pb remained <1, indicating that only small amount of Pb was translocated from roots to aerial parts.

Keywords

Heavy metals; Phytoextraction; Chlorophyll content; Translocation factor

Ali, S. Y., S. N. Banerjee and S. Chaudhury. 2016. Phytoextraction of cadmium and lead by three vegetable-crop plants. *Plant Science Today* 3(3): 298-303. <http://dx.doi.org/10.14719/pst.2016.3.3.247>

Introduction

Environmental pollution caused by heavy metals is a growing concern now-a-days due to its ill effects on plants and animals including human beings (John et al., 2008). Various agricultural and industrial activities is on the rise to cope up with the increasing demand of modern civilization, leading to an accumulation of these metals in the water bodies and soil, from which they can easily enter into the food-chain and pose health risks to the human population. Heavy metals are very hazardous pollutants because they are non-biodegradable, extremely toxic even at low concentrations and can change their mobility under different physico-chemical conditions (Mathew, 2001).

Some heavy metals act as essential elements for animals and plants in trace amounts, such as Cobalt (Co), Copper (Cu), Manganese (Mn), Molybdenum (Mo), Vanadium (V), Nickel (Ni) and Zinc (Zn) (He et al., 2005; Falusi and Olanipekun, 2007). These metals are used for redox processes as components of various enzymes and for regulation of osmotic pressure in cells (Bruins et al., 2000). Other heavy metals such as Cadmium (Cd), Lead (Pb), Chromium (Cr), Mercury (Hg), Aluminum (Al), Gold (Au) and Silver (Ag) have no biological function/beneficial effects (Chang et al., 1996). All these heavy metals may be detrimental to living organisms when present in excess amounts in soil and water bodies. Of all toxic heavy metals, cadmium (Cd) ranks the highest in terms of its

to cause damage to plant growth and pose serious risk to human health via food chain (Shah and Dubey, 1998).

The levels of Cadmium may rise to be toxic in soils due to mining activities, smelting, fuel combustion, as well as the use of phosphate fertilizers, sewage sludge, batteries, pigments, metal coatings, and plastics (ATSDR, 2011; Goswami and Das, 2015). Lead is another most abundant toxic metal in the earth crust. Presence of excessive amounts of Cd and Pb in soils affect various plant processes such as growth reduction, especially root growth, chlorosis, disturbances in mineral nutrition and carbohydrate metabolism, photosynthesis, water absorption, and cause wilting of leaves (Moya *et al.*, 1993, John *et al.*, 2008).

Phytoextraction, an indispensable component of phytoremediation, is widely in use to remove heavy metals from soil by growing selective plants and then harvesting those plants for management the contaminated soil (Jadia and Fulekar, 2009). Both aquatic and terrestrial plants have been used to reclaim contaminated water bodies and soil (Rahmani and Sternberg, 1999; Prasad *et al.*, 2001; John *et al.*, 2008). The ideal plant used for the purpose of phytoextraction should possess the ability to accumulate metal(s) intended to be extracted, preferably in the aboveground parts, tolerate high metal concentrations in soil, grow at an appreciably fast rate to be considered as an agricultural crop (Goswami and Das, 2015). In the present study, three vegetable-crop plants, *Momordica charantia*, *Vigna unguiculata* and *Solanum melongena*, were studied to estimate their bioaccumulation potential for cadmium and lead present in the artificially contaminated soils and to assess their ability to translocate the metals in shoots and leaves.

Materials and Methods

Soil sample were collected from field at 20 cm depth, dried, sieved and was then thoroughly mixed. Various soil parameters such as soil texture, pH, EC, organic Carbon, water soluble sodium, potassium, calcium, exchangeable sodium, potassium, calcium, total nitrogen, water soluble phosphorus, chloride and sulfate, available phosphorus, cation exchange capacity were analyzed. Grain size of the soils was analyzed by 'pipette method'. Soil pH was determined by potentiometric method using pH meter (Sytonic pH-meter 361) in 1:2.5 soil:water suspension. Electrical conductivity was measured in 1:2 soil:water extract using conductivity meter (Thermo Scientific conductivity cell, Orion 013605MD). Organic carbon of the soil was determined by method of Walkely and Black, (1934). Available nitrogen was determined using Kjeldahl instrument (PlicanKelplus-Distyl) following the method of Subbiah and Asija, (1956).

Available phosphorus was measured by Bray's method (Bray and Kurtg, 1945). Water soluble anion and cation were determined using Ion Chromatography (Metrohm 797 VA Computrace). For exchangeable sodium, potassium, and calcium, soil samples were extracted by ammonium acetate and the liquid extracts were measured by flame photometry (ELICO CL 361). Cation exchangeable capacity of the soil was determined by following the method of Harada and Inoko, (1980).

After analysis of soil parameters plastic pots with holes at the bottom, were filled with approximately 2kg air-dried and sieved soil. The soil was fertilized using 200gm of cow dung and 5gm of urea and then artificially contaminated by lead [Lead nitrate, $Pb(NO_3)_2$] and cadmium [Cadmium nitrate, $Cd(NO_3)_2 \cdot 4H_2O$] at 50mg of each metal/kg of soil following procedures described by Turan and Esringü, (2007). For control pots, similar processes were followed except the addition of metals in the soil. Three edible plant species, *Momordica charantia*, *Vigna unguiculata* and *Solanum melongena* were cultivated in separate pots (in triplicate) in metal contaminated and normal (control) soils. Both the control and metal contaminated pots were watered 3 to 4 times per week. Pot experiments were conducted under ambient climatic conditions from May to August, 2013. The maximum and minimum temperature recorded were 29-34°C and the relative humidity ranged from 30-80%. All plant species were harvested after 3 months.

After three months plants were collected from the pots, thoroughly washed in running tap water and rinsed with deionized water to remove each and every soil particle remaining attached with the plant surfaces. Then plant parts (roots, shoots and leaves) were oven dried at 70°C for 48 hours. The dried samples were weighed and powdered. For the soil samples, upper most soil portion from the pots were discarded. Then the remaining soils were mixed thoroughly and sieved at 2mm. Sieved soil samples were collected in the zipper bag for metal analysis.

Soil and dry plant samples (1gm) were digested after adding 15 ml of tri-acid mixture (HNO_3 , H_2SO_4 , and $HClO_4$ in 5:1:1 ratio) at 80°C until a transparent solution was obtained (Allen *et al.*, 1986). After cooling, the digested sample was filtered using Whatman No. 42 filter paper and the filtrate was finally raised to 50 ml with distilled water. Metals were analyzed by using Anodic Stripping Voltammetry (Metrohm VA 797 Computraces).

The chlorophyll content was determined according to the method of Arnon, (1949). Fresh leaf samples (500mg) were homogenized in 80% cold acetone and centrifuged at 5000rpm for 15 min (Remi Cooler centrifuge C-4), the absorbance was read at 645nm and 663 nm (Cyberlab UV-100). Values were expressed as mg chlorophyll/gm leaves (fresh weight).

Table 3: Heavy Metals (in ppm) concentration of soil, root, shoot and leaves of three harvested plants

Pt	Cd root	Pb root	Cd shoot	Pb Shoot	Cd leaves	Pb leaves
<i>Momordica charantia</i>	36.91±1.87 ^a	53.72±1.55 ^a	22.22±1.18 ^a	34.29±1.53 ^a	20.63±1.05 ^a	2.48±0.30 ^a
<i>Vigna unguiculata</i>	29.15±0.98 ^b	78.54±1.83 ^b	26.35±0.83 ^b	4.41±1.12 ^b	40.43±1.52 ^b	7.06±0.61 ^b
<i>Solanum melongena</i>	28.48±1.72 ^{ab}	41.99±1.57 ^c	24.97±1.53 ^{bc}	10.87±1.47 ^c	11.90±1.63 ^{ab}	6.33±0.58 ^{ab}

The same letter within columns are not significantly different at the 5% probability level by least significant range.

Table 4: BCF and TF value of harvested plants

	Bioconcentration factors		Translocation factors	
	Cd	Pb	Cd	Pb
<i>Momordica charantia</i>	1.60±0.04	1.29±0.04	1.16±0.03	0.68±0.005
<i>Vigna unguiculata</i>	1.92±0.01	1.29±0.03	2.29±0.04	0.15±0.03
<i>Solanum melongena</i>	1.31±0.03	0.85±0.04	1.29±0.07	0.41±0.05




Fig. 1: Cd and Pb concentration in roots, shoots and leaves of three harvested plants

The results were statistically analyzed with the help of Microsoft Excel 2013 using one way ANOVA to determine the degree of significance for the studied plants.

Results and Discussion

The physico-chemical characteristics of uncontaminated soil samples are listed in Table 1. The three plant species were found to grow well in the heavy metal contaminated soils. There were neither any visible change in the morphology of the plants under the influence of metal stress as compared to control plants nor any sign of chlorosis in the treated plants (Table 2). Earlier observers (Unni *et al.*, 1992; Kopittke *et al.*, 2007; Siddhu *et al.*, 2008; Daniel *et al.*, 2009; Yilmaz *et al.*,

2009; Gautam *et al.*, 2014) have reported such changes as plant growth, germination and peroxidase activity at various doses of Pb/Cd applied separately on the same plants. It may be said that combination of these two metals did not alter the morphology of these plants which is well evident in the chlorophyll content of the treated and control plants. Experiment conducted by Al-Subu *et al.*, (1993), on marrow vegetables showed that the toxicity of cadmium and lead was mostly antagonistic and sometimes irregular on the parts of root-treated or foliar-treated plants.

The value of Cd concentration applied to the soil was higher than the Indian Standard (3 to 6 mg/kg) but the Pb concentration was much lower than the tolerable limits (250 to 300 mg/kg) as stated by Kabata-Pendias, (2001).

Table 1: Physio-chemical characteristics of uncontaminated soil

Parameters	Quantity
Soil type	Sandy Loam
pH	6.3
Electron Conductivity	1mS/cm
Organic Carbon	1.1%
Water soluble Cations:	
Sodium	46.9mg/kg
Potassium	60.4mg/kg
Calcium	31.7mg/kg
Exchangeable Cations:	
Sodium	213.2mg/kg
Potassium	17.1mg/kg
Calcium	1537mg/kg
Water Soluble Anions:	
Chloride	350mg/kg
Phosphorus	0.56mg/kg
Sulfate	48mg/kg
Total Nitrogen	209mg/kg
Available Phosphorus	2.23mg/kg
Cation Exchange Capacity	11.2meq/100gm
Heavy Metals	
Zinc	80mg/kg
Copper	55mg/kg
Lead	20.5mg/kg
Cadmium	ND

Table 2: Total Chlorophyll content in control and treated experimented plants

Species Name	Total Chlorophyll mg/g FW	
	Control Plants	Treated Plants
<i>Momordica charantia</i>	1.23	1.18
<i>Vigna unguiculata</i>	0.46	0.42
<i>Solanum melongena</i>	0.93	1.01

Phytoextraction studies showed that the three plant species accumulated higher levels of Cd and Pb in their roots, but their ability for translocation of these heavy metals to the aerial plant parts were considerably different. The levels of Cd were higher in roots of *Momordica charantia* and *Solanum melongena* (36.91 mg/kg and 28.48 mg/kg) followed by the shoot (22.22 mg/kg and 24.97 mg/kg) and leaves (20.63 mg/kg, 11.90 mg/kg). The trend was opposite in case of *Vigna unguiculata* which accumulated higher levels of Cd in leaves (40.43 mg/kg) as compared to root system (29.15 mg/kg). In case of Pb, the highest accumulation was found to occur in root of all harvested plants. *Vigna unguiculata* accumulated highest amounts of Pb in root (78.54mg/kg), followed by *Momordica charantia*

(53.72 mg/kg) and *Solanum melongena* (41.99mg/kg). In contrast to Cd, translocation of Pb from root to aerial parts was found to be very poor in *Vigna* and *Solanum* with the exception of *Momordica* which accumulated higher levels of Pb in shoot (34.29 mg/kg). Translocation of Pb from root to the leaves was insignificant in all the three plant species studied (in the range of 2.48 mg/kg to 7.06 mg/kg).

The accumulation of heavy metals in a plant can be quantified by two factors: the Bioconcentration Factor (BCF) (Eqn. 1) and the Translocation Factor (TF) ((Eqn. 2) (Selamat *et al.*, 2014). The BCF is calculated as the ratio of the element concentration in plant tissues to the concentration of the element in the soil (Zayed *et al.*, 1998).

$$\text{Bioconcentration Factor (BCF)} = \frac{\text{element concentration in plant tissues} \dots\dots\dots 1}{\text{element concentration in soil}}$$

TF is calculated as the ratio of metal concentrations in aerial part of the plant to those in roots, indicating the ability of the plant to translocate metals from the roots to the shoots (Roongtanakiat, 2009).

$$\text{Translocation Factor (TF)} = \frac{\text{element concentration in aerial part of the plant} \dots 2}{\text{element concentration in root these plant}}$$

The BCF values of the Cd and Pb of all the three harvested plants in the metal contaminated soil were >1 except *Solanum melongena* for Pb. The highest BCF value of Cd was observed in *Vigna unguiculata* about 1.92 followed by *Momordica charantia* and *Solanum melongena* ranging about 1.60 and 1.29 respectively (Table 4). Furthermore, the Cd concentrations in the aerial parts of harvested plants were higher than that of the corresponding roots, i.e. the translocation factors (TF) were >1. The highest TF values of Cd were observed in *Vigna unguiculata*, ranging from 2.18 to 2.38, followed by *Solanum melongena* ranging from 1.17 to 1.38 and *Momordica charantia* ranging from 1.1 to 1.23 (Table 4).

From the result observed it is clear that BCF and TF values of Pb were less than the values of the Cd. Highest BCF value of Pb was observed in *Momordica charantia* and *Vigna unguiculata* about 1.29 and lowest in *Solanum melongena* about 0.85 (Table 4). Furthermore, the Pb concentrations in the aerial parts of harvested plants were lower than that of their roots, i.e. the translocation factors (TF) were <1. The TF values of the three harvested plants *Momordica charantia*, *Vigna unguiculata* and *Solanum melongena* were 0.67, 0.13 and 0.41 respectively.

The present study showed that the distribution of Cd in the aerial plant parts was found to be significantly higher than roots. The BCF and TF values of Cd were always >1 in all harvested plants. High BCF values indicated that the plants could accumulate Cd from soil. On the other hand, high TF values indicated that plants could take up Cd from the soil and store it in its above ground parts (Selamat *et al.*, 2014). Highest BCF and TF values were observed in the *Vigna unguiculata*, which indicated that this plant was capable of accumulating Cd to a greater extent from soil than the other two plants and store the metal in the aerial plant parts. Cadmium was stored in the *Vigna unguiculata* according to following order Leaves>Roots>shoots. Mengel and Kirkby, (1982) stated that Cadmium is readily transported from the soil to the upper parts of plants. Transport of Cd from the soil to plant parts of agricultural crops is significantly greater than other heavy metals except Zinc (Moolenaar and Lexmond, 1999).

TF values of Pb was always <1 in all experimental plants, indicating that Pb concentration in the plant root was always greater than the aerial parts. According to Grill *et al.*, (1985), this may be due to low mobility of Pb, when it is found in low quantities in the soil, given that the root systems prevents the migration of Pb towards the above ground part of the plant and it only reaches this part if it is found in high concentration.

In the present study phytoextraction and bioaccumulation potential of the three plant species *Momordica charantia*, *Vigna unguiculata* and *Solanum melongena* against Cd and Pb were found to be favorable. Translocation of Cd within the plant system was, however, much higher than that of Pb for all the three plant species studied. Most of the Pb extracted from contaminated soil was localized within the root system while a considerable fraction of extracted Cd was found to be translocated to the aerial parts of the three plant species. Kopittke *et al.*, (2007), obtained similar result while working on *Vigna unguiculata*. They observed that the tissue concentrations of Pb were 10 to 50 times greater in the roots than in the shoots, with the critical Pb concentration being 330 µg/g for the roots and 49 µg/g for the shoots. Another study reported by Daniel *et al.*, (2009) that the Cd²⁺ accumulation was greater than Pb in seeds of bitter-gourd. Qadir *et al.*, (2000) and Chauhan, (2014) observed twofold Cd accumulation in leafy vegetables as compared to others.

Conclusion

The results clearly indicate that phytoextraction studies of edible plants are important not only from nutritional point of view, but also from the differential abilities of an individual plant to

remove heavy metals from soil and localize them in specific plant parts.

Competing Interest

The authors declare that they have no competing interests.

Authors' contributions

All authors have contributed significantly. They have performed the laboratory works and prepared the manuscript.

Acknowledgements

Reviewers and editor of the article are greatly acknowledged for their tremendous effort in reviewing the manuscript. A special mention is important for Mr. Bikash Chakroborty and Mrs. Rani Kisku who helped to maintain the plants.

References

- Allen, S. E., H. M. Grimshaw, and A. P. Rowland. 1986. Chemical analysis. In: *Methods in Plant Ecology*, P.D. Moore and S. B. Chapman, Ed. Blackwell, Scientific Publication, Oxford, London. p. 285-344.
- Al-Subu, M. M., R. Salim, A. Douleh, and A. Atallah. 1993. Combined Effects of Cadmium, Lead and Copper on the growth and on metal uptake of Broad Beans, Carrots, Radishes and Marrow vegetables, *Rev. Int. Contam. Ambient* 9: 1-9.
- Arnon, D.I. 1949. Copper enzymes in isolated chloroplast: Polyphenoloxidase in *Beta vulgaris*. *Plant Physiol* 24: 1-15.
- ATSDR, 2011. CERCLA Priority List of Hazardous Substances, Agency for Toxic Substances and Disease Control (Online), available URL: <http://www.atsdr.cdc.gov/spl/index.html> (accessed on 2nd October 2012).
- Bray, R. H., and L. T. Kurtz. 1945. Determination of total, organic and available forms of phosphorus in soil. *Soil Sci* 59: 39-45.
- Bruins, M., S. Kapil, and F. Oehme. 2000. Microbial resistance to metals in the environment. *Ecotoxicol Environ Saf* 45: 198-207. doi: 10.1006/eesa.1999.1860
- Chang, L., L. Magos, and T. Suzuki. 1996. *Toxicology of Metals*. CRC Press, Boca Raton, FL, USA.
- Chauhan, G. 2014. Toxicity study of metals contamination on vegetables grown in the vicinity of cement factory. *International Journal of Scientific and Research Publication* 4: 1-8.
- Daniel, G., K. Hussain, and N. Salim. 2009. Bioaccumulation and effect of lead, chromium and cadmium on peroxidase activity in bittergourd (*Momordica charantia* L. walp) cv. PRIYA. *International Journal of Plant Sciences* 4: 239-244.
- De la Cruz-Landero, N., V. E. Hernandez, E. Guevara, M. A. Lopez-Lopez, A. T. Santos, E. Ojeda-Trejo, and A. Alderete-Chavez. 2010. *Lupinus versicolor* Response in Soils Contaminated with Heavy Metals from a Petroleum Extraction Field. *Journal of Applied Sciences* 10: 694-698. doi: 10.3923/jas.2010.694.698
- Falusi, B.A., and E. O. Olanipekun. 2007. Bioconcentration Factors of Heavy Metals in Tropical Crab (*Carcinus* sp) From River Aponwe, Ado-Ekiti, Nigeria. *Journal*

of *Applied Sciences and Environmental Management* 11: 51-54.

- Gautam, M., A. K. Singh, and R. M. Johri. 2014. Influence of Pb toxicity on yield, yield attributing parameters and photosynthetic pigment of tomato (*Solanum lycopersicum*) and eggplant (*Solanum melongena*). *Indian Journal of Agricultural Sciences* 84: 808-815.
- Goswami, S., and S. Das. 2015. A Study on Cadmium Phytoremediation Potential of Indian Mustard, *Brassica juncea*. *International Journal of Phytoremediation* 17: 583-588. doi: 10.1080/15226514.2014.935289
- Grill, E., E. L. Winnacker, and M. H. Zenk. 1985. Phytochelatins: The principal heavy-metal complexing peptides of higher plants. *Sciences* 230: 674-676. doi: 10.1126/science.230.4726.674
- Harada, Y., and A. Inoko. 1980. The Measurement of the Cation-Exchange Capacity of Composts for the Estimation of the Degree of Maturity. *Soil Science and Plant Nutrition* 26: 127-134.
- He, Z., X. Yang, and P. Stoffella. 2005. Trace elements in agroecosystems and impacts on the environment. *Journal of Trace Elements and medicine and biology* 19: 125-40. doi: 10.1016/j.jtemb.2005.02.010
- Jadia, C., and M. Fulekar. 2009. Phytoremediation of heavy metals: Recent techniques. *African Journal of Biotechnology* 8: 921-928.
- John, R., P. Ahmad, K. Gadgil, and S. Sharma. 2008. Effect of cadmium and lead on growth, biochemical parameters and uptake in *Lemna polyrrhiza* L. *Plant Soil and Environment* 54: 262-270.
- Kabata Pendias A. 2001. *Trace Elements in soils and Plants*. CRC Press, Boca Raton, FL.
- Kamnev, A. A., and D. ven der Lelie. 2000. Chemical and Biological Parameters as Tools to Evaluate and Improve Heavy Metal Phytoremediation. *Bioscience Reports* 20: 279-258. doi: 10.1023/A:1026436806319
- Kopittke, P. M., C. J. Asher, R. A. Kopittke, and N. W. Menzies. 2007. Toxic Effects of Pb²⁺ on Growth of Cowpea (*Vigna unguiculata*). *Environmental Pollution* 150: 280-287. doi: 10.1016/j.envpol.2007.01.011
- Mathew, A. R. 2001. *Phytoremediation of heavy metal contaminated soil*. <http://digital.library.okstate.edu/etd/umi-okstate-1445.pdf>
- Mengel, K., and E. Kirkby. 1982. *Principles of plant nutrition*. International Potash Institute, Switzerland.
- Moolenaar, S., and I. Lexmond. 1999. Heavy metal balances. Part 1. General aspects of cadmium, copper, zinc and lead balance studies in agroecosystems. *Journal of Industrial Ecology* 2: 4-8. doi: 10.1162/jiec.1998.2.4.45
- Moya, J. L., R. Ros, and I. Picazo. 1993. Influence of cadmium and nickel on growth, net photosynthesis and carbohydrate distribution in rice plants. *Photosynthesis* 36: 75-80. doi: 10.1007/BF00016271
- Selamat, S. N., R. S. Abdullah, and M. Idris. 2014. Phytoremediation of lead (Pb) and Arsenic (As) by *Melastoma malabathricum* L. from Contaminated Soil in Separate Exposure. *International Journal of Phytoremediation* 16: 694-703. doi: 10.1080/15226514.2013.856843
- Olsen, S., C. Cole, F. Watanabe, and L. Dean. 1954. *Estimation of available phosphorus in soils by extraction with sodium bicarbonate*. USDA Circular 939. U.S. Washington D.C., Government Printing Office.
- Prasad, M. N. 2003. Phytoremediation of Metal-Polluted Ecosystems: Hype for Commercialization. *Russian Journal of Plant Physiology* 50: 686-700. doi: 10.1023/A:1025604627496
- Prusty, B., K. Sahu, and G. Godgul. 1994. Metal contamination due to mining and milling activities at the Zawar zinc mine, Rajasthan, India. *Contamination of stream sediments: Chemical Geology* 112: 275-292. doi: 10.1016/0009-2541(94)90029-9
- Qadir, M., A. Ghafoor, and G. Murtaza. 2000. Cadmium concentration in Vegetables grown on Urban soils irrigated with untreated municipal sewage. *Environment, Development and Sustainability* 2: 11-19. doi: 10.1023/A:1010061711331
- Rahmani, G. N. H., and S.P. K. Sternberg. 1999. Bioremoval of lead from water using *Lemna minor*. *Bioresource Technology* 70: 225-230. doi: 10.1016/S0960-8524(99)00050-4
- Roongtanakiat, N. 2009. *Vetiver Phytoremediation for Heavy Metal Decontamination*. Technical Bulletin no. 2009/1. Pacific Rim Vetiver Network, Office of the Royal Development Projects Board, Bangkok, Thailand.
- Shah, K., and R.S. Dubey. 1998. Effect of cadmium on proline accumulation and ribonuclease activity in rice seedlings: Role of proline as a possible enzyme protectant. *Biologia Plantarum* 40: 121-130. doi: 10.1023/A:1000956803911
- Siddhu, G., D. S. Sirohi, K. Kashyap, I. A. Khan, and M. A. Khan. 2008. Toxicity of cadmium on the growth and yield of *Solanum melongena* L. *Journal of Environmental Biology* 29: 853-857.
- Subbiah, B., and Asija, G. 1956. A rapid procedure for the determination of available nitrogen in soil. *Current Science* 25: 259-260.
- Turan, M., and M. T. Esringü. 2007. Phytoremediation based on canola (*Brassica napus* L.) and Indian mustard (*Brassica juncea* L.) planted on spiked soil by aliquot amount of Cd, Cu, Pb, and Zn. *Plant Soil and Environment* 53: 7-15.
- Unni, P. N., G. Daniel, and S. R. Nair. 1992. Effect of heavy metals on field crops I. effect of lead on germination, growth and chlorophyll content in bitter gourd (*Momordica charantia* L.). *International Journal of Environmental Studies* 41: 115-119. doi: 10.1080/00207239208710750
- Walkley, A. and I. Black. 1934. An estimation of the Degtjareff method for determining soil organic matter, and a proposed modification of the Chromic acid titration method. *Soil Science* 34: 29-38.
- Yilmaz, K., Akinci, I. E., and S. Akinci. 2009. Effect of lead accumulation on growth and mineral composition of eggplant seedlings (*Solanum melongena*). *New Zealand Journal of crop and horticultural Science* 37: 189-199. doi: 10.1080/01140670909510264
- Zayed, A., S. Gowthaman, and N. Terry. 1998. Phytoaccumulation of trace elements by wetland plants. *Journal Environmental Quality* 27: 715-721. doi: 10.2134/jeq1998.00472425002700030032x

