



RESEARCH ARTICLE

Double phase culture system mediated enhanced protocol for shoot proliferation of *Vanilla andamanica* Rolfe. - an endemic wild relative of commercial Vanilla from the Andaman and Nicobar Islands

Shaina Jerald¹, A.Gangaprasad^{1,2*} & Sam P. Mathew³

¹Department of Botany, University of Kerala, Karyavattom P.O., Thiruvananthapuram, Kerala 695581, India

²Centre for Biodiversity Conservation, Department of Botany, University of Kerala, Karyavattom P.O., Thiruvananthapuram, Kerala 695581, India

³Jawaharlal Nehru Tropical Botanic Garden and Research Institute, Thiruvananthapuram, Kerala 695562, India

*Email: agangaprasad@yahoo.com



ARTICLE HISTORY

Received: 22 March 2023

Accepted: 01 June 2023

Available online

Version 1.0 : 28 July 2023



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care etc. See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Shaina J, Gangaprasad A, Mathew SP. Double phase culture system mediated enhanced protocol for shoot proliferation of *Vanilla andamanica* Rolfe. - an endemic wild relative of commercial Vanilla from the Andaman and Nicobar Islands. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.2475>

Abstract

A liquid overlay culture system in micropropagation improved shoot proliferation in many species. In the present investigation, we have developed an enhanced shoot proliferation protocol using the double phase culture system (DPS) for *Vanilla andamanica*, an endemic species of the Andaman and Nicobar Islands recognized under the vulnerable category by the IUCN. *In vitro* generated nodal explants were used for shoot proliferation experiments and were tested in Murashige and Skoog medium augmented with various cytokinins (BAP, KIN, 2iP) and auxins (IBA, NAA, IAA) to produce a maximum of 3.24 ± 0.08 shoots per explant in 5 μ M BAP and 5.23 ± 0.67 roots per shoot in basal MS medium. A remarkable increase in shoot regeneration was observed when the nodal explant was cultured on DPS system with optimum cytokinin (5 μ M BAP). On evaluation of the influence of DPS and conventional single-phase systems (SPS), (solid medium) exhibits an improved multiplication rate on a DPS with optimal BAP (5 μ M) with an average of 8.66 ± 0.17 shoots per explant, which represents a 2-fold increase over the rate of SPS + 5 μ M BAP. The plantlets were rooted at 100% frequency in half-strength MS medium devoid of auxins and acclimatized with 100% success. A double-phase mediated enhanced shoot proliferating procedure could be employed for large scale multiplication and commercial breeding trials for better results with regard to *V. andamanica*.

Keywords

Vanilla andamanica; endemic; double phase culture; liquid medium; micropropagation; shoot proliferation

Introduction

The genus *Vanilla* Plum. ex-Mill., belonging to the family Orchidaceae and comprising more than 110 species, is distributed throughout the tropics (1, 2). Interestingly, *Vanilla* is the only commercial orchid taxon found to be an expensive flavouring material across the world after saffron (3). All the *vanilla* species except commercial *vanilla* are under the threat of serious habitat destruction and massive exploitation, highlighting the relevance of their conservation (3). The commercial crop improvement programmes are apparently meagre owing to limited clonal sources and the availability of a wide range of genetic variability. It highlights the necessity of *ex-situ* conservation in different geographical habitats of the taxon.

V. andamanica is a climbing, hemiepiphytic shrub whose occurrence

in the Andaman and Nicobar Islands was reported by Rolfe in 1918. The genetic distance of *V. andamanica* from other wild relative species of *vanilla* showed that it had a different origin, which may have arisen during past historical introduction events in the Asian sub-continent along with India (3). This evidently emphasises the importance and rarity of *V. andamanica*.

According to Lubinsky (4), the primary gene pool of *vanilla* species is evidently threatened, and therefore the secondary gene pool contains wild relatives of commercial *vanilla* that serves as a source of many desirable characteristics which can be used for crop improvement programmes (5). Disease resistance was considered one of the most important crop breeding traits. The foot rot and wilt diseases commonly found in cultivated species of *vanilla* by *Fusarium oxysporum* and *Phytophthora meadii*, which are rather least infected with *V. andamanica*, are a hallmark of this wild endemic taxon in the breeding programmes among existing germplasm of *vanilla*. Successful isolation followed by protoplast fusion of *V. planifolia* and *V. andamanica* was also reported in order to achieve gene transfer of desirable traits and hybrid production (6).

Habitat destruction coupled with frequently occurring natural calamities in the Andaman and Nicobar group of islands severely threatens many endemic and rare species, highlighting the immediate requirement of *ex-situ* conservation of plant species outside the islands. Micropropagation facilitates enhanced rapid multiplication of plant species from the limited plant material and is the best propagation methodology for the rare species concerned. In the present investigation, an evaluation of a double phase culture system over a conventional system of propagation was employed. A successful protocol for enhanced shoot multiplication for the conservation of *V. andamanica* through a double phase culture system was employed for the first time.

Materials and Methods

Explant source and culture condition

Plant materials for *in vitro* culture were originally collected from South Andaman Island and were well acclimatised in the greenhouse of the Department of Botany, University of Kerala, India (Fig. 1.A). They served as a source of explant. Young top shoots having two to three nodes from healthy plants grown in the greenhouse were defoliated, excised into nodal segments, and washed in tap water for 10-20 min, followed by treatment in the 10% Labolene solution (Galaxo India Ltd., Mumbai) for 10-15 min with constant shaking. Nodal segments were then subjected to washing in running tap water for 30-45 min, were further rinsed thrice using double-distilled water, and were surface sterilised using 0.1% HgCl₂ (SRL, Pvt. Ltd., India) for 5-6 min also. The surface-sterilized explants were further washed with sterile double-distilled water three to four times and inoculated into culture medium after the excision of nodal explants into appropriate sizes (1-1.5 cm). Surface sterilisation and inoculation were carried out under an

aseptically maintained laminar airflow hood (Klenzaid, India).

For initiating cultures, MS medium (7) containing salts and vitamins augmented with 3% sucrose (w/v) (Himedia, India) and gelled with 0.8% agar (w/v) (Himedia, India) was used, and the pH was adjusted to 5.8 ± 0.2 using 1N NaOH or 1N HCl. Sterilization of a culture bottle (13 x 6 cm) was carried out in an autoclave (Nat Steel, Mumbai) at 120°C for 18 minutes under 15 lbs. pressure. The inoculated cultures were carefully maintained in a culture room provided with 25 ± 2 °C under a 16/8 h photoperiod with a light intensity of 2500 lux (μmol⁻²S⁻¹) provided with white LED tubes (Crompton Ltd., Mumbai) and 55- 60 % relative humidity.

Culture initiation and *in vitro* multiplication

The surface-sterilised nodal segment was inoculated into MS medium without plant growth regulators. Nodes containing a petiole attached to a portion of an internode (nodal segments) of appropriate size were excised from proliferated shoots in cultures and subcultures into fresh hormone-free medium for further proliferation. Nodal segments obtained from subcultures served as explants for further *in vitro* experiments. *In vitro*-developed nodal explants were cultured on MS medium supplemented with various concentrations of different cytokinin individually, viz. 6-Benzyl Amino Purine (BAP 1-5 μM), kinetin (KIN 1-5 μM), and 2-isopentenyl adenine (2 iP 1-5 μM) was tested for shoot multiplication and medium devoid of plant growth regulators was used as a control.

Shoot proliferation in Double Phase Culture System

To evaluate the effect of double phase culture systems and their duration, MS medium supplemented with optimum cytokinin (5 μM BAP) for shoot multiplication was selected. A nodal segment derived from stock cultures with one axillary bud was excised and inoculated into the double phase medium (DPS). The explants inoculated on 60 mL of solid medium and 15 mL of liquid medium with optimum BAP (5μM) were added after 30 days of culture without removing the solid medium. For solid phase medium (SPS), the shoots were retained on the same medium itself. The parameters were evaluated at 30, 60, and 90 days after the culture inoculation. The total shoot number, shoot length and percentage of response were recorded.

In vitro rooting

Healthy shoots of about four to five leaves were transferred to half strength-MS medium with different concentrations (0.5-2 μM) of auxins: (IBA (Indole -3- butyric acid), IAA (Indole-3- acetic acid) and NAA (1- Naphthalene acetic acid) to produce *in vitro* roots.

Acclimatization and field establishment of rooted plantlets

In vitro-rooted cultures were taken from the culture room, kept at room temperature for one week, and deflasked and washed using tap water to remove the traces of agar. Rooted plantlets were transferred to plastic pots (6 x 7.5 cm) filled with a potting mixture of garden soil and river sand (1:1) and maintained humidity by covering the

planted pots with a transparent polythene bag. The polythene bags were removed from the pots after 30 days, and the pots were transferred to greenhouse conditions. Hardened plants were planted in a potting mixture (garden soil, sand, and cow dung in a 1:1:1 ratio) containing plastic pots (17 x 15 cm) and then in the field.

Design of experiment and data collection

A completely randomized experimental design was employed for conducting experiments with three replications and three explants per flask. Experimental data were recorded after 30, 60, and 90 days in the case of the double phase culture system experiment. After 60 days, data were recorded for *in vitro* rooting and shooting experiments using PGRs and subjected to ANOVA, and the data means were compared by Tukey's test ($p \leq 0.05$) (Minitab ver. 20.4.0., Minitab, LLC., Pennsylvania, US).

Result and Discussion

Influence of PGRs on shoot multiplication

The effect of plant growth regulators supplemented with MS medium on nodal explant response for shoot multiplication after 60 days of culture is represented in Table 1. *In vitro* derived nodal segments were inoculated with MS medium supplemented with various concentrations of cytokinins like BAP, KIN, and 2iP to induce multiple shoots. Among the cytokinins used, the maximum multiple shoot induction (3.24 ± 0.08) and shoot length (2.52 ± 0.10 cm) were obtained on 5 μ M BAP augmented MS medium after 60 days with a 79.20 ± 2.77 percentage of response (Table 1) (Fig. 1.B). A reduced number of shoots per explant was given by other cytokinins tested (KIN and 2iP), which also did not contribute much shoot length when compared to BAP.

The present investigation reveals that the MS medium augmented with 5 μ M BAP showed a better

regeneration response in *V. andamanica* when compared to other tested cytokinins. BAP induced multiple shoot regeneration was also reported in *V. planifolia* by Abede *et al.* (8) and de Oliveira *et al.* (9). However, Diwakaran *et al.* (10) report a combinational medium (4.44 μ M BAP and 2.4 μ M IBA) for enhanced shoot production in *vanilla* species, which was completely absent in the present study.

Influence of double phase culture system on shoot multiplication

In order to enhance multiple shoot induction, the explants were inoculated with double phase medium containing optimum cytokinin (5 μ M BAP) supplemented with agar-gelled solid MS medium and a layer of liquid MS medium. The results obtained on experiments with SPS and DPS with different culture durations are shown in Table 2. It was observed that the use of DPS with 5 μ M BAP induces enhanced shoot multiplication (8.66 ± 0.17) and significantly improves the rate of shoot multiplication and shoot elongation (2.14 ± 0.09) (Fig. 1.C) compared to other systems (SPS, SPS + BAP, DPS) with a higher percentage of response (84.70 ± 0.25) after 90 days of culture. The DPS culture system improves the shoot multiplication efficiency at least two times higher than that of a conventional system of culture with a culture duration of 90 days. Rapid development of explants as well as the healthy development of multiple shoots was enabled by the DPS system.

Several plant species, such as *Malus sylvestris* (11), pineapple (12, 13), *Cymbidium giganteum* (14), *Pyrus calleryana* (15), potato (16), and *Gynerium sagittatum* (17), provide evidence for the use of liquid medium for improved shoot regeneration. Usually, liquid medium based culture systems use a support such as a bridge or paper raft. In the present DPS system, the solid phase basement acts as an anchor, whereas the liquid phase replenishes plant growth regulators and nutrients, which

Table 1. The effect of various cytokinins & augmented - with MS medium on shoot multiplication from *in vitro* generated nodal explants of *Vanilla andamanica*.

PGRs (μ M)			Means no. of shoots/explant (cm) \pm SE	Mean shoot length in (cm) \pm SE	Percentage of Response
BAP	KIN	2ip			
0	0	0	1.00 \pm 0.00 ^f	1.53 \pm 0.01 ^{ef}	42.85 \pm 0.00 ⁱ
1			0.90 \pm 0.08 ^{gh}	1.65 \pm 0.04 ^{de}	61.27 \pm 1.09 ^e
2			1.62 \pm 0.08 ^{de}	1.99 \pm 0.03 ^{bc}	66.23 \pm 0.25 ^d
3			2.14 \pm 0.14 ^{bc}	2.13 \pm 0.06 ^b	70.94 \pm 0.82 ^c
4			2.42 \pm 0.14 ^b	2.10 \pm 0.04 ^b	75.96 \pm 0.05 ^b
5			3.24 \pm 0.08 ^a	2.52 \pm 0.10 ^a	79.20 \pm 2.77 ^a
	1		1.00 \pm 0.14 ^{gh}	1.55 \pm 0.02 ^{ef}	47.20 \pm 0.20 ^h
	2		1.14 \pm 0.14 ^{fgh}	1.38 \pm 0.07 ^c	52.33 \pm 0.45 ^g
	3		1.38 \pm 0.08 ^{ef}	1.66 \pm 0.05 ^{de}	56.70 \pm 0.60 ^f
	4		1.61 \pm 0.16 ^{de}	1.79 \pm 0.09 ^{cd}	70.80 \pm 0.72 ^c
	5		1.85 \pm 0.14 ^{cd}	1.84 \pm 0.03 ^{cd}	66.04 \pm 0.14 ^d
		1	0.80 \pm 0.08 ^h	1.35 \pm 0.24 ^c	42.85 \pm 0.00 ⁱ
		2	1.19 \pm 0.08 ^{fg}	1.33 \pm 0.03 ^a	52.42 \pm 0.42 ^g
		3	1.42 \pm 0.14 ^{ef}	1.63 \pm 0.02 ^{de}	60.43 \pm 2.12 ^e
		4	1.38 \pm 0.08 ^{ef}	1.74 \pm 0.01 ^{de}	60.80 \pm 0.73 ^e
		5	1.66 \pm 0.08 ^{de}	1.85 \pm 0.02 ^{cd}	57.14 \pm 0.00 ^f
		Treatment Df(n-1)=15	94.50***	32.32***	364.76***
		Cytokinin(T) Df(n-1)=2	4.59*	6.67**	15.51***
		Cytokinin con.(C) Df(n-1)=4	481.41***	130***	157.71***
		TxCDF (n-1)=14	62.09***	3.84*	2.27 ^{NS}

Means followed by same letter within a column are not significantly different ($p < 0.05$) as determined by Tukey's test
NS - Non significant, *** F value is highly significant at $p < 0.001$, ** significant at $p < 0.01$ and * significant at $p < 0.05$

Table 2. The effect of various *in vitro* culture systems (SPS - solid phase system and DPS - double phase system) with 5 µM BAP on shoot proliferation from *in vitro* generated nodal explants of *Vanilla andamanica*.

Culture System	Culture Duration (No. of days)	Mean Shoot Number Per explant ±SE	Mean Shoot Length ±SE	Percentage of Response
SPS	30	1.00±0.00 ^f	1.25±0.02 ^f	40.63±1.19 ^f
	60	1.00±0.00 ^f	1.40±0.01 ^e	41.90±1.65 ^f
	90	1.00±0.00 ^f	1.47±0.01 ^e	46.66±1.65 ^{ef}
SPS+BAP	30	2.19±0.08 ^d	1.76±0.04 ^{cd}	73.73±4.03 ^c
	60	3.28±0.14 ^c	1.86±0.03 ^c	79.63±1.48 ^{abc}
	90	4.29±0.15 ^b	2.03±0.01 ^b	81.43±1.25 ^{ab}
DPS	30	1.14±0.14 ^f	1.49±0.02 ^e	50.10±0.26 ^e
	60	1.52±0.08 ^e	1.71±0.01 ^d	51.67±2.89 ^{de}
	90	1.48±0.08 ^e	2.01±0.01 ^c	56.53±0.92 ^d
DPS+BAP	30	2.14±0.15 ^d	1.47±0.02 ^e	77.77±3.78 ^{bc}
	60	4.14±0.14 ^b	1.74±0.03 ^d	81.57±1.40 ^{ab}
	90	8.66±0.17 ^a	2.14±0.09 ^a	84.70±0.25 ^a
Treatment Df(n-1)=11		1182.8 ^{***}	197.04 ^{***}	206.41 ^{***}
Culture system(S) Df(n-1)=3		17.05 ^{***}	40.74 ^{***}	95.95 ^{***}
Culture Duration(D) Df(n-1)=2		233.75 ^{***}	656.07 ^{***}	61.68 ^{***}
DxSdf (n-1)=11		105.06 ^{***}	42.35 ^{***}	0.17 ^{NS}

Means followed by same letter within a column are not significantly different ($p < 0.05$) as determined by Tukey's test

NS - Non significant, *** F value is highly significant at $p < 0.001$, ** significant at $p < 0.01$ and *significant at $p < 0.05$

Table 3. The effect of various auxins augmented - with half strength MS medium on rooting from *in vitro* generated shoots of *Vanilla andamanica*.

PGRs (µM)			Means no. of Roots/explant ±SE	Mean Root length in (cm) ±SE	Percentage of Response
IBA	IAA	NAA			
0	0	0	5.23±0.67 ^a	1.36±0.11 ^{cd}	100
0.5			4.61±0.36 ^{ab}	1.27±0.02 ^{cde}	100
1			4.57±0.53 ^{ab}	1.59±0.01 ^b	100
2			4.76±0.16 ^{ab}	1.92±0.02 ^a	100
	0.5		4.71±0.14 ^{ab}	1.17±0.06 ^{cde}	100
	1		4.80±0.16 ^{ab}	1.15±0.03 ^{de}	100
	2		4.80±0.08 ^{ab}	1.26±0.01 ^{cde}	100
		0.5	4.19±0.08 ^{ab}	1.13±0.02 ^e	100
		1	4.38±0.08 ^b	1.13±0.06 ^e	100
		2	4.42±0.38 ^b	1.38±0.17 ^{bc}	100
Treatment Df(n-1)=9			2.95 ^{NS}	34.94 ^{***}	-

Means followed by same letter within a column are not significantly different ($p < 0.05$) as determined by Tukey's test

NS - Non significant, *** F value is highly significant at $p < 0.001$, ** significant at $p < 0.01$ and *significant at $p < 0.05$

enhance shoot proliferation (18). Reduction of subculture with maintenance of explants for several months in a culture system had another advantage in double phase culture system, as reported earlier (13). A DPS system devoid of physical barriers increases nutrient availability and facilitates enhanced absorption by the explants, thereby promoting shoot growth. The DPS system ensures a higher quantity of nutrients in the culture system. From the present investigation, the evaluation of the efficiency of DPS in shoot enhancement of *V. andamanica* provides a cost-effective and reliable method for routine micropropagation and thereby ensures *in vitro* conservation strategies for this endangered and endemic species.

Rooting

Generally, influences of auxin quality and concentration showed a significant influence on rooting in the micropropagation technique. The observed results for *in vitro* rooting of micro stems with various concentrations of auxins are recorded in Table 3. Significant improvement in rooting was not observed in the addition of auxins, and the basal half strength MS medium provided the maximum

result in an average of 5.23±0.67 roots per explant (Fig. 1.D) with 100% rooting after 60 days of culture. However, IBA showed a maximum root length of 1.92±0.02cm (Fig. 1.E).

Spontaneous rooting in basal MS medium was exhibited by *V. andamanica* micro stems. Endogenous regulators for rooting at the optimum level were present in the nodal explant and are suggested as the reason for the spontaneous rooting of microstems even in the basal MS medium of *Vanilla* species (10). Several herbaceous species produce roots easily in auxin devoid or mediums with low levels of auxin, as already reported (8, 19, 20).

Hardening and acclimatization

In vitro produced rooted plants were taken out, washed thoroughly, and planted in plastic pots with river sand and garden soil (1:1). The potted plants were transferred to a greenhouse for acclimatization. It was observed that the plants raised from both SPS and DPS culture systems exhibited 100% survival rates (Fig. 1.F). A high percentage survival rate of *in vitro* regenerated *Vanilla* species was reported earlier by de Oliveira *et al.* (9).



Fig. 1. **A.** Habitat of *V. andamanica* **B.** *In vitro* produced multiple shoots with SPS after 60 days **C.** *In vitro* produced multiple shoots using DPS system (5 μ M BAP) after 90 days **D.** Basal MS medium producing roots without any PGRs **E.** IBA induced *in vitro* rooting **F.** Hardened *in vitro* plants of *V. andamanica*

Conclusion

The present investigation evaluated the efficiency of a double phase system and revealed that enhanced shoot multiplication of *V. andamanica* could be achieved through this novel technique rather than conventional methods. As regards large-scale propagation of *V. andamanica*, the DPS system can be used as a cost-effective and reproducible method that ensures its conservation, which may be applicable for other orchid members as well. This is the first report of the usage of DPS-mediated shoot proliferation in *V. andamanica*, an endangered endemic species on the road to extinction, from the Andaman and Nicobar Islands in the Bay of Bengal.

Acknowledgements

The authors are deeply indebted to University Grants Commission (UGC) for financial assistance for the

successful completion of work in the form of Junior Research Fellowship (F.No.16-6(DEC.2017)/2018(NET/CSIR)) and Head, Department of Botany, University of Kerala, India for providing the necessary facilities for doing the work.

Authors' contributions

SJ carried out the whole experiment and drafted the manuscript. AGP and SPM incorporated corrections to the manuscript. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None.

References

- Rodolphe G, Séverine B, Michel G, Pascale B. Biodiversity and evolution in the *Vanilla* genus. The dynamical processes of biodiversity-case studies of evolution and spatial distribution. 2011 Dec;2:1-27. <https://doi.org/10.5772/24567>
- Bory S, Grisoni M, Duval MF, Besse P. Biodiversity and preservation of *vanilla*: Present state of knowledge. 2008;551-71. <https://doi.org/10.1007/s10722-007-9260-3>
- Verma PC, Chakrabarty D, Jena SN, Mishra DK, Singh PK, Sawant SV, Tuli R. The extent of genetic diversity among *Vanilla* species: Comparative results for RAPD and ISSR. *Industrial Crops and Products*. 2009 Mar 1;29(2-3):581-89. <https://doi.org/10.1016/j.indcrop.2008.11.006>
- Lubinsky P. Conservation of wild *vanilla*. In *First International Congress on the Future of the Vanilla Business*. Princeton, NJ, USA. 2003 Nov 11; pp. 11-12.
- Minoo DI, Jayakumar VN, Veena SS, Vimala J, Basha A, Saji KV, Nirmal Babu K, Peter KV. Genetic variations and interrelationships in *Vanilla planifolia* and few related species as expressed by RAPD polymorphism. *Genetic Resources and Crop Evolution*. 2008 May;55:459-70. <https://doi.org/10.1007/s10722-007-9252-3>
- Divakaran M, Pillai GS, Babu KN, Peter KV. Isolation and fusion of protoplasts in *Vanilla* species. *Current Science*. 2008 Jan; 10:115-20.
- Murashige T, Skoog F. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia Plantarum*. 1962 Jul;15(3):473-97. <https://doi.org/10.1111/j.1399-3054.1962.tb08052.x>
- Abebe Z, Mengesha A, Teressa A, Tefera W. Efficient *in vitro* multiplication protocol for *Vanilla planifolia* using nodal explants in Ethiopia. *African Journal of Biotechnology*. 2009;8 (24). <https://doi.org/10.4314/AJB.V8I24.68674>
- de Oliveira SO, Sayd RM, Balzon TA, Scherwinski-Pereira JE. A new procedure for *in vitro* propagation of vanilla (*Vanilla planifolia*) using a double-phase culture system. *Scientia Horticulturae*. 2013 Sep 24;161:204-09. <https://doi.org/10.1016/j.scienta.2013.06.039>
- Divakaran M, Babu KN, Peter KV. Conservation of *Vanilla* species, *in vitro*. *Scientia Horticulturae*. 2006 Oct 9;110(2):175-80. <https://doi.org/10.1016/j.scienta.2006.07.003>
- Sota V, Benelli C, Çuko B, Kongjika E. Effect of a double phase culture system and activated charcoal on *in vitro* propagation of *Malus sylvestris* (L.) Mill. *Advances in Horticultural Science*. 2022 Oct 1;35(4):361. <https://doi.org/10.36253/ahsc-11825>
- Feuser S, Meler K, Daquinta M, Guerra MP, Nodari RO. Genotypic fidelity of micropropagated pineapple (*Ananas comosus*) plantlets assessed by isozyme and RAPD markers. *Plant Cell, Tissue and Organ Culture*. 2003 Mar;72:221-27. <https://doi.org/10.1023/A:1022321405683>
- Scherwinski-Pereira JE, Lima ED, da Silva TL, Mesquita AG, Maciel SD, da Silva Costa FH. Double-phase culture system for large scale production of pineapple. *Plant Cell, Tissue and Organ Culture*. 2012 May;109:263-69. <https://doi.org/10.1007/s11240-011-0091-8>
- Roy AR, Sajeev S, Pattanayak A, Deka BC. TDZ induced micropropagation in *Cymbidium giganteum* Wall. Ex Lindl. and assessment of genetic variation in the regenerated plants. *Plant Growth Regulation*. 2012;68:435-45. <https://doi.org/10.1007/s10725-012-9732-0>
- Moraes LK, Felisbino C, Crestani L, Silva AL. *In vitro* establishment and multiplication of *Pyrus calleryana* D-6 on double-phase culture system. *Revista Brasileira de Fruticultura*. 2004;26:403-05. <https://doi.org/10.1590/S0100-29452004000300008>
- Pereira JE, Fortes GR. Protocol for the production of potato propagating material in liquid medium. *Brazilian Agricultural Research*. 2003;38:1035-43. <https://doi.org/10.1590/S0100-204X2003000900003>
- López D C, Suárez IE. *In vitro* arrow cane (*Gynerium sagittatum* Aubl.) multiplication in double phase medium. *Revista de Ciencias Agrícolas*. 2018 Dec;35(2):5-13. <https://doi.org/10.22267/rcia.183502.86>
- Sandhyarani N, Kishor R, Sharma GJ. Clonal propagation of triploid *Acorus calamus* Linn. using dual-phase culture system. *Journal of Crop Science and Biotechnology*. 2011 Sep;14:213-17. <https://doi.org/10.1007/s12892-011-0028-0>
- Kalimuthu K, Senthilkumar R, Murugalatha N. Regeneration and mass multiplication of *Vanilla planifolia* Andr.–A tropical orchid. *Current Science*. 2006 Nov; 25:1401-03.
- Gurel E, Yucesan B, Aglic E, Gurel S, Verma SK, Sokmen M, Sokmen A. Regeneration and cardiotoxic glycoside production in *Digitalis davisiana* Heywood (*Alanya foxglove*). *Plant Cell, Tissue and Organ Culture (PCTOC)*. 2011 Feb;104:217-25. <https://doi.org/10.1007/s11240-010-9824-3>