



MINI REVIEW ARTICLE

# The role of polyamines in plants: A review

Aya Ihsan Rshan<sup>1</sup>, Rana T. Altaee<sup>2</sup>, Farah Sameer Salh<sup>3</sup>, Omar Younis AL-Abbasy<sup>4</sup> & Nashwan Al-Lehebe<sup>5\*</sup>

<sup>1,2,4,5</sup>Chem. Dept. College of Education for pure science. University of Mosul, Mosul, Iraq

<sup>3</sup>Branch of Basic Science, College of Agriculture and Forestry University of Mosul, Mosul, Iraq

\*Email: [nashwan78ibrahem@uomosul.edu.iq](mailto:nashwan78ibrahem@uomosul.edu.iq)



OPEN ACCESS

## ARTICLE HISTORY

Received: 21 March 2023

Accepted: 26 June 2023

Available online

Version 1.0 : 06 August 2023

Version 2.0 : 07 October 2023



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care etc. See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Rshan A I, Altaee R T, Salh F S, AL-Abbasy O Y, Al-Lehebe N. The role of polyamines in plants: A review. Plant Science Today. 2023; 10(sp2): 164-171.  
<https://doi.org/10.14719/pst.2520>

## Abstract

Polyamines (PAs) are linear alkali nitrogenous with two, three or more amino groups that have a low molecular weight. PAs are created by organisms through metabolism and can be found in nearly wholly cells. They are considered a unique type of plant stimulant because they act significant roles in many kinds of plant growth and developmental stages as well as stress responses. More evidence suggests that the PA can be produced endogenously or exogenously applied to improve plant reproduction, productivity, and bear stress. While the mechanism of polyamines influencing plant growth and stress responses is still not clear. We are making an attempt to supply an inclusive review of the available literature that explained the association between PAs and flowering, stress responses and senescence. This review aimed at this is focused and abbreviates how PAs enhance and increase plant productivity and then serve as a foundation for forthcoming study on the mechanisms of PAs exploit in plant reproduction and growth.

## Keywords

polyamine, polyamine oxidase, growth, proliferation

## Introduction

Polyamines (PAs) compounds like that spermine (Spm), spermidine (Spd), cadaverine (Cad) and putrescine (Put) are positively charged aliphatic molecules, play an essential role in the nucleic acid regulation, protein synthesis and structures, interactions of nucleic acid, protein, oxidative balance, and proliferation of the cell. (1,2). They can be found in prokaryotic as well as eukaryotic cells. (3,4). PAs are mostly found in free form in higher plants. Many physiological processes are regulated by them, including flower development(5), cell proliferation embryogenesis, morphology, organogenesis, senescence, root system formation, fruit maturation, architecture, induce a biosynthesis of osmotic alteration substances and responses to abiotic and biotic stresses and as an antioxidant (6,7,8,9), salinity and fruit maturity and development (10) also recognized as being crucial for plant growth (11). They are likewise elaborate in abiotic and biotic stress responses (12). Meanwhile, free polyamines contribute form covalent bonds with biological substances such as nucleic acids and proteins also lignin or phenolic compounds (13). Plants such as potatoes contain numerous phytochemicals like PAs, which are highly required in the diet because of their useful effects on health (14). In addition, PAs can maintain the stability of the membrane and aid in the scavenging of reactive oxygen species (ROS) in chloroplasts (15). It seems that PAs boost the activity of antioxidant enzymes and noticeably lessen the oxidative effects of ROS on membranes (16).

We review the latest research of PAs effects on plant growth, from flowering to salinity, and investigate their characteristics in response to various stresses.

## Methodology

The methodology of this article was designed to search and diagnose PAs effectiveness, which depended on the information and evidence available and published on the sciences platforms, Web of Science, Scopus, John Thompson Science Direct etc., for the approximately last 40 years. The article dealt with these compounds from several aspects focusing on the papers on plant kingdom only inclusion while excluding all tissues and organisms of humans and animals.

PAs compounds were discussed in terms of the chemical distribution compositions, biosynthesis, degradation and the effects of the products of these vital processes on plant cell divisions, and proliferation. The criteria of the effect of PAs in this article was discussed by searching and collecting information that indicates and confirms the effect of these compounds' free or combined with other negative compounds or ions on the sensitivity of plants towards environmental influences such as temperature, physiological pH or internal influences such as exposure to oxidative stress and their role as antioxidants by combined with membrane phospholipid of the cell to protect it. For obtaining all the information of this review, using some keywords were adopted, namely PAs compounds, types, the relationship of PAs to plant life, and the functions of PAs. It was found that they have an important role in the development and flowering, in addition to their significant role in enhancing plant resistance to environmental conditions.

## Results

### Distribution of PAs

Polyamines are abundant in eukaryotic and prokaryotic cells (17). PAs are distributed in plants in tissue and organ-specific forms and participate play in photosynthesis (18). The best common PAs in plants are Spm, Spd and Put, with high intracellular concentrations, and the least common are Cad and Dap (19,20). Many studies indicate the presence of polyamine compounds in rice, pistachio, green tea, mangoes, and mushrooms (21-23) noticed clear differences in the concentration of polyamine compounds in a variety of plants, and indicated that some plants contain high levels of Put and Spm in corn, citrus, and peas. Spd was also found at a higher level in soybeans, beans and castor tissues and these levels of plant leaves changed during the life of the leaf (24). Immature citrus seed species also were found to contain high levels of cadaverine (25,26).

On the other hand, an inverse relationship was observed between ascorbic acid with of Put and Spm levels in green peppers and tomatoes respectively (27). High levels of endogenous Put were found to be unfavourable for the accumulation of GA, and endogenous Put was discovered to be strictly interrelated to gibberellin

contents (28). The patterns of PA distribution may be linked to their unique functions. Generally, greater PA contents are associated with more robust plant growth and metabolism (29).

### Chemical nature of polyamines

The majority of PAs have a minimum of two positively charged amino or guanidinium groups, distinct by a carbon backbone with a range of lengths (Figure 1) (46). Different from other cations like  $Mg^{2+}$  or  $Ca^{2+}$ , PAs have no less than two charged groups linked by a flexible carbon chain. By doing this, electrostatic forces of amino or guanidine groups are enhanced by hydrophobic effects. Agmatine (AGM), which is a chemical compound, is a member of the PA family (47).





Fig 1. Chemical structure of polyamines (46)

### Biosynthesis of Polyamines

In all the kingdoms of living things, the PA biosynthetic pathway has been thoroughly investigated (48,49). The common PA biosynthetic pathway's main output is putrescine, in general. Through three distinct arginine-dependent pathways, plants can produce PAs from the proteinogenic amino acid L-Arginine. (Figure 2)(12). The first pathway is comparable to that of animals and fungi and involves the conversion of L-arginine by the mitochondrial arginase into ornithine, that is next decarboxylated via ornithine decarboxylase (ODC) to yield putrescine. Additionally, plants can produce agmatine by using the arginine decarboxylase (ADC) enzyme, which then catalyzes the conversion of agmatine to N-carbamoyl putrescine and putrescine using the enzymes agmatine iminohydrolase and N-carbamoyl putrescine amidohydrolase, respectively. ADC and agmatine ureohydrolase, respectively, can catalyze the reactions that result in L-arginine and agmatine contributing to the generation of Put (50-53).

The third pathway starts with the conversion of Arg to citrulline, that is before decarboxylated via citrulline decarboxylase to produce Put (54,55). The first two pathways are more prevalent in plants because the Cit pathway has only been discovered so far in sesame. Spermidine synthase (SPDS) generates spermidine from Put and amino-propyl, that is progressively supplied by L-methionine (56). Additionally, putrescine could be transformed into spermidine via the carboxy norspermidine decarboxylase and carboxy norspermidine dehydrogenase, respectively, using amide group provided



**Fig 2.** Biosynthesis pathway of main PAs in plants (12)

through the L-aspartate- $\beta$ -semialdehyde from the glycine catabolism pathway. In the presence of homospermidine synthase, Put and Spd both promote a synthesis of the homospermidine. Finally, by means of the enzymes spermidine synthase and thermospermidine synthase, respectively, spermidine is transformed into Spm or thermospermidine through a combination of an additional amino-propyl group (57,58). Additionally, some other PAs may be derived from other amino acids that are proteinogenic. For example, lysine decarboxylase (LDC) catalyzes the synthesis of the diamine cadaverine from L-lysine (59). The amino acids L- glycine, L-proline, L- glutamate, and L-methionine are effective sources in the biosynthesis of polyamines (60,61).

### Catabolism of Polyamines

The action of amine oxidases is crucial in the degradation of PAs in plants. Polyamine oxidase (PAO) provokes the formation of PA as metabolic end products. Wheat PAO, oxidizes Spm and Spd to produce 3-aminopropyl-4-amino butanal, 1,3-diamino propane, 4-amino butanal and hydrogen peroxide. The flavin adenine dinucleotide-dependent enzymes known as polyamine oxidases, which work in PA degradation, are essential in the pathway. (62,63). Tassoni (64) discovered that exo-Spd smeared to kinds of sunflower tuberos resulted in Put transformation. The generated  $H_2O_2$  plays a role in plant indication transduction through abiotic and biotic stress responses (65,66). The function of polyamines in helping plants adapt to salt stress conditions has been demonstrated. Under salt stress conditions, *Calendula officinalis* L. plants were given exogenous Spm and Spd to see what effects they had on their development and

physiological characteristics (67). Furthermore, PA metabolism is linked to the creation of NO (68), which is an important signalling constituent for plant growth (69). It has been demonstrated that polyamines and plant growth regulators are crucial for the start and control of the programmed cell death process. A variety of secondary metabolites known as phenol amides are produced when a phenolic moiety is conjugated with polyamines or deaminated aromatic amino acids. (70,71). The catabolism of PA compounds is a non-reversible process through which the cells can be supplied with carbon and nitrogen, which are used as precursors for cellular components (72). In order to better understand the functions of PAs in plant growth and development and the mechanisms underlying them, it is important to understand how PA metabolism interacts with plant hormones and how it affects signalling molecules.

### Physiological action of polyamines

#### Polyamines and the Flowering

Higher plants go through a phase of vegetative growth before beginning a phase of proliferation and growth, then leaf and bud tissues undergo differentiate physiological turnover to the tissues of flower bud, which then cultivates into a floral part of plants, it's referred to as flower bud reproduction (73). Putrescine, Spermidine, and Spermamine levels were elevated at prior developmental periods, whereas Putrescine and Spermamine levels stayed constant (74). Polyamines, which are important and ubiquitous molecules required for cell growth and differentiation, are oxidatively de-aminated by copper amine oxidases and amine oxidases that include flavin (75). PA is thought to be a plant growth regulator (10).

Polyamines and reproductive development appear to be closely related, according to a number of lines of experimental data, it was found that PAs were more prevalent in *Arabidopsis* flowers in any other organ, and adding invitros PAs to plants that were not flowering significantly increased their flowering response (76). Gibberellic acid (GA3) and spermamine (SPM), two plant growth regulators, have been characterized as extending the postharvest life of flowers (77) Spm enhanced the quality of the flower and vase life of cut rose blossoms (78). On the other hand, feeding of Spd is directly correlated to the time of flowering in *Arabidopsis thaliana* (79).

**Table 1.** Distribution of Polyamines

|   |               |  |                          |
|---|---------------|--|--------------------------|
| 1 | Put           | broccoli, cauliflower, eggplant, jiló (scarlet eggplant), tomato, green onion, parsley, spinach, capers, cassava, heart of palm and bean sprouts                                       | 30                       |
| 2 | Put, spd, spm | Soybeans   | 31                       |
| 3 | Spm, Spd, put | Maize grain, Common wheat  | 32                       |
| 4 | Spm, Spd, put | Fruits: Apple – bananas – cherry – kiwi – pineapple – orange - mandarin - avocados, strawberry   | 33,34,35,36              |
| 5 | Spm, Spd, put | Vegetables: Broccoli – cucumber – cauliflower – celeriac – courgette – eggplant - cabbage green beans – lettuce – mushroom - green pepper – onion - potato – carrot – spinach – tomato | 37,38                    |
| 6 | Spm, Spd, put | Legumes: lentils - soybean - red beans – peas - soybean milk - tofu – sprouts – miso   | 34,36,37, 39,40,41,42,43 |
| 7 | Spm, Spd, put | Cereals: wheat germ – Rice - white bread   | 34, 35, 37, 44           |
| 8 | Spm, Spd, put | Nuts: chestnuts - Almonds - pistachios   | 34, 39                   |
| 9 | Spm, Spd, put | spinach leaf cells - intact chloroplasts- thylakoid membranes  | 45                       |

### Polyamines and Plant Senescence

PAs change as the plant grows. Endogenous PAs were found to be lowest in the senescent tissues in whole plants (80). A decrease in levels of PA appears to be an important precursor to senescence signals (81). Delayed leaf senescence was linked to higher Spm levels and higher NO levels (82). (83) discovered that chlorophyll degraded quickly and Put collected through senescence, whereas exogenous Spm or Spd inhibited protein breakdown and reduced chlorophyll harm. PA synthesis inhibitors in peonies extended the lifespan and delayed flower senescence, but PAs reduced faster flower senescence (84). Even during nematode pathogenesis, the tomato plant's physiological reactions to the polyamines were found to be improved (85).

### Polyamines and Temperature Stress

PAs can stimulate photosynthesis while also rising plant antioxidant ability and osmotic adjustment ability under high-temperature stress (86). PAs serve numerous roles in all plants, and the primary physiological processes underlying the tolerance of high-temperature vary between plant species. PAs have the ability to bind to the cell membrane's phospholipid site, increasing cold resistance and preventing cytolysis (87). When the chilling temperature was applied to sweet pepper and zucchini fruits, the Put content expanded rapidly, accompanied by terrifying deterioration (88). Put accumulation, on the other hand, enhanced Spm was a defence reaction to cold damage, which caused chilling damage. They found that the spermidine, spermine, and Put levels eventually rose when loquat fruit was kept at low temperatures (89).

### The Role of Polyamines with Oxidative Stress

PAs can activate a variety of enzymes known as antioxidants in plants, permitting them to successful defence against reactive oxygen species and then decrease oxidative stress, which is caused by a diversity of environmental triggers. Spm and Put pretreatment of maize leaves improved their resistance to the oxidative damage caused by paraquat (90). To combat the detrimental impact of different environmental stressors on plant processes polyamines have been advocated as a viable option. (91) Under hypoxia stress, exogenous Spd elevated the amount of Spd and Spm while reducing the amount of Putrescin in cucumber seedling rootstock. (92). It was also discovered that cold-treated or cold polyamine-treated groups had higher specific antioxidant enzyme activity than comparable controls, including catalase and ascorbate peroxidase. The treated plants' H<sub>2</sub>O<sub>2</sub> and malondialdehyde levels were significantly reduced as a result (93). The impacts of Cd<sup>2+</sup> and Cu<sup>2+</sup> on the peroxidation of lipids were reduced by Spm treatment. (94). PAs have the potential to harm cells under conditions of stress because their degradation generates the powerful oxidizers hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and acrolein. (95). H<sub>2</sub>O<sub>2</sub> is, on the other hand, a molecule that signals that may pass through a stressful signal transduction pathway.

As a result, PAs appear to be redox regulators of homeostasis in plants that show a dual role in oxidative stress (96). Numerous studies revealed that polyamine

oxidase (PAO) genes are essential for a variety of abiotic and biotic stress responses (97).

### Discussion

All living organisms contain polyamines, which are involved in a variety of metabolic processes in plants like cell division, proliferation, expansion, differentiation, and programmed cell death to protect them from various stresses (98). They are important for cells and are known as endogenous growth regulators (99).

The capability of polyamines to interact via negatively charged sites in molecules like lipids, proteins, and nucleic acids is one of the reasons for their wide range of functions, also some natural compounds like phenolic compounds. Additionally, it has been discovered that a few byproducts of polyamine catabolism play important signalling functions in a variety of physiological, cellular, and developing processes (100). For example, transglutaminases (TGases) covalently link polyamines to particular proteins. TGases play a central role in photosynthesis in plants (101). According to Belda-Palazón et al. (102), the spermidine-dependent hypusination pathway may have novel roles in plant improvement and responses to hormonal, dietary, and environmental signals.

According to research by Nambeesan et al. (103), PAs have also been revealed to enhance longevity in a number of organisms, including tomato fruit, and to extend the shelf life of tomato fruit. They also participate in both transcriptional and translational gene regulation (104). In previous research, tomato fruits with high levels of the anabolic biogenic amines Spm and Spd, which are produced as a result of yeast S-adenosyl methionine decarboxylase expression, would increase an expression of SA biosynthesis genes, resulting in increased SA creation.

Putrescine N-methyltransferase is a crucial enzyme of the second metabolism and is involved in the biosynthesis of tropane alkaloids, nicotine and calystegines. An in-silico study demonstrated structural as well as functional characterization and developed three-dimensional computational models of putrescine N-methyltransferase proteins. Recently the number of reports illustrating that crosstalk between PA and phytohormones in plants to respond to abiotic stresses has increased and exposed the favorable influence of the co-application of exogenous PA and phytohormones (105).

### Conclusion

In this review, we conclude that polyamines are active natural compounds having at least two positively charged amino groups that play important roles in many biological purposes in plants. The involvement of polyamines in a variety of plant processes, including DNA replication, gene transcription, organ development, cell division, fruit development and leaves senescence, ripening, and abiotic stresses, is supported by a substantial body of research. As a result, the main genes ADC, ODC, or SAMDC have been overexpressed or down regulated, altering the amount of

PA in cells. Overexpression of heterologous ADC or ODC typically results in increased Put levels without affecting Spd or Spm. In contrast, overexpression of the pathway's genes downstream (SAMDC or Spd synthase) typically results in higher levels of Spd and Spm, demonstrating that the levels of Spm and Spd are tightly regulated by cellular homeostasis. The phytochemical, polyamines have an important impact on the contents of bioactive compounds with antioxidant activity besides in the activity of the key antioxidant enzymes in fruits during the ripening and senescence processes. It can be inferred that polyamines have wide applications due to their effect on the properties of phenolic compounds and flavonoids.

## Acknowledgements

The authors are appreciative of those who provided the resources needed to finish this work. Special thanks go out to the Chemistry Department at the College of Education for Pure Sciences, University of Mosul.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interest to declare.

**Ethical issues:** None .

## References

1. Vuosku J, Karppinen K, Muilu-Mäkelä R, Kusano T, Sagor GHM, Avia K. Scots pine aminopropyltransferases shed new light on evolution of the polyamine biosynthesis pathway in seed plants. *Ann Bot.* 2018;121:1243-56. <https://doi.org/10.1093/aob/mcy012>.
2. Zahedi K, Barone S, Soleimani M. Polyamines and their metabolism: From the maintenance of physiological homeostasis to the mediation of disease. *Med Sci.* 2022;10:38. <https://doi.org/10.3390/medsci10030038>.
3. Liu W, Tan M, Zhang C, Al E. Functional characterization of murB-potABCD operon for polyamine uptake and peptidoglycan synthesis in *Streptococcus suis*. *Microbiol Res.* 2017;207:177-87. <https://doi.org/10.1016/j.micres.2017.11.008>.
4. Mustafavi SH, Badi HN, Sekara A, Al E. Polyamines and their possible mechanisms involved in plant physiological processes and elicitation of secondary metabolites. *Acta Physiol Plant.* 2018;40:102. <https://doi.org/10.1007/s11738-018-2671-2>.
5. Chen D, Shaho Q, Yin L, Younis A. Polyamine function in plants: Metabolism, regulation on development and roles in abiotic stress responses. *Plant Abiotic stress.* 2018;9. <https://doi.org/10.3389/fpls.2018.01945>.
6. Reis RS, Vale EM, Heringer AS, Al E. Putrescine induces somatic embryo development and proteomic changes in embryogenic callus of sugarcane. *J Proteomics.* 2016;130:170-79. <https://doi.org/10.1016/j.jprot.2015.09.029>.
7. Tiburcio AF, Alcazar R. Potential applications of polyamines in agriculture and plant biotechnology. *Methods Mol Biol.* 2018;1694:489-508. [https://doi.org/10.1007/978-1-4939-7398-9\\_40](https://doi.org/10.1007/978-1-4939-7398-9_40).
8. Miyamoto M, Shimao S, Tong W, Motose H, Takahashi T. Effect of thermospermine on the growth and expression of polyamine-related genes in rice seedlings. *Plants.* 2019;269. <https://doi.org/10.3390/plants8080269>.
9. Mohammadi H, Ghorbanpour M, Brestic M. Exogenous putrescine changes redox regulations and essential oil constituents in field-grown *Thymus vulgaris* L. under well-watered and drought stress conditions. *Ind Crops Prod.* 2018;122:119-32. <https://doi.org/10.1016/j.indcrop.2018.05.064>.
10. Xu L, Xing ST, Sun X. Effects of polyamines on hormones contents and the relationship with the flower bud differentiation in chrysanthemum. *Plant Physiol.* 2014;50:1195-202. <https://doi.org/10.13592/j.cnki.ppj.2014.0212>.
11. de Oliveira LF, Elbl P, Navarro BV, Al E. Elucidation of the polyamine biosynthesis pathway during Brazilian pine (*Araucaria angustifolia*) seed development. *Tree Physiol.* 2016;37:116-30. <https://doi.org/10.1093/treephys/tpw107>.
12. Anshika T, Sajad A, Goriparthi R, Anupam S, Suvin P, Henda M, Hanhong B. Revisiting the role of polyamines in plant growth and abiotic stress resilience: Mechanisms, crosstalk and future perspectives. *Journal of Plant Growth Regulation.* 2022; <https://doi.org/10.1007/s00344-022-10847-3>.
13. Holbert CE, Foley JR, Yu A, Murray Stewart T, Phanstiel OIV, Oupicky D, Casero RA Jr. Polyamine-based nanostructures share polyamine transport mechanisms with native polyamines and their analogues: Significance for polyamine-targeted therapy. *Med Sci.* 2022;10:44. <https://doi.org/10.3390/medsci10030044>.
14. Ezekiel PR, Singh N, Sharma S, Kaur A . Beneficial phytochemicals in potato a review. *Food Research International.* 2013;50(2):487-96. <https://doi.org/10.1016/j.foodres.2011.04.025>
15. Kubís J, Floryszak-Wieczorek J, Arasimowicz-Jelonek M. Polyamines induce adaptive responses in water deficit stressed cucumber roots. *J Plant Res.* 2014;127:151-58. <https://doi.org/10.1007/s10265-013-0585-z>.
16. Li J, Hu L, Zhang L, Pan X, Hu X. Exogenous spermidine is enhancing tomato tolerance to salinity-alkalinity stress by regulating chloroplast antioxidant system and chlorophyll metabolism. *BMC Plant Biol.* 2015;15:303. <https://doi.org/10.1186/s12870-015-0699-7>.
17. Dandan C, Qingsong S, Lianghong Y, Adnan Y, Bingsong Z. Polyamine function in plants: Metabolism, regulation on development and roles in abiotic stress responses. *Front Plant Sci.* 2019;9:1945. <https://doi.org/10.3389/fpls.2018.01945>.
18. Bograh A, Gingras Y, Tajmir RH, Carpentier R. The effects of spermine and spermidine on the structure of photo system II proteins in relation to inhibition of electron transport. *FEBS Lett.* 1997;402(1):41-44. [https://doi.org/10.1016/S0014-5793\(96\)01453-6](https://doi.org/10.1016/S0014-5793(96)01453-6).
19. Takahashi Y, Tahara M, Yamada Y, Al E. Characterization of the polyamine biosynthetic pathways and salt stress response in *Brachypodium distachyon*. *J Plant Growth Regul.* 2017;37:625-34. <https://doi.org/10.1007/s00344-017-9761-z>
20. Del Duca S, Serafini-Fracassini D, Cai G. Senescence and programmed cell death in plants: polyamine action mediated by transglutaminase. *Plant Polyamines in Stress and Development.* 2014;98. <https://doi.org/10.3389/fpls.2014.00120>.
21. Nishimura K, Kashiwagi SK, Igarashi K. Decrease in polyamines with aging and their ingestion from food and drink. *J Bioche.* 2006;139:81-90. <https://doi.org/10.1093/jb/mvj003>.
22. Igarashi K, Kashiwagi K. Polyamines: mysterious modulators of cellular functions. *Biochem Biophys Res Commun.* 2000;19:271 (3):559-64. <https://doi.org/10.1006/bbrc.2000.2601>.
23. Goni S, Pathak AK, Sharma RK, Rastogi A. Polyamines: As futuristic feed additives. *Appro Poult Dairy and Vet Sci.* 2018;4 (5). <https://doi.org/10.31031/APDV.2018.04.000600>
24. Antognoni F, Fornale S, Grimmer C, Komor E, Bagni N. Long-distance translocation of polyamines in phloem and xylem of *Ricinus communis* L. plants. *Planta.* 1998;204(4):520-27. <https://doi.org/10.1007/s004250050287>.

25. Tassoni A, Germanà MA, Bagni N. Free and conjugated polyamine content in *Citrus sinensis* Osbeck, cultivar Brasileiro NL. A navel orange, at deferent maturation stages. *Food Chem.* 2004;87:537-41. <https://doi:10.1016/j.foodchem.2004.01.001>.
26. Shahid MA, Balal RM, Khan N, Rossi L, Rathinasabapathi B, Liu G, Khan J, Cámara-Zapata JM, Martínez-Nicolas JJ, Garcia-Sanchez F. Polyamines provide new insights into the biochemical basis of Cr-tolerance in Kinnow mandarin grafted on diploid and double-diploid rootstocks. *Environ Exp Bot.* 2018;156:248-60. <https://doi:10.1016/j.envexpbot.2018.09.015>.
27. Yahia EM, Contreras M, Gonzalez-Aguilar G. Ascorbic acid content in relation to ascorbic acid oxidase activity and polyamine content in tomato and bell pepper fruits during development, maturation and senescence. *Lebensm Wiss Technol Food Sci Technol.* 2001;34(7):452-57. <https://doi.org/10.1006/fstl.2001.0790>
28. Xu C, Wu X, Zhang H. Impact of D-Arg on drought resistance and endogenous polyamines in mycorrhizal *Pinus massoniana*. *J Nanjing Forestry Univ.* 2009;33:019-023. <https://doi:10.3969/j.issn.1000-2006.2009.04.004>.
29. Cai Q, Zhang J, Guo C, Al E. Reviews of the physiological roles and molecular biology of polyamines in higher plants. *J Fujian Educ Coll.* 2006;7:118-24. <https://doi:10.3969/j.issn.1673-9884.2006.10.039>.
30. Bruno MDala-Paula B, Maria de Fátima V Starling AC, M Beatriz AG. Vegetables consumed in Brazilian cuisine as sources of bioactive amines. *Food Bioscience.* 2021;40(6):100856. <https://doi.org/10.1016/j.fbio.2020.100856>.
31. Tatsuya S, Dhaka RB, Bernhard S, Johann V. Spermidine and other functional phytochemicals in soybean seeds: Spatial distribution as visualized by mass spectrometry imaging. *Food Sci Nutr.* 2019;8:675-82. <https://doi.org/10.1002/fsn3.1356>
32. Koc EC, Bagga S, Songstad DD, Betz SR, Kuehn GD, Phillips GC. Occurrence of uncommon polyamines in cultured tissues of maize. *In Vitro Cellular & Developmental Biology- Plant.* 1998;34:252-55. <https://doi:10.1104/pp.94.3.855>.
33. Eliassen KA, Reistad R, Risoen U, Ronning HF. Dietary polyamines. *Food Chem.* 2002;78:273-80. [https://doi:10.1016/S0308-8146\(01\)00405-8](https://doi:10.1016/S0308-8146(01)00405-8).
34. Nishimura K, Shiina R, Kashiwagi K, Igarashi K. Decrease in polyamines with aging and their ingestion from food and drink. *J Biochem.* 2006;139:81-90. <https://doi:10.1093/jb/mvj003>.
35. Okamoto A, Sugi E, Koizumi Y, Yanadiga F, Udaka S. Polyamine content of ordinary foodstuffs and various fermented foods. *Biosci Biotechnol Biochem.* 1997;61:1582-84. <https://doi:10.1271/bbb.61.1582>.
36. Brian MD, Jeff SR. Determination of biogenic amines in alcoholic beverages by ion chromatography with suppressed conductivity detection and integrated pulsed amperometric detection. *J Chrom A.* 2007;1155(1):22-30. <https://doi:10.1016/j.chroma.2007.01.114>.
37. Byun BY, Bai X, Mah JH. Occurrence of biogenic amines in Doubanjiang and Tofu. *Food Sci Biotech.* 2013;22:55-62. <https://doi:10.1007/s10068-013-0008-x>.
38. Kalač P, Krížek M, Pelikánová T, Langová M, Veskrna O. Contents of polyamines in selected foods. *Food Chem.* 2005;90:561-64. <https://doi:10.1016/j.foodchem>.
39. Nishibori N, Fujihara S, Akatuki T. Amounts of polyamines in foods in Japan and intake by Japanese. *Food Chem.* 2007;100:491-97. <https://doi:10.1016/j.foodchem.2005.09.070>.
40. Moret S, Smela D, Populin T, Conte L. A survey on free biogenic amine content of fresh and preserved vegetables. *Food Chem.* 2005;89:355-61. <https://doi:10.1016/j.foodchem.2004.02.050>.
41. Ziegler W, Hahn M, Wallnöfer PR. Changes in biogenic amine contents during processing of several plant foods. *Deut Lebensm Rundsch.* 1994;90:108-12. <https://doi:10.3390/food7120205>.
42. Kalac P, Svecová S, Pelikánová T. Levels of biogenic amines in typical vegetable products. *Food Chem.* 2002;77:349-51. [https://doi:10.1016/S0308-8146\(01\)00360-0](https://doi:10.1016/S0308-8146(01)00360-0).
43. Cipolla BG, Havouis R, Moulinoux JP. Polyamine contents in current foods: A basis for polyamine reduced diet and a study of its long-term observance and tolerance in prostate carcinoma patients. *Amino Acids.* 2007;33:203-12. <https://doi:10.1007/s00726-007-0524-1>.
44. Preti R, Rapa M, Vinci G. Effect of Steaming and boiling on the antioxidant properties and biogenic amines content in Green Bean (*Phaseolus vulgaris*) varieties of different colours. *J Food Qual.* 2017;5329070. <https://doi:10.1155/2017/5329070>.
45. Toro Funes N, Bosch Fusté J, Latorre Moratalla ML, Veciana Nogués MT, Vidal Carou MC. Biologically active amines in fermented and non-fermented commercial soybean products from the Spanish market. *Food Chem.* 2015;173:1119-24. <https://doi:10.1016/j.foodchem.2014.10.118>.
46. Kabir A, Kumar GS. Binding of the biogenic polyamines to deoxyribonucleic acids of varying base composition: Base specificity and the associated energetics of the interaction. *PLoS ONE.* 2013;8:e70510. <https://doi.org/10.1371/journal.pone.0070510>
47. Sagar NA, Tarafdar S, Agarwa S, Tarafdar A, Sharma S. Polyamines: functions, metabolism and role in human disease management. *Med Sci (Basel).* 2021;9(2):44. <https://doi:10.3390/medsci9020044>
48. Koc EC, Bagga S, Songstad DD, Betz SR, Kuehn GD, Phillips GC. Occurrence of uncommon polyamines in cultured tissues of maize. *In Vitro Cellular & Developmental Biology- Plant.* 1998;34:252-55. <https://doi:10.1104/pp.94.3.855>.
49. Eliassen KA, Reistad R, Risoen U, Ronning HF. Dietary polyamines. *Food Chem.* 2002;78:273-80. [https://doi:10.1016/S0308-8146\(01\)00405-8](https://doi:10.1016/S0308-8146(01)00405-8).
50. Xu C, Wu X, Zhang H. Impact of D-Arg on drought resistance and endogenous polyamines in mycorrhizal *Pinus massoniana*. *J Nanjing Forestry Univ.* 2009;33:019-023. <https://doi:10.3969/j.issn.1000-2006.2009.04.004>.
51. Pegg AE. Functions of polyamines in mammals. *J Biol Chem.* 2016;291:14904-912. <https://doi:10.1074/jbc.R116.731661>.
52. Nishimura K, Shiina R, Kashiwagi K, Igarashi K. Decrease in polyamines with aging and their ingestion from food and drink. *J Biochem.* 2006;139:81-90. <https://doi:10.1093/jb/mvj003>.
53. Okamoto A, Sugi E, Koizumi Y, Yanadiga F, Udaka S. Polyamine content of ordinary foodstuffs and various fermented foods. *Biosci Biotechnol Biochem.* 1997;61:1582-84. <https://doi:10.1271/bbb.61.1582>.
54. Paolo P, Gabriella F, Denis B, Andrea T, Silvano C, Giovanna S. Determination of biogenic amines in fruit, vegetables and chocolate using ion chromatography with suppressed, conductivity and integrated pulsed amperometric detections. *Appl Update.* 2005;162:1-8. <https://doi:10.1016/j.chroma.2005.08.065>.
55. Ouyang J, Song C, Chen D. Research progress on heat-tolerance mechanism and transports of polyamines in plant. *Mol Plant Breed.* 2017;15:3286-94. <https://doi:10.13271/j.mpb.015.003286>.
56. De Oliveira LF, Navarro BV, Cerruti G, Al E. Polyamines and amino acid related metabolism: the roles of arginine and ornithine are associated with the embryogenic potential. *Plant Cell Physiol.* 2018;59:1084-98. <https://doi:10.1093/pcp/pcy049>.

57. Hanzawa Y, Takahashi T, Michael AJ, Burtin D, Long D, Pineiro M, Coupland G, Komeda Y. ACAULIS5, an Arabidopsis gene required for stem elongation, encodes a spermine synthase. *EMBO J.* 2000;19:4248-56. <https://doi.org/10.1093/emboj/19.16.4248>.
58. Byun BY, Bai X, Mah JH. Occurrence of biogenic amines in Doubanjiang and Tofu. *Food Sci Biotech.* 2013;22:55-62. <https://doi.org/10.1007/s10068-013-0008-x>.
59. Kalac P, Krizek M, Pelikánová T, Langová M, Veskrna O. Contents of polyamines in selected foods. *Food Chem.* 2005;90:561-64. <https://doi.org/10.1016/j.foodchem.2005.09.070>.
60. Nishibori N, Fujihara S, Akatuki T. Amounts of polyamines in foods in Japan and intake by Japanese. *Food Chem.* 2007;100:491-97. <https://doi.org/10.1016/j.foodchem.2005.09.070>.
61. Moret S, Smela D, Populin T, Conte L. A survey on free biogenic amine content of fresh and preserved vegetables. *Food Chem.* 2005;89:355-61. <https://doi.org/10.1016/j.foodchem.2004.02.050>.
62. Hao Y, Huang B, Jia D, Al E. Identification of seven polyamine oxidase genes in tomato (*Solanum lycopersicum* L.) and their expression profiles under physiological and various stress conditions. *J Plant Physiol.* 2018;228:1-11. <https://doi.org/10.1016/j.jplph.2018.05.004>.
63. Liu T, Kim DW, Niitsu M, Al E. Polyamine oxidase 7 is a terminal catabolism-type enzyme in *Oryza sativa* and is specifically expressed in anthers. *Plant Cell Physiol.* 2014;55:1110-22. <https://doi.org/10.1093/pcp/pcu047>.
64. Tassoni A, Buuren MV, Franceschetti M, Al E. Polyamine content and metabolism in *Arabidopsis thaliana* and effect of spermidine on plant development. *Plant Physiol Biochem.* 2000;38:383-93. [https://doi.org/10.1016/S0981-9428\(00\)00757-9](https://doi.org/10.1016/S0981-9428(00)00757-9).
65. Freitas VS, Miranda RDS, Costa JH, Al E. Ethylene triggers salt tolerance in maize genotypes by modulating polyamine catabolism enzymes associated with H<sub>2</sub>O<sub>2</sub> production. *Environ Exp Bot.* 2017;145:75-86. <https://doi.org/10.1016/j.envexpbot.2017.10.022>.
66. Sen S, Ghosh D, Mohapatra S. Modulation of polyamine biosynthesis in *Arabidopsis thaliana* by a drought mitigating *Pseudomonas putida* strain. *Plant Physiol Biochem.* 2018;129:180-88. <https://doi.org/10.1016/j.plaphy.2018.05.034>.
67. Baniasadi F, Saffari VR, Moud AAM. Physiological and growth responses of *Calendula officinalis* L. plants to the interaction effects of polyamines and salt stress. *Sci Horticult.* 2018;234:312-17. <https://doi.org/10.1016/j.scienta.2018.02.069>.
68. Pál M, Szalai G, Janda T. Speculation: polyamines are important in abiotic stress signaling. *Plant Sci.* 2015;237:16-23. <https://doi.org/10.1016/j.plantsci.2015.05.003>.
69. Agurla S, Gayatri G, Raghavendra AS. Polyamines increase nitric oxide and reactive oxygen species in guard cells of *Arabidopsis thaliana* during stomatal closure. *Protoplasma.* 2017;255:153-62. <https://doi.org/10.1007/s00709-017-1139-3>.
70. Bassard JE, Ullmann P, Bernier F, Al E. Phenolamides: bridging polyamines to the phenolic metabolism. *Phytochemistry.* 2010;71:1808-24. <https://doi.org/10.1016/j.phytochem.2010.08.003>.
71. Cai G, Sobieszczuknowicka E, Aloisi I, Al E. Polyamines are common players in different facets of plant programmed cell death. *Amino Acids.* 2015;47:27-44. <https://doi.org/10.1007/s00726-014-1865-1>.
72. Bachrach U, Abu-Elheiga L, Talmi M, Schnur LF, El-On J, Greenblatt CL. Polyamines and the growth of leishmanial parasites. *Med Biol.* 1981Dec;59(5-6):441-47. <https://doi.org/10.3390/medsci10020024>.
73. Guo J, Tian L, Sun XZ, Al E. Relationship between endogenous polyamines and floral bud differentiation in *Chrysanthemum morifolium* under short-day conditions. *Wonye kwahak kisluchi.* 2015;33:31-38. <https://doi.org/10.7235/hort.2015.14043>.
74. Agudelo-Romero P, Bortolotti C, Pais MS, Al E. Study of polyamines during grape ripening indicate an important role of polyamine catabolism. *Plant Physiol Biochem.* 2013;67:105-19. <https://doi.org/10.1016/j.plaphy.2013.02.024>.
75. Cona A, Rea G, Angelini R, Al E. Functions of amine oxidases in plant development and defence. *Trends Plant Sci.* 2006;11:80-88. <https://doi.org/10.1016/j.tplants.2005.12.009>.
76. Applewhite PB, Kaur-Sawhney R, Galston AW. A role for spermidine in the bolting and flowering of *Arabidopsis*. *Physiol Plant.* 2003;108:314-20. <https://doi.org/10.1034/j.1399-3054.2000.108003314.x>.
77. Simões ADN, Diniz NB, Vieira MRDS, Al E. Impact of GA3 and spermine on postharvest quality of anthurium cut flowers (*Anthurium andraeanum*) cv. Arizona. *Sci Horticult.* 2018;241:178-86. <https://doi.org/10.1016/j.scienta.2018.06.095>.
78. Tenreiro S, Nunes PA, Viegas CA, Neves MS, Teixeira MC, Cbral MG, Sa-Correia I. AQR1 gene (ORF YNL065w) encodes a plasma membrane transporter of the major facilitator superfamily that confers resistance to short-chain monocarboxylic acids and quinidine in *Saccharomyces cerevisiae*. *Biochem Biophys Res Commun.* 2002;292:741-48. <https://doi.org/10.1006/bbrc.2002.6703>.
79. Ahmed S, Ariyaratne M, Patel J, Al E. Altered expression of polyamine transporters reveals a role for spermidine in the timing of flowering and other developmental response pathways. *Plant Sci.* 2017;258:146-55. <https://doi.org/10.1016/j.plantsci.2016.12.002>.
80. Takahashi T. Plant polyamines. *Plants.* 2020;9:511. <https://doi.org/10.3390/plants9040511>.
81. Duan G, Huang Z, Lin H. The role of polyamines in the ontogeny of higher plants. *Acta Agric Boreali Occidentalis Sinica.* 2006;15:190-94. <https://doi.org/10.7606/j.issn.1004-1389.2006.02.050>.
82. Sobieszczuk-Nowicka E. Polyamine catabolism adds fuel to leaf senescence. *Amino Acids.* 2017;49:49-56. <https://doi.org/10.1007/s00726-016-2377-y>.
83. Cai G, Sobieszczuknowicka E, Aloisi I, Al E. Polyamines are common players in different facets of plant programmed cell death. *Amino Acids.* 2015;47:27-44. <https://doi.org/10.1007/s00726-014-1865-1>.
84. Nakanishi S, Cleveland JL. Polyamine homeostasis in development and disease. *Med Sci.* 2021;9:28. <https://doi.org/10.3390/medsci9020028>.
85. Khajuria A, Ohri P. Exogenously applied putrescine improves the physiological responses of tomato plant during nematode pathogenesis. *Sci Horticult.* 2018;230:35-42. <https://doi.org/10.1016/j.scienta.2017.11.021>.
86. Anna F, Bekebrede JK, Walter JJ, Vincent CJ. The molecular and physiological effects of protein-derived polyamines in the intestine. *Nutrients.* 2020;12(1):197. <https://doi.org/10.3390/nu12010197>.
87. Li Y, He J. Advance in metabolism and response to stress of polyamines in plant. *Acta Agric Boreali Sinica.* 2012;27:240-45. <https://doi.org/10.3969/j.issn.1000-7091.2012.z1.048>.
88. Wu M, Yuan L. Research progress in the relationship between polyamine and plant resistance. *J Anhui Agric Sci.* 2008;36:3516-18. <https://doi.org/10.3969/j.issn.0517-6611.2008.09.020>.
89. Anwar R, Fatima Sh, Mattoo AK, K Handa A. Fruit architecture in polyamine-rich tomato germplasm is determined via a medley of cell cycle, cell expansion and fruit shape genes. *Plants.* 2019;8:387. <https://doi.org/10.3390/plants8100387>.

90. Durmu N, Kadioglu A. Spermine and putrescine enhance oxidative stress tolerance in maize leaves. *Acta Physiol Plant.* 2005;27:515-22. <https://doi:10.1007/s11738-005-0057-8>.
91. Mohammadi H, Ghorbanpour M, Brestic M. Exogenous putrescine changes redox regulations and essential oil constituents in field-grown *Thymus vulgaris* L. under well-watered and drought stress conditions. *Ind Crops Prod.* 2018;122:119-32. <https://doi:10.1016/j.indcrop.2018.05.064>
92. Wu J, Shu S, Li C, Sun J, Guo S. Spermidine-mediated hydrogen peroxide signaling enhances the antioxidant capacity of salt-stressed cucumber roots. *Plant Physiol Biochem.* 2018;128:152-62. <https://doi:10.1016/j.plaphy.2018.05.002>.
93. Peynevandi KM, Razavi SM, Zahri S. The ameliorating effects of polyamine supplement on physiological and biochemical parameters of *Stevia rebaudiana* Bertoni under cold stress. *Plant Prod Sci.* 2018;21:123-31. <https://doi:10.1080/1343943X.2018.1437756>
94. María D Groppa, María L Tomaro, María P Benavides. Polyamines as protectors against cadmium or copper-induced oxidative damage in sunflower leaf discs. *Plant Sci.* 2001;161:481-88. [https://doi.org/10.1016/S0168-9452\(01\)00432-0](https://doi.org/10.1016/S0168-9452(01)00432-0).
95. Minocha R, Majumdar R, Minocha SC. Polyamines and abiotic stress in plants: A complex relationship. *Front Plant Sci.* 2014;5:175. <https://doi:10.3389/fpls.2014.00175>.
96. Saha J, Brauer EK, Sengupta A, Al E. Polyamines as redox homeostasis regulators during salt stress in plants. *Front Environ Sci.* 2015;3:21. <https://doi:10.3389/fenvs.2015.00021>.
97. Cheng XQ, Zhu XF, Tian WG, Al E. Genome-wide identification and expression analysis of polyamine oxidase genes in upland cotton (*Gossypium hirsutum* L.). *Plant Cell Tissue Organ Cult.* 2017;129:237-49. <https://doi:10.1007/s11240-017-1172-0>.
98. Handa AK, Mattoo AK. Differential and functional interactions emphasize the multiple roles of polyamines in plants. *Plant Physiology and Biochemistry.* 2010;48(7):540-46. <https://doi:10.1016/j.plaphy.2010.02.009>.
99. Srivastava A, Handa AK. Hormonal regulation of tomato fruit development: A molecular perspective. *Journal of Plant Growth Regulation.* 2005;24(2):67-82. <https://doi:10.1093/jxb/eru347>
100. Masson PH, Takahashi T, Angelini R. Editorial: molecular mechanisms underlying polyamine functions in plants *Front Plant Sci.* 2017;8:14. Published online 2017 Jan 24. <https://doi:10.3389/fpls.2017.00014>.
101. Ioannidis NE, Malliarakis DM, Torné MT, Santos M, Kotzabasis K. The over-expression of the plastidial transglutaminase from maize in *Arabidopsis* increases the activation threshold of photoprotection. *Front Plant Sci.* 2016;7. <https://doi.org/10.3389/fpls.2016.00635>
102. Borja Belda P, Carla A, Esmeralda M, Juan C, Alejandro F. Relevance of the axis spermidine/elF5A for plant growth and development. *Front Plant Sci.* 2016; <https://doi.org/10.3389/fpls.2016.00245>.
103. Nambeesan S, Datsenka T, Ferruzzi MG, Malladi A, Mattoo AK, Handa AK. Overexpression of yeast spermidine synthase impacts ripening, senescence and decay symptoms in tomato. *Plant Journal.* 2010;63(5):836-47. <https://doi:10.1111/j.1365-313X.2010.04286.x>.
104. Handa AK, Fatima T, Mattoo AK. Polyamines: Bio-molecules with diverse functions in plant and human health and disease. *Frontiers in Chemistry.* 2018;6:10-28. <https://doi.org/10.3389/fchem.2018.00010>.
105. Napieraj N, Janicka M, Reda M. Interactions of polyamines and phytohormones in plant response to abiotic stress. *Plants.* 2023;12(5):1159. <https://doi.org/10.3390/plants12051159>.