



MINI REVIEW ARTICLE

Physiological and AI-based study of endophytes on medicinal plants: A mini review

Saloni Kunwar¹, Aditya Joshi², Prateek Gururani¹, Deepa Pandey³ & Neha Pandey^{1*}

- ¹Department of Biotechnology, Graphic Era Deemed to be University, Dehradun 248002, India
- ²Department of Computer Science and Engineering, Graphic Era Deemed to be University, Dehradun 248002, India
- ³Department of Zoology, Govt. P. G. College Ranikhet, 263647, India
- *Email: neha.pandey@geu.ac.in, neha.pandey221@gmail.com

3 OPEN ACCESS

ARTICLE HISTORY

Received: 31 March 2023 Accepted: 05 August 2023

Available online Version 1.0 : 31 August 2023 Version 2.0 : 22 September 2023



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care etc. See https://horizonepublishing.com/journals/ index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an openaccess article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (https://creativecommons.org/licenses/by/4.0/)

CITE THIS ARTICLE

Kunwar S, Joshi A, Gururani P, Pandey D, Pandey N. Physiological and Al-based study of endophytes on medicinal plants: A mini review. Plant Science Today. 2023; 10(sp2): 53 -60. https://doi.org/10.14719/pst.2555

Abstract

Research on the natural resources found in medicinal plants and endophytes makes important contributions to a wide variety of fields, including drug development, agribusiness, biotechnology, and sustainable development. Endophytes are a group of microorganisms that can be discovered in the rhizosphere of plants being used in medical treatment. These microorganisms have the capability of producing a wide range of primary and secondary metabolites by utilizing a variety of distinct biosynthetic pathways. Several different technologies, such as genetic modification and artificial intelligence (AI), play a significant role in the acceleration of endophytic research. These methods aid in the discovery and synthesis of novel compounds with medicinal promise, the predictive analysis of bioactive compounds, the identification and classification of endophytes, and the optimization of potential bioactive compounds. In light of this, the current review focuses on providing a concise comprehension of the influence of bioactive compounds secreted by specific endophytes on medicinal plants through the application of significant technologies in the field of endophytic research.

Keywords

Artificial Intelligence; bacteria; bioactive compounds; Endophytes; fungus; medicinal plants

Introduction

Microbes residing in plant roots tend to be beneficial for plants, showing plant-microbe interaction. Microbes acquire good nutritional values from plants and serve to produce various bioactive compounds (1). In return, they provide essential nutrients to plants and help them survive in extreme climatic conditions, as well as perform a crucial role in a sustainable environment (2).

Bioactive compounds such as quinols, phenols, steroids, alkaloids, peptides, flavonoids, polyketones, and terpenoids (3) are produced by the endophytes, including bacteria, actinobacteria, and fungi, that reside in symbiotic association with plants and are in great demand (4, 5, 6). Additionally, endophytes, which are found in the roots of plants used for medicinal purposes, are a crucial source of nutrients for plant growth and development. According to reports, three main kinds of exchanges that are advantageous exist between endophytes and host plants: (i) enhancement of the growth of the medicinal plant, (ii) enhancement of the medicinal plant's resistance to stress, and (iii) high

-value secondary metabolites (2). There is various advanced development against pathogenic species inhibiting their growth and modulating the nutrient balance in pathogens and providing benefits to existing endophytes (7). Various species of medicinal plants and endophytes produce similar groups of bioactive compounds Table 1 such as alkaloids, terpenoids, flavonoids, steroids, phenolics, etc. example from Catharanthus roseus, Alternaria sp. was isolated and found to produce vinblastine (also called vinca alkaloid), It was discovered that Fusarium oxysporum, another endophytic fungus, produces vincristine (vinca alkaloids) when it is isolated from Catharanthus roseus (1) possessing numerous advantages (3) as shown in Fig. 1. These secondary metabolites possess beneficial, multifunctional activity which promotes human health and wellness, and also used for drug development (1, 3) shown in Fig. 2.

However, research showed that genome sequencing of numerous endophytes exhibits the absence of comprehensive biosynthetic pathways for the production of phytochemicals (4). In addition, plant secondary metabolites (PSMs) themselves perform several types of roles in plant development and growth, including defence response signalling (8), innate immunity (9), and response to environmental stress (10). They also provide resistance to certain pathogens and pesticides, have significance over

plant-microbe relations, and transform the microbial communities with plants (11). Rhizobacteria has been reported to encourage plant development by improving root hair branching, promoting early nodulation, improving nodule function, increasing leaf surface area and biomass, and most importantly, increasing nutrient use efficiency (12), however, decreasing the application of chemical fertilizers and pesticides (13). Furthermore, the secondary metabolites from fungal and bacterial endophytes contribute to the enhancement of the growth of medicinal plants. Additionally, these endophytes have antibacterial, anticancer, antioxidant, and inflammatory properties as well (14) (Fig. 2). Moreover, there are certain biotechnological techniques such as CRISPR-Cas9, TALEN, and ZFN for the genetic improvement in endophytes which helps to enhance the production of targeted bioactive compounds and also helps to enhance the immune memory in the medicinal plants (15, 16, 17). The genetically modified endophytes provide the capacity to survive adverse situations and tolerate drought, which enhances the durability, yield, fertility, and health of to host plant (8) Fig. 3 depicts all phases of endophytic research on medicinal plants, from the identification of potential endophytic species to the commercialization of endophytic-derived medications.

Table 1. Enodpytes produce various bioactive compounds by using medicinal host plants.

S.No.	Bioactive Compound	Endophyte	Plant Species	References
1	Galanthamine	Paenibacillus lautus	Leucojum aestivum	(18)
2	Vinblastine	Curvularia verruculosa	Catharanthus roseus	(19)
3	Taxol (paclitaxel)	<i>Annulohypoxylon</i> sp. MUS1	Taxus wallichiana	(20)
4	Saikosaponin	Fusarium acuminatum Fusarium oxysporum	Bupleurum scorzonerifolium	(21)
5	Resveratrol	Aspergillus stellifer AB4 Quambalaria cyanescens	Vitis vinifera	(22, 26)
6	Ginsenoside Rg3	Chaetomium sp.	Panax ginseng	(24)
7	Emodin	Epicoccum nigrum	Hypericum perforatum	(25)
8	Dendrobine	Trichoderma Longibrachiatum MD33	Dendrobium nobile	(26)
9	Colchicine	Diaporthe perseae	Glorosa superb	(27)
10	Asiaticoside	Colletotrichum Gloeosporioides	Centella asiactica	(28)




Fig. 1. Various secondary metabolites present in medicinal plants.

Fig. 2. The Schematic diagram of AI with its relation to ML and DL.




Fig. 3. Influence of bioactive compounds from entophytes in medicinal plants. (Bio Render software).

To analyze and interpret the vast amounts of data produced by such studies, artificial intelligence (AI) can be extremely important. Artificial intelligence could identify and classify endophytic microbes using their genetic and biochemical profiles (29). Machine learning algorithms can detect specific endophytes and their medicinal properties from large DNA and RNA sequence datasets (30). Artificial intelligence can help facilitate the discovery of promising medicinal substances (31). To predict the activity of newly identified endophytic bioactive compounds, machine learning algorithms can be trained on large databases of chemical compounds and their properties. This can aid in the discovery of novel agents and medications (32). Endophytic microorganism growth can also be optimized with AI. Machine learning algorithms can maximize endophytic growth by analyzing environmental factors like temperature, pH, and nutrient availability (33).

Role of endophytes on medicinal plants

Fungal endophytes

Endophytic fungi are responsible for the production of growth-promoting plant hormones (4), for example; *Azospirillum* spp. enhances the growth of *Triticum aestivum L.*, and they also produce bioactive compounds which increase the stress resistance in host plants (34, 35). By secreting indole acetic acid, *Mycena dendrobii* could

Table 2. Effects of Fungal endophytes on medicinal plants.

encourage seed germination in addition to the growth of the host plant *Gastrodia elata* (36). Furthermore, it has been reported that by promoting the production of nitrate reductase and the starch-degrading enzymes, *Piriformospora indica* accelerated the growth in the roots of tobacco (37). The host plants of endophytic fungus belonging to the genus *Neotyphodium* may benefit from alkaloids produced by these organisms boosting their resistance to insect attacks (9). Table 2 shows the influence of fungal endophytes on various medicinal plants.

Bacterial endophytes

Bacterial endophytes are found in various plant components, including the seed, root, stem, leaves, and flowers (43-45). Endophytic bacteria enhance the biomass and seed germination in the host plants, they present at the different stages of plant development (46). Through a variety of methods, including the production of phytohormones, nitrogen fixation, and phosphorus solubilization, endophytes aid in the growth of plants (47, 48). Bacteria such as Streptomyces, Bacillus, Paenibacillus, and Pseudomonas (49-51) reported to promote the growth of plants, induce stress resistance, and enhance the metabolism of the medicinal plant (14). Actinomycetes support the ecology due to their environmentally favourable nature. They have been recognized to generate a range of antibiotics, biocontrol agents, and compounds that encourage plant growth (52-55). Actinomycetes enhance the soil fertility and production of the plant and also increase the nitrogen level (15). Table 3 shows the effect of bacterial endophytes on various medicinal plants.

Artificial intelligence and its subsets

Artificial intelligence (AI) refers to the mimicking of human intelligence processes by machines, particularly computer systems. It trains machines how to mimic human characteristics including learning, judging, and making decisions (61). Artificial intelligence (AI) seeks to replicate human intellectual activity by using knowledge as the object, obtaining knowledge, researching and analyzing the various ways in which knowledge is expressed, and then using these expressions (62).

S.No.	Medicinal Plant	Fungal Endophyte	Impact on Plants	References
1	Cymbidium sinese	Mycena archdicola	Increase hormones in the host plant.	(38)
2	Dendrobium nobile, D. chrysanthum	Epulorhiza sp., Mycena sp., Tulasnellales, Sebacinales,Cantharellales	Improve absorption of nutrients and promote seed germination in medicinal plants.	(39)
3	Dendrobium candidum	Mycena dendrobii	Increase hormones in the host plant.	(38)
4	Sesbaniasesban	Funneliformis mosseae, Claroideoglomus etunicatum	Secret the plant hormones	(40)
5	Cucumis sativus	Phoma glomerata, Penicillium sp.	Secret phytohormones	(41)

Table 3. Effects of bacterial endophytes on medicinal plants.

S.No.	Host Plant	Bacterial Endophytes	Impact on Host Plant	References
1	Coriandrum Sativum	Bacillus siamensis	Enhance the growth of stem and roots.	(56)
2	Teucrium polium	Bacillus subtilis	Improve the area, weight, and length of the root.	(57)
3	Panax ginseng	Micrococcus luteus	Enhanced biomass of seeds.	(58)
4	Atractylodes Macrocephala	Pseudomonas fluorescens	Increase yield of metabolites.	(59)
5	Lonicera japonica	Paenibacillus	Length of shoot, root, and fresh and dry weight increased.	(60)

The ability of a machine to learn by itself from data, improve its efficiency depending on prior experiences, and make predictions is known as machine learning, which is a subset of artificial intelligence (AI). The four basic categories of machine learning i.e., supervised, unsupervised, semi-supervised, and reinforcement learning are determined by the learning processes and methods (63). Deep learning (DL) is a member of a larger group of machine learning techniques built on artificial neural networks and representation learning (64). The Schematic diagram of AI and its relation to ML and DL is represented in Fig. 2.

Al and its functioning

The field of artificial intelligence is vast and includes numerous approaches and methodologies. Below is a general summary of how AI works and some of the important principles involved:

Data Collection: AI systems learn and make choices using massive volumes of data. The initial stage is to collect appropriate information from multiple data sources, such as images, text, audio, video, or numerical data (64).

Data Pre-processing: After gathering the data, it is frequently necessary to pre-process it, which involves cleaning, converting, and arranging it to make it appropriate for AI algorithms (64).

Machine learning: The procedure for ML modelling is to input data for training, use a machine learning algorithm for learning; and fine-tune the learning algorithm's parameters to improve accuracy. After the learning is completed, the model is now formed (63).

Neural networks: Modern AI and machine learning rely heavily on neural networks. They are inspired by the organization of the human brain and are made up of interconnected nodes or artificial neurons. Because neural networks can understand complex patterns and connections within data, they are highly effective for difficult jobs (65).

Deep learning: It emphasizes the application of multilayered deep neural networks. These deep networks may automatically learn hierarchical data representations, allowing them to solve more complex problems (66).

Inference and decision-making: Following training, the AI model can make predictions or judgments on previously unknown data. This is referred to as inference. (67). For example, an AI system trained on diseased or healthy plants may be used to categorize newly acquired images as diseased or healthy.

Natural language processing: The field of artificial intelligence known as "natural language processing" (NLP) is concerned with designing algorithms to understand and interact with human language (68). Techniques such as language translation, digit recognition, handwriting recognition, and chatbots fall under this category.

Computer vision: Another important field of AI is computer vision, which enables intelligent machines to evaluate and comprehend visual data from images or videos (69). It is

used in domains such as object detection, facial identification, autonomous vehicles, medicinal plant recognition, plant disease detection, and so on.

Role of AI in plant-based endophytic research

In endophytic research on medicinal plants, artificial intelligence (AI) has assumed an increasingly significant role. Here, are some examples of AI applications in this field shown in Fig. 4.




Fig. 4. Al assists in Endophytic research phases.

Prediction of bioactive compounds

Artificial intelligence can help to predict the bioactive compounds developed via endophytic microorganisms in medicinal plants. Predicting the production of bioactive metabolites by endophytic fungi in the medicinal plant *Panax ginseng* using machine learning algorithms is studied (70). Transcriptomics and artificial intelligence techniques for bioactive compound prediction and selection are also studied (71). On the extraction of bioactive chemicals from unripe *M. acuminata* peel using microwave-assisted extraction (MAE), the association between response surface methodology (RSM) and an artificial neural network prediction model was examined (72).

Discovery of new endophytes and natural products

Artificial Intelligence can aid in the discovery of novel endophytes and natural products with potential medical applications. Al is utilized to analyze the genome of the endophytic fungus *Aspergillus terreus* isolated from the medicinal plant *Huperzia serrata* to identify new natural products with anti-inflammatory and anti-tumor properties (73). In a review, emerging methods for discovering natural products from plant microbiomes were examined, with a focus on the potential of deep learning as a tool for bioprospecting, biochemical novelty prediction, and regulatory control of endophytes (74). For in-depth data analysis and forecasting in the search for anticancer leads, machine learning has provided potent

approaches to identify anticancer lead compounds with an emphasis on natural products from plants and their related microbes (75). Bioprospecting different species and running laboratory screening techniques are traditionally employed to identify natural compounds, producing complex data. This strategy, meanwhile, frequently confronts difficulties and high rates of loss. To solve these issues, machine learning (ML) and artificial intelligence (AI) are being used. The area of natural product-based drug discovery algorithms has undergone a revolution as a result of recent advances in AI, notably in ML. AI tools have shown success in classifying things, finding hidden patterns, and grouping compounds according to their properties (76, 77).

Optimization of endophyte-based production

Artificial Intelligence can improve medicinal plant endophytic microorganism bioactive compound production. Artificial Intelligence optimized bioactive compound synthesis by the endophytic fungus Fusarium solani isolated from the medicinal plant Lithospermum erythrorhizon (78). Response surface methodology (RSM) and an artificial neural network-genetic algorithm (ANN-GA) were investigated using a single-factor analysis of the efficacy of culture situations upon the quantity of cellulase produced by endophytic fungi of T. cuspidata to obtain factors for improving the enzyme activity process of production (79). For optimizing and predicting the 6gingerol concentration in Zingiber officinale samples from 60 distinct locations around the state with varying agricultural and climatic circumstances, two deep learning algorithms of the Artificial Neural Network (ANN) model were used (80).

Identification of endophyte-host interactions

Artificial intelligence can be used to identify the associations between endophytic microorganisms and their host plants. It is used to examine the transcriptomes of the host plant *Camellia sinensis* and the endophytic fungus *Colletotrichum tofieldiae* and discover genes involved in their interaction (81).

Using genetic, morphological, and biochemical traits, artificial intelligence (AI) can analyze large datasets of plant samples to pinpoint possible endophytic species (73). It can assist in standardizing the process of isolation and maintaining culture conditions for endophytes to ensure their viability and growth (82). It is capable of assessing the metabolic profiles of endophytes to find bioactive substances with potential medicinal applications (75). Artificial Intelligence can aid in the analysis of genetic and biochemical data generated by endophytic research to identify novel drug development targets (83). Artificial Intelligence can be used in the design and optimization of drugs based on endophytic compounds, resulting in the creation of novel medicines (84). In addition, Al also analyses clinical trials for endophyte-derived pharmaceuticals, thereby enhancing their efficacy and safety. Apart from this, AI can assist in the commercialization of endophytic research by identifying potential markets, developing marketing strategies, and optimizing supply chains (85).

Conclusion

Endophytes existing as eco-friendly bioagents help to achieve a sustainable environment in the future. Bacterial and fungal endophytes aid in the greater development of medicinal plants by increasing the rate of plant hormones. Correspondingly, they increase medicinal properties by improving the seed germination stage and nutrient content. Besides, focus can be given to various techniques regeneration, genetic modification, electroporation which give a high yield of targeted bioactive compounds which gives numerous benefits to medicinal plants. Furthermore, the advancement of technologies such as AI has been incorporated to revolutionize the field of endophytic research in medicinal plants. The capability of AI to analyze large datasets of genomic and metabolomic data can help identify suitable endophytes with pharmaceutical potential and aid in the development of new drugs.

Indeed, the research of endophytes on medicinal plants has shown AI to be of enormous value to human society. Overall, the study of endophytes on medicinal plants in conjunction with artificial intelligence has enormous potential to advance drug discovery, develop more potent therapies, and advance human health. Researchers can speed up the discovery and development of novel medicinal compounds from these natural sources by utilizing AI.

Acknowledgements

The authors are grateful to IERP-GBPNIHE and Graphic Era (Deemed to be University) for providing general support.

Authors' contributions

Saloni Kunwar and Aditya Joshi: Writing-original draft and Data curation. Prateek Gururani and Deepa Pandey: Conceptualization and Investigation. Neha Pandey: Supervision, Resources. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None.

References

- Jamwal VL, Gandhi SG. Endophytes as a source of High-value phytochemicals: Present scenario and future outlook. Endophytes Second. Metabolites Reference Ser. Phytochemistry. 2019; 571-90. https://doi.org/10.1007/978-3-319-90484-9_14
- Akula R, Ravishankar GA. Influence of abiotic stress signals on secondary metabolites in plants. Plant Signal. Behav. 2011;6(11):1720-31. https://doi.org/10.4161/psb.6.11.17613
- 3. Singh M, Kumar A, Singh R, Pandey KD. Endophytic bacteria: a new source of bioactive compounds. 3 Biotech. 2017; 7:1-4. https://doi.org/10.1007/s13205-017-0942-z

- 4. Li J, Zhao GZ, Chen HH, Wang HB, Qin S, Zhu WY, Xu LH, Jiang CL, Li WJ. Antitumour and antimicrobial activities of endophytic streptomycetes from pharmaceutical plants in rainforest. Lett Appl Microbio. 2008; 47(6):574-80. https://doi.org/10.1111/j.1472-765X.2008.02470.x
- Kusari S, Verma VC, Lamshoeft M, Spiteller M. An endophytic fungus from Azadirachtaindica A. Juss. that produces azadirachtin. World J Microbiol Biotechnol. 2012; 28(3):1287-94. https://doi.org/10.1007/s11274-011-0876-2
- Molina G, Pimentel MR, Bertucci TC, Pastore GM. Application of fungal endophytes in biotechnological processes. Chem Eng Trans. 2012;27:1-6. https://doi.org/10.3303cet/1227049
- Pang Z, Chen J, Wang T, Gao C, Li Z, Guo L, Xu J, Cheng Y. Linking plant secondary metabolites and plant microbiomes: a review. Front Plant Sci. 2021;12:621276. https://doi.org/10.3389/fpls.2021.621276b
- Isah T. Stress and defense responses in plant secondary metabolites production. Biol Res. 2019;52. http:// dx.doi.org/10.1186/s40659-019-0246-3
- Piasecka A, Jedrzejczak-Rey N, Bednarek P. Secondary metabolites in plant innate immunity: conserved function of divergent chemicals. New Phytolo. 2015; 206(3):948-64. https://doi.org/10.1111/nph.13325
- Yang L, Wen KS, Ruan X, Zhao YX, Wei F, Wang Q. Response of plant secondary metabolites to environmental factors. Molecules. 2018; 23(4):762. https://doi.org/10.3390/ molecules23040762
- Guerrieri A, Dong L, Bouwmeester HJ. Role and exploitation of underground chemical signaling in plants. Pest management science. 2019;75(9):2455-63. https:// doi.org/10.1002/ps.5507
- Vocciante M, Grifoni M, Fusini D, Petruzzelli G, Franchi E. The role of plant growth-promoting rhizobacteria (PGPR) in mitigating plant's environmental stresses. Appl Sci. 2022;12 (3):1231. https://doi.org/10.3390/app12031231
- Liu K, Newman M, McInroy JA, Hu CH, Kloepper JW. Selection and assessment of plant growth-promoting rhizobacteria for biological control of multiple plant diseases. Phytopathology. 2017;107(8):928-36. https://doi.org/10.1094/ phyto-02-17-0051-r.
- 14. Divekar PA, Narayana S, Divekar BA, Kumar R, Gadratagi BG, Ray A, Singh AK, Rani V, Singh V, Singh AK, Kumar A. Plant secondary metabolites as defense tools against herbivores for sustainable crop protection. Int J Mol Sci. 2022; 23 (5):2690. https://doi.org/10.3390/ijms23052690
- Chen K, Gao C. TALENs: customizable molecular DNA scissors for genome engineering of plants. J Genet Genom. 2013; 40 (6):271-79. https://doi.org/10.1016/j.jgg.2013.03.009
- Li D, Tang Y, Lin J, Cai W. Methods for genetic transformation of filamentous fungi. Microbial Cell Factories. 2017;16:1-3. https://doi.org/10.1186/s12934-017-0785-7
- 17. El-Sayed AS, Abdel-Ghany SE, Ali GS. Genome editing approaches: manipulating of lovastatin and taxol synthesis of filamentous fungi by CRISPR/Cas9 system. Appl Microbiol Biotechnol. 2017;101:3953-76. https://doi.org/10.1007/s00253-017-8263-z
- Gupta S, Bhatt P, Chaturvedi P. Determination and quantification of asiaticoside in endophytic fungus from Centellaasiatica (L.) Urban. World J Microbiol Biotechnol. 2018; 34:1-10. https://doi.org/10.1007/s11274-018-2493-9
- Deepika VB, Vohra M, Mishra S, Dorai K, Rai P, Satyamoorthy K, Murali TS. DNA demethylation overcomes attenuation of colchicine biosynthesis in an endophytic fungus Diaporthe. J Biotechnol. 2020; 323:33-41. https://doi.org/10.1016/j.jbiotec.2020.07.019.

- Sarsaiya S, Jain A, Fan X, Jia Q, Xu Q, Shu F, Zhou Q, Shi J, Chen J. New insights into detection of a dendrobine compound from a novel endophytic Trichodermalongibrachiatum strain and its toxicity against phytopathogenic bacteria. Front Microbiol. 2020; 11:337. https://doi.org/10.3389/fmicb.2020.00337
- Vigneshwari A, Rakk D, Németh A, Kocsubé S, Kiss N, Csupor D, Papp T, Škrbić B, Vágvölgyi C, Szekeres A. Host metabolite producing endophytic fungi isolated from Hypericum perforatum. PLoS One. 2019;14(5):e0217060. https://doi.org/10.1371/journal.pone.0217060
- 22. Xu XD, Liang WX, Yao L, Paek KY, Wang J, Gao WY. Production of ginsenoside by Chaetomium sp. and its effect on enhancing the contents of ginsenosides in *Panax ginseng* adventitious roots. Biochem Engin J. 2021;174:108100. https://doi.org/10.1016/j.bej.2021.108100
- Roat C, Saraf M. Isolation and characterization of t-resveratrol and α-viniferin, a bioactive secondary metabolite of an endophytic fungus Aspergillus stellifer ab4, from Vitisvinifera. The J Microbiol BiotechnolFood Sci. 2020;19 (4):708. https://doi.org/10.15414/jmbfs.2020.9.4.708-713
- 24. Meshram V, Kapoor N, Dwibedi V, Srivastava A, Saxena S. Extracellular resveratrol producing endophytic fungus, *Quambalaria cyanescens*. S Afr J Bot. 2022;146:409-16. https://doi.org/10.1016/j.sajb.2021.11.026
- Cheng Y, Liu G, Li Z, Zhou Y, Gao N. Screening saikosaponin d (SSd)-producing endophytic fungi from *Bupleurum* scorzonerifolium Willd. World J Microbiol Biotechnol. 2022; 38 (12):242. https://doi.org/10.1007/s11274-022-03434-x
- 26. Gauchan DP, Vélëz H, Acharya A, Östman JR, Lundén K, Elfstrand M, García-Gil MR. Annulohypoxylon sp. strain MUS1, an endophytic fungus isolated from Taxus wallichiana Zucc., produces taxol and other bioactive metabolites. 3 Biotech. 2021;11:1-6. https://doi.org/10.1007/s13205-021-02693-z
- Parthasarathy R, Shanmuganathan R, Pugazhendhi A. Vinblastine production by the endophytic fungus *Curvularia verruculosa* from the leaves of *Catharanthus roseus* and its in vitro cytotoxicity against HeLa cell line. Analy Biochem. 2020;593:113530. https://doi.org/10.1016/j.ab.2019.113530
- Ptak A, Morańska E, Warchoł M, Gurgul A, Skrzypek E, Dziurka M, Laurain-Mattar D, Spina R, Jaglarz A, Simlat M. Endophytic bacteria from in vitro culture of Leucojumaestivum L. a new source of galanthamine and elicitor of alkaloid biosynthesis. Scienti Rep. 2022;12(1):13700. https://doi.org/10.1038/s41598-022-17992-5
- Sagita R, Quax WJ, Haslinger K. Current state and future directions of genetics and genomics of endophytic fungi for bioprospecting efforts. Front Bioengin Biotechno. 2021; 9:649906. https://doi.org/10.3389/fbioe.2021.649906
- Busato S, Gordon M, Chaudhari M, Jensen I, Akyol T, Andersen S, Williams C. Compositionality, sparsity, spurious heterogeneity, and other data-driven challenges for machine learning algorithms within plant microbiome studies. Curr Opin Plant Biol. 2023; 71:102326. https://doi.org/10.1016/ j.pbi.2022.102326
- 31. Mak KK, Pichika MR. Artificial intelligence in drug development: present status and future prospects. Drug Discovery Today. 2019; 24(3):773-80. https://doi.org/10.1016/j.drudis.2018.11.014
- 32. Helmy M, Elhalis H, Yan L, Chow Y, Selvarajoo K. Perspective: multi-omics and machine learning help unleash the alternative food potential of microalgae. Advances in Nutrition. 2022. https://doi.org/10.1016/j.advnut.2022.11.002
- 33. Kudjordjie EN, Sapkota R, Steffensen SK, Fomsgaard IS, Nicolaisen M. Maize synthesized benzoxazinoids affect the host-associated microbiome. Microbiome. 2019; 7(1):1-7.

- https://doi.org/10.1186/s40168-019-0677-7
- 34. Maggini V, Mengoni A, Bogani P, Firenzuoli F, Fani R. Promoting model systems of microbiota-medicinal plant interactions. Trends Plant Sci. 2020; 25(3):223-25. https://doi.org/10.1016/j.tplants.2019.12.013
- Shunxing G, Qiuying W. Character and action of good strain on stimulating seed germination of Gastrodiaelata. Mycosystema. 2001;20(3):408-12.
- 36. Sherameti I, Shahollari B, Venus Y, Altschmied L, Varma A, Oelmuller R. The endophytic fungus Piriformosporaindica stimulates the expression of nitrate reductase and the starch -degrading enzyme glucan-water dikinase in tobacco and Arabidopsis roots through a homeodomain transcription factor that binds to a conserved motif in their promoters. J Biolo Chem. 2005; 280(28):26241-47. https://doi.org/10.1074/jbc.M500447200
- 37. Qin S, Li J, Chen HH, Zhao GZ, Zhu WY, Jiang CL, Xu LH, Li WJ. Isolation, diversity, and antimicrobial activity of rare actinobacteria from medicinal plants of tropical rain forests in Xishuangbanna, China. Appl Environ Microbio. 2009; 75 (19):6176-86. https://doi.org/10.1128/AEM.01034-09
- 38. Zhang J, Wang C, Guo S, Chen J, Xiao P. Studies on the plant hormones produced by 5 species of endophytic fungi isolated from medicinal plants (Orchidacea). Acta Academiae Medicinae Sinicae. 1999;21(6):460-65.
- Chen XM, Guo SX. Effects of four species of endophytic fungi on the growth and polysaccharide and alkaloid contents of *Dendrobium nobile*. China J Chinese Materia Med. 2005;30 (4):253-57.
- Abd_Allah EF, Hashem A, Alqarawi AA, Bahkali AH, Alwhibi MS. Enhancing growth performance and systemic acquired resistance of medicinal plant Sesbaniasesban (L.) Merr using arbuscular mycorrhizal fungi under salt stress. Saudi J of Biol Sci. 2015; 22(3):274-83. https://doi.org/10.1016/j.sjbs.2015.03.004
- 41. Waqas M, Khan AL, Kamran M, Hamayun M, Kang SM, Kim YH, Lee IJ. Endophytic fungi produce gibberellins and indoleacetic acid and promotes host-plant growth during stress. Molecules. 2012;17(9):10754-73. https://doi.org/10.3390/molecules170910754
- 42. Compant S, Mitter B, Colli-Mull JG, Gangl H, Sessitsch A. Endophytes of grapevine flowers, berries, and seeds: identification of cultivable bacteria, comparison with other plant parts, and visualization of niches of colonization. Microbial Ecol. 2011;62:188-97. https://doi.org/10.1007/s00248-011-9883-y
- 43. Elmagzob AA, Ibrahim MM, Zhang GF. Seasonal diversity of endophytic bacteria associated with Cinnamomumcamphora (L.) Presl. Diversity. 2019;11(7):112. https://doi.org/10.3390/d11070112
- 44. Mocali S, Bertelli E, Di Cello F, Mengoni A, Sfalanga A, Viliani F, Caciotti A, Tegli S, Surico G, Fani R. Fluctuation of bacteria isolated from elm tissues during different seasons and from different plant organs. Res in Microbiology. 2003;154(2):105-14. https://doi.org/10.1016/S0923-2508(03)00031-7
- 45. Qin S, Miao Q, Feng WW, Wang Y, Zhu X, Xing K, Jiang JH. Biodiversity and plant growth promoting traits of culturable endophytic actinobacteria associated with Jatropha curcas L. growing in Panxi dry-hot valley soil. Appl Soil Ecol. 2015;93:47-55. https://doi.org/10.1016/j.apsoil.2015.04.004
- Borah A, Das R, Mazumdar R, Thakur D. Culturable endophytic bacteria of Camellia species endowed with plant growth promoting characteristics. J Appl Microbiol. 2019;127 (3):825-44. https://doi.org/10.1111/jam.14356
- 47. Gao Y, Liu Q, Zang P, Li X, Ji Q, He Z, Zhao Y, Yang H, Zhao X,

- Zhang L. An endophytic bacterium isolated from Panax ginseng CA Meyer enhances growth, reduces morbidity, and stimulates ginsenoside biosynthesis. Phytochem Lett. 2015;11:132-38. https://doi.org/10.1016/j.phytol.2014.12.007
- 48. Zhao L, Xu Y, Lai XH, Shan C, Deng Z, Ji Y. Screening and characterization of endophytic Bacillus and Paenibacillus strains from medicinal plant Lonicera japonica for use as potential plant growth promoters. Braz J Microbiol. 2015;46:977-89. https://doi.org/10.1590/S1517-838246420140024
- 49. Qi Y, Li X, Wang J, Wang C, Zhao S. Efficacy of plant growth-promoting bacteria *Streptomyces werraensis* F₃ for chemical modifications of diseased soil of ginseng. Biocontrol Science and Technology. 2021; 31(2):219-33. https://doi.org/10.1080/09583157.2020.1843598
- Bhatti AA, Haq S, Bhat RA. Actinomycetes benefaction role in soil and plant health. Microbial Pathogenesis. 2017;111:458-67. https://doi.org/10.1016/j.micpath.2017.09.036
- 51. Abdallah ME, Haroun SA, Gomah AA, El-Naggar NE, Badr HH. Application of actinomycetes as biocontrol agents in the management of onion bacterial rot diseases. Arch Phytopathol Pflanzenschutz 2013;46(15):1797-808. https://doi.org/10.1080/03235408.2013.778451
- Behie SW, Bonet B, Zacharia VM, McClung DJ, Traxler MF. Molecules to ecosystems: Actinomycete natural products in situ. Front in Microbiol. 2017;7:2149. https://doi.org/10.3389/ fmicb.2016.02149
- Abd-Alla MH, El-Sayed ES, Rasmey AH. Indole-3-acetic acid (IAA) production by Streptomyces atrovirens isolated from rhizospheric soil in Egypt. J Biol Earth Sci. 2013; 3(2):182-93.
- 54. Zhang C, Lu Y. Study on artificial intelligence: The state of the art and future prospects. Journal of Industrial Information Integration. 2021;23:100224. https://doi.org/10.1016/j.jii.2021.100224
- 55. Duan L, Da Xu L. Business intelligence for enterprise systems: a survey. IEEE Transactions on Industrial Informatics. 2012;8 (3):679-87. https://doi.org/10.1109/tii.2012.2188804
- 56. Sarker IH. Machine learning: Algorithms, real-world applications and research directions. SN Computer Sci. 2021;2(3):160. https://doi.org/10.1007/s42979-021-00592-x
- 57. Ibrahim E, Fouad H, Zhang M, Zhang Y, Qiu W, Yan C, Li B, Mo J, Chen J. Biosynthesis of silver nanoparticles using endophytic bacteria and their role in inhibition of rice pathogenic bacteria and plant growth promotion. RSC Advances. 2019;9(50):29293-99. https://doi.org/10.1039/c9ra04246f
- 58. Hassan SE. Plant growth-promoting activities for bacterial and fungal endophytes isolated from medicinal plant of *Teucrium polium* L. J Advanc Res. 2017;8(6):687-95. https://doi.org/10.1016/j.jare.2017.09.001
- 59. Vendan RT, Yu YJ, Lee SH, Rhee YH. Diversity of endophytic bacteria in ginseng and their potential for plant growth promotion. J Microbiol. 2010;48:559-65. https://doi.org/10.1007/s12275-010-0082-1
- Yang HR, Yuan J, Liu LH, Zhang W, Chen F, Dai CC. Endophytic Pseudomonas fluorescens induced sesquiterpenoid accumulation mediated by gibberellic acid and jasmonic acid in Atractylodes macrocephala Koidz plantlets. Plant Cell, Tissue and Organ Culture. 2019; 138:445-57. https://doi.org/10.1007/s11240-019-01640-4
- 61. Zhao L, Xu Y, Lai XH, Shan C, Deng Z, Ji Y. Screening and characterization of endophytic *Bacillus* and *Paenibacillus* strains from medicinal plant *Lonicera japonica* for use as potential plant growth promoters. Brazilian Journal of Microbiology. 2015;46:977-89.

- Fujioka T, Mori M, Kubota K, Oyama J, Yamaga E, Yashima Y, Katsuta L, Nomura K, Nara M, Oda G, Nakagawa T. The utility of deep learning in breast ultrasonic imaging: a review. Diagnostics. 2020;10(12):1055. https://doi.org/10.3390/diagnostics10121055
- 63. Chen B. Wineinformatics: A New Data Science Application. Springer Nature; 2022 1-70p.
- 64. Kujawa S, Niedbała G. Artificial neural networks in agriculture. Agriculture. 2021;11(6):497. https://doi.org/10.3390/agriculture11060497
- Gupta A, Anpalagan A, Guan L, Khwaja AS. Deep learning for object detection and scene perception in self-driving cars: Survey, challenges, and open issues. Array. 2021;10:100057. https://doi.org/10.1016/j.array.2021.100057
- 66. Agrawal A, Gans JS, Goldfarb A. Exploring the impact of artificial intelligence: Prediction versus judgment. Inform Econo Pol. 2019;47:1-6. https://doi.org/10.1016/j.infoecopol.2019.05.001
- 67. Baclic O, Tunis M, Young K, Doan C, Swerdfeger H, Schonfeld J. Artificial intelligence in public health: Challenges and opportunities for public health made possible by advances in natural language processing. Can Communi Dis Rep. 2020;46 (6):161. https://doi.org/10.14745%2Fccdr.v46i06a02
- Bhatt D, Patel C, Talsania H, Patel J, Vaghela R, Pandya S, Modi K, Ghayvat H. CNN variants for computer vision: History, architecture, application, challenges, and future scope. Electronics. 2021;10(20):2470. https://doi.org/10.3390/ electronics10202470
- Li Z, Chen Y, Chen H, Zhang X, Liu Y. Predicting the biosynthetic potential of endophytic fungi isolated from Panax ginseng with artificial intelligence. BMC Bioinformatics. 2021;2(1):1-14.
- Zameer R, Tariq S, Noreen S, Sadaqat M, Azeem F. Role of Transcriptomics and Artificial Intelligence Approaches for the Selection of Bioactive Compounds. Drug Design Using Machine Learning. 2022; 10:283-317. https:// doi.org/10.1002/9781394167258.ch10
- 71. Said FM, Gan JY, Sulaiman J. Correlation between response surface methodology and artificial neural network in the prediction of bioactive compounds of unripe *Musa acuminata* peel. Engineering Sci Technol. 2020; 23(4):781-77. https://doi.org/10.1016/j.jestch.2019.12.005
- Tiwari P, Bae H. Endophytic fungi: key insights, emerging prospects, and challenges in natural product drug discovery. Microorganisms. 2022; 10(2), p.360. https:// doi.org/10.3390/microorganisms10020360
- Aghdam SA, Brown AM. Deep learning approaches for natural product discovery from plant endophytic microbiomes.
 Environ Microbiome. 2021; 16(1):1-20. https://doi.org/10.1186/s40793-021-00375-0
- Li G, Lin P, Wang K, Gu CC, Kusari S. Artificial intelligence-guided discovery of anticancer lead compounds from plants and associated microorganisms. Trends in Cancer. 2022; 8 (1):65-80. https://doi.org/10.1016/j.trecan.2021.10.002

- 75. Yang X, Wang Y, Byrne R, Schneider G, Yang S. Concepts of artificial intelligence for computer-assisted drug discovery. Chem Rev. 2019; 119(18):10520-94. https://doi.org/10.1021/acs.chemrev.8b00728
- 76. Gupta A, Meshram V, Gupta M, Goyal S, Qureshi KA, Jaremko M, Shukla KK. Fungal Endophytes: Microfactories of Novel Bioactive Compounds with Therapeutic Interventions; A Comprehensive Review on the Biotechnological Developments in the Field of Fungal Endophytic Biology over the Last Decade. Biomolecules. 2023; 13(7):1038. https://doi.org/10.3390/biom13071038
- 77. Chen Y, Chen H, Zhang X, Liu Y. Optimization of bioactive compounds production by endophytic fungus *Fusarium solani* using artificial intelligence. J Biotechno. 2020; 310:57-65
- 78. Li H, Dou M, Wang X, Guo N, Kou P, Jiao J, Fu Y. Optimization of cellulase production by a novel endophytic fungus *Penicillium oxalicum* R4 isolated from *Taxus cuspidata*. Sustainability. 2021;13(11):6006. https://doi.org/10.3390/su13116006
- 79. Sahoo M, Dey S, Sahoo S, Das A, Ray A, Nayak S, Subudhi E. MLP (multi-layer perceptron) and RBF (radial basis function) neural network approach for estimating and optimizing 6-gingerol content in *Zingiber officinale* Rosc. in different agroclimatic conditions. Ind Crop Pro. 2023; 198:116658. https://doi.org/10.1016/j.indcrop.2023.116658
- 80. Zhang C, Wang Y, Zhang X, Huang H, He X. Identification of genes involved in the interaction between *Colletotrichum tofieldiae* and *Camellia sinensis* using machine learning algorithms. Genomics. 2020;112(6):4197-04.
- 81. YoosefzadehNajafabadi M, Hesami M, Eskandari M. Machine learning-assisted approaches in modernized plant breeding programs. Genes. 2023;14(4):777. https://doi.org/10.3390/genes14040777
- 82. Paul D, Sanap G, Shenoy S, Kalyane D, Kalia K, Tekade RK.

 Artificial intelligence in drug discovery and development.

 Drug Discovery Today. 2021;26(1):80. https://doi.org/10.1016%2Fj.drudis.2020.10.010
- 83. Saldívar-González FI, Aldas-Bulos VD, Medina-Franco JL, Plisson F. Natural product drug discovery in the artificial intelligence era. Chem Sci. 2022;13(6):1526-46. https://doi.org/10.1039/d1sc04471k
- 84. Askin S, Burkhalter D, Calado G, El Dakrouni S. Artificial Intelligence Applied to clinical trials: opportunities and challenges. Health and Techno. 2023;13(2):203-13. https://doi.org/10.1007/s12553-023-00738-2
- 85. Shankar A, Sharma KK. Fungal secondary metabolites in food and pharmaceuticals in the era of multi-omics. Appl Microbiol Biotechno. 2022;106(9-10):3465-88. https://doi.org/10.1007/s00253-022-11945-8