



REVIEW ARTICLE

The potential of bacterial endophytes on orchids

Aneesa Ansiya¹, Rajeevan Parvathi², Appukkuttannair Gangaprasad³ & Sugathan Shiburaj^{1*}

¹Department of Botany, University of Kerala, Karyavattom, Thiruvananthapuram, Kerala, India, 695 581

²Department of Biotechnology, University of Kerala, Karyavattom, Thiruvananthapuram, Kerala, India, 695 581

³Centre for Biodiversity Conservation, Department of Botany, University of Kerala, Karyavattom, Thiruvananthapuram, Kerala, India, 695 581

*Email: drshiburaj@keralauniversity.ac.in



ARTICLE HISTORY

Received: 02 April 2023

Accepted: 13 December 2023

Available online

Version 1.0 : 28 July 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Ansiya A, Parvathi R, Gangaprasad A, Shiburaj S. The potential of bacterial endophytes on orchids. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.2573>

Abstract

Orchids are one of the most exquisite and diverse plant species in nature. The seeds of orchids are non-endospermic and, therefore, dependent on endophytes for germination, growth, and adaptability. Orchids are prized for their beauty and therapeutic and culinary qualities by naturalists and the general public. Many orchid species are now endangered or vulnerable due to collectors' eagerness to take them. Current studies on orchids have concentrated on isolating and identifying the mycorrhizal and non-mycorrhizal endophytes that either directly or indirectly help orchids to grow, develop, and produce beneficial secondary metabolites. In orchids, bacterial endophytes play an essential role in the formation of mycorrhizae and the stability of relationships between plants and fungi. Endophytic bacteria can grow in orchids by producing phytohormones, doing photosynthesis, fixing nitrogen, promoting the mineral nutrition cycle, forming siderophores, and producing diverse beneficial metabolites to improve biomass production, stress tolerance, and biocontrol of potential phytopathogenic fungal species. This article examines how endophytic bacteria are associated with orchids and their potential growth-promoting abilities.

Keywords

Orchids; Orchid endophytes; Bacterial endophytes; PGPR

Introduction

The orchid family represents the second-largest angiosperm family, with over 900 genera and 27,000 species worldwide (1). They have evolved in various captivating ways to survive extreme weather conditions, acquire nutrients, and attract pollinators (2). Due to their decorative, culinary, and therapeutic uses, orchids are widely grown in domestic and foreign cities. In particular, the orchid plant's roots, foliage, and pseudo-bulbs can treat conditions including rheumatism, respiratory illnesses, nausea, piles, inflammation, viral infections, and even cancer (3).

Orchids produce many highly reduced, microscopic seeds per capsule, but only 0.2–0.3% germinate in nature, while thousands die (4). A symbiotic relationship between orchid seeds and mycorrhizae is required for orchid propagation (5). As mycorrhizae provide most minerals, nutrients, vitamins, and water for orchid germination and seedling development, they differ in how they relate to their orchid hosts (6). *In vitro*, symbiotic germination has been used for orchid propagation. Sustainability is crucial in preventing declines in orchid biodiversity, scenic attractiveness, and therapeutic benefits. In orchids, microbial communities provide nutrients

and enhance immunity through symbiotic relationships. In nature, most plants work in symbiotic harmony with other species; some depend on bacterial or fungal interactions for survival. Still, successful propagation requires extensive knowledge of orchid-mycorrhizal interaction and the ability to isolate and grow the fungus, which can be time-consuming (7, 8, 9).

Orchids have not been extensively explored for their endophytic bacteria, even though some commercially valuable epiphytic orchids, such as *Vanilla* spp. (10), *Dendrobium* spp. (11), and *Cymbidium* spp. (12), contain endophytes. These microorganisms can promote plant growth, particularly in acclimating micro-propagated seedlings (13). Endophytic bacteria may promote plant growth by producing indole acetic acid (14) and solubilizing inorganic phosphate (15).

This review article explores the potentiality of bacterial endophytes on orchids and their importance, roles, and implications in orchid ecology, growth, and conservation. It also discusses the limited research in the field of orchid-associated bacterial endophytes and points out the need for further investigations in this area. The review also explores the potential of bacterial endophytes to benefit orchids in various ways, such as promoting plant growth, enhancing stress tolerance, and possibly contributing to the discovery of bioactive compounds with pharmaceutical applications.

Orchid endophytes

Endophytic bacteria have been discovered in various environments, including tropical, temperate, aquatic, xerophytic, deserts, Antarctica, geothermal soils, rainforests, mangrove swamps, and coastal forests (16, 17). Microbial endophytes have also been extensively examined in orchids (10,11,12). Orchids and mycorrhizal fungi interact much, and the distribution of these fungi and orchids is interrelated (18). Mycorrhizal associations are essential for orchid survival. Most microbiological studies have focused on them while disregarding the possible contributions of other microbial partners (19,20). There are also endophytic bacteria that can form symbiotic relationships (21). Some bacteria that inhabit orchids include *Streptomyces* spp. (22), *Bacillus* spp. (23), *Erwinia* spp., *Pseudomonas* spp. (24), *Flavobacterium* spp. (25), *Sphingomonas paucimobilis* ZJSH1, and *Streptosporangium oxazolinicum* (24). Only a few studies on the interactions between bacteria and orchids have shown the potential of bacteria as a biological tool for conserving orchids and generating bioactive compounds (24, 26). However, some bacterial and fungal species were thought to be the best sources for researching the interactions between orchids and microorganisms and how they affect the evolution of orchid species (27).

Bacterial endophytes

Few studies focus on orchid bacterial endophytes. In orchid roots and leaves, bacterial endophytes have been isolated after surface sterilization; some isolated bacteria increased seed germination (28, 29). Root associated bacteria (RAB) interact with plants, colonize root

compartments, and control plant host function (30). These are recruited based on host genotype, metabolic profiles, root exudates, and soil physicochemical properties (30). They facilitate plant growth by acquiring nutrients, supplying hormones, protecting against infections, and reducing abiotic stressors (Fig. 1) (31). Duponnois and Garbaye published insight into mycorrhizal and bacterial associations in orchid root tissues (32). They demonstrated that *Pseudomonas fluorescens* BBC6 aids in the formation of mycorrhizae and gave the name "mycorrhiza helper bacteria" (MHB). MHB releases active substances or biomolecules in stress response, which promotes mycelia development, root colonization, and root growth (33). Endophytic rhizobacteria, including *Paenibacillus lentimorbus* and *P. macerans*, were isolated from the meristems of *Cymbidium eburneum* (26). Strains of *Sphingomonas* sp., *Agrococcus* sp., and *Mycobacterium* sp. promoted orchid germination and were isolated from *Dendrobium moschatum* (24). The study by Yu *et al.* amplified endophytic bacteria from the roots of *D. officinale* belonging to the genera *Burkholderia*, *Rhodanobacter*, *Pseudomonas*, and *Sphingomonas* (34). The major endophytic bacteria of *Cymbidium* orchid are *Bacillus thuringiensis*, *Burkholderia cepacia*, *B. gladioli*, *Herbaspirillum frisingense*, *Pseudomonas stutzeri*, *Rhizobium cellulosilyticum*, *R. radiobacter*, and *Stenotrophomonas maltophilia* (12). Endophytic bacteria from *Spiranthes spiralis*, *Serapias vomeracea*, and *Neottia ovata* were identified using the 16S rRNA gene. These include *Pseudomonas*, *Pantoea*, *Rahnella*, *Staphylococcus*, *Sphingomonas*, *Microbacterium*, *Streptomyces*, *Fictibacillus*, and *Bacillus* (35). The endophytes like *Paenibacillus taichungensis*, *P. pabuli*, *Enterobacter* sp., *Rhizobium* sp, and *Pseudomonas* sp. were isolated from *Cattleya walkeriana* (37).

Importance of endophytic bacteria

In orchids, endophytic bacteria can colonize symbiotically without harming their hosts. In addition to exchanging metabolites and sharing physiological processes with the host plant, these microorganisms can reside within plant




Fig. 1. The beneficial role of bacterial endophytes on orchid

tissues, including roots and leaves (38). In orchid roots, bacteria interact with fungi and orchid root cells to maintain symbiosis by producing plant growth-promoting metabolites and indirectly by inhibiting fungal growth that could harm the plant (31). There are several plant growth-promoting rhizobacteria (PGPR), such as *Acinetobacter*, *Arthrobacter*, *Azospirillum*, *Azotobacter*, *Bacillus*, *Burkholderia*, *Clostridium*, *Enterobacter*, *Erwinia*, *Flavobacterium*, *Klebsiella*, *Pseudomonas*, *Serratia* and *Rhizobium* (38). These bacteria may be able to stimulate plant host growth through the production of phytohormones and through the production of ACC deaminase, which effectively lowers plant ethylene concentrations (39) (Figure 2). By reducing the inhibitory effects of various phytopathogens, PGPR facilitates plant growth and adaptive capacities through nitrogen fixation (40), assistance in acquiring phosphorus (41) and essential minerals (42), improved water uptake, or acting as a biocontrol agent (Fig.2)(43). Endophytic bacteria are less investigated for their metabolic potential (44, 45) (Table 1). Endophytes synthesize bioactive compounds such as paclitaxel from *Taxomyces andreanae*, endophytic fungi isolated from pacific yew (*Taxus brevifolia*) (46) help the host plant to develop resistance against pathogens that are used in pharmaceutical industries as antibiotics (47), and anticancer agents (48), anti-virals (48), diabetes medications (48). Endophytic bacteria isolated from *Vanda cristata* contain bioactive metabolite ethyl iso-allocholate with antimicrobial properties that contribute to plant fitness by overcoming biotic stress (23).

Conservation implications of orchid by orchid-associated bacteria

It is challenging to conserve orchids due to their strong biotic relationships and tiny seeds that require mycorrhiza to germinate. Arbuscular mycorrhizal fungi (AMF), one of these advantageous microbes, have symbiotic relationships with over 80% of the species of vascular plants (42) and are regarded as an essential class of soil

microorganisms that enhance plants' uptake of phosphorous (49). AMF may also increase a plant's resistance to abiotic stressors (50) and its ability to defend itself against infections. Mycorrhiza helper bacteria (MHB) facilitate the establishment and the functioning of the symbiotic association of AMF by stimulating spore germination (51), mycelial growth (52), root colonization (53), sporulation and also by reducing stresses that could impact AMF symbiosis. Also, they can stimulate root exudates, activating AM hyphae and, therefore, a higher rate of mycorrhizal root colonization (54).

Orchid plant growth promoting rhizobacteria

Plant growth - promoting rhizobacteria (PGPR) has been continuously reported as a plant root colonizing endophytic bacterium that helps plants absorb nutrients and minerals from the soil and plays a vital role in mineralizing organic phosphorus (55) and iron, fixing di-nitrogen (56) and producing phytohormones and bio regulators (57). An endophytic bacteria isolated from *Cymbidium* orchid contributes to plant growth and biomass production (12). A study by Alibrandi *et al.* analyzed the diversity of endophytic bacteria associated with *Neottia ovata*, *Serapias vomeracea*, and *Spiranthes spiralis* and determined the importance of these bacteria for orchid development (35). In cultivated tropical orchids genera of *Calanthe*, *Acampe*, and *Dendrobium*, heterotrophic and phototrophic bacteria were abundant. The isolated strains were shown to fix nitrogen from the orchid's surrounding cyanobacterial community and to produce Indole acetic acid (IAA) from heterotrophic bacteria (58). It was also found that some strains of orchid-associated bacteria could promote symbiotic germination of *Cymbidium goeringii* (59). Plants obtain iron via root-mediated degradation by producing siderophores and chelating agents (60). In addition, some beneficial RABs, *Bacillus*, *Rhizoctonia*, etc., make phosphorus more available by acidifying (61), chelating, or releasing phosphatases (62). Nitrogen is fixed by nitrogenase activity, which makes it easier for plants to use (63) (Table 2).

Table 1. Bioactive metabolites from endophytic bacterial source

Sl.No.	Name of the endophytic bacteria	Name of the metabolites produced	Name of the host plant	References
1	<i>Streptomyces</i> sp.	Eicosanoic acid	<i>Vanda corulea</i>	22
2	<i>Streptomyces</i> sp.	Pyrazole-5-carboxylic acid	<i>V. corulea</i>	22
3	<i>Streptomyces</i> sp.	phenol,2,4-bis-(1,1-dimethyl ethyl)	<i>V. corulea</i>	22
4	<i>Streptomyces</i> sp.	Cyclononasiloxane	<i>V. corulea</i>	22
5	<i>Streptomyces</i> sp.	1-Deoxy-d-mannitol	<i>Renanthera imschootiana</i>	22
6	<i>Streptomyces</i> sp.	phenol,2,4-bis-(1,1-dimethyl ethyl)	<i>R. imschootiana</i>	22
7	<i>Streptomyces</i> sp.	Benzoic acid, 3-(2-methoxyethyl) heptyl ester	<i>R. imschootiana</i>	22
8	<i>Streptomyces</i> sp.	1-Nonadecene.	<i>R. imschootiana</i>	22
9	<i>Streptomyces</i> sp.	1,2-Benzenedicarboxylic acid, bis (2-methyl propyl) ester	<i>R. imschootiana</i>	22
10	<i>Bacillus</i> sp.	6-acetyl-beta-d-mannose	<i>V. cristata</i>	23
11	<i>Bacillus</i> sp.	ethyl iso-allocholate	<i>V. cristata</i>	23

Table 2. Role of endophytic bacteria in orchids.

Sl.No	Orchid species	Endophytic bacteria	Role in plant growth and development	References
1	<i>Paphiopedilum appletonianum</i>	<i>Streptomyces</i>	IAA production	7
		<i>Bacillus</i>	●The biological action of microbial IAA was demonstrated by treating kidney bean cuttings with bacterial supernatants, significantly stimulating root development and growth.	
2	<i>Pholidota articulata</i>	<i>Pseudomonas</i>	IAA production	7
		<i>Burkholderia</i>	●The biological action of microbial IAA was demonstrated by treating kidney bean cuttings with bacterial supernatants, which significantly stimulated root development and growth.	
		<i>Erwinia</i>		
		<i>Pseudomonas</i>		
3	<i>Dendrobium moschatum</i>	<i>Flavobacterium</i>	IAA production	14
		<i>Stenotrophomonas</i>	●The biological action of microbial IAA was demonstrated by treating kidney bean cuttings with bacterial supernatants, which significantly stimulated root development and growth.	
		<i>Pantoea</i>		
4	<i>Cymbidium eburneum</i>	<i>Chryseobacterium</i>	IAA production	13
		<i>Sphingomonas</i>	●Inoculation of <i>D. moschatum</i> seeds with <i>Sphingomonas</i> sp. and <i>Mycobacterium</i> sp. significantly improved orchid seed germination even in the absence of plant growth stimulators or mycorrhizal fungi	
5	<i>Cymbidium sp.</i>	<i>Mycobacterium</i>	Plant growth promoters	15
		<i>Paenibacillus lemtimorbus</i>	●Endophytic <i>P. lemtimorbus</i> and <i>P. macerans</i> strains significantly increased biomass in <i>Cattleya</i> seedling shoots and roots compared to non-inoculated seedlings.	
6	<i>Neottia ovata</i>	<i>P. macerans</i>	Plant growth promoters	16
		<i>Bacillus</i>	●Isolated <i>Herbaspirillum</i> and <i>Stenotrophomonas</i> increased the dry matter in <i>Cymbidium</i> sp. plants by 26% and 29%, respectively, compared to the control.	
		<i>Burkholderia</i>	Plant growth promoters	
		<i>Rhodanobacter</i>	●Most bacterial isolates demonstrated some potential Plant Growth Promoting characteristics, such as nutrient solubilization, ACC deaminase activity, and IAA production.	
7	<i>Serapias vomareceae</i>	<i>Pseudomonas</i>	Plant growth promoters	16
		<i>Sphingomonas</i>	●Most bacterial isolates demonstrated some potential Plant Growth Promoting characteristics, such as nutrient solubilization, ACC deaminase activity, and IAA production.	
		<i>Cutibacterium</i>	Plant growth promoters	
8	<i>Spiranthes spiralis</i>	<i>Pseudomonas</i>	●Most bacterial isolates demonstrated some potential Plant Growth Promoting characteristics, such as nutrient solubilization, ACC deaminase activity, and IAA production.	16
		<i>Sphingomonas</i>	Plant growth promoters	
9	<i>Cattleya walkeriana</i>	<i>Leutibacter</i>	●Most bacterial isolates demonstrated some potential Plant Growth Promoting characteristics, such as nutrient solubilization, ACC deaminase activity, and IAA production.	17
		<i>Cutibacterium</i>	Plant growth promoters	
		<i>Paenibacillus</i>	●These endophytes were inoculated in acclimatizing seedlings of <i>C. walkeriana</i> obtained from micropropagation and have a potential role as plant growth promoters, as well as in morphological alterations, nutrient uptake, and improved antioxidant enzyme activity.	
		<i>Enterobacter</i>	Plant growth promoters	
10	<i>Dendrobium officinale</i>	<i>Rhizobium</i>	●Nitrogen fixation and the secretion of plant growth regulators such as IAA, Zeatin, Salicylic acid, and ABA offer <i>S. paucimobilis</i> growth-promoting capabilities that could be used for commercial cultivation of <i>D. officinale</i> .	25
		<i>Pseudomonas</i>	Plant growth promoters	
11	<i>Dendrobium sp.</i>	<i>Sphingomonas paucimobilis</i>	Antimicrobial activity	26
		<i>Burkholderia</i>	●Antimicrobial activity against <i>Athelia rolfsii</i> , <i>Myrothecium roridum</i> , and <i>Pectobacterium carotovorum</i> by the Kirby-Bauer method showed activity against at least one pathogen in this study.	
		<i>Bacillus</i>		
		<i>Paraburkholderia</i>		
		<i>Kluyvera</i>		

		Plant growth promoters		
12	<i>Epipactis</i> sp.	<i>Bacillus</i> <i>Pseudomonas</i>	●Endophytic bacteria, particularly those categorized as prototrophic organisms, can boost the growth and development of <i>Epipactis</i> orchids, particularly those grown in nutrient-poor soils.	27
13	<i>Epipactis</i> sp.	<i>Bacillus</i> <i>Clostridium</i> <i>Pseudomonas</i> <i>Stenotrophomonas</i>	●Endophytic bacteria may stimulate the growth and development of <i>Epipactis</i> orchids, especially those that grow in nutrient-poor soil.	28
14	<i>Dendrobium nobile</i>	<i>Pseudomonas fluorescens</i> <i>Klebsiella oxytoca</i>	● <i>K. oxytoca</i> and <i>P. fluorescens</i> produced the highest amounts of auxin in nitrogen-limiting and NO ₃ -containing environments. Bacterization of <i>D. nobile</i> seeds increased their in vitro germination.	29
15	<i>Doritaenopsis Jiuhiboo</i>	<i>Mycobacterium</i>	● <i>Mycobacterium</i> can synthesize and secrete the plant growth hormone indole-3-acetic acid (IAA). In <i>Doritaenopsis</i> , <i>Mycobacterium</i> inoculation enhanced root number and length, plant height, leaf number, and leaf length. It also increases seed germination in <i>Doritaenopsis</i> .	30




Fig. 2. Schematic diagram about the mode of action of PGPR in promoting plant growth

Several strains of *Sphingomonas*, *Agrococcus*, *Mycobacterium*, and *Bacillus* produce IAA, which facilitates the germination of host orchids (31). *Sphingomonas paucimobilis*, an endophytic bacterium from *Dendrobium officinale* roots, fixes atmospheric nitrogen and increases seedling growth. Seedlings inoculated with *S. paucimobilis* contain high amounts of salicylic acid, IAA, and abscisic acid, which may promote growth or provide systemic resistance to biotic stresses (64).

Conclusion

The article focuses on the importance of endophytic bacteria in orchid ecology, particularly their symbiotic relationships and potential contributions to orchid

growth, development, and conservation. Orchids are fascinating plants with remarkable adaptations to survive in various environments, and their interactions with mycorrhizal fungi have been extensively studied. However, the role of bacterial endophytes has been relatively understudied, despite some notable research highlighting their significance, the diversity of endophytic bacteria found in various orchid species, including important genera like *Bacillus*, *Streptomyces*, *Pseudomonas*, and *Sphingomonas*. These endophytic bacteria have been shown to facilitate plant growth by providing nutrients, promoting hormonal balance, and enhancing the resistance to biotic and abiotic stresses.

Moreover, some endophytic bacteria have been found to produce bioactive metabolites with potential

pharmaceutical applications, including antibiotics, anti-cancer compounds, anti-virals, and diabetes medications. These findings indicate that orchid-associated bacteria play essential roles in plant health and hold promise for future drug discovery and biotechnological applications. Conservation implications are also discussed, as orchids face challenges in their conservation due to their complex relationships with mycorrhizal fungi and the low germination rates of their tiny seeds. Understanding the role of endophytic bacteria in these interactions may offer new avenues for conserving orchids, enhancing their germination success, and improving their adaptability to changing environmental conditions. This article highlights the often-overlooked role of endophytic bacteria in orchid biology and ecology. Further research could lead to valuable insights into orchid conservation, sustainable propagation methods, and the potential discovery of novel bioactive compounds with critical pharmaceutical applications. Orchids, with their captivating beauty and therapeutic qualities, deserve continued attention and research to ensure their preservation and sustainable use in the future.

Acknowledgments

The authors thank the Department of Botany, University of Kerala, for constant support. Aneesa Ansiya acknowledges the University Grants Commission (UGC) and Rajeevan Parvathi, the Department of Science and Technology (DST-INSPIRE), Govt. of India, for financial support through the award of Junior Research Fellowship.

Authors' contributions

Conceptualization - Sugathan S.; writing - original draft preparation -Aneesa A., Rajeevan P.; resources - Sugathan S.; writing - review and editing - Sugathan S., Gangaprasad A.; supervision - Sugathan S.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None.

References

- Popova E, Kim HH, Saxena PK, Engelmann F, Pritchard HW. Frozen beauty: The cryobiotechnology of orchid diversity. *Biotechnology advances*. 2016;34(4):380-403. <https://doi.org/10.1016/j.biotechadv.2016.01.001>
- Teo ZW, Zhou W, Shen L. Dissecting the function of MADS-box transcription factors in orchid reproductive development. *Frontiers in Plant Science*. 2019;10:1474. <https://doi.org/10.3389/fpls.2019.01474>
- Gutiérrez RM. Orchids: A review of uses in traditional medicine, its phytochemistry and pharmacology. *J. Med. Plants Res*. 2010;4(8):592-638.
- Kaur S. Mycorrhiza in Orchids. *Orchids Phytochemistry, Biology, and Horticulture: Fundamentals and Applications*. 2020:1-4. https://doi.org/10.1007/978-3-030-11257-8_7-1
- Liu H, Luo Y, Liu H. Studies of mycorrhizal fungi of Chinese orchids and their role in orchid conservation in China-a review. *The Botanical Review*. 2010;76:241-62. <https://doi.org/10.1007/s12229-010-9045-9>
- Herrera H, García-Romera I, Meneses C, Pereira G, Arriagada C. Orchid mycorrhizal interactions on the Pacific side of the Andes from Chile. A review. *Journal of Soil Science and Plant Nutrition*. 2019;19:187-202. <https://doi.org/10.1007/s42729-019-00026-x>
- Pujasatria GC, Miura C, Kaminaka H. *In vitro* symbiotic germination: A revitalized heuristic approach for orchid species conservation. *Plants*. 2020;9(12):1742. <https://doi.org/10.3390/plants9121742>
- Li T, Wu S, Yang W, Selosse MA, Gao J. How mycorrhizal associations influence the orchid distribution and population dynamics. *Frontiers in Plant Science*. 2021;12:647114. <https://doi.org/10.3389/fpls.2021.647114>
- McCormick MK, Whigham DF, Canchani-Viruet A. Mycorrhizal fungi affect orchid distribution and population dynamics. *New Phytologist*. 2018;219(4):1207-15. <https://doi.org/10.1111/nph.15223>
- White Jr JF, Torres MS, Sullivan RF, Jabbour RE, Chen Q, Tadych M, Irizarry I, Bergen MS, Havkin-Frenkel D, Belanger FC. Occurrence of *Bacillus amyloliquefaciens* as a systemic endophyte of vanilla orchids. *Microscopy research and technique*. 2014;77(11):874-85. <https://doi.org/10.1002/jemt.22410>
- Yang S, Zhang X, Cao Z, Zhao K, Wang S, Chen M, Hu X. Growth-promoting *Sphingomonas paucimobilis* ZJSH 1 associated with *Dendrobium officinale* through phytohormone production and nitrogen fixation. *Microbial biotechnology*. 2014;7(6):611-20. <https://doi.org/10.1111/1751-7915.12148>
- Gontijo JB, Andrade GV, Baldotto MA, Baldotto LE. Bioprospecting and selection of growth-promoting bacteria for *Cymbidium* sp. orchids. *Scientia Agricola*. 2018 Sep;75:368-74. <https://doi.org/10.1590/1678-992x-2017-0117>
- Dias AC, Costa FE, Andreote FD, Lacava PT, Teixeira MA, Assumpção LC, Araújo WL, Azevedo JL, Melo IS. Isolation of micropropagated strawberry endophytic bacteria and assessment of their potential for plant growth promotion. *World Journal of Microbiology and Biotechnology*. 2009;25:189-95. <https://doi.org/10.1007/s11274-008-9878-0>
- Tsavkelova EA, Cherdyntseva TA, Klimova SY, Shestakov AI, Botina SG, Netrusov AI. Orchid-associated bacteria produce indole-3-acetic acid, promote seed germination, and increase their microbial yield in response to exogenous auxin. *Archives of Microbiology*. 2007;188:655-64. <https://doi.org/10.1007/s00203-007-0286-x>
- Malboobi MA, Behbahani M, Madani H, Owlia P, Deljou A, Yakhchali B, Moradi M, Hassanabadi H. Performance evaluation of potent phosphate solubilizing bacteria in potato rhizosphere. *World Journal of Microbiology and Biotechnology*. 2009;25:1479-84. <https://doi.org/10.1007/s11274-009-0038-y>
- Strobel G, Ford E, Worapong J, Harper JK, Arif AM, Grant DM, Fung PC, Chau RM. Isopestacin, an isobenzofuranone from *Pestalotiopsis microspora*, possessing antifungal and antioxidant activities. *Phytochemistry*. 2002;60(2):179-183. [https://doi.org/10.1016/S0031-9422\(02\)00062-6](https://doi.org/10.1016/S0031-9422(02)00062-6)
- Suryanarayanan TS, Murali TS. Incidence of *Leptosphaerulina crassiasca* in symptomless leaves of peanut in southern India. *Journal of Basic Microbiology*. 2006;46(4):305-309. <https://doi.org/10.1002/jobm.200510126>
- McCormick MK, Whigham DF, Canchani-Viruet A. Mycorrhizal fungi affect orchid distribution and population dynamics. *New Phytologist*. 2018;219(4):1207-15. <https://doi.org/10.1111/nph.15223>

19. Tsavkelova EA, Cherdyntseva TA, Netrusov AI. Bacteria associated with the roots of epiphytic orchids. *Microbiology*. 2004;73:710-715. <https://doi.org/10.1007/s11021-005-0013-z>
20. Tsavkelova EA, Lobakova ES, Kolomeitseva GL, Cherdyntseva TA, Netrusov AI. Localization of associative cyanobacteria on the roots of epiphytic orchids. *Microbiology*. 2003 Jan;72:86-91. <https://doi.org/10.1023/A:1022286225013>
21. Wu W, Chen W, Liu S, Wu J, Zhu Y, Qin L, Zhu B. Beneficial relationships between endophytic bacteria and medicinal plants. *Frontiers in plant science*. 2021 ;12:646146. <https://doi.org/10.3389/fpls.2021.646146>
22. Saikia J, Mazumdar R, Thakur D. Phylogenetic affiliation of endophytic actinobacteria associated with selected orchid species and their role in growth promotion and suppression of phytopathogens. *Frontiers in Plant Science*. 2022;13:1058867. <https://doi.org/10.3389/fpls.2022.1058867>
23. Shah S, Chand K, Rekadwad B, Shouche YS, Sharma J, Pant B. A prospectus of plant growth promoting endophytic bacterium from orchid (*Vanda cristata*). *BMC biotechnology*. 2021;21(1):1-9. <https://doi.org/10.1186/s12896-021-00676-9>
24. Tsavkelova EA, Cherdyntseva TA, Botina SG, Netrusov AI. Bacteria associated with orchid roots and microbial production of auxin. *Microbiological research*. 2007;162(1):69-76. <https://doi.org/10.1016/j.micres.2006.07.014>
25. Herrera, H., Fuentes, A., Soto, J., Valadares, R. and Arriagada, C., 2022. Orchid-associated bacteria and their plant growth promotion capabilities. In *Orchids phytochemistry, biology, and horticulture: fundamentals and applications* (pp. 175-200). Cham: Springer International Publishing. https://doi.org/10.1007/978-3-030-38392-3_35
26. Faria DC, Dias AC, Melo IS, de Carvalho Costa FE. Endophytic bacteria isolated from orchid and their potential to promote plant growth. *World Journal of Microbiology and Biotechnology*. 2013;29:217-21.
27. Dearnaley JD. Further advances in orchid mycorrhizal research. *Mycorrhiza*. 2007;17(6):475-86.
28. Araújo WL, Marcon J, Maccheroni Jr W, Van Elsas JD, Van Vuurde JW, Azevedo JL. Diversity of endophytic bacterial populations and their interaction with *Xylella fastidiosa* in citrus plants. *Applied and Environmental Microbiology*. 2002;68(10):4906-14. <https://doi.org/10.1128/AEM.68.10.4906-4914.2002>
29. Alibrandi P, Cardinale M, Rahman MM, Strati F, Ciná P, de Viana ML, Giamminola EM, Gallo G, Schnell S, De Filippo C, Ciaccio M. The seed endosphere of *Anadenanthera colubrina* is inhabited by a complex microbiota, including *Methylobacterium* spp. and *Staphylococcus* spp. with potential plant-growth-promoting activities. *Plant and Soil*. 2018;422:81-99.
30. Bulgarelli D, Schlaeppi K, Spaepen S, Van Themaat EV, Schulze-Lefert P. Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology*. 2013 ;64:807-38. <https://doi.org/10.1146/annurev-arplant-050312-120106>
31. Tsavkelova EA, Egorova MA, Leontieva MR, Malakho SG, Kolomeitseva GL, Netrusov AI. *Dendrobium nobile* Lindl. Seed germination in co-cultures with diverse associated bacteria. *Plant Growth Regulation*. 2016;80:79-91.
32. Duponnois R, Garbaye J. Mycorrhization helper bacteria associated with the Douglas fir-*Laccaria laccata* symbiosis: effects in aseptic and in glasshouse conditions. In *Annales des sciences forestières* 1991 (Vol. 48, No. 3, pp. 239-251). EDP Sciences. <https://doi.org/10.1051/forest:19910301>
33. Frey-Klett P, Garbaye JA, Tarkka M. The mycorrhiza helper bacteria revisited. *New phytologist*. 2007;176(1):22-36.
34. Yu J, Zhou XF, Yang SJ, Liu WH, Hu XF. Design and application of specific 16S rDNA-targeted primers for assessing endophytic diversity in *Dendrobium officinale* using nested PCR-DGGE. *Applied microbiology and biotechnology*. 2013;97:9825-36.
35. Alibrandi P, Lo Monaco N, Calevo J, Voyron S, Puglia AM, Cardinale M, Perotto S. Plant growth promoting potential of bacterial endophytes from three terrestrial Mediterranean orchid species. *Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology*. 2021;155(6):1153-64. <https://doi.org/10.1080/11263504.2020.1829731>
36. Andrade GV, Rodrigues FA, Nadal MC, Caroline MD, Martins AD, Rodrigues VA, dos Reis Ferreira GM, Pasqual M, Buttros VH, Dória J. Plant-endophytic bacteria interactions associated with root and leaf microbiomes of *Cattleya walkeriana* and their effect on plant growth. *Scientia Horticulturae*. 2023;309:111656.
37. Compant S, Cambon MC, Vacher C, Mitter B, Samad A, Sessitsch A. The plant endosphere world-bacterial life within plants. *Environmental Microbiology*. 2021;23(4):1812-29. <https://doi.org/10.1111/1462-2920.15240>
38. Sudhakar P, Chattopadhyay GN, Gangwar SK, Ghosh JK. Effect of foliar application of *Azotobacter*, *Azospirillum*, and *Beijerinckia* on leaf yield and quality of mulberry (*Morus alba*). *The Journal of Agricultural Science*. 2000;134(2):227-34. <https://doi.org/10.1017/S0021859699007376>
39. Glick BR. Bacteria with ACC deaminase can promote plant growth and help to feed the world. *Microbiological Research*. 2014;169(1):30-9. <https://doi.org/10.1016/j.micres.2013.09.009>
40. Glick BR. Plant growth-promoting bacteria: mechanisms and applications. *Scientifica*. 2012;2012. <https://doi.org/10.6064/2012/963401>
41. Rodríguez H, Fraga R. Phosphate solubilizing bacteria and their role in plant growth promotion. Department of Microbiology, Cuban Research Institute on Sugarcane By-Products (ICIDCA), Havana, Cuba. Havana. Cuba. 1999.
42. Simon L, Bousquet J, Lévesque RC, Lalonde M. Origin and diversification of endomycorrhizal fungi and coincidence with vascular land plants. *Nature*. 1993;363(6424):67-9. <https://doi.org/10.1038/363067a0>
43. Singh SP, Gaur R. Evaluation of antagonistic and plant growth promoting activities of chitinolytic endophytic actinomycetes associated with medicinal plants against *Sclerotium rolfsii* in chickpea. *Journal of Applied Microbiology*. 2016;121(2):506-18. <https://doi.org/10.1111/jam.13176>
44. Brader G, Compant S, Mitter B, Trognitz F, Sessitsch A. Metabolic potential of endophytic bacteria. *Current Opinion in Biotechnology*. 2014;27:30-7. <https://doi.org/10.1016/j.copbio.2013.09.012>
45. Wilson D. Endophyte: the evolution of a term, and clarification of its use and definition. *Oikos*. 1995:274-6. <https://doi.org/10.2307/3545919>
46. Stierle A, Strobel G, Stierle D. Taxol and taxane production by *Taxomyces andreanae*, an endophytic fungus of Pacific yew. *Science*. 1993;260(5105):214-6. <https://doi.org/10.1126/science.8097061>
47. Christina A, Christopher V, Bhore SJ. Endophytic bacteria as a source of novel antibiotics: an overview. *Pharmacognosy reviews*. 2013;7(13):11.
48. Guo B, Wang Y, Sun X, Tang K. Bioactive natural products from endophytes: a review. *Applied Biochemistry and Microbiology*. 2008;44:136-42.
49. Garcia-Garrido JM, Tribak M, Rejon-Palomares A, Ocampo JA, Garcia-Romera I. Hydrolytic enzymes and ability of arbuscular mycorrhizal fungi to colonize roots. *Journal of Experimental Botany*. 2000;51(349):1443-8. <https://doi.org/10.1093/jexbot/51.349.1443>
50. Lies A, Delteil A, Prin Y, Duponnois R. Using mycorrhiza helper microorganisms (MHM) to improve the mycorrhizal efficiency on plant growth. *Role of rhizospheric microbes in soil: volume 1:*

- Stress Management and Agricultural Sustainability. 2018;277-98.
51. Barea JM, Pozo MJ, Azcon R, Azcon-Aguilar C. Microbial co-operation in the rhizosphere. *Journal of Experimental Botany*. 2005;56(417):1761-78. <https://doi.org/10.1093/jxb/eri197>
 52. Garbaye J. Helper bacteria: a new dimension to the mycorrhizal symbiosis. *New Phytologist*. 1994;128(2):197-210.
 53. Yadav BK, Sidhu AS. Dynamics of Potassium and Their Bioavailability for Plant Nutrition. In: Meena V, Maurya B, Verma J, Meena R. (eds) *Potassium Solubilizing Microorganisms for Sustainable Agriculture*. Springer, New Delhi. 2016 https://doi.org/10.1007/978-81-322-2776-2_14
 54. Sangwan S, Prasanna R. Mycorrhizae helper bacteria: unlocking their potential as bioenhancers of plant-arbuscular mycorrhizal fungal associations. *Microbial Ecology*. 2022 Jul;84(1):1-0. <https://doi.org/10.1007/s00248-021-01831-7>
 55. Rashid M, Khalil S, Ayub N, Alam S, Latif F. Organic acids production and phosphate solubilization by phosphate solubilizing microorganisms (PSM) under *in vitro* conditions. *Pak J Biol Sci*. 2004;7(2):187-96. <https://doi.org/10.3923/pjbs.2004.187.196>
 56. Mabood F, Zhou X, Smith DL. Microbial signaling and plant growth promotion. *Canadian Journal of Plant Science*. 2014;94(6):1051-63. <https://doi.org/10.4141/cjps2013-148>
 57. Borriss R. Use of plant-associated *Bacillus* strains as biofertilizers and bio control agents in agriculture. *Bacteria in agrobiology: Plant Growth Responses*. 2011:41-76. https://doi.org/10.1007/978-3-642-20332-9_3
 58. Tsavkelova EA, Cherdyntseva TA, Netrusov AI. Auxin production by bacteria associated with orchid roots. *Microbiology*. 2005;74:46-53.
 59. Sun L, Shao H, Liu L, Zhang R, Zhao L, Li L, Yao N. Diversity of siderophore-producing endophytic bacteria of *Cymbidium goeringii* roots. *Acta Microbiologica Sinica*. 2011;51(2):189-95
 60. Molnár Z, Solomon W, Mutum L, Janda T. Understanding the Mechanisms of Fe Deficiency in the Rhizosphere to Promote Plant Resilience. *Plants*. 2023;12(10):1945. <https://doi.org/10.3390/plants12101945>
 61. Santoyo G, Moreno-Hagelsieb G, del Carmen Orozco-Mosqueda M, Glick BR. Plant growth-promoting bacterial endophytes. *Microbiological Research*. 2016;183:92-9. <https://doi.org/10.1016/j.micres.2015.11.008>
 62. Rasul M, Yasmin S, Zubair M, Mahreen N, Yousaf S, Arif M, Sajid ZI, Mirza MS. Phosphate solubilizers as antagonists for bacterial leaf blight with improved rice growth in phosphorus deficit soil. *Biological Control*. 2019;136:103997. <https://doi.org/10.1016/j.biocontrol.2019.05.016>
 63. Afzal I, Shinwari ZK, Sikandar S, Shahzad S. Plant beneficial endophytic bacteria: Mechanisms, diversity, host range and genetic determinants. *Microbiological Research*. 2019;221:36-49. <https://doi.org/10.1016/j.micres.2019.02.001>
 64. Yang S, Zhang X, Cao Z, Zhao K, Wang S, Chen M, Hu X. Growth-promoting *Sphingomonas paucimobilis* ZJSH 1 associated with *Dendrobium officinale* through phytohormone production and nitrogen fixation. *Microbial Biotechnology*. 2014;7(6):611-20. <https://doi.org/10.1111/1751-7915.12148>