



REVIEW ARTICLE

Beneficial effects of biochar application on mitigating the drought and salinity stress implications on plants

Anil Patani¹, Dharmendra Prajapati¹, Sachidanand Singh¹, Yuri Enakiev², Snezhan Bozhkov², Dilfuza Jabborova^{3,4,5}, Chinmayi Joshi^{1*}

1 Department of Biotechnology, Smt. S. S. Patel Nootan Science and Commerce College, Sankalchand Patel University, Visnagar-384315, Gujarat, India.

2 Nikola Pushkarov Institute of Soil Science, Agrotechnologies and Plant Protection

3, Shose Bankya St., Sofia, 1331, Bulgaria

3 Institute of Genetics and Plant Experimental Biology, Uzbekistan Academy of Sciences, Kibray 111208, Uzbekistan;

4 Faculty of Biology, National University of Uzbekistan, Tashkent 100174, Uzbekistan

5 Department of Biotechnology, School of Biological Engineering & Sciences, Shobhit Institute of Engineering & Technology, Meerut 250110, India

*Email: ckjoshi.fsh@spu.ac.in



ARTICLE HISTORY

Received: 19 April 2023

Accepted: 23 June 2023

Available online

Version 1.0 : 30 September 2023



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Patani A, Prajapati D, Singh S, Enakiev Y, Bozhkov S, Jabborova D, Joshi C. Beneficial effects of biochar application on mitigating the drought and salinity stress implications on plants. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.2591>

Abstract

Biochar, an amorphous and highly porous carbonaceous substance derived from the thermal decomposition of organic matter, has been empirically proven to enhance soil water retention capacity, mitigate soil salinity, and augment nutrient bioavailability. Consequently, these improvements exert a stimulating influence on the growth and development of medicinal plants. Numerous scientific investigations have corroborated that the incorporation of biochar into the cultivation of medicinal flora can lead to increased plant biomass, heightened photosynthetic efficiency, and augmented accumulation of bioactive compounds. Furthermore, the utilization of biochar exhibits the potential to curtail the necessity for chemical fertilizers, which can otherwise have deleterious effects on soil health and the environment. A comprehensive comprehension of biochar's prospective role as a sustainable, long-term strategy for augmenting the productivity and resilience of medicinal plant cultivation in arid and saline environments holds paramount importance for ensuring a consistent supply of medicinal plants in the forthcoming years. This review aims to delve into the mechanistic foundations underpinning the beneficial impacts of biochar on plant development and the accumulation of bioactive constituents. It also explores the feasibility of biochar as a sustainable instrument for enhancing the cultivation of medicinal plants under adverse environmental conditions.

Keywords

Biochar, medicinal plants, salinity stress, drought stress

Introduction

Biochar is a carbon-rich substance produced through the pyrolysis of organic materials, such as wood or agricultural residues, under controlled, low-oxygen conditions (1). This process results in a highly porous and chemically stable material with versatile applications. One prominent application involves utilizing biochar as a soil amendment in agriculture to augment soil quality and fertility significantly. Extensive research has demonstrated its capacity to increase crop yields, reduce the reliance on synthetic fertilizers and pesticides, and enhance soil water retention (2). Importantly, biochar's exceptional stability allows it to persist in the soil for extended periods, potentially spanning hundreds to thousands of years, enabling it to serve as a carbon sequestration tool. Consequently, the integration of biochar into agricultural practices yields a dual benefit: it

enhances soil health while simultaneously contributing to climate change mitigation through long-term carbon retention (3). Figure 1 illustrates the various types of biochar with applications relevant to soil health improvement.




Fig. 1. Types of biochar

Drought and salinity stress are significant environmental factors that can profoundly impact the growth, development, and chemical composition of medicinal plants, potentially leading to diminished yield, compromised quality, and reduced potency of bioactive phytochemicals, frequently employed in both traditional medicinal practices and contemporary pharmaceuticals (4, 5). Drought stress manifests when plants are subjected to prolonged periods of diminished water availability, resulting in decreased vegetative growth, desiccation of leaves, and diminished photosynthetic activity (6). Various types of biochar have been documented to exhibit favorable effects (Table 1).

As a consequence, the synthesis of secondary metabolites in medicinal plants, including alkaloids and flavonoids, can be significantly reduced, leading to compromised therapeutic efficacy (12-14). Conversely, salinity stress occurs when plants are exposed to elevated salt levels in the soil or water, resulting in ion toxicity and osmotic stress. This can manifest as leaf chlorosis, inhibited plant growth, and ultimately, plant mortality. Excessive salt levels can also disrupt the uptake and transport of essential nutrients, causing imbalances in plant metabolism and a reduction in the production of bioactive compounds (15, 16). To mitigate the adverse effects of drought and salinity stress on medicinal plants, various strategies can be implemented, such as utilizing drought and salt-tolerant plant varieties, incorporating biochar, employing plant growth-promoting rhizobacteria, amending soil to enhance its structure and water-holding capacity, and adopting more efficient irrigation methods

(17). Additionally, several plant growth regulators and natural products, such as extracts from seaweed, have demonstrated the ability to stimulate plant growth and promote the synthesis of secondary metabolites in medicinal plants under conditions of drought and salinity stress (18). Biochar possesses notable water retention capacity, enabling it to support medicinal plants in enduring drought conditions. Furthermore, biochar has the capacity to absorb and retain nutrients, thereby making them accessible to medicinal plants even in saline environments. Studies have revealed that biochar can enhance the production of secondary metabolites in plants, including medicinal species, consequently enhancing their therapeutic potential (19).

This review aims to explore the advantageous ramifications of biochar application in ameliorating the adverse effects of drought and salinity stress on medicinal plants. The primary focus of this review is to delve into the underlying mechanisms that drive the beneficial influence of biochar on plant growth and the accumulation of bioactive compounds. Gaining insights into the potential of biochar as a sustainable agronomic approach for augmenting the productivity and resilience of medicinal plant cultivation in arid and saline environments is imperative for securing a consistent supply of medicinal plants in the forthcoming years.

The effects of biochar on plant growth and physiology

Biochar exerts significant effects on plant development and physiological processes. A primary advantage of biochar lies in its ability to enhance soil structure and augment water retention capacity, thereby fostering optimal conditions for plant growth and development. Moreover, biochar serves as a reservoir of essential nutrients, including nitrogen, phosphate, and potassium, which are gradually released into the soil over time, ensuring a sustained supply of plant nutrition (20, 21). Additionally, biochar exhibits the capacity to stimulate soil microbial activity, thereby conferring benefits upon plant health. It has been empirically demonstrated to promote the proliferation of beneficial microorganisms, such as mycorrhizal fungi, which enhance plant nutrient uptake and bolster resistance to diseases (22). Notably, the carbon-sequestering property of biochar in soil holds promise for environmental conservation, as it aids in carbon storage, reduces greenhouse gas emissions, and contributes to mitigating climate change effects (23).

In sum, biochar emerges as a valuable tool for advancing sustainable agriculture and environmental management, given its profound influence on plant growth and physiological processes. However, it is

Table 1. Effect of different types of biochar on soil and plant health

Type of biochar	Reported for	Reference
Manure-based biochar	Improvement of the water holding capacity of the soil and improve crop yield by conserving the rainfall water in arid regions.	(7)
Cow manure-based biochar	Soil amendment for reducing cadmium availability and accumulation by <i>Brassica chinensis</i> L. in acidic red soil	(8)
Soft wood biochar	Reduction in N ₂ O emission in long-term.	(9)
Sewage sludge biochar	Improvement in corn fruit yield	(10)
Bamboo biochar	Improvement of soil health	(11)

imperative to acknowledge that the appropriate rate and timing of biochar application may vary contingent upon soil type, plant species, and prevailing weather conditions. Hence, further research is imperative to comprehensively elucidate its multifaceted impacts on plant growth and physiology (24).

Biochar's ability to mitigate drought stress

Biochar has been identified as a potential mitigation strategy for alleviating drought-induced stress in plants. It achieves this by enhancing soil's water retention capacity, thereby serving as a pivotal mechanism for aiding plants in coping with water scarcity (25). Biochar's efficacy in ameliorating drought stress stems from its high porosity and water-retention capabilities. Consequently, it plays a vital role in maintaining adequate soil moisture levels, mitigating stress on plants. Furthermore, aside from its water-holding capacity, biochar has been demonstrated to improve soil structure and aeration. These improvements augment the soil's pore space, which in turn enhances water infiltration and retention (26). The enhanced aeration facilitated by biochar is also instrumental in increasing oxygen availability to plant roots, a critical factor for optimal plant growth and development, especially in arid conditions. Additionally, the presence of biochar in the environment has been associated with the proliferation of mycorrhizal fungi, which assist plants in more efficiently absorbing nutrients and water. This microbial interaction further contributes to enhanced plant growth and heightened resistance to water stress (27).

In summary, one noteworthy advantage of utilizing biochar as a soil amendment is its potential to alleviate drought-induced stress in plants. Nonetheless, it is crucial to acknowledge that the effectiveness of biochar in reducing drought stress may vary depending on factors such as application rate, timing, and prevailing soil conditions. Consequently, further research is imperative to gain a comprehensive understanding of the underlying mechanisms governing biochar's ability to mitigate drought stress and to identify optimal strategies for leveraging biochar to enhance plant resilience in water-limited environments (28).

Biochar's ability to mitigate salinity stress

Biochar has emerged as a promising solution for mitigating salinity-induced stress on plants, albeit its effectiveness is contingent upon several factors, including application rate, timing, and soil conditions. This natural carbonaceous material, when incorporated into soil, can enhance its structural integrity and water retention capacity, thereby ameliorating the adverse effects of salinity stress. Biochar's high porosity and water-absorbing capabilities become particularly valuable during periods of water scarcity induced by excessive salinity. By reducing the detrimental impacts of salt on plants, biochar aids in preserving soil moisture levels (29). In soil environments afflicted with excessive salinity and waterlogging, biochar proves instrumental in improving soil aeration, which, in turn, enhances the availability of

oxygen to plant roots. This oxygen supply is pivotal for optimal plant growth and development, especially in conditions of water shortage (30).

Furthermore, biochar has been demonstrated to augment the cation exchange capacity (CEC) of soil, complementing its influence on soil structure and water availability. In this regard, biochar exhibits the capacity to attract and retain positively charged ions such as calcium and magnesium, thereby mitigating the detrimental consequences of excessive soil salinity (31, 32). Additionally, biochar elevates soil pH, which, in turn, reduces the availability of harmful ions in the soil solution, such as salt and chloride. This alteration in soil pH contributes to its overall efficacy in alleviating salinity-induced stress (33, 34). However, it is important to note that the effectiveness of biochar in reducing salinity stress may vary depending on the specific application rate, timing, and prevailing soil conditions (33, 34). Further research is essential to ascertain the optimal methods for applying biochar in salt-affected soils and to gain a deeper understanding of the underlying mechanisms responsible for its salinity stress-reducing capabilities. For instance, Zulfikar et al. (35) reported notable improvements in *Alpinia zerumbet* plants, including a more than 28% increase in net photosynthetic rates, a 92% and 78% rise in chlorophyll a and b contents, respectively, and a 50% enhancement in carboxylation efficiency when biochar (BC) and biochar enriched with cobalt (BC+Co) were introduced, compared to those grown in unamended sandy loam soil.

Efficacy of biochar for the improvement of various plants

Biochar has emerged as a promising tool for enhancing the growth and development of medicinal plants while also promoting the synthesis of secondary metabolites with therapeutic properties (35). This is primarily attributed to its multifaceted impact on soil quality, encompassing improved fertility, enhanced soil structure, and increased water retention capacity, all of which collectively accelerate the growth of medicinal plants. Notably, biochar acts as a slow-release reservoir of essential nutrients, including nitrogen, phosphorus, and potassium, crucial for both plant growth and the biosynthesis of secondary metabolites (36). Biochar additionally fosters the establishment of beneficial soil microorganisms, particularly mycorrhizal fungi, which play a pivotal role in enhancing nutrient uptake and overall plant health. The presence of these microorganisms has been found to facilitate the synthesis of secondary metabolites, which medicinal plants typically produce in response to biotic and abiotic stressors (37).

Furthermore, biochar has been observed to exert a positive influence on the production of secondary metabolites in medicinal plants, such as phenolic compounds, flavonoids, and essential oils. These phytochemicals possess pharmacological properties, making them valuable in both traditional and contemporary medicinal applications (38). In sum, the

incorporation of biochar into soil holds the potential to significantly benefit medicinal plants by promoting their growth and the production of medicinally relevant secondary metabolites. However, it is important to note that the optimal application rate and timing of biochar may vary depending on the specific medicinal plant species, soil type, and environmental conditions, necessitating further research (39).

Numerous studies have reported that biochar enhances root morphological traits (40) and physiological characteristics (41,42) in medicinal plants. Furthermore, biochar amendments have been shown to reduce plant oxidative stress, as evidenced by reduced levels of leaf free proline and glycine betaine (43). Additionally, the biomass of *Salvia miltiorrhiza* increased with the proportion of biochar in the soil mix (ranging from 0% to 32%). Importantly, biochar supplementation resulted in a significant reduction in cadmium (Cd) content in both leaves and roots of *S. miltiorrhiza*, with decreases of 52.8% and 43.6%, respectively. Moreover, extensive research has demonstrated the efficacy of biochar in mitigating the adverse effects of drought and salt stress on various plant species (Table 2).

Conclusion

In summary, biochar has demonstrated its efficacy in mitigating the adverse effects of drought and salinity stress on medicinal plants. It enhances soil's water retention capacity, diminishes soil salinity levels, and augments nutrient accessibility, thereby fostering plant growth and development. Numerous studies have reported that the application of biochar to medicinal plants results in increased plant biomass, enhanced photosynthetic activity, and heightened accumulation of bioactive compounds. Moreover, the utilization of biochar can curtail the reliance on chemical fertilizers, which can be detrimental to both soil health and the environment. In conclusion, biochar presents itself as a promising strategy for enhancing the productivity and resilience of medicinal plant cultivation in arid and saline environments. Nonetheless, further research is imperative to fine-tune biochar application methods tailored to specific plant species and environmental conditions, as well as to assess its long-term effects on soil health and plant growth.

Authors contributions

AP: Writing–Original Draft, Review and Editing. CJ: Writing, Review, Editing. DP: Writing–Original Draft, Review and Editing. SS: Conceptualization, Review, Editing. YE: Conceptualization, Review, Editing. SB: Conceptualization, Review, Editing. DJ: Conceptualization, Review, Editing. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None.

References

1. Yaashikaa PR, Kumar PS, Varjani S, Saravanan A. A critical review on the biochar production techniques, characterization, stability and applications for circular bioeconomy. *Biotechnol Rep.* 2020; 28 (1): e00570. <https://doi.org/10.1016/j.btre.2020.e00570>
2. Ding Y, Liu Y, Liu S, Li Z, Tan X, Huang X, Zeng G, Zhou L, Zheng B. Biochar to improve soil fertility. A review. *Agron Sustain Dev.* 2016; (36). <https://doi.org/10.1007/s13593-016-0372-z>
3. Gross A, Bromm T, Glaser B. Soil Organic Carbon Sequestration after Biochar Application: A Global Meta-Analysis. *Agron.* 2021; (11): 2474. <https://doi.org/10.3390/agronomy11122474>
4. Seleiman MF, Al-Suhaibani N, Ali N, Akmal M, Alotaibi M, Refay Y, Dindaroglu T, Abdul-Wajid HH, Battaglia ML. Drought Stress Impacts on Plants and Different Approaches to Alleviate Its Adverse Effects. *Plants (Basel).* 2021; 10(2):259. <https://doi.org/10.3390/plants10020259>
5. Patani AV, Patel RR, Prajapati DH, Patel MS, Patel AD. Evaluation of Plant Growth Promoting Rhizobacteria to Improve Tomato (*Lycopersicon esculentum* Mill.) Productivity and Resilience under Salinity Stress. *J Cell Tissue Res.* 2020;20(2):6953-64.
6. Patani A, Prajapati D, Ali D, Kalasariya H, Yadav VK, Tank J, Bagatharia S, Joshi M, Patel A. Evaluation of the growth-inducing efficacy of various *Bacillus* species on the salt-stressed tomato (*Lycopersicon esculentum* Mill.). *Frontiers Plant Sci.* 2023;14:1018. <https://doi.org/10.3389/fpls.2023.1168155>
7. Rehman A, Nawaz S, Alghamdi HA, Alrumman S, Yan W, Nawaz MZ. Effects of manure-based biochar on uptake of nutrients and water holding capacity of different types of soils. *Case Studies Chem Environ Eng.* 2020;2:100036. <https://doi.org/10.1016/j.cscee.2020.100036>

Table 2. Efficacy of biochar in agriculture to improve plants under drought and salt stress

Biochar and its benefit	Plant(s) used for the study	Reference
Improvement of plant growth under drought condition	Spinach	(44)
Improvement of root morphological traits and growth of under salinity stress	Soybean	(45)
Enhancement of plant growth and root morphological traits under drought conditions	Okra	(46)
Improvement in plant yield in salt stress	Tomato	(47)
Mitigate drought and salt stress	Quinoa	(48)
Mitigating Drought and Salt Stress	Soybean	(49)
Improvement in plant yield and physiological properties in salt stress	Forage sorghum	(50)
Induction of drought-tolerance	Basil	(51)

8. Kiran YK, Barkat A, CUI XQ, Ying FE, PAN FS, Lin TA, YANG XE. Cow manure and cow manure-derived biochar application as a soil amendment for reducing cadmium availability and accumulation by *Brassica chinensis* L. in acidic red soil. *J Integr Agric*. 2017;16(3):725-34. [https://doi.org/10.1016/S2095-3119\(16\)61488-0](https://doi.org/10.1016/S2095-3119(16)61488-0)
9. Kalu S, Sijojoki A, Karhu K, Tammeorg P. Long-term effects of softwood biochar on soil physical properties, greenhouse gas emissions and crop nutrient uptake in two contrasting boreal soils. *Agric Ecosyst Environ*. 2021;316:107454. <https://doi.org/10.1016/j.agee.2021.107454>
10. Xie S, Yu G, Jiang R, Ma J, Shang X, Wang G, et al. Moderate sewage sludge biochar application on alkaline soil for corn growth: a field study. *Biochar*. 2021;3:135-47. <https://doi.org/10.1007/s42773-021-00085-3>
11. Cruz-Méndez AS, Ortega-Ramírez E, Lucho-Constantino CA, Arce-Cervantes O, Vázquez-Rodríguez GA, Coronel-Olivares C, et al. Bamboo biochar and a nopal-based biofertilizer as improvers of alkaline soils with low buffer capacity. *Appl Sci*. 2021;11(14):6502. <https://doi.org/10.3390/app11146502>
12. Yeshi K, Crayn D, Ritmejeriyé E, Wangchuk P. Plant Secondary Metabolites Produced in Response to Abiotic Stresses Has Potential Application in Pharmaceutical Product Development. *Molecules*. 2022 Jan 5;27(1):313. <https://doi.org/10.3390/molecules27010313>
13. Zuidersma EI, Ausma T, Stuiver CEE, Prajapati DH, Hawkesford MJ, De Kok LJ. Molybdate toxicity in Chinese cabbage is not the direct consequence of changes in sulphur metabolism. *Plant Biol*. 2020;22(2):331-6. <https://doi.org/10.1111/plb.13065>
14. Prajapati DH, Ausma T, De Boer J, Hawkesford MJ, De Kok LJ. Nickel toxicity in *Brassica rapa* seedlings-Impact on sulfur metabolism and mineral nutrient content. *J Kulturpflanzen-J Cult Plants*. 2020;72(9S):473-8. <https://doi.org/10.5073/JfK.2020.09.03>
15. Acosta-Motos JR, Ortuño MF, Bernal-Vicente A, Diaz-Vivancos P, Sanchez-Blanco MJ, Hernandez JA. Plant Responses to Salt Stress: Adaptive Mechanisms. *Agron*. 2017;7:18. <https://doi.org/10.3390/agronomy7010018>
16. Neves MI, Prajapati DH, Parmar S, Aghajanzadeh TA, Hawkesford MJ, De Kok LJ. Manganese toxicity hardly affects sulfur metabolism in *Brassica rapa*. In: *Sulfur Metabolism in Higher Plants-Fundamental, Environmental and Agricultural Aspects*. Springer International Publishing; 2017. p. 155-162.
17. Yang A, Akhtar SS, Li L, Fu Q, Li Q, Naeem MA, et al. Biochar Mitigates Combined Effects of Drought and Salinity Stress in Quinoa. *Agron*. 2020;10(6):912. <https://doi.org/10.3390/agronomy10060912>
18. Ali O, Ramsubhag A, Jayaraman J. Biostimulant Properties of Seaweed Extracts in Plants: Implications towards Sustainable Crop Production. *Plants*. 2021;10(3):531. <https://doi.org/10.3390/plants10030531>
19. Murtaza G, Ahmed Z, Eldin SM, Ali B, Bawazeer S, Usman M, et al. Biochar-Soil-Plant interactions: A cross talk for sustainable agriculture under changing climate. *Frontiers in Environmental Science*. 2023;11:1059449. <https://doi.org/10.3389/fenvs.2023.1059449>
20. Rawat J, Saxena J, Sanwal P. Biochar: A Sustainable Approach for Improving Plant Growth and Soil Properties. *IntechOpen*; 2019. <https://doi.org/10.5772/intechopen.82151>
21. Gabhane JW, Bhange VP, Patil PD, Bankar ST, Kumar S. Recent trends in biochar production methods and its application as a soil health conditioner: a review, *SN Appl Sci*. 2020;2:1-21. <https://doi.org/10.1007/s42452-020-3121-5>
22. Jabborova D, Annapurna K, Paul S, Kumar S, Saad HA, Desouky S, Ibrahim MF, Elkelish A. Beneficial features of biochar and arbuscular mycorrhiza for improving spinach plant growth, root morphological traits, physiological properties, and soil enzymatic activities. *J Fungi*. 2021 Jul 17;7(7):571. <https://doi.org/10.3390/jof7070571>
23. Woolf D, Amonette JE, Street-Perrott FA, Lehmann J, Joseph S. Sustainable biochar to mitigate global climate change. *Nature Commun*. 2010;1(1):56. <https://doi.org/10.1038/ncomms1053>
24. Carter S, Shackley S, Sohi S, Suy TB, Haefele S. The impact of biochar application on soil properties and plant growth of pot grown lettuce (*Lactuca sativa*) and cabbage (*Brassica chinensis*). *Agronomy*. 2013;3(2):404-18. <https://doi.org/10.3390/agronomy3020404>
25. Hasnain M, Munir N, Abideen Z, Zulfiqar F, Koyro HW, El-Naggar A, Caçador I, Duarte B, Rinklebe J, Yong JW. Biochar-plant interaction and detoxification strategies under abiotic stresses for achieving agricultural resilience: A critical review. *Ecotoxicol Environ Saf*. 2023;249:114408. <https://doi.org/10.1016/j.ecoenv.2022.114408>
26. Heydari M, Hajinia S, Jafarian N, Karamian M, Mosa Z, Asgharzadeh S, Rezaei N, Guidi L, Valkó O, Prévosto B. Synergistic use of biochar and the plant growth-promoting rhizobacteria in mitigating drought stress on oak (*Quercus brantii* Lindl.) seedlings. *For Ecol Manage*. 2023 ;531:120793. <https://doi.org/10.1016/j.foreco.2023.120793>
27. Begum N, Qin C, Ahanger MA, Raza S, Khan MI, Ashraf M, Ahmed N, Zhang L. Role of arbuscular mycorrhizal fungi in plant growth regulation: implications in abiotic stress tolerance. *Front Plant Sci*. 2019;10:1068. <https://doi.org/10.3389/fpls.2019.01068>
28. Kim YJ, Hyun J, Yoo SY, Yoo G. The role of biochar in alleviating soil drought stress in urban roadside greenery. *Geoderma*. 2021;404:115223. <https://doi.org/10.1016/j.geoderma.2021.115223>
29. Parkash V, Singh S. Potential of biochar application to mitigate salinity stress in eggplant. *Hort Science*. 2020;55(12):1946-55. <https://doi.org/10.21273/HORTSCI15398-20>
30. Razzaghi F, Obour PB, Arthur E. Does biochar improve soil water retention? A systematic review and meta-analysis. *Geoderma*. 2020 Mar 1;361:114055. <https://doi.org/10.1016/j.geoderma.2019.114055>
31. Oni BA, Oziegbe O, Olawole OO. Significance of biochar application to the environment and economy. *Ann Agric Sci*. 2019;64(2):222-236. DOI: 10.1016/j.aoas.2019.08.003
32. Alkharabsheh HM, Seleiman MF, Battaglia ML, Shami A, Jalal RS, Alhammad BA, et al. Biochar and its broad impacts in soil quality and fertility, nutrient leaching and crop productivity: A review. *Agron*. 2021;11(5):993. <https://doi.org/10.3390/agronomy11050993>
33. Dahlawi S, Naeem A, Rengel Z, Naidu R. Biochar application for the remediation of salt-affected soils: Challenges and opportunities. *Sci Total Environ*. 2018;625:320-335. <https://doi.org/10.1016/j.scitotenv.2017.12.257>
34. Kul R, Arjumend T, Ekinci M, Yildirim E, Turan M, Argin S. Biochar as an organic soil conditioner for mitigating salinity stress in tomato. *Soil Sci Plant Nutr*. 2021;67(6):693-706. <https://doi.org/10.1080/00380768.2021.1998924>
35. Zulfiqar F, Chen J, Younis A, Abideen Z, Naveed M, Koyro HW, et al. Biochar, compost, and biochar-compost blend applications modulate growth, photosynthesis, osmolytes, and antioxidant system of medicinal plant *Alpinia zerumbet*. *Front Plant Sci*. 2021;12:707061. DOI: 10.3389/fpls.2021.707061
36. Khan Z, Zhang K, Khan MN, Fahad S, Xu Z, Hu L. Coupling of

- biochar with nitrogen supplements improve soil fertility, nitrogen utilization efficiency and rapeseed growth. *Agron.* 2020;10:1661. <https://doi.org/10.3390/agronomy10111661>
37. Xiang L, Harindintwali JD, Wang F, Redmile-Gordon M, Chang SX, Fu Y, et al. Integrating biochar, bacteria, and plants for sustainable remediation of soils contaminated with organic pollutants. *Environ Sci Technol.* 2022 Dec 6;56(23):16546-16566. <https://doi.org/10.1021/acs.est.2c02976>
 38. Oladipo A, Enwemiwe V, Ejeromedoghene O, Adebayo A, Ogunyemi O, Fu F. Production and functionalities of specialized metabolites from different organic sources. *Metabolites.* 2022;12(6):534. <https://doi.org/10.3390/metabo12060534>
 39. Saha A, Basak BB, Gajbhiye NA, Kalariya KA, Manivel P. Sustainable fertilization through co-application of biochar and chemical fertilizers improves yield, quality of *Andrographis paniculata* and soil health. *Ind Crops Prod.* 2019;140:111607. <https://doi.org/10.1016/j.indcrop.2019.111607>
 40. Jabborova D, Ma H, Bellingrath-Kimura SD, Wirth S. Impacts of biochar on basil (*Ocimum basilicum*) growth, root morphological traits, plant biochemical and physiological properties and soil enzymatic activities. *Sci Hort.* 2021;290:110518. <https://doi.org/10.1016/j.scienta.2021.110518>
 41. Jabborova D, Wirth S, Halwani M, Ibrahim MF, Azab IH, El-Mogy MM, Elkelish A. Growth response of ginger (*Zingiber officinale*), its physiological properties and soil enzyme activities after biochar application under greenhouse conditions. *Hortic.* 2021 ;7(8):250. <https://doi.org/10.3390/horticulturae7080250>
 42. Jabborova D, Annapurna K, Choudhary R, Bhowmik SN, Desouky SE, Selim S, Azab IH, Hamada MM, Nahhas NE, Elkelish A. Interactive impact of biochar and arbuscular mycorrhizal on root morphology, physiological properties of fenugreek (*Trigonella foenum-graecum* L.) and soil enzymatic activities. *Agron.* 2021;11(11):2341. <https://doi.org/10.3390/agronomy11112341>
 43. Liu A, Tian D, Xiang Y, Mo H. Biochar improved growth of an important medicinal plant (*Salvia miltiorrhiza* Bunge) and inhibited its cadmium uptake. *J Plant Biol Soil Health.* 2013;3(2):1-6.
 44. Jabborova D, Ziyadullaeva N, Enakiev Y, Narimanov A, Dave A, Sulaymanov K, et al. Growth of Spinach As Influenced By Biochar and *Bacillus Endophyticus* IGPEB 33 In Drought Condition. *Pak. J. Bot.* 2023;55(SI). [https://doi.org/10.30848/PJB2023-SI\(6\)](https://doi.org/10.30848/PJB2023-SI(6))
 45. Jabborova D, Kannepalli A, Azimov A, Tyagi S, Pengani KR, Sharma P, et al. Co-inoculation of Biochar and Arbuscular Mycorrhizae for Growth Promotion and Protein and Enzymes Fortification in Soybean Under Drought Conditions. *Front Plant Sci.* 2022 Jul 22;13:947547.
 46. Jabborova D, Annapurna K, Al-Sadi AM, Alharbi SA, Datta R, Zuan AT. Biochar and Arbuscular mycorrhizal fungi mediated enhanced drought tolerance in Okra (*Abelmoschus esculentus*) plant growth, root morphological traits and physiological properties. *Saudi J Biol Sci.* 2021 Oct;28(10):5490-9. <https://doi.org/10.1016/j.sjbs.2021.08.016>
 47. Wu Z, Fan Y, Qiu Y, Hao X, Li S, Kang S. Response of yield and quality of greenhouse tomatoes to water and salt stresses and biochar addition in Northwest China. *Agric Water Manage.* 2022;270:107736.
 48. Yang A, Akhtar SS, Li L, Fu Q, Li Q, Naeem MA, et al. Biochar mitigates combined effects of drought and salinity stress in quinoa. *Agronomy.* 2020;10(6):912. <https://doi.org/10.3390/agronomy10060912>
 49. Zhang Y, Ding J, Wang H, Su L, Zhao C. Biochar addition alleviate the negative effects of drought and salinity stress on soybean productivity and water use efficiency. *BMC Plant Biol.* 2020;20:1-11. <https://doi.org/10.1186/s12870-019-2232-5>
 50. Hussien Ibrahim ME, Adam Ali AY, Zhou G, Ibrahim Elsiddig AM, Zhu G, Ahmed Nimir NE, Ahmad I. Biochar application affects forage sorghum under salinity stress. *Chilean J Agric Res.* 2020;80(3):317-25.
 51. Kordi S, Saidi M, Ghanbari F. Induction of drought tolerance in sweet basil (*Ocimum basilicum* L.) by salicylic acid. *Int J Agric Food Res.* 2013;2(2):18-26.