



RESEARCH COMMUNICATION

# *Cheirostylis* Blume (Orchidaceae): a new generic addition to the orchid flora of Manipur state, Northeast India

Kazhuhrii Eshuo

Department of Botany, D.M. College of Science, Dhanamanjuri University, Imphal-795001, India

\*Email: [kazhuhrii@gmail.com](mailto:kazhuhrii@gmail.com)



## ARTICLE HISTORY

Received: 19 April 2023  
Accepted: 23 September 2023  
Available online  
Version 1.0 : 10 January 2024  
Version 2.0 : 24 January 2024



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care etc. See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Eshuo K. *Cheirostylis* Blume (Orchidaceae): a new generic addition to the orchid flora of Manipur state, Northeast India. *Plant Science Today*. 2024; 11(1): 564–567. <https://doi.org/10.14719/pst.2609>

## Abstract

The genus *Cheirostylis*, belonging to the Orchidaceae family, is represented by numerous species distributed worldwide. In India, the genus is represented by 10 species, with two species reported from Western Ghats and Eastern Ghats, and the remaining eight species from the Northwestern Himalayas and Northeastern regions. This study reports the occurrence of *Cheirostylis griffithii* Lindl. in Manipur, Northeast India, which is a new generic record for the state. The taxon was discovered in Pudunamei village and identified based on herbarium specimens and available literature. Detailed morphological descriptions and measurements were made using living plant specimens. The taxonomic treatment includes information on plant characteristics, distribution, and a comprehensive list of Indian *Cheirostylis* species. The species is a terrestrial orchid with fleshy stems, lanceolate bracts, pubescent flowers, connate sepals forming a tube, and a 3-lobed lip with deeply lacinate lobes. The occurrence of *Cheirostylis* in Manipur expands its distributional range within India. The study contributes to the documentation of the orchid flora of Manipur, which encompasses a diverse range of orchid taxa.

## Keywords

*Cheirostylis griffithii*; generic record; extended distribution; Manipur

## Introduction

*Cheirostylis* Blume, belonging to the flowering plants of the family Orchidaceae, tribe Cranichideae, and sub-tribe Goodyerinae, is represented by 60 species worldwide (1, 2). It is distributed mostly in tropical Africa through Southeast Asia, Japan, Indonesia, and Pacific Islands to Australia (3–5). *Cheirostylis* are characterized by a monoliform to terete fleshy stem, connate sepals forming a sapeline tube, a hypochile of the lip with a saccate base, a column with two deeply bifid stledia, and rostellum arms (3, 4). In India, the genus *Cheirostylis* is represented by 10 species (6, 7) namely: *Cheirostylis flabellata* Wight, *C. Griffithii* Lindl., *C. gunnarii* A.N.Rao, *C. Monoliformis* (Griff.) Seidenf., *C. Parvifolia* Lindl., *C. Pusilla* Lindl., *C. sessanica* A.N.Rao, *C. tabiyahanensis* (Hayata) N.Pearce & P.J.Cribb, *C. tippica* A.N.Rao and *C. yunnanensis* Rolfe. Out of the 10 species, 2 species namely *C. flabellata* and *C. parvifolia* have been reported from the Western Ghats (8–10), Eastern Ghats (11), while the other 8 species are reported from the Northwestern Himalayas and Northeast region of India (4, 6, 7, 11–13). Three species, namely *C. gunnarii*, *C. sessanica* and *C. tippica*, are endemic to India and are reported from Arunachal Pradesh (7).


In Manipur, the orchid flora is represented by 389 taxa (including 5 varieties) belonging to 90 genera (14). The orchid flora of the state has been explored by several plant collectors and botanists (15–23). The orchid flora of Manipur state is well documented, as many orchid lovers and taxonomists have explored it extensively. However, orchid literature in the Mao areas of Senapati district, Manipur, is limited, with only fragmentary works available (24–26). During a field visit at Pudunamei village, Mao Senapati district, Manipur, the author encountered an interesting species of terrestrial orchid. Terrestrial orchid specimens were collected, and upon further investigation and consultation with the herbarium and available literature, the collected species has been identified as *Cheirostylis griffithii* Lindl., a previously unknown species in Manipur. The occurrence of this taxon is a new generic record for the state and has an extended distributional range to Manipur, Northeast India. The total number of Indian *Cheirostylis* species and their distribution are listed in Table 1.

### Materials and Methods

The terrestrial orchid plants were collected, pressed, and prepared as herbarium specimens following conventional herbarium techniques (27). The herbarium specimen was deposited at the Herbarium, Botany Department of D.M. College of Science, Imphal, and at the Herbarium Botanical Survey of India, ERC, Shillong (ASSAM), for future reference. Photos of the live plants were captured using Sony SLT-A58 and Canon SX120 digital cameras, respectively. All illustrations, morphological descriptions, and measurements were based on living plant specimens.

### Taxonomic Treatment


***Cheirostylis griffithii* Lindl.**, J. proc. Linn. Soc., Bot. 1: 188. 1857. (Fig. 1 & Fig. 2)



**Fig. 1.** *Cheirostylis griffithii* Lindl. A. Plant habit with a single flower; B. Plant habit with three flowers; C. Sepal dorsal view; D. Labellum; E. Pollinia.

**Table 1.** List of Indian *Cheirostylis* species and their distribution (7).

S. No.	Name	Distribution
1	<i>Cheirostylis flabellata</i> Wight	Karnataka, Kerala, Maharashtra & Tamil Nadu.
2	<i>C. griffithii</i> Lindl.	Arunachal Pradesh, Meghalaya, Mizoram, Nagaland, Sikkim, Uttarakhand, West Bengal, Manipur [present report].
3	<i>C. gunnarii</i> A.N.Rao	Arunachal Pradesh (Endemic to India).
4	<i>C. monoliformis</i> (Griff.) Seidenf.	Arunachal Pradesh, Sikkim & West Bengal.
5	<i>C. parvifolia</i> Lindl.	Kerala, Maharashtra, Tamil Nadu & Odisha.
6	<i>C. pusilla</i> Lindl.	Arunachal Pradesh, Meghalaya & Nagaland.
7	<i>C. sessanica</i> A.N.Rao	Arunachal Pradesh (Endemic to India).
8	<i>C. tabiyahanensis</i> (Hayata) N.Pearce & P.J.Cribb	Nagaland.
9	<i>C. tippica</i> A.N.Rao	Arunachal Pradesh (Endemic to India).
10	<i>C. yunnanensis</i> Rolfe	Arunachal Pradesh, Sikkim & West Bengal.



**Fig. 2.** *Cheirostylis griffithii* Lindl. A. Habit; B. Sepal dorsal view; C. Column with ovary; D. Labellum; E-F. Petals; G. Anther cap; H. Pollinia.

Plants 15–30 cm tall; rhizome creeping, terete to torose; roots hair like at the rhizome internode. Stem erect, fleshy, with withered leaves at anthesis. Inflorescence ca. 15 cm long, 2–3 sterile bracts, rachis 1–2 cm, with 1–3 flowers in terminal raceme, bracts lanceolate, 0.6–0.9 cm, flower whitish, large pubescent. Sepals 1–1.5 cm, lower portion connate in  $\frac{1}{2}$  forming a tube, upper portion spreading, surface pubescent, apex acute. Petals are oblong, oblique, falcate, with obtuse to round apex. The lip or labellum is 3-lobed with deeply lacinate lobes having 9–11 divisions. The column measures 0.2–0.3 cm and has straight, linear stelia, rostellum arms, and an anther with two sectile pollinia. The pollinia are attached with a tegula that has viscidia at the distal end.

### Flowering

February–March.

### Fruit

Not seen.

### Distribution

INDIA: (Arunachal Pradesh, Meghalaya, Mizoram, Nagaland, Sikkim, Uttarakhand, West Bengal, Manipur [present report]).

### Specimen examined

India, Manipur, Pudunamei-Mao, 1650–1750 m, 25°31'77.40"N and 94°09'10.18"E, KE100005.

### Ecology

The orchid plants were found growing in the deciduous forest at an altitude ranging between 1650–1750 m.

## Discussion

The orchid flora of Manipur is well documented, with several researchers having studied the orchids of Manipur state. The most recent work, the “Updated Checklist of Orchid Flora of Manipur” carried out (14), documented 389 taxa belonging to 90 genera. They classified the orchid genera of Manipur into four subfamilies, thirteen tribes, and fourteen sub-tribes based on the classification of Orchidaceae (28). According to this classification, the genus *Cheirostylis* Blume falls under the sub-family Orchidoideae, tribe Cranichideae, and sub-tribe Goodyerinae. Within the sub-tribe Goodyerinae, six genera, namely *Anoectochilus* Blume, *Erythrodes* Blume, *Goodyera* R. Br., *Myrmechis* Blume, *Odontochilus* Blume and *Zeuxine* Lindl. were reported from Manipur state. The occurrence of the genus *Cheirostylis* Blume has not been reported by any workers from Manipur state so far. Therefore, the occurrence of a taxon *Cheirostylis* represents a new generic record for the state and extends its distributional range to Manipur.

## Acknowledgements

The author is grateful to N. Eshuo Mao and Adahe Pfoziro for their help during field work. The author is also thankful to Dr. Santanu Dey and Lovikali V Kiho for providing relevant literature. The facility rendered by D.M. College of Science, Dhanamanjuri University, Imphal is duly acknowledged.

## Compliance with ethical standards

**Conflict of interest :** Author does not have any conflict of interest to declare.

**Ethical issues:** None.

## References

- Govaerts R, Pfahl J, Campacci MA, Baptista DH, Tigges H, Shaw J, Cribb P, George A, Kreuz K, Wood J. World Checklist of Orchidaceae. The Board of Trustees of the Royal Botanic Gardens, Kew; 2018. Published on the Internet; <http://www.kew.org/wcsp/> (Accessed on 23<sup>rd</sup> January, 2023)
- POWO. “Plants of the world Online. Facilitated by the Royal Botanic Gardens, Kew. Published on the internet; <http://www.plantsoftheworldonline.org/> Accessed on 23<sup>rd</sup> January 2023.
- Pridgeon AM, Cribb PJ, Chase MW, Rasmussen FN. Genera Orchidacearum 3, Orchidoideae (part 2), Vanilloideae. Oxford University Press; 2003.
- Bhattacharjee A. On the status of some species of *Cheirostylis* Blume (Orchidaceae) from India. *Candollea*. 2013;67(1):31-35. <https://doi.org/10.15553/c2012v671a3>
- Chen S, Gale SW, Cribb PJ, Ormerod P. *Cheirostylis* Blume. In: Wu ZY, Raven PH, Hong DY, editors. *Flora of China*. Science Press and Missouri Botanical Garden Press. 2009; p. 57-63.
- Odyuo N, Roy DK, Deori C, Daimary R. *Cheirostylis tabiyahanensis* (Hayata) N. Pearce & P.J. Cribb (Orchidaceae). A new addition to the Orchid flora of India. *The MIOS Journal*. 2018;19(10):2-6.

7. BSI. "Cheirostylis" on <https://efloralindia.gov.in>. Botanical Survey of India, Kolkata. (Accessed on 23<sup>rd</sup> January, 2023)
8. Puneekar S. Report of rare terrestrial orchid species *Cheirostylisparvifolia* Lindl. from Dajipur Wildlife Sanctuary, Kohlapur District of Maharashtra State, India. Journal of Economic and Taxonomic Botany. 2002;26(1):105-07.
9. Shigwan BK, Kulkarni A. Extended distribution of rare orchid *Cheirostylisparvifolia* from Northern Western Ghats of Maharashtra, India. Plantasia #5, in: Zoo's Print. 2019;34(11):49-51.
10. Dash PK, Bhattacharjee A, Kumar P, Mohapatra P. *Cheirostylis* Blume (Orchadaceae), a generic record for the Eastern Ghats, India. Journal of Threatened Taxa. 2015;7(1):6826-6829. <http://dx.doi.org/10.11609/JoTT.o4139.6826-9>
11. Rao NA. Two new species of *Cheirostylis* (Orchidaceae) from Arunachal Pradesh India. Nordic Journal of Botany. 1988;7:339-40.
12. Rao NA. *Cheirostylis pinnata* Nagaswana Rao A New species of Orchid from Arunachal Pradesh, India. Arunachal Forest News. 1991;9(2):17-20. <https://doi.org/10.1111/j.1756-1051.1988.tb00508.x>
13. Rao NA. *Cheirostylis gunnarii* (Orchidaceae)—a new orchid from Arunachal Pradesh, India. Nordic Journal of Botany. 2008;18(1):23-25. <https://doi.org/10.1111/j.1756-1051.1998.tb001092.x>.
14. Rao AN, Kumar V. Updated checklist of orchid flora of Manipur. *Turczaninowia*. 2018;21(4):109-34. DOI: 10.14258/turczaninowia.21.4.12
15. Watt G. The forests of Manipur. Ind For. 1888;14:291-99; 338-44; 387-94.
16. Clarke CB. On the plants of Kohima and Muneypore. The Journal of Linnean Society. 1889;25:1-74. <https://doi.org/10.1111/j.1095-8339.1889.tb00793.x>
17. Deb DB. Monocotyledonous plants of Manipur Territory. The Bulletin of the Botanical Survey of India. 1961;3:126-29.
18. Mukerjee SK. An enumeration of the orchids of Ukhrul, Manipur. Notes from the Royal Botanic Garden, Edinburgh. 1953;21:149-54.
19. Ghatak J, Devi RK J. Orchids of Manipur. In: Vij SP, editor. Biology, Conservation and Culture of Orchids. Affiliated East West Press Pvt. Ltd., New Delhi. 1986;p. 357-62.
20. Chauhan AS. A conspectus of orchids of Manipur: Their status and conservation. In: Pathak P, Sehgal RN, Shekar N, Sharma M, Sood A, editors. Orchids Science and Commerce. Bishen Singh mahendra Pal Singh, Dehra Dun. 2001;p. 81-99.
21. Kumar CS, Kumar PCS. An Orchid Digest of Manipur, Northeastern India. Rheedeia. 2005;15(1):1-70.
22. Rao AN. Orchid flora of North East India- an upto date analysis. Bulletin of Arunachal Forest Research. 2007;23(1-2):6-38.
23. Rao AN, Chowlu K, Kumar V, Sharma HB, Devi YN. Orchids of North East India. Vol. 1. FEEDS, Hengbung, Manipur, India; 2016.
24. Mao AA. Notes on the orchids of Senapati and surrounding hills in the state of Manipur, India. The Journal of the Orchid Society of India. 1999;13(1-2):55-58.
25. Phukan SJ, Mao AA. *Armadorumsenapatianum*—a new species from India. Orchid Review. 2002;110:298-300.
26. Phukan SJ, Mao AA. Additions to the Indian Orchid Flora. Orchid Review. 2004;112:115- 18.
27. Jain SK, Rao RR. A handbook of field and herbarium methods. Today & Tomorrow's Printers and Publishers, New Delhi; 1976.
28. Chase MW, Cameron KM, Freudenstein JV, Pridgeon AM, Salazar G, Berg CVD, Schuiteman A. An updated classification of Orchidaceae. Botanical Journal of the Linnean Society. 2015;177:151-74. <https://doi.org/10.1111/boj.12234>