



RESEARCH ARTICLE

# Effect of phosphate fertilizer-coated Dicarboxylic Acid Polymer on rice yield and components under greenhouse conditions

Tran Van Hung<sup>1</sup>, Ngo Ngoc Hung<sup>2</sup> & Vo Quang Minh<sup>1\*</sup>

<sup>1</sup>Department of Land Resources, College of Environment and Natural Resources, Can Tho University, Can Tho City, 90000, Viet Nam

<sup>2</sup>Department of Soil Science, College of Agricultural, Cantho University, Can Tho City, 90000, Viet Nam

\*Email: [vqminh@ctu.edu.vn](mailto:vqminh@ctu.edu.vn)



## ARTICLE HISTORY

Received: 18 April 2023  
Accepted: 07 December 2023  
Available online  
Version 1.0 : 28 January 2024



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care etc. See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Hung TV, Hung NN, Minh VQ. Effect of phosphate fertilizer-coated Dicarboxylic Acid Polymer on rice yield and components under greenhouse conditions. *Plant Science Today* (Early Access). <https://doi.org/10.14719/pst.2610>

## Abstract

A significant amount of phosphorus (P) becomes fixed by aluminium (Al) and iron (Fe) in acidic soils, leading to decreased efficiency in P utilization and subsequently lowering crop yield. Enhanced P fertilization offers a potential solution, as the dicarboxylic acid polymer (DCAP) coating on P fertilizer promotes increased plant productivity and more effective P utilization. The improvement achieved through enhanced P fertilization can contribute to higher rice yields in acidic soils, accompanied by an increase in P solubility. The study aimed to determine the impact of DCAP-mixed phosphate fertilizer on P uptake by plants, absorption efficiency, and rice yield. The results demonstrated a significant increase in available P (about 3.5 mg P/kg) when DCAP was used in a greenhouse setting, resulting in elevated yields and total P absorption (ranging from 0.03 to 0.05 grams/pot). However, the addition of 60 kg of phosphate mixed with DCAP has not yet demonstrated a significant increase in available phosphorus in the soil compared to adding just 60 kg of phosphate. The application of phosphate at a dose of 30 kg of P<sub>2</sub>O<sub>5</sub> mixed with DCAP for growth and phosphorus absorption yield results equivalent to using 60 kg of P<sub>2</sub>O<sub>5</sub> without DCAP. Furthermore, the use of DCAP in conjunction with 50% P fertilizer increased P availability by the same amount as that achieved with 100% P fertilizer. Consequently, DCAP reduced chemical P fertilizer in the soil by approximately 50%. However, it is essential to evaluate the effectiveness of mixed phosphate fertilizer (DCAP) under field conditions before recommending its widespread use.

## Keywords

Dicarboxylic Acid Polymer (DCAP); available phosphorus; rice; phosphate fertilizer; phosphorus absorption

## Introduction

One of the nutrients that globally influences agricultural output is phosphorus (P) (1–3). While tropical and subtropical soils can naturally contain high total P content, reaching values as high as 1800 mg kg<sup>-1</sup>(4), the strong P adsorption of iron (Fe) and aluminium (Al) oxides, along with 1:1 clay minerals, means that only a minimal amount of this P is readily available for plants (5). In these soils, phosphate fertilizers becomes the primary means to meet the crop's P requirements (6). According to Pal and Adhikary (7), phosphorus is a secondary requirement that often poses the most significant limitation for crops and forage output after nitrogen.

The primary function of phosphorus in plants is to store and transport the energy generated by photosynthesis for use in growth and reproduction.

Phosphorus in the soil exists in forms such as phosphates ( $\text{PO}_4^{3-}$ ), mono- and di-hydrogen phosphates ( $\text{HPO}_4^{2-}$ ), and  $\text{H}_2\text{PO}_4^-$ . The predominance of these anions depends on the pH of the solution or soil, which controls their eager interconversion. Plants most readily absorb phosphorus when the pH is between 6.5 and 7.5. Due to its high mobility, phosphorus can move from older, actively growing plant tissue to younger, more actively growing parts when it is insufficient. Consequently, early vegetative responses to phosphorus deficiency are frequently observed.

In acid sulphate soil, the efficiency of phosphorus uptake by plants can be limited, with only about 5-25% of the applied fertilizer being utilized. This is due to the high iron and aluminium content in acid sulphate soil, where phosphorus becomes fixed into insoluble compounds that are not easily dissolved by plants, making absorption more challenging (8). Consequently, approximately 75% to 95% of the phosphorus applied to the soil remains unavailable for crops. Moreover, this unutilized phosphorus, located near or on the soil's surface, can significantly impact the environment through erosion and leaching into rivers and lakes (9). It's noteworthy that phosphorus is most available to the plant within a few days to two weeks after fertilizer addition, gradually decreasing over time (20). Therefore, enhancing the efficiency of phosphorus uptake by plants not only improves economic efficiency but also reduces environmental pollution. Recent research and development have explored the additive effect of "Dicarboxylic Acid Polymer" (DCAP) in phosphate fertilizer. DCAP aims to protect phosphorus particles from immobilization by iron and aluminium in low pH conditions and by calcium and magnesium in high soil pH conditions. Results indicate that DCAP combined with phosphorus application increases the efficiency of phosphorus uptake and the yield of some crops (10–12). However, findings from Curtis research (13) on phosphate fertilizer application at a dose of 67 kg  $\text{P}_2\text{O}_5$ /ha mixed with DCAP in five experimental potato growing sites revealed that four sites exhibited higher yields.

In contrast, one site yielded less than the control group under an unmixed condition. This can be explained by the mixing of phosphorus with DCAP, which increases the available soil phosphorus content while reducing certain trace elements such as copper, zinc, and manganese in the soil, subsequently leading to a decrease in crop yield (14–17). Therefore, the effective use of DCAP depends on factors such as the amount of phosphate fertilizer applied, the response of each soil, and the specific types of crops being cultivated. The active ingredient "dicarboxylic acid polymer" is utilized to enhance the efficiency of phosphate fertilizer in various locations. However, the response to this active ingredient varies across different soil areas and crop types. In the Mekong Delta, acid sulfate soils cover approximately 1.6 million hectares, and while the phosphorus content in these soils is high, the efficiency of phosphorus use remains low. Hence, there is a pressing need to improve the efficiency of phosphate fertilizer use in acidic soils. DCAP, as an active ingredient, is known to increase the efficiency of

phosphate fertilizer use and rice productivity. Consequently, this study aims to assess the response of mixed phosphate fertilizer (DCAP) in a greenhouse setting. This preliminary evaluation is crucial, providing a foundation for conducting extended trials in the acidic soil regions of the Mekong Delta before recommending its widespread use.

## Materials and Methods

### Material

**Place and time of performance:** The experiment took place from February 2020 to May 2021 in the net house located at the Soil Science Department, College of Agriculture, Can Tho University, Can Tho City, Vietnam. (Latitude: 10° 01'43.30"N, Longitude: 105°45'59.71"E)

**Experimental soil** was sourced from the 0–20 cm layer in Hoa An commune, Phung Hiep district, Hau Giang province. The collected soil underwent a thorough process, including drying, mixing to ensure homogeneity, and removal of any debris. Subsequently, the soil was subjected to comprehensive analysis, covering key criteria such as pH, electric conductivity (EC), available phosphorus (P), free iron (Fe), and exchangeable aluminium (Al).

**Fertilizers:** urea (46% N), DAP (18% N–46%  $\text{P}_2\text{O}_5$ ), and KCl (60%  $\text{K}_2\text{O}$ ).

**Experimental pots:** For the experiment, pots with a height of 35cm, width of 40cm, and a soil capacity of 15 kg soil were utilized. Each pot was planted with five germinating rice seeds.

**Rice variety and its characteristics:** The rice variety chosen for the experiment was OM2517. This variety typically has a growth cycle of approximately 90–95 days, reaching a height of about 90–100 cm and displaying robust tillering ability. The number of panicles or spikelets ranges from 9–12, while the grain length measures between 7.0–7.3 mm. A thousand seeds weigh around 26–28 g, with an amylase content of about 24–25%. The chalkiness of rice, denoting the silver belly of rice grain, falls within grades 1 to 5. For this variety, the expected yields are 5 tons/ha for the summer-autumn crop and 8 tons/ha for the winter-spring crop.

### Method

#### Experimental layout

The experiment was arranged in a Completely Randomized Design (CRD), comprising 5 treatments, each with six replicates, where each replicate corresponds to an individual pot. The experimental treatments are shown in Table 1.

#### Soil and plant analysis

Soil analysis was conducted at the Soil Physics - Chemistry Analysis Laboratory, situated within the Soil Science Department, College of Agriculture, Can Tho University. The specific methods employed for soil analysis are mentioned in Table 2a, while the methods for crop analysis are in Table 2b.

**Table 1.** The treatments of the experiment

No	Experiment	Describe
1	0 P <sub>2</sub> O <sub>5</sub>	No phosphate fertilizer (control)
2	30 P <sub>2</sub> O <sub>5</sub>	Apply 100% phosphate as recommended with DAP
3	30 P <sub>2</sub> O <sub>5</sub> + DCAP	Apply 100% phosphate as recommended with DAP fertilizer mixed with DCAP
4	60 P <sub>2</sub> O <sub>5</sub>	Apply 100% phosphate as recommended with DAP
5	60 P <sub>2</sub> O <sub>5</sub> + DCAP	Apply 100% phosphate as recommended with DAP fertilizer mixed with DCAP

**Note:** Use 2 litres of mixed DCAP solution for one ton of DAP fertilizer to achieve a concentration of 2‰ (as recommended by the manufacturer); DCAP: Dicarboxylic Acid Polymer

**Table 2a.** Method of soil analysis at the beginning of the crop

Parameters	Unit	Method
pH <sub>H2O</sub>		Extraction with distilled water, ratio 1:2.5 (soil/water), measured by pH meter.
EC	mS/cm	Extraction with distilled water, ratio 1:2.5 (soil/water), measured by EC meter.
Avai P	mg P/kg	Bray II method: soil extraction with HCl 0.1N + NH <sub>4</sub> F0.03N, ratio 1:7 (soil: extract solution), then measured by colorimetric method on a spectrophotometer at wavelength 880 nm.
Fe <sup>3+</sup>	%Fe <sub>2</sub> O <sub>3</sub>	extraction with oxalate - oxalic acid, Fe determination on atomic absorber
Al <sup>3+</sup>	meq/100g	Extract with 1N KCl, titrate with 0.01N NaOH, complex with NaF and titrate with 0.01N H <sub>2</sub> SO <sub>4</sub>

**Table 2a.** Method of soil analysis at the beginning of the crop

Parameters	Unit	Destruction of samples	Method*
P total	%P	6g salicylic acid + 18ml demineralized water + 100ml H <sub>2</sub> SO <sub>4</sub> 96%, H <sub>2</sub> O <sub>2</sub> used for oxidation	Colorimeter on a spectrophotometer

**Notes:** Walsh and Beaton (1973)

### Time and dose of fertilizer

The fertilizer formula for rice consists of 100 N–30 K<sub>2</sub>O (kg/ha). The application schedule and corresponding fertilizer doses are planned at 10, 20, and 40 days after seeding (DAS) (Table 3).

**Table 3.** Period and dose of fertilizer for the experiment

Fertilization period	Amount of fertilizers (%)		
	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
1st application (10 DAS)	30	50	50
2nd application (20 DAS)	40	50	0
3rd application (40 DAS)	30	0	50

**DAS:** days after sowing

### Data collection and analysis

Soil samples were collected for available P analysis at both the beginning and ending of the cropping season.

The data on height and the number of rice shoots were collected at specific intervals: 20, 45, 65, and 90 days after sowing (DAS).

Data collection included yield components such as the number of flowers per pot, the percentage of firm seeds (%), the weight of a thousand seeds (at 14% moisture content), and the number of seeds per panicle. The grain yield/pot at 14% moisture was determined by harvesting the entire rice plant from each pots. The yield was then calculated in grams.

For the straw and seed sampling method, 5 plants were randomly chosen from each treatment. The biomass of straw and grains was measured by weighing them in the pots at the harvest stage and subsequently drying them at 70 °C for 72 hours.

Absorbance = biomass (straw and seeds) x the content of each part.

The analysis of variance (ANOVA) was done using SPSS software version 16.0 to compare the mean differences, using the Duncan test.

## Results

### Initial soil properties of the experiment


The initial soil characteristics at a depth of 0–20cm are presented in Table 4. The soil was initially in a state of perennial rice cultivation. The impact of soil chemical parameters on plant growth was assessed using the rating scale developed by Có dấu giữa 2 tác giả (19). The soil selected for the experiment was identified as acid sulfate soil (Epi-Orthi Thionic Fluvisols), characterized by a low pH (3.8) and high levels of exchangeable aluminium (Table 4). The prevalence of high iron and aluminium content in the soil led to phosphorus fixation, resulting in low soil phosphorus content (20).

**Table 4.** Initial experimental soil properties

Targets	Value	Evaluation for plant growth
pH <sub>H2O</sub>	3.82 ± 0.05	Low, slightly acid
EC <sub>1:2.5</sub> (mS/cm)	4.52 ± 0.08	High
Available P (mg/kg)	9.63 ± 0.14	Relatively low
Exchange Al (meqAl <sup>3+</sup> /100g)	13.3 ± 0.11	Pretty high
Free Fe (%Fe <sub>2</sub> O <sub>3</sub> )	0.95 ± 0.01	Medium

### The concentration of available soil phosphorus after mixing phosphorus with DCAP

The addition of 30 kg of P<sub>2</sub>O<sub>5</sub> mixed with DCAP to the soil resulted in an increase in available phosphorus compared to the control (Figure 1). The peak available phosphorus content was observed four days after application. Interestingly, when 60 kg of P<sub>2</sub>O<sub>5</sub> mixed with DCAP was introduced to the soil, there was no additional increase in available phosphorus, similar to the case with 30 kg of P<sub>2</sub>O<sub>5</sub> mixed with DCAP.



**Fig 1.** The content of available phosphorus doses

The absence of added phosphorus in fertilizers leads to a decrease in soil phosphorus content due to the uptake by plants during their growth and development. Application of mixed phosphate fertilizer DCAP plays a crucial role in reducing the fixation of  $\text{Fe}^{2+}$ ,  $\text{Fe}^{3+}$ , and  $\text{Al}^{3+}$  ions with  $\text{HPO}_4^{2-}$  and  $\text{H}_2\text{PO}_4^-$ , consequently elevating the available phosphorus content in the soil. This observation aligns with findings from Noble's study (21), where the application of mixed phosphate fertilizer DCAP was associated with an increase in the soil's available phosphorus content.

#### **Effects of mixed phosphate fertilizer DCAP on growth parameters of rice growing under Greenhouse Conditions**

**Height of rice plants:** The height of rice plants in treatments without any phosphate application was significantly shorter at 45, 65 and 90 DAS compared to those receiving 30  $\text{P}_2\text{O}_5$  + DCAP, 60  $\text{P}_2\text{O}_5$ , and 60  $\text{P}_2\text{O}_5$  + DCAP. This difference was found to be statistically significant at a 5% level of significance. Specifically, the application of 30 kg of  $\text{P}_2\text{O}_5$  mixed with DCAP resulted in a precise height of rice, while 60 kg of  $\text{P}_2\text{O}_5$  was applied without DCAP at the 45 and 90 DAS stages (Table 5).

Given that phosphorus plays a crucial role in photosynthesis, respiration, energy storage and transfer, cell division, and various other plant processes, including growth promotion and enhancement of water use efficiency (22), the absence of phosphorus application led to a reduction in the height of rice plants compared to those receiving other treatments. Notably, the soil

**Table 5.** Effect of DCAP phosphate fertilizer on rice plant height

Experiment	Rice plant height (cm)			
	20 DAS	45 DAS	65 DAS	90 DAS
0 $\text{P}_2\text{O}_5$	34.9	53.1 b	76.8 b	80.7 b
30 $\text{P}_2\text{O}_5$	34.8	55.3 b	82.4 a	85.1 b
30 $\text{P}_2\text{O}_5$ + DCAP	35.8	60.2 a	82.0 a	94.6 a
60 $\text{P}_2\text{O}_5$	33.7	59.7 a	82.3 a	92.2 a
60 $\text{P}_2\text{O}_5$ + DCAP	34.1	61.5 a	81.9 a	94.4 a
F	Ns	*	*	*
CV(%)	5.04	6.75	7.10	5.24

**Note:** In the same column, numbers followed by different letters have statistical significance at 5% (\*); ns: no statistically significant difference

treatment involving 30 kg  $\text{P}_2\text{O}_5$ /ha for rice cultivation showed reduces height at stages 45 and 90 DAS. This reduction might be attributed to the fixation of the applied phosphorus applied (13), preventing the rice plants from utilizing it fully or partially.

**The number of rice shoots:** Table 6 indicates no significant difference in the number of rice shoots among treatments at the 20 and 45 DAS stages. However, variations emerged at the 65 DAS stage. Specifically, treatments without phosphate and 30 kg of  $\text{P}_2\text{O}_5$  without DCAP exhibited the lowest number of rice shoots compared to the other treatments at the 65 DAS stage. Notably, the number of rice shoots in the treatment without phosphate significantly decreased from 45 to 65 DAS. This decline could be attributed to the plant's need for sufficient phosphorus to maintain the number of flowers at this stage. These findings align with the results of Khuong (23), which demonstrated an increase in rice shoots in soils mixed with DCAP.

**Table 6.** Effect of DCAP mixed phosphate fertilizer on the rice shoot numbers

Experiment	Rice shoot's number per pot		
	20 DAS	45 DAS	65 DAS
0 $\text{P}_2\text{O}_5$	10.0	25.7	20.8 b
30 $\text{P}_2\text{O}_5$	11.8	24.3	22.1 b
30 $\text{P}_2\text{O}_5$ + DCAP	11.0	27.0	26.2 a
60 $\text{P}_2\text{O}_5$	11.5	25.9	26.4 a
60 $\text{P}_2\text{O}_5$ + DCAP	10.1	26.8	26.6 a
F	Ns	ns	*
CV(%)	4.12	8.17	5.89

**Note:** In the same column, numbers followed by different letters have statistical significance at 5% (\*); ns: no statistically significant difference

#### **Effect of mixed phosphate fertilizer DCAP on rice yield components and yield under greenhouse conditions**

The application of phosphate at a dosage of 30 kg  $\text{P}_2\text{O}_5$  mixed with DCAP showed that the number of panicles per pot, the ratio of firm seeds, and yield (grams/pot) were comparable to the treatment of 60 kg  $\text{P}_2\text{O}_5$  and 60 kg  $\text{P}_2\text{O}_5$  mixed with DCAP. However, a significantly higher value was observed in the treatment without any additional phosphate and in the treatment with 30 kg of  $\text{P}_2\text{O}_5$  without DCAP (Table 7).



**Table 7.** Effect of DCAP phosphate fertilizer on rice yield components and yield

Experiment	Number of panicles/pot	Number of seeds per panicle	Percentage of firm seeds (%)	Weight 1,000 seeds (grams)	Yield (gram/pot)
0 P <sub>2</sub> O <sub>5</sub>	19.5 b	91.0	75.7 b	26.6	38.1 b
30 P <sub>2</sub> O <sub>5</sub>	20.8 b	95.5	79.9 b	27.0	40.6 b
30 P <sub>2</sub> O <sub>5</sub> + DCAP	25.1 a	93.5	84.4 a	26.9	56.6 a
60 P <sub>2</sub> O <sub>5</sub>	26.1 a	95.7	84.5 a	26.8	55.7 a
60 P <sub>2</sub> O <sub>5</sub> + DCAP	24.3 a	97.9	83.5 a	26.9	54.4 a
F	**	Ns	*	ns	**
CV(%)	5.12	5.27	4.06	3.24	6.22

**Note:** In the same column, numbers followed by different letters have statistical significance at 1% (\*\*) and 5% (\*); ns: no statistically significant difference

Interestingly, the application of 60kg of P<sub>2</sub>O<sub>5</sub> mixed with DCAP showed no difference in yield and yield components (grams/pot) compared to the treatment with 60kg of P<sub>2</sub>O<sub>5</sub> alone. This suggests that rice plants may only require approximately 60kg of P<sub>2</sub>O<sub>5</sub>/ha for optimal growth and development, and the addition of DCAP at this dose does not increase phosphorus uptake efficiency.

Contrary to this research, research indicates that while high doses of phosphate application combined with DCAP did not result in a difference in rice yield compared to treatments without DCAP (11), the combination of low doses of phosphate with DCAP increased the phosphorus uptake efficiency and boosted yields in certain crops (12). According to the research results of Dang (24), phosphate fertilizer application at a dose of 30 kg P<sub>2</sub>O<sub>5</sub>/ha mixed with DCAP produced the yield of sweet potato and cassava tubers equivalent to those obtained with 60kg P<sub>2</sub>O<sub>5</sub>/ha without DCAP, a notable finding considering the potential reduction in yield when applying only 30kg P<sub>2</sub>O<sub>5</sub>/ha to the soil.


#### Effect of mixed phosphate fertilizer DCAP on phosphorus content and absorption by rice

No statistically significant differences were observed between phosphorus fertilization treatments in straw and seed phosphorus content. However, the total phosphorus absorption among the treatments was statistically different at the 5% significance level, potentially attributed to variations in the biomass of straw and grain (Table 8). The absence of phosphate application resulted in the lowest phosphorus absorption. The phosphorus absorption in treatments involving 30kg P<sub>2</sub>O<sub>5</sub>/ha mixed with DCAP was equivalent to that in the treatment with 60 kg of P<sub>2</sub>O<sub>5</sub>/ha without DCAP. These findings align with Wiatrak (25), where the application of phosphate at a dose of 45 kg P<sub>2</sub>O<sub>5</sub>/ha mixed with DCAP increased phosphorus uptake and wheat yield, comparable to the treatment with 90 kg P<sub>2</sub>O<sub>5</sub>/ha, which also enhanced phosphorus uptake and grain yield. The phosphorus content in seeds ranged

from 0.20-0.24%, while in straw, it ranged from 0.10- 0.12%. These findings are consistent with Fairhurst (26), which stated that the average P content in seeds was around 0.20%, and in straw, it was 0.10%.

#### Effect of phosphate fertilizer mixed with DCAP on the content of available soil phosphorus at the end of the crop.

The application of phosphorus mixed with DCAP at 30kg P<sub>2</sub>O<sub>5</sub>/ha resulted in soil available phosphorus content equivalent to that achieved by applying 60kg P<sub>2</sub>O<sub>5</sub>/ha by the end of the season, as illustrated in Figure 2. Treatments without phosphate and with phosphate applied at a dose of 30kg P<sub>2</sub>O<sub>5</sub>/ha exhibited low soil phosphate content at the end of the crop, significantly differing at the 5% significance level compared to other treatments. Phosphate application at the dose of 60kg P<sub>2</sub>O<sub>5</sub>/ha mixed with DCAP did not demonstrate a difference in available phosphorus content in the soil compared to the application of 60kg P<sub>2</sub>O<sub>5</sub>/ha alone. Despite the increase in available phosphorus content resulting from the application of DCAP mixed phosphate fertilizer, the levels remained low, measuring less than 20mg P/kg (18).

**Fig 2.** Effect of mixed phosphate fertilizer DCAP on the content of available soil phosphorus at the end of the crop**Table 8.** Effect of mixed phosphate fertilizer DCAP on phosphorus content and phosphorus absorption of rice

Experiment	Biomass (gram/pot)		Phosphate content (%P)		Total P absorption (grams/pot)
	Seed	Straw	Seed	Straw	
0 P <sub>2</sub> O <sub>5</sub>	27.2 b	24.1 b	0.20	0.11	0.08 c
30 P <sub>2</sub> O <sub>5</sub>	29.0 b	28.5 b	0.22	0.12	0.10 bc
30 P <sub>2</sub> O <sub>5</sub> + DCAP	40.4 a	39.9 a	0.22	0.11	0.13 ab
60 P <sub>2</sub> O <sub>5</sub>	39.7 a	42.0 a	0.24	0.12	0.15 a
60 P <sub>2</sub> O <sub>5</sub> + DCAP	39.8 a	40.3 a	0.23	0.10	0.13 ab
F	*	*	ns	ns	*
CV(%)	5.11	4.32	8.19	7.10	4.15

**Note:** In the same column, numbers followed by different letters have statistical significance at 1% (\*\*) and 5% (\*); ns: no statistically significant difference

## Conclusion

In summary, the experiments demonstrated that the utilization of DCAP-coated P fertilizer significantly increased the concentration of available phosphorus (~22%) in the soil, total phosphorus absorption (~23%), and yield (~28%) compared to the use of uncoated P fertilizer. Nevertheless, the addition of 60 kg of phosphate mixed with DCAP did not show a notable increase in phosphorus availability compared to the application of 60 kg of phosphate alone.

Furthermore, the study highlights the adverse effects of not using phosphate fertilizer, leading to reduced plant height, the number of shoots, and the percentage of firm seeds, ultimately lowering overall yield. The application of phosphate at a dose of 30 kg of P<sub>2</sub>O<sub>5</sub> mixed with DCAP demonstrated an equivalent phosphorus absorption yield to that of 60 kg of P<sub>2</sub>O<sub>5</sub> without DCAP.

The incorporation of DCAP in phosphate fertilizer proved effective in protecting phosphate particles from fixation by iron and aluminium toxins in low pH conditions and calcium and magnesium in high pH conditions. This not only enhances economic efficiency but also contributes to environmental sustainability by reducing pollution.

The study establishes that adding DCAP to phosphate fertilizer has the potential to improve phosphorus uptake and growth parameters, particularly in rice cultivation in acidic soils with limited phosphorus availability. These findings suggest the promising application of DCAP in agricultural practices, especially in regions with similar soil conditions, such as acid sulfate soils. However, further research and field trials are imperative to fully comprehend the extent of DCAP's effectiveness in real-world agricultural settings.

## Acknowledgment

This study was supported by greenhouses, equipment, and laboratories from Can Tho University and partly by scientific research funds from Vietnam's State-level project code, BDKH 57"

## Authors' contributions

Tran Van Hung: Field experiment data collection, processing, and drafting the manuscript.

Ngo Ngoc Hung: Topic development, field experiment design, final approval of the manuscript.

Vo Quang Minh\*: Correct, revise the manuscript, and corresponding.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interest to declare.

**Ethical issues:** "None".

## References

- Rodrigues M, Pavinato PS, Withers PJA, Teles APB, Herrera WFB. Legacy phosphorus and no-tillage agriculture in tropical oxisols of the Brazilian savanna. *Science of The Total Environment*.2016;542:1050-61. <https://doi.org/10.1016/j.scitotenv.2015.08.118>
- Du E, Terrer C, Pellegrini AFA, Ahlström A, van Lissa CJ, Zhao X, Xia N, Wu X, Jackson RB. Global patterns of terrestrial nitrogen and phosphorus limitation. *Nat Geosci*. 2020;13:221-26. <https://doi.org/10.1038/s41561-019-0530-4>
- Oliveira LEZ de, Nunes R de S, Figueiredo CC de, Rein TA. Spatial distribution of soil phosphorus fractions in a clayey oxisol submitted to long-term phosphate fertilization strategies. *Geoderma*. 2022;418:115847. <https://doi.org/10.1016/j.geoderma.2022.115847>
- Pavinato PS, Merlin A, Rosolem CA. Phosphorus fractions in Brazilian Cerrado soils as affected by tillage. *Soil and Tillage Research*. 2009;105:149-55. <https://doi.org/10.1016/j.still.2009.07.001>
- Fink JR, Inda AV, Tiecher T, Barrón V. Iron oxides and organic matter on soil phosphorus availability. *Ciênc agrotec*. 2016;40:369-79. <https://doi.org/10.1590/1413-70542016404023016>
- Mumbach GL, Gatiboni LC, Dall'Orsoletta DJ, Schmitt DE, Grando DL, Junior AA de S, Brignoli FM, Iochims DA. Refining phosphorus fertilizer recommendations based on buffering capacity of soils from Southern Brazil. *Rev Bras Ciênc Solo*. 2021;45. <https://doi.org/10.36783/18069657rbcs20200113>
- Pal A, Adhikary R. Improving of phosphorus use efficiency in acid and alkaline soil: A critical review study. 2020;10:18558-62.
- Sanders JL, Murphy LS, Noble A, Melgar R, Perkins J. Improving phosphorus use efficiency with polymer technology. *Procedia Engineering*. 2012;46:178-84. <https://doi.org/10.1016/j.proeng.2012.09.463>
- Murphy L, Agro M, Sanders L. Improving N and P use efficiency with polymer technology: Proceedings of Indiana Certified Crop Adviser Conference; 2007 Dec 18-19; Purdue University, Indianapolis, IN.
- Mooso G, Tindall T, Jackson G, Zhang H. Increasing the efficiency of MAP and urea applied to winter wheat in Montana with AVAIL and NutriSphere-N.
- Dunn DJ, Stevens G. Response of rice yields to phosphorus fertilizer rates and polymer coating. *Crop Management*. 2008;7:1-4. doi: 10.1094/CM-2008-0610-01-RS.
- Tindall T, Mooso G. Nitrogen and phosphorus: Mechanisms of loss from the soil system and effect to slow the losses and increase plant availability.
- Curtis J, Hill M, Hopkins B. Improving phosphorus use efficiency with Carbond® P and Dicarboxylic Acid Polymer (Avail®) fertilizer additives.
- Barben SA, Hopkins BG, Jolley VD, Webb BL, Nichols BA. Optimizing phosphorus and zinc concentrations in hydroponic chelator-buffered nutrient solution for russet burbank potato. *Journal of Plant Nutrition*. 2010;33:557-70. <https://doi.org/10.1080/01904160903506282>
- Barben SA, Hopkins BG, Jolley VD, Webb BL, Nichols BA. Phosphorus and manganese interactions and their relationships with zinc in chelator-buffered solution grown russet burbank potato. *Journal of Plant Nutrition*.2010;33:752-69. <https://doi.org/10.1080/01904160903575964>
- Barben SA, Hopkins BG, Jolley VD, Webb BL, Nichols BA. Phosphorus and zinc interactions in chelator-buffered solution grown russet burbank potato. *Journal of Plant Nutrition*. 2010;33:587-601. <https://doi.org/10.1080/01904160903506308>

17. Hopkins BG, Ellsworth JW, Bowen TR, Cook AG, Stephens SC, Jolley VD, Shiffler AK, Eggett D. Phosphorus fertilizer timing for russet burbank potato grown in calcareous soil. *Journal of Plant Nutrition*. 2010;33:529-40. <https://doi.org/10.1080/01904160903506266>
18. Horneck D, Sullivan D, Owen Jr J, Hart J. *Soil Test Interpretation Guide*; 2011.
19. Hung NN, Ve NB, Minh VQ, Hiep NH, Khuong NQ. Managing rice soil fertility in the Mekong Delta.
20. Zhang M, Kumar A, Calvert D. Aluminum and iron fractions affecting phosphorus solubility and reactions in selected sandy soils. *Soil Science*. 2001;166:940-48. <https://doi.org/10.1097/00010694-200112000-00008>
21. Noble A, Murphy L, Murray C. The use of polymer technology to improve the production of nitrogen and phosphorus fertilizer and organic efficiency. In: *Annual Meeting, Crop Protection Southern Britain, Peterborough, Cambridgeshire*. 2012;p. 27-28.
22. Mooso G, Tindall T, Hettiarachchi G. Phosphorus use efficiency in crop production. *Proceedings of Western Nutrient Management Conference*; 2013 Mar 7-8; Vol.10. Reno, Nevada, USA
23. Khuong NQ, Nghia NV, Toan LP, Hung TV, Hung NN. Effect of mixed phosphate fertilizer "Dicarboxylic Acid Polymer - DCAP" on growth and yield of rice on acid soil in the Mekong Delta. *Journal of Science Can Tho University*.2015;41:63-70.
24. Dang LV, Huu TN, Phuong LN. Effects of Dicarboxylic Acid Polymer (DCAP) mixed phosphate fertilizer application on sweet potato, cassava and yams yield on acid soil. *Journal of Science and Development*. 2016;8:1138-44.
25. Wiatrak P. Evaluation of phosphorus application with avail on growth and yield of winter wheat in Southeastern coastal plains. *American Journal of Agricultural and Biological Sciences*. 2013;8:222-29. <https://doi.org/10.3844/ajabssp.2013.222.229>
26. Fairhurst T, Witt C, Buresh R, Dobermann A. *Rice: A practical guide to nutrient management* (2nd edition). International Rice Research Institute, International Plant Nutrition Institute and International Potash Institute; 2007.