



RESEARCH ARTICLE

# Influence of some chemicals and solvents on the lytic activity and the adsorption of bacteriophages on *Pectobacterium carotovorum* subsp. *carotovorum*

Mohammed H. Abdel-Aal<sup>1</sup>, Yasmeen A. Hasanien<sup>1</sup>, Nahed A. Younis<sup>1</sup>, Gamal El Didamony<sup>2</sup>, Ahmed Askora<sup>2</sup>, Naglaa M. Balabel<sup>3,4</sup> & Khaled Abdelaal<sup>5\*</sup>

<sup>1</sup> Plant Research Department, Nuclear Research Center (NRC), Egyptian Atomic Energy Authority (EAEA), Cairo 13759, Egypt

<sup>2</sup> Department of Microbiology and Botany, Faculty of Science, Zagazig University, Zagazig 44519, Egypt

<sup>3</sup> Bacterial Diseases Research Department, Plant Pathology Research Institute, Agricultural Research Center, Giza 12619, Egypt

<sup>4</sup> Potato Brown Rot Project, Ministry of Agriculture and Land Reclamation, Dokki, Egypt

<sup>5</sup> Agricultural Botany Department, EPCRS Excellence Center, Plant Pathology and Biotechnology Laboratory, Faculty of Agriculture, Kafr Elsheikh University, 33516, Egypt

\*Email: Khaled\_elhaies@yahoo.com

 OPEN ACCESS

ARTICLE HISTORY

Received: 06 October 2023

Accepted: 13 December 2023

Available online

Version 1.0 : 15 January 2024



Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Abdel-Aal M H, Hasanien Y A, Younis N A, Didamony G E, Askora A, Balabel N M, Abdelaal K. Influence of some chemicals and solvents on the lytic activity and the adsorption of bacteriophages on *Pectobacterium carotovorum* Subsp. *carotovorum*. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.2998>

Abstract

Recently, bacteriophages have been used to control hazardous bacterial soft rot disease on crops. However, agricultural plants are frequently treated with different chemicals (fertilizers, pesticides and solvents), so we assessed the effect of some commonly used chemicals and solvents on the lytic activity of tested bacteriophages and their adsorption potential. This study reports the isolation of three specific phages against the *Pectobacterium carotovorum* subsp. *carotovorum* DSM 30170 strain, designated as  $\phi$ PC1,  $\phi$ PC2 and  $\phi$ PC3, then partially characterized using electron microscopy and genome size. The 3 isolated phages belong to the Myoviridae family. The results obtained were based on the plaque-forming unit observed after incubation. By increasing the chemical concentrations (from 0.1 to 0.5 mM), calcium chloride (CaCl<sub>2</sub>) and potassium chloride (KCl) showed a significant increase in the lytic activity of the phages. Copper sulphate (CuSO<sub>4</sub>) and copper chloride (CuCl<sub>2</sub>) showed a substantial decrease in the activity of  $\phi$ PC3; however, such a decrease was insignificant for  $\phi$ PC1 and  $\phi$ PC2. By increasing the solvent concentrations (from 30 % v/v to 70 % v/v), propanol, ethanol and methanol showed a significant decrease in the count of the three isolated phages,  $\phi$ PC1,  $\phi$ PC2 and  $\phi$ PC3, compared to the control. Chloroform was the only solvent that did not reduce the phage titer. Our findings offer significant information for developing a strategy to combat the *P. carotovorum* subsp. *carotovorum* caused bacterial soft rot disease. avoiding copper compounds and alcoholic solvents such as propanol, ethanol and methanol in plots where phages are applied seems advisable.

Keywords

*Pectobacterium carotovorum*; soft rot; bacteriophages; pesticides; fertilizers

Introduction

Numerous commercially significant vegetables, including carrot, cucumber, cabbage, eggplant, garlic, pepper, potato, onion, radish, squash, sweet potato and tomato, are susceptible to the destructive soft rot disease caused by *P. carotovorum* subsp. *carotovorum* (Pcc). At any point along the agricultural production cycle, including during planting, the growth season, transportation and storage, the diseases can be highly destructive and result in catastrophic losses. (Pcc), the most common soft rot pathogen, is a gram nega-

tive, necrotrophic, facultative anaerobe of the family *Enterobacteriaceae*, motile via flagella. The symptoms of bacterial soft rot include fast softening of the tuber tissues which could be reduced to a watery mass within a week under humid conditions and optimum temperatures.

The pathogen invades the roots through stomata, lenticels, wounds and the stolen mother plant, then multiplies quickly causing maceration of tuber tissues by secretion of different extracellular degrading enzymes (4). The ability of *Pectobacterium* species to produce and secrete large amounts of a variety of extracellular plant cell wall-degrading enzymes (PCWDE), such as pectate lyases (Pel), polygalacturonases (Peh), proteases (Prt) and cellulases (Cel), which cause extensive tissue maceration, rotting and ultimately the death of the entire plant, is essential to their virulence and pathogenicity (5). These bacterial rots cause extensive damage to crops in agriculture and in the potato sector, a total of EUR 46 million of annual losses in the European Union (6).

Phages, viruses that attack bacteria, are believed to play a significant ecological role in regulating bacterial activity and population. They represent a potentially significant biotic component that could impact bacterial soil populations and agricultural output. Phages infect their host through a range of biochemically diverse host surface receptors, such as carbohydrates, lipopolysaccharides and proteins. The phage host range is usually based on how precisely it interacts with the host receptor. For example, if the phage recognises a very specific region, it may only be able to infect a single host species or strain (8). When the lytic phage genome is injected into its host, it initiates processes that hijack the host metabolic systems to ultimately produce multiple viral progenies, which are then released from the host utilizing phage encoded lytic enzymes. These progenies are able to infect new hosts and repeat the lytic cycle. In the right circumstances, they can proliferate in the field and possibly achieve a level of control that chemicals cannot.

The use of bacteriophages in agriculture has many advantages, but their high specificity is the most significant. Because they cannot infect humans or animals, their usage in food is safe. They are persistent as long as their hosts are around and they multiply effectively in the environment. They do not need to be used repeatedly due to the population's rapid growth (12).

Chemicals used as insecticides, fertilizers and solvents significantly rely on agricultural productivity. Intentional releases of pesticides into the environment are made to eradicate weeds, insects, plant diseases and other pests that negatively impact farm or livestock output. Fertilizers are supplementary materials that are given to crops to boost productivity. Farmers use these daily to increase crop productivity. Solvents make sure that these compounds (fertilizers) dry precisely when needed and are efficiently applied to the desired crops. This guarantees that the least product is required to ensure food security.

Previous studies discussed the effect of agricultural chemicals (agrichemicals) on phages' lytic activity

(infectivity) and their adsorption on their bacterial hosts. The changes in phage titer were seen *in vitro* (16) after mixing phages into fresh formulations of some of the most widely used agrochemicals. Only one of the tested chemicals, Kocide 2000 (active component copper-hydroxide), decreased the phage titer. It was found that copper sulphate and sodium chloride showed a significant increase in the activity of *Streptomyces* phages infecting *Streptomyces scabies*, causing potato scab disease (17).

To the best of our knowledge, however, no research discusses the impact of agrichemicals and solvents on the lytic activity and the adsorption of bacteriophages on Pcc. The main objective of the study was to find out how some commonly used agrichemicals affected the ability of Pcc phages to break down cells and attach to the Pcc DSM 30170 strain.

## Materials and Methods

### Bacterial strains and culture conditions

Ain Shams University in Egypt's Faculty of Agriculture provided the Pcc DSM 30170 strain. The bacterial cells were grown at 28 °C with shaking at 200 to 300 rpm in a Casamino acid Peptone Glucose (CPG) medium that contained 0.1% Casamino Acids, 1% peptone and 0.5% glucose.

### Isolation and characterization of bacteriophages

#### Sampling and isolation of bacteriophages

Bacteriophages were isolated according to standard procedure (18) with some modifications from sewage water samples collected from a Sewage water treatment plant in Gabal Asfar, Khanka, Qalubeia, Egypt. For phage enrichment, 5 mm of mixed sewage water samples were put on hold in 50 ml of CPG medium previously inoculated with Pcc DSM 30170 strain with a  $2 \times 10^8$  CFU/ml. The inoculated flasks were incubated for 24 h at 28 °C and 150 rpm in a shaker incubator (VIS-180, Taisite Lab Sciences Inc., USA). Centrifugation was performed on the resultant cell suspension for 15 min. at 6000 rpm. The supernatant was filtered through a membrane filter with a 0.22 µm pore size. The plaque assay was carried out using the double-layer agar (DLA) overlay method. 200 µl each of the log phase host bacteria Pcc DSM 30170 strain and the phage filtrate were combined with 0.3-0.4% soft agar and put onto a 1.5% hard CPG agar plate. For 24 h, the plates were incubated at 28 °C. After incubation, the plaques' size, shape and number were noted.

#### Preparation of the phage lysates

By using the single plaque isolation approach, phage lysates were created (19). Different plaques were chosen and added to 3 ml of the liquid culture of the Pcc DSM 30170 strain based on the morphological shape of the plaques' characters. The cultures were incubated for 24 h at 150 rpm per min and 28 °C. To get the progeny phages, the cells were centrifuged. At least 3 repetitions were required to complete the procedure to achieve homogeneous plaque morphology.

### Propagation of phages

A large amount of high titer phage stock was generated by propagating the phage isolates on their liquid Pcc DSM 30170 strain culture. A single plaque was chosen and added to an Pcc DSM 30170 log phase culture (24-h culture). The phage-host mixture was incubated at 28 °C for 24 h before being centrifuged in a chilled centrifuge (INO-BRC Micro, INOVIA technology, Turkey) for 10 min at 6000 rpm. The phage particles were then precipitated in the presence of 0.5 M NaCl and 5% polyethylene glycol 6000 after the filtrate had been passed through a membrane filter with a 0.22- $\mu$ m pore size. Moreover, SM buffer, including 100 mM NaCl, 10 mM MgSO<sub>4</sub> and 0.01% gelatin, was used to dissolve the pellet after it had been centrifuged at 6000 rpm for 30 min at 4 °C. Phages that had been purified were stored at 4 °C until use (20).

### Transmission electron microscopy

On 200 mesh carbon-coated copper grids, a drop of (10<sup>12</sup> pfu/mL) phage suspension was placed, and the excess was drawn off with filter paper as described (21). Next, the grids were covered with a 2% uranyl acetate solution and the excess was drawn off as before and then examined using a transmission electron microscope (Model JEOL JEM- 2100) at Nano Tech Egypt, Dreamland, 6th October, Cairo, Egypt.

### Phages DNA extraction

Each phage's nucleic acid was extracted using a modified phenol/chloroform technique (22). Bacteriophage suspension in a volume of 947.5  $\mu$ L was gently combined with 2.5  $\mu$ L of proteinase K (20 mg. mL<sup>-1</sup>) and 50  $\mu$ L of 10% (w/v) SDS and then incubated for one h at 56 °C. The solution was then topped off with an equal volume of 25:24:1 phenol-chloroform-isoamyl alcohol (Merck, UK). The liquid was thoroughly mixed before being centrifuged at room temperature for 15 min at 12000 g. The aqueous phase was gathered. This phenol-chloroform-isoamyl alcohol extraction step was repeated twice. Nucleic acids precipitated in ice-cold 70% ethanol and the pellet were resuspended in 50  $\mu$ L of TE buffer pH 8 and stored at -20 °C.

### DNA Quantification

A spectrophotometer (NanoDrop 2000, Thermo Fisher Scientific) was used to measure the nucleic acid concentration and agarose gel 1% electrophoresis was used to verify the integrity of the nucleic acids.

### Phages DNA digestion with restriction enzymes

A mixture of *Hind*III (10 U/L) and *Eco*RI (10 U/L) was utilized for DNA fragmentation as per the manufacturer's protocol (Thermo Fisher Scientific). Also used as a DNA ladder was Quick-Load® Purple 2-log DNA Ladder (0.1 - 10.0 kb) - New England Biolabs, Inc.: N0550S. Finally, data interpretation was done using Totallab (Ver.1.1) software. MultiSUB Maxi, Maxi Horizontal Electrophoresis System Cleaver Scientific was used for fragmentation. DNA fragment sizes derived from restriction enzyme digestion profiles were used to determine the phage genome size.

### Thermal stability of phages

The thermal inactivation point of Pcc bacteriophages *in vitro* was carried out with some modifications (24). Phages suspensions were incubated at various temperatures for 10 min: 30, 40, 50, 60, 70, 80 and 90 °C. As previously described, the plaque assay was carried out 3 times for each examined temperature.

### Phages stability at different pH

With certain adjustments, the *in vitro* testing of phage stability under various pH settings was carried out in accordance with the procedures (25). The stability of the three isolated phages was tested in CPG broth at pH levels 2, 5, 7, 9 and 11. After one h of incubation at 28 °C, the surviving phage particles were immediately counted using the double-layer agar plate method.

### Phages stability to UV

The phage stability under UV radiation was tested with a modification (26). Petri dishes with 5 ml of high titer phage suspensions were individually positioned 50 cm from an ultraviolet light source ( $\lambda = 365$  nm) and exposed for 10, 20, 30, 40, 50 and 60 min. As previously described, the double-layer agar plate method counted the phage particles that had survived after each exposure interval.

### Effect of different agricultural chemicals on isolated phages

Potassium chloride (KCl), calcium chloride (CaCl<sub>2</sub>), copper sulphate (CuSO<sub>4</sub>) and copper chloride (CuCl<sub>2</sub>) were chosen to be assessed on the isolated phages. Each salt was produced as stock solutions (100 mM). Each salt was diluted to a concentration of 0, 1, 2, 3, 4 and 0.5 mM. The phage solution with a final concentration of 10<sup>8</sup> pfu/ml and 1.5 ml of the host bacteria (Pcc DSM 30170 strain) were combined with 500  $\mu$ L of each salt dilution. At room temperature, the mixture was incubated for 10 min. The double-layer method was used to calculate the number of plaques. A bacterial solution and phage were combined to create the control test without using the substances under investigation. The experiment was conducted in triplicates to minimize manual error.

### Effect of different solvents on the isolated phages

The concentration effects of different solvents (chloroform, propanol, methanol and ethanol) on the isolated phages were studied with different concentrations, such as 30, 40, 50, 60 and 70% (v/v) for each selected solvent. Phages (10<sup>8</sup> pfu/ml) in a 0.1M phosphate buffer solution (20 ml, pH7, supplemented with alcohol) were incubated at room temperature. After one h, the samples' plaque formation (units) was detected .

### Statistical analysis

Data for phage stability to different temperature degrees, pH, UV radiation and different chemicals were analyzed by One-Way ANOVA using PASW STATISTICS 18 software. Duncan's Significant Difference test included in the software was used to compare means. Differences were considered statistically significant if  $p < 0.05$ .


## Results



**Isolation of bacteriophages**

Three lytic phages with infection potential against *Pcc* DSM 30170 strain were isolated and designated as  $\phi$ PC1,  $\phi$ PC2 and  $\phi$ PC3.  $\phi$ PC1 was clear circular 3 mm plaque size.  $\phi$ PC2 was a clear circular 1 mm plaque size.  $\phi$ PC3 was a clear circular 2 mm plaque size (Fig. 1) (Table 1).

**Transmission electron microscopy**




**Fig. 1** Plaque Morphology of isolated phages

**Table 1.** Plaque morphology of the isolated phages

| Plaque character | Phage      |            |            |
|------------------|------------|------------|------------|
|                  | $\phi$ PC1 | $\phi$ PC2 | $\phi$ PC3 |
| Shape            | Circle     | Circle     | Circle     |
| Diameter in mm   | 3          | 1          | 2          |
| Transparency     | Clear      | Clear      | Clear      |
| Presence of halo | Absent     | Absent     | Absent     |

The transmission electron microscope revealed that all the isolated phages belong to the Myoviridae family.  $\phi$ PC1 was to be approximately 284.6 nm in size, comprising the head (123.07 nm) and tail (161.53 nm). The phage  $\phi$ PC2 had an icosahedral head (114.28 nm) and tail (74.1 nm) with a visible base plate (11.4 nm) and tail fibers. The average length of the phage  $\phi$ PC2 from the top of the capsid to the base plate is 199.8 nm. The phage  $\phi$ PC3 was to be approximately 273.8 nm in size, comprising the head (118.4 nm) and tail (155.4) (Fig. 2) (Table 2).



**Fig. 2** Electron micrograph of isolated phage particles under TEM


**Table 2.** Morphological characters of the isolated phages as illustrated by transmission electron microscopic examination

| phage      | Head diameter | Tail length | Tail width | Base plates |
|------------|---------------|-------------|------------|-------------|
| $\phi$ PC1 | 123.07        | 161.53      | 26.923     | -           |
| $\phi$ PC2 | 114.28        | 74.1        | 31.42      | 11.42       |
| $\phi$ PC3 | 118.4         | 155.4       | 29.6       | -           |

**Restriction endonuclease analysis of Phages DNA**

All phages include dsDNA, as shown by the effective digestion of the DNA (Fig. 3).  $\phi$ PC1's genomic DNA was digested into 11 different-sized restriction bands, whereas  $\phi$ PC2 and  $\phi$ PC3's genomic DNA was digested into seventeen distinct restriction bands. The bacteriophage genome for  $\phi$ PC1, the presence of  $\phi$ PC2 and  $\phi$ PC3 is estimated to be 14, 20 and 22 kbp in length based on the total of the restriction fragment sizes.


**Thermal stability of phages**



**Fig. 3** Fragmentation pattern for phages with a mix of Hind III and Eco RI

The sensitivity of phages  $\phi$ PC1,  $\phi$ PC2 and  $\phi$ PC3 to various temperatures is represented in Fig. 4. The activity of the 3 phages remained stable in a range of temperatures up to 70 °C. The activity of Phages  $\phi$ PC2 and  $\phi$ PC3 was lost when subjected to a temperature of 80 °C for 10 min.  $\phi$ PC1 was totally lost when subjected to 90 °C for 10 min. Interestingly,  $\phi$ PC1,  $\phi$ PC2 and  $\phi$ PC3 phages survived at 50 °C with no significant loss in phage particle number.

**Phages stability at different pH**



**Fig.4** Phages stability at different temperatures

Fig. 5 illustrates how sensitive the phages  $\phi$ PC1,  $\phi$ PC2 and  $\phi$ PC3 are to different pH levels. When incubated at 28 °C,  $\phi$ PC1 and  $\phi$ PC2 maintained their capacity to lyse the *Pcc* DSM 30170 strain for up to 1 h at pH levels ranging from 3 to 9, with the highest stability at pH 6, 7 and 8. When incubated at 28 °C for an h, the activity of  $\phi$ PC3 remained stable for up to an h at pH levels ranging from 3 to 8 and it had the most significant stability at pH 6 and 7.

**Phages stability to UV radiation**

By increasing the exposure time from 0 to 60 min, UV radiation significantly reduced the number of plaques for each of  $\phi$ PC1,  $\phi$ PC2 and  $\phi$ PC3.  $\phi$ PC1,  $\phi$ PC2 and  $\phi$ PC3 retained their lytic capability after exposure to ultraviolet radiation for 60 min at 50 cm (Fig. 6).

**Effect of different agricultural chemicals on the isolated phages**




Fig. 5 Phages stability at different pH




Fig. 6 Phages stability to UV

By increasing the chemical concentrations (from 0.1 to 0.5 mM), calcium chloride ( $\text{CaCl}_2$ ) showed a significant increase in the activity of  $\phi\text{PC1}$ ,  $\phi\text{PC2}$  and  $\phi\text{PC3}$ , compared to the control. Potassium chloride (KCl) showed a significant increase in the activity of  $\phi\text{PC1}$  and  $\phi\text{PC2}$ . Potassium chloride (KCl) increased the activity of  $\phi\text{PC3}$ ; however, such an increase was insignificant. Copper sulphate

( $\text{CuSO}_4$ ) and copper chloride ( $\text{CuCl}_2$ ) showed a significant decrease in the activity of  $\phi\text{PC3}$ ; however, such a decrease was insignificant for  $\phi\text{PC1}$  and  $\phi\text{PC2}$  (Table 3).

#### Effect of different solvents on the isolated phages

The only solvent tested that didn't cause the phage titer to decrease was chloroform. The count of the three isolated phages,  $\phi\text{PC1}$ ,  $\phi\text{PC2}$  and  $\phi\text{PC3}$ , was significantly reduced by increasing the solvent concentrations (from 30% v/v to 70% v/v) in propanol, ethanol and methanol when compared to the control. Following exposure to 60% propanol, 70% ethanol or 70% methanol, no  $\phi\text{PC1}$  plaques were discovered. Following exposure to 50% propanol, 70% ethanol or 70% methanol, no  $\phi\text{PC2}$  plaques were discovered. Following exposure to 50% propanol, 60% ethanol or methanol, no  $\phi\text{PC3}$  plaques were discovered (Table 4).

#### Discussion

We successfully isolated three phages from sewage samples collected from the Sewage water treatment plant in Gabal Asfar, Khanka, Qalubeia, Egypt. to provide biological disease prevention strategies. It was not the first study to isolate bacteriophages lytic against *Pectobacterium carotovorum* as isolated bacteriophage PP16 from wastewater in the Moscow region using the highly virulent Pcc strain F002 (PB69) as a bacterial host.

DNA or RNA, either double- or single-stranded, makes up the genomes of phages. These genetic components are contained in a capsid, which can be polyhedral (Microviridae, Corticoviridae, Tectiviridae, Leviviridae and

Table 3. Effect of different agricultural chemicals on the isolated phages

| Chemicals       | Effect                     | Concentration (mM) | Phage count*( $10^7$ pfu/ml) |                          |                           |
|-----------------|----------------------------|--------------------|------------------------------|--------------------------|---------------------------|
|                 |                            |                    | $\phi\text{PC1}$             | $\phi\text{PC2}$         | $\phi\text{PC3}$          |
| $\text{CaCl}_2$ | Increase phages activity ↑ | 0.1                | 18.00 ± 0.06 <sup>e</sup>    | 13 ± 0.17 <sup>e</sup>   | 15 ± 0.29 <sup>e</sup>    |
|                 |                            | 0.2                | 20 ± 0.58 <sup>d</sup>       | 16 ± 0 <sup>d</sup>      | 17 ± 0.06 <sup>d</sup>    |
|                 |                            | 0.3                | 25.00 ± 0 <sup>c</sup>       | 20 ± 0.58 <sup>c</sup>   | 18 ± 0.58 <sup>c</sup>    |
|                 |                            | 0.4                | 28.00 ± 0.11 <sup>b</sup>    | 22 ± 0.35 <sup>b</sup>   | 19 ± 0.12 <sup>b</sup>    |
|                 |                            | 0.5                | 30.00 ± 0.11 <sup>a</sup>    | 25 ± 0.29 <sup>a</sup>   | 20 ± 0.0 <sup>a</sup>     |
| KCl             | Increase phages activity ↑ | 0.1                | 11.20 ± 0.64 <sup>d</sup>    | 10 ± 0.46 <sup>b</sup>   | 10.2 ± 0.12 <sup>c</sup>  |
|                 |                            | 0.2                | 13.00 ± 0.29 <sup>c</sup>    | 11 ± 0.23 <sup>b</sup>   | 10.5 ± 0.29 <sup>bc</sup> |
|                 |                            | 0.3                | 14.00 ± 0.35 <sup>b</sup>    | 11.3 ± 0.75 <sup>b</sup> | 12 ± 1.15 <sup>ab</sup>   |
|                 |                            | 0.4                | 14.10 ± 0.06 <sup>b</sup>    | 13 ± 0.58 <sup>a</sup>   | 12.3 ± 0.40 <sup>a</sup>  |
|                 |                            | 0.5                | 15.30 ± 0.17 <sup>a</sup>    | 13.3 ± 0.06 <sup>a</sup> | 12.8 ± 0.12 <sup>a</sup>  |
| $\text{CuSO}_4$ | Decrease phages activity ↓ | 0.1                | 10.00 ± 0.40 <sup>a</sup>    | 10 ± 0.58 <sup>a</sup>   | 9 ± 0.58 <sup>a</sup>     |
|                 |                            | 0.2                | 9.70 ± 0.17 <sup>a</sup>     | 9.9 ± 0.06 <sup>a</sup>  | 8 ± 0.35 <sup>ab</sup>    |
|                 |                            | 0.3                | 9.00 ± 0.17 <sup>ab</sup>    | 9.9 ± 0.11 <sup>a</sup>  | 7 ± 0.29 <sup>bc</sup>    |
|                 |                            | 0.4                | 8.70 ± 0.75 <sup>ab</sup>    | 10 ± 0.0 <sup>a</sup>    | 7.2 ± 0.12 <sup>bc</sup>  |
|                 |                            | 0.5                | 8.00 ± 0.58 <sup>b</sup>     | 9.8 ± 0.11 <sup>a</sup>  | 6 ± 0.58 <sup>d</sup>     |
| $\text{CuCl}_2$ | Decrease phages activity ↓ | 0.1                | 9.00 ± 0.06 <sup>a</sup>     | 10 ± 0.58 <sup>a</sup>   | 8.8 ± 0.12 <sup>a</sup>   |
|                 |                            | 0.2                | 8.50 ± 0.29 <sup>a</sup>     | 9.5 ± 0 <sup>a</sup>     | 7.4 ± 0.23 <sup>b</sup>   |
|                 |                            | 0.3                | 8.20 ± 0.17 <sup>a</sup>     | 9 ± 0.17 <sup>a</sup>    | 6.6 ± 0.69 <sup>bc</sup>  |
|                 |                            | 0.4                | 7.00 ± 0.58 <sup>b</sup>     | 8 ± 0.17 <sup>b</sup>    | 6.4 ± 0.35 <sup>bc</sup>  |
|                 |                            | 0.5                | 6.40 ± 0.06 <sup>b</sup>     | 7.5 ± 0.40 <sup>b</sup>  | 6 ± 0.35 <sup>c</sup>     |

**Table 4.** Effect of different solvents on the isolated phages

| Solvents   | Effect                        | Concentration % (v/v) | Phage count*(10 <sup>7</sup> pfu/ml) |                           |                           |
|------------|-------------------------------|-----------------------|--------------------------------------|---------------------------|---------------------------|
|            |                               |                       | φPC1                                 | φPC2                      | φPC3                      |
| Chloroform | No effect                     | 30                    | 9.83 ± 0.09 <sup>b</sup>             | 10.06 ± 0.09 <sup>a</sup> | 10.03 ± 0.13 <sup>a</sup> |
|            |                               | 40                    | 9.93 ± 0.20 <sup>ab</sup>            | 10 ± 0.06 <sup>a</sup>    | 9.96 ± 0.09 <sup>a</sup>  |
|            |                               | 50                    | 10.16 ± 0.03 <sup>ab</sup>           | 10.16 ± 0.03 <sup>a</sup> | 10.10 ± 0.06 <sup>a</sup> |
|            |                               | 60                    | 10.23 ± 0.07 <sup>a</sup>            | 10.16 ± 0.09 <sup>a</sup> | 10.10 ± 0.10 <sup>a</sup> |
|            |                               | 70                    | 10.06 ± 0.07 <sup>ab</sup>           | 10.06 ± 0.07 <sup>a</sup> | 9.96 ± 0.14 <sup>a</sup>  |
| Propanol   | Decrease phages activity<br>↓ | 30                    | 10.06 ± 0.07 <sup>ab</sup>           | 10.06 ± 0.07 <sup>a</sup> | 9.96 ± 0.14 <sup>a</sup>  |
|            |                               | 40                    | 5.3 ± 0.15 <sup>a</sup>              | 4.76 ± 0.14 <sup>a</sup>  | 2.94 ± 0.03 <sup>a</sup>  |
|            |                               | 50                    | 3.06 ± 0.12 <sup>b</sup>             | 2.46 ± 0.26 <sup>b</sup>  | 1.01 ± 0.09 <sup>b</sup>  |
|            |                               | 60                    | 0.16 ± 0.01 <sup>c</sup>             | 0.09 ± 0.00 <sup>c</sup>  | 0 ± 0 <sup>c</sup>        |
|            |                               | 70                    | 0.00 ± 0.00 <sup>c</sup>             | 0 ± 0 <sup>c</sup>        | 0 ± 0 <sup>c</sup>        |
| Methanol   | Decrease phages activity<br>↓ | 30                    | 9.8 ± 0.06 <sup>a</sup>              | 9.13 ± 0.07 <sup>a</sup>  | 9.9 ± 0.06 <sup>a</sup>   |
|            |                               | 40                    | 9.36 ± 0.09 <sup>b</sup>             | 7.96 ± 0.09 <sup>b</sup>  | 9.66 ± 0.14 <sup>b</sup>  |
|            |                               | 50                    | 7.9 ± 0.06 <sup>c</sup>              | 6.13 ± 0.14 <sup>c</sup>  | 6.06 ± 0.12 <sup>c</sup>  |
|            |                               | 60                    | 2.23 ± 0.14 <sup>d</sup>             | 2.76 ± 0.12 <sup>d</sup>  | 0.07 ± 0.03 <sup>d</sup>  |
|            |                               | 70                    | 0 ± 0 <sup>e</sup>                   | 0 ± 0 <sup>e</sup>        | 0 ± 0 <sup>d</sup>        |
| Ethanol    | Decrease phages activity<br>↓ | 30                    | 9.66 ± 0.09 <sup>a</sup>             | 9.9 ± 0.06 <sup>a</sup>   | 9.9 ± 0.15 <sup>a</sup>   |
|            |                               | 40                    | 9.36 ± 0.09 <sup>b</sup>             | 9.16 ± 0.09 <sup>b</sup>  | 9.03 ± 0.03 <sup>b</sup>  |
|            |                               | 50                    | 7.50 ± 0.29 <sup>c</sup>             | 7 ± 0.29 <sup>c</sup>     | 3.23 ± 0.18 <sup>c</sup>  |
|            |                               | 60                    | 0.13 ± 0.03 <sup>d</sup>             | 0.25 ± 0.02 <sup>d</sup>  | 0 ± 0 <sup>d</sup>        |
|            |                               | 70                    | 0.00 ± 0 <sup>d</sup>                | 0 ± 0 <sup>d</sup>        | 0 ± 0 <sup>d</sup>        |

Cystoviridae), filamentous (Inoviridae), pleomorphic (Plasmaviridae), or attached to a tail (Caudovirales). The Caudovirales order contains the Myoviridae family with a contractile tail and the Podoviridae family with a short tail, making up around 96% of the documented bacteriophages. The transmission electron microscope and genomic characteristics of the 3 isolated phages in this study revealed that they all belong to the Myoviridae family. φPC1 was to be approximately 284.6 nm in size, comprising head (123.07 nm) and tail (161.53 nm). The phage φPC2 had an icosahedral head (114.28 nm) and tail (74.1 nm) with a visible base plate (11.4 nm) and tail fibers. The average length of the phage φPC2 from the top of the capsid to the base plate is 199.8 nm. The phage φPC3 was to be approximately 273.8 nm in size, comprising of head (118.4 nm) and tail (55.4). Likewise, several *Myoviridae* bacteriophages infecting *Pcc* have been described: Pc1, Wc2, Wc3, Wc4 vB\_PcaM-D1 and vB\_PcaM-J3 infecting *Pcc*. All isolated phages include dsDNA, as shown by the effective digestion of the DNA. The genomic DNA of φPC1 digested in eleven restriction bands, whereas φPC2 and φPC3 digested in seventeen restriction bands of different sizes. The genome for φPC1, φPC2 and φPC3 is estimated to be 14, 20 and 22 kbp in length based on the total of the restriction fragment sizes. Their stability significantly influences agriculture's acceptance of biocontrol agents.

The isolated phages exhibited thermostability at elevated temperatures (up to 70 °C for φPC2 and φPC3 and 80 °C for φPC1). This is in accordance with, who suggested

that bacteriophage PP1 1, with lytic activity against *Pcc*, was stable at relatively high temperatures. Similarly, found that Phage P7\_Pc could maintain its viability against *Pcc* when stored at 45 °C and 55 °C for 1 h (19).

Phages were contagious in both acidic and alkaline settings, but excessive alkalinity completely suppressed them (pH 10 for φPC1, pH 11 for φPC2 and φPC3). The optimum infectivity of φPC1 and φPC2 was obtained at pH range 6-8 and 6-7 for φPC3. This is in agreement with, who stated that the optimal pH for phages is neutral and the rise in acidity and alkalinity has an inverse relationship with the infectivity of the phage (36). As reported, the denaturation of phages' proteins is to blame for this.

φPC1, φPC2 and φPC3 retained their lytic capability after exposure to ultraviolet radiation for 60 min at 50 cm. Inversely, Myoviridae bacteriophages isolated could not survive 10 min of UV exposure, although 5 min of exposure cut their concentration in half as measured by a plaque assay (38).

Agricultural plants are frequently treated with different chemicals. To uncover potential interferences, we mixed phages with varying concentrations of some of the most commonly used agrochemicals and monitored the changes in phage titer after each treatment. By increasing the chemical concentrations (from 0.1 to 0.5 mM), calcium chloride (CaCl<sub>2</sub>) and potassium chloride (KCl) showed a significant increase in the lytic activity of the 3 phages φPC1, φPC2 and φPC3. This rise in phage activity is likely due to higher rates of adsorption and penetration. Our

results are in accordance with, who showed that calcium and potassium ions increase the process of phage infectivity (39). Copper sulphate (CuSO<sub>4</sub>) and copper chloride (CuCl<sub>2</sub>) showed a significant decrease in the activity of  $\phi$ PC3; however, such a decrease was insignificant for  $\phi$ PC1 and  $\phi$ PC2. Despite its obvious efficacy, it is still unclear how copper inactivates phages. According to the majority of publications, redox cycling between the various copper species produces hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and/or reactive oxygen species (ROS), which in turn causes phage inactivation. These molecules are known to damage DNA and proteins, the components of bacteriophages. Thus, avoiding using copper compounds in plots where phages are applied seems advisable.

Propanol, ethanol and methanol showed a significant decrease in the count of the 3 isolated phages,  $\phi$ PC1,  $\phi$ PC2 and  $\phi$ PC3, by increasing the concentrations (from 30 % v/v to 70 % v/v).  $\phi$ PC1,  $\phi$ PC2 and  $\phi$ PC3 were more tolerant to ethanol and methanol than propanol as 50-60% (v/v) propanol resulted in total loss in the phages count, but 60-70% (v/v) ethanol, or methanol decreased the phages count to zero. Below these concentrations, the killing effect of alcohols against bacteriophages became much weaker. On the other hand, chloroform was the only one among the solvents tested that did not significantly reduce the phage titer. Other phages were found to tolerate chloroform, such as *Pectobacterium* phage CB7 isolated (42). After 1 h of exposure to chloroform, the phage was tolerable, with no discernible log-fold decrease in phage titer. In contrast, *Pectobacterium* phages isolated earlier were found to be sensitive to chloroform (43). 1 h of incubation with chloroform at room temperature caused a 100-fold titer drop.

Our findings offer significant information for developing a strategy to combat the *P. carotovorum* subsp. *carotovorum* caused bacterial soft rot disease.

## Conclusion

Three phages lytic against *P. carotovorum* subsp. *carotovorum* were isolated, designated as ( $\phi$ PC1,  $\phi$ PC2,  $\phi$ PC3) and classified based on morphological and genomic characteristics as Myoviridae. Before applying phages to treat the soft rot disease caused by Pcc, we tested if the lytic activity of phages could be affected by some common chemicals and solvents generally used in agriculture. From these chemicals tested, calcium chloride and potassium chloride were found to increase the lytic activity of the phages and their adsorption on Pcc. On the other hand, copper sulphate, copper hydroxide and alcohols decreased the lytic activity of the phages against Pcc. Thus, it seems advisable to avoid using copper compounds and alcoholic solvents in plots where phages are applied.

## Authors contributions

MHA performed the experiments and wrote the manuscript and participated in manuscript revising and editing.

YAH conducted experimental methodology, participated in data analysis and representation and participated in manuscript revising and editing. NAY conceived and designed research, provided the used chemicals, provided practical guidance and participated in manuscript revising and editing. NMB provided the used chemicals, provided practical guidance and participated in manuscript revising and editing. AA conceived and designed the research, conducted experimental methodology, participated in data analysis and representation. GD suggested the research point, investigated the article, conceived and designed the research, conducted experimental methodology. KA participated in manuscript revising, editing and publishing. All authors read and approved the article.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** None.

## References

1. Abd-El-Khair H, Abdel-Gaied TG, Mikhail MS, Abdel-Alim AI, El-Nasr HIS. Biological control of *Pectobacterium carotovorum* subsp. *carotovorum*, the causal agent of bacterial soft rot in vegetables, *in vitro* and *in vivo* tests. Bulletin of the National Research Centre. 2021;45:1-9. <https://doi.org/10.1186/s42269-021-00491-4>
2. Abu-Obeid IM. Soft rot disease in Jordan: A review. Advances in Environmental Biology. 2019;13(6):1-6. <http://doi.org/10.22587/aeb.2019.13.6.1>
3. Youdkes D, Helman Y, Burdman S, Matan O, Jurkevitch E. Potential control of potato soft rot disease by the obligate predators *Bdellovibrio* and like organisms. Applied and Environmental Microbiology. 2020;86(6):e02543-02519. <https://doi.org/10.1128/AEM.02543-19>
4. Khedr AA. Management of soft rot disease, caused by *Erwinia carotovora* subsp. *carotovora* in potato tubers. African Journal of Biological Sciences. 2019;15(1):211-18. <https://doi.org/10.21608/ajbs.2019.72268>
5. Agyemang PA, Kabir MN, Kersey CM, Dumenyo CK. The bacterial soft rot pathogens, *Pectobacterium carotovorum* and *P. atrosepticum*, respond to different classes of virulence-inducing host chemical signals. Horticulturae. 2020;6(1):13. <https://doi.org/10.3390/horticulturae6010013>
6. Beňo F, Horsáková I, Kmoch M, Petrzik K, Krátká G, Ševčík R. Bacteriophages as a strategy to protect potato tubers against *Dickeya dianthicola* and *Pectobacterium carotovorum* soft rot. Microorganisms. 2022;10(12):2369. <https://doi.org/10.3390/microorganisms10122369>
7. Liang X, Radosevich M, DeBruyn JM, Wilhelm SW, McDearis R, Zhuang J. Incorporating viruses into soil ecology: A new dimension to understand biogeochemical cycling. Critical Reviews in Environmental Science and Technology. 2023:1-21. <https://doi.org/10.1080/10643389.2023.2223123>
8. Batinovic S, Wassef F, Knowler SA, Rice DT, Stanton CR, Rose J, Tucci J, Nittami T, Vinh A, Drummond GR. Bacteriophages in natural and artificial environments. Pathogens. 2019;8(3):100.
9. Howard-Varona C, Hargreaves KR, Abedon ST, Sullivan MB. Lysogeny in nature: Mechanisms, impact and ecology of temperate phages. The ISME journal. 2017;11 (7):1511-20. <https://doi.org/10.1038/ismej.2017.16>



10. D'Accolti M, Soffritti I, Mazzacane S, Caselli E. Bacteriophages as a potential 360-degree pathogen control strategy. *Microorganisms*. 2021;9(2):261. <https://doi.org/10.3390/microorganisms9020261>
11. Czajkowski R, Perombelon MC, van Veen JA, van der Wolf JM. Control of blackleg and tuber soft rot of potato caused by *Pectobacterium* and *Dickeya* species: A review. *Plant Pathology*. 2011;60(6):999-1013. <https://doi.org/10.1111/j.1365-3059.2011.02470.x>
12. Kering KK, Kibii BJ, Wei H. Biocontrol of phyto-bacteria with bacteriophage cocktails. *Pest Management Science*. 2019;75(7):1775-81. <https://doi.org/10.1002/ps.5324>
13. Shang Y, Hasan MK, Ahammed GJ, Li M, Yin H, Zhou J. Applications of nanotechnology in plant growth and crop protection: A review. *Molecules*. 2019;24(14):2558. <https://doi.org/10.3390/molecules24142558>
14. Itelima J, Bang W, Onyimba I, Sila M, Egbere O. Bio-fertilizers as key player in enhancing soil fertility and crop productivity: A review. *Direct Research Journal of Agriculture and Food Science*. 2018;6(3):73-83. <http://hdl.handle.net/123456789/1999>
15. Xi L, Zhang M, Zhang L, Lew TT, Lam YM. Novel materials for urban farming. *Advanced Materials*. 2022;34(25):2105009. <https://doi.org/10.1002/adma.202105009>
16. Balogh B, Jones J, Momol M, Olson S. Persistence of bacteriophages as biocontrol agents in the tomato canopy. In 1<sup>st</sup> International Symposium on Tomato Diseases. 2004;695:299-302.
17. AlKhazindar M, Sayed E, Khalil M, Zahran D. Isolation and characterization of two phages infecting *Streptomyces scabies*. *Research Journal of Pharmaceutical Biological and Chemical Sciences*. 2016;7(2):1351-63.
18. Abo-Senna A, El-Fouly M, Hussein H, Swailam H, Abdel-Aal M. Biocontrol of food borne *Salmonella* using bacteriophages. *Arab Journal of Nuclear Sciences and Applications (Online)*. 2018;51(1):100-09. <https://doi.org/10.5771/0010-3497-2018-1-109>
19. Naligama KN, Halmillawewa AP. *Pectobacterium carotovorum* phage vB\_PcaM\_P7\_Pc is a new member of the genus *Certrevirus*. *Microbiology Spectrum*. 2022;10(6):e03126-32. <https://doi.org/10.1128/spectrum.03126-22>
20. Jurczak-Kurek A, Gąsior T, Nejman-Faleńczyk B, Bloch S, Dydecka A, Topka G *et al.* Biodiversity of bacteriophages: Morphological and biological properties of a large group of phages isolated from urban sewage. *Scientific Reports*. 2016;6(1):1-17. <https://doi.org/10.1038/srep34338>
21. Kalatzis PG, Bastías R, Kokkari C, Katharios P. Isolation and characterization of two lytic bacteriophages,  $\phi$ St2 and  $\phi$ Grn1; phage therapy application for biological control of *Vibrio alginolyticus* in aquaculture live feeds. *PloS one*. 2016;11(3):e0151101. <https://doi.org/10.1371/journal.pone.0151101>
22. Phothichaisri W, Ounjai P, Phetruen T, Janvilisri T, Khunrae P, Singhakaew S *et al.* Characterization of bacteriophages infecting clinical isolates of *Clostridium difficile*. *Frontiers in Microbiology*. 2018;9:1701. <https://doi.org/10.3389/fmicb.2018.01701>
23. Akhwale JK, Rohde M, Rohde C, Bunk B, Spröer C, Boga HI, Klenk H-P, Wittmann J. Isolation, characterization and analysis of bacteriophages from the haloalkaline lake Elmenteita, Kenya. *PLoS One*. 2019;14(4):e0215734. <https://doi.org/10.1371/journal.pone.0215734>
24. Othman B, Askora A, Abo-Senna AS. Isolation and characterization of a *Siphoviridae* phage infecting *Bacillus megaterium* from a heavily trafficked holy site in Saudi Arabia. *Folia Microbiologica*. 2015;60:289-95. <https://doi.org/10.1007/s12223-015-0375-1>
25. Xue Y, Zhai S, Wang Z, Ji Y, Wang G, Wang T *et al.* The yersinia phage X1 administered orally efficiently protects a murine chronic enteritis model against *Yersinia enterocolitica* infection. *Frontiers in Microbiology*. 2020; 11:351. <https://doi.org/10.3389/fmicb.2020.00351>
26. Lee S, Vu N-T, Oh E-J, Rahimi-Midani A, Thi T-N, Song Y-R *et al.* Biocontrol of soft rot caused by *Pectobacterium odoriferum* with bacteriophage phiPccP-1 in Kimchi cabbage. *Microorganisms*. 2021;9(4):779. <https://doi.org/10.3390/microorganisms9040779>
27. El-Afifi S, Hammad A. Biochemical and molecular characteristics of Pc1 virulent phage isolate infecting *Pectobacterium carotovorum*. *Pakistan Journal of Biological Sciences: PJBS*. 2020;23(11):1481-86. <https://doi.org/10.3923/pjbs.2020.1481.1486>
28. Chandrarathna H, Nikapitiya C, Dananjaya S, De Silva B, Heo G-J, De Zoysa M, Lee J. Isolation and characterization of phage AHP-1 and its combined effect with chloramphenicol to control *Aeromonas hydrophila*. *Brazilian Journal of Microbiology*. 2020;51:409-16. <https://doi.org/10.1007/s42770-019-00178-z>
29. Voronina M, Bugaeva E, Vasiliev D, Kabanova A, Barannik A, Shneider M *et al.* Characterization of *Pectobacterium carotovorum* subsp. *carotovorum* bacteriophage PP16 prospective for biocontrol of potato soft rot. *Microbiology*. 2019;88:451-60. <https://doi.org/10.1134/S0026261719040118>
30. Dion MB, Oechslein F, Moineau S. Phage diversity, genomics and phylogeny. *Nature Reviews Microbiology*. 2020;18(3):125-38. <https://doi.org/10.1038/s41579-019-0311-5>
31. Kaliniene L, Šimoliūnas E, Truncaitė L, Zajančauskaitė A, Nainys J, Kaupinis A, Valius M, Meškys R. Molecular analysis of *Arthrobacter myovirus* vB\_ArtM-ArV1: We blame it on the tail. *Journal of Virology*. 2017;91(8):e00023-00017. <https://doi.org/10.1128/jvi.00023-17>
32. Parena AJS, Silva BBI, Mercado RML, Sendon AAA, Signabon FB, Balidion JF, Encabo JR. Lytic phages display protective effects against soft rot-causing *Pectobacterium* sp. *Biocontrol Science and Technology*. 2022;32(11):1326-45. <https://doi.org/10.1080/09583157.2022.2122403>
33. Marei E, El-Afifi SI, Elsharouny T, Hammad AM. Biological control of *Pectobacterium carotovorum* via specific lytic bacteriophage. *J Basic Appl Sci Res*. 2017;7:1-9.
34. Muturi P, Yu J, Maina AN, Kariuki S, Mwaura FB, Wei H. Bacteriophages isolated in China for the control of *Pectobacterium carotovorum* causing potato soft rot in Kenya. *Virologica Sinica*. 2019;34:287-94. <https://doi.org/10.1007/s12250-019-00091-7>
35. Svircev A, Roach D, Castle A. Framing the future with bacteriophages in agriculture. *Viruses*. 2018;10(5):218. <https://doi.org/10.3390/v10050218>
36. Lim J-A, Jee S, Lee DH, Roh E, Jung K, Oh C, Heu S. Biocontrol of *Pectobacterium carotovorum* subsp. *carotovorum* using bacteriophage PP1. *Journal of Microbiology and Biotechnology*. 2013;23(8):1147-53. <http://dx.doi.org/10.4014/jmb.1304.04001>
37. Sharma M, Kumar D, Poluri KM. Elucidating the pH-dependent structural transition of T7 bacteriophage endolysin. *Biochemistry*. 2016;55(33):4614-25. <https://doi.org/10.1021/acs.biochem.6b00240>
38. Czajkowski R, Ozymko Z, Lojkowska E. Isolation and characterization of novel soilborne lytic bacteriophages infecting *Dickeya* spp. biovar 3 ('D. solani'). *Plant Pathology*. 2014;63(4):758-72. <https://doi.org/10.1111/ppa.12157>
39. Szermer-Olearnik B, Drab M, Mąkosa M, Zembala M, Barbasz J, Dąbrowska K, Boratyński J. Aggregation/dispersion transitions of T4 phage triggered by environmental ion availability. *Journal of Nanobiotechnology*. 2017;15(1):1-15. <https://doi.org/10.1186/s12951-017-0266-5>
40. Grass G, Rensing C, Solioz M. Metallic copper as an antimicrobial surface. *Applied and Environmental Microbiology*. 2011;77(5):1541-47. <https://doi.org/10.1128/AEM.02766-10>



41. Li J, Dennehy JJ. Differential bacteriophage mortality on exposure to copper. *Applied and Environmental Microbiology*. 2011;77(19):6878-83. <https://doi.org/10.1128/AEM.05661-11>
42. Buttner C, Lynch C, Hendrix H, Neve H, Noben J-P, Lavigne R, Coffey A. Isolation and characterization of *Pectobacterium* phage vB\_PatM\_CB7: New insights into the genus *Certrevirus*. *Antibiotics*. 2020; 9(6):352. <https://doi.org/10.3390/antibiotics9060352>
43. Lukianova AA, Evseev PV, Shneider MM, Dvoryakova EA, Tokmakova AD, Shpirt AM *et al.* *Pectobacterium versatile* bacteriophage possum: A complex polysaccharide-deacetylating tail fiber as a tool for host recognition in pectobacterial schitoviridae. *International Journal of Molecular Sciences*. 2022; 23(19):11043. <https://doi.org/10.3390/ijms231911043>