



RESEARCH ARTICLE

Phenolic contents, antioxidant activity and proximate analysis of *Ziziphus oxyphylla* Edgew. (Angrezi ber) from Soon Valley, Salt Range, Pakistan

Sarvat Rahim¹, Javaria Nargis¹, Amin Shah^{1*}, Muhammad Nadeem², Muhammad Kabir³, Gulfam Shahzad¹, Tehreem Tahir¹, Asif Abbas Shah¹, Nuzhat Rasul¹ & Umna Zainab¹

¹Department of Botany, University of Sargodha, Sargodha-40100, Pakistan

²Institute of Food Science and Nutrition, University of Sargodha, Sargodha-40100, Pakistan

³Department of Biological Sciences, Thal University Bhakkar (University of Sargodha, Ex-Sub Campus Bhakkar), Bhakkar-30000, Punjab, Pakistan

*Email: aminullah.amin@uos.edu.pk



ARTICLE HISTORY

Received: 05 December 2023

Accepted: 27 March 2024

Available online

Version 1.0 : 25 June 2024

Version 2.0 : 01 July 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Rahim S, Nargis J, Shah A, Nadeem M, Kabir M, Shahzad G, Tahir T, Shah AA, Rasul N, Zainab U. Phenolic contents, antioxidant activity and proximate analysis of *Ziziphus oxyphylla* Edgew. (Angrezi ber) from Soon Valley, Salt Range, Pakistan. Plant Science Today. 2024; 11(3): 136-141. <https://doi.org/10.14719/pst.3163>

Abstract

The purpose of this research was to investigate the phenolic composition, antioxidant activity and proximate composition of *Ziziphus oxyphylla* Edgew. (Angrezi ber) through the use of ethanol and aqueous extracts. The flavonoid concentration of the extracts exhibited significant variation as a result of employing two distinct extraction methods, specifically sun-drying and shade-drying. Shade-dried aqueous extracts of leaves yielded the highest TPC of 599 ± 1.00 mg/100 g GAE (where GAE stands for Gallic acid equivalent), while shade-dried ethanol extracts of leaves exhibited a minimum value of TPC 281 ± 1.00 mg/100 g GAE. In leaves, the highest flavonoid content was observed in sun-dried ethanol extracts at 496.296 ± 3.70 mg/100 g CE (where CE stands for catechin equivalent), while the minimum was observed in shade-dried aqueous extracts at 159.259 ± 48.15 mg/100 g CE. Sun-dried ethanol extract 3711.35 ± 2.70 µg/100 g AAE (where AAE stands for ascorbic acid equivalent) exhibits a higher level of antioxidants as compared to shade-dried aqueous extracts valued at 462.70 ± 2.70 µg/100 g AAE. Aqueous and ethanol extracts of fruit yielded 758.000 mg/100 g GAE and 739.333 mg/100 g GAE respectively. At the same time, aqueous and ethanol extracts of fruit yielded 405.045 µg/100 g AAE and 329.369 µg/100 g AAE antioxidants respectively. Proximate analysis of *Z. oxyphylla* fruit yielded the following results: Moisture 88.4 ± 0.02081 %, Fiber 5.03 ± 0.02 %, ash 5.6666 ± 0.1527 %, protein 2.0366 ± 0.0152 % and NFE (Nitrogen-Free Extract) 79.233 ± 0.02081 %. The results obtained from this study, which focused on the total phenolic contents and proximate compositions of *Z. oxyphylla*, can serve as a foundation for further investigations into its antioxidant capabilities also.

Keywords

Ziziphus oxyphylla? Soon Valley? nutritional analysis? Folin-Ciocalteu reagent; food source

Introduction

Food serves as a vital source of energy and essential building blocks for the restoration of damaged tissues as well as the production of crucial physiological components. Hence, it is imperative to ensure the provision of appropriate quality and amount of food to an individual. The assessment of food quality can be based on its nutritive values, which are determined by analyzing the composition of food in terms of its carbohydrate, protein, fat, mineral and vitamin content. The caloric content of food is contingent upon

the quantity of these constituents. The presence of insufficient quantities of one or more essential components can lead to the manifestation of signs and symptoms associated with a range of disorders. These several components collaborate in order to maintain the proper functioning of the human body. Nevertheless, the nutritional benefit of a diverse dietary regimen is solely advantageous if the essential components are sufficiently accessible in appropriate amounts and compositions (1-3).

Antioxidants serve the purpose of mitigating cellular damage caused by free radicals, which are molecules produced as a result of the regular metabolic process of oxidation. Several types of free radicals are present in biological systems, such as reactive hydroxyl radicals, reactive oxygen free radicals species, the superoxide anion radical, peroxy and hydrogen peroxides. These molecules are capable of generating metabolic products that can initiate attacks on DNA or lipids in cell membranes. These factors are implicated in several forms of biological harm, including the development of cancer, cellular deterioration associated with aging, DNA damage and their contribution to heart disease and arthritis (4).

Currently, there exists a global liking for the utilization of aromatic and medicinal plants as sources of antioxidants in the context of food (5). The primary source of antioxidative properties in plants can be traced to phenolic compounds, including phenolic diterpenes, flavonoids and phenolic acids. These compounds possess the capacity to effectively neutralize free radicals, either by

scavenging them, donating hydrogen atoms or electrons, or chelating metal cations (6). A strong positive connection was observed between the antioxidant capability and total phenol levels of several plant species (7-9). In addition to their antioxidant capacity, phenolic compounds demonstrate a diverse array of physiological activities, encompassing antiallergenic, anti-inflammatory, antiarthrogenic, antibacterial, antithrombotic, vasodilatory and cardioprotective actions (10). Hence, the addition of antioxidant plant phenols to a food product may potentially confer a positive impact on human health.

Ziziphus oxyphylla also known as Indian jujube or Ber, belongs to the family Rhamnaceae (known as the buckthorn family) and is native to Southeast Asia and South Asia. The tree grows in scrublands, wastelands and dry deciduous woods. In Southeast and South Asia, it is found in India, Pakistan, Bangladesh and Sri Lanka. The plant has been utilized in traditional medicine for the treatment of many ailments i.e., diabetes (11), jaundice (12, 37), hypertension (13) as well as gas troubles (14) for a long.

Materials and Methods

Plant Material: *Ziziphus oxyphylla* was collected from Soon Valley, Khushab District, Punjab, Pakistan (Fig. 1). Our collection site was precisely located at N 32°30'05.7" and S 072°09'30.2", situated at an elevation of 856 m above sea level.



Fig. 1. Collection site (Soon Valley).

Furthermore, the plant was collected during fruiting season; the identified and scientific name of the plant specimen was cross-referenced using the World Flora Online (WFO) (<https://worldfloraonline.org>) that was deposited in the Herbarium of the Department of Botany, University of Sargodha, Pakistan under voucher number GA-640 for future reference.

Sample Preparation: Sun and shade-dried leaves were crushed into a fine powder, and this powder was used to make solutions for different assays. Using 5 g of plant materials (leaves and fruits), 2 types of solutions were made (100 % aqueous and 20 % ethanol and 80 % aqueous) for quantification of total phenolic content and antioxidants.

Plant material (5 g) from sun and shade dried leaves and fruits was added in 100 mL of solvent using different beakers. Solutions were thoroughly mixed with a vortex mixer and then left to rest for a period of 24 h. Then, this solution was filtered using Whatman filter paper no. 42, these solutions were used for further analysis.

For the proximate analysis, the fruit sample was dried in the shade for a period of 3 weeks; then, this sample was further dried using the hot air-drying method.

Determination of Total Phenolic Content: TPC was calculated using a slightly adjusted version of the Folin-Ciocalteu reagent technique (15). Total phenols were measured using ethanol with Gallic acid as a reference and were represented as (GAE) Gallic acid equivalents per 100 g of fresh weight sample. All experiments were carried out 3 times, whereas the experiments were conducted out twice.

Determination of total Flavonoid content: The approach given was used to compute FC (16). The Catechin was used as the standard; results were represented as mg of (+) -catechin equivalent per 100 g of fresh weight. All experiments were conducted in triplicate as were all determinations.

Determination of Total Antioxidants: The technique was employed to assess the overall antioxidant activity in water-solubilized samples (17). The results are presented both as micrograms of Ascorbic acid equivalent per mL of pulp and as mol TE (Trolox equivalent) per mL of pulp, utilizing AA (Ascorbic acid) for generating standard calibration curves. The process was repeated 3 times (in triplicate) to ensure accuracy in all computations and experiments.

Proximate analysis: Proximate analysis was conducted in IFSN (Institute of Food Science and Nutrition) at the University of Sargodha. This analysis encompassed proximate factors, including crude protein, crude fat, moisture, ash and NFE, crude fiber, ash and NFE, by following the procedure outlined by (18). All the values obtained were converted into %.

Results and Discussion

Phytochemical screening of *Zizyphoxyphylla* yielded the following results. Shade-dried aqueous extracts of leaves

yielded the highest TPC, 599 ± 1.00 mg/100 g GAE (where GAE stands for Gallic acid equivalent), followed by sun-dried aqueous extracts 589 ± 1.00 mg/100 g GAE, sun-dried and shade dried ethanol extracts showed minimum amount of TPC 367 ± 1.00 mg/100 g GAE and 281 ± 1.00 mg/100 g GAE respectively (Table 1).

Table 1. Mean values of Total phenolic contents (leaves)

Solvent	Drying method		Mean
	Sun dry	Shade	
Ethanol	367 ± 1.00^C	281 ± 1.00^D	324 ± 47.11^B
Aqueous	589 ± 1.00^B	599 ± 1.00^A	594 ± 5.55^A
Mean	478 ± 121.60^A	440 ± 174.18^B	

Units= mg/100 g GAE (GAE stands for Gallic acid equivalent). Significantly different mean values do not share a letter.

The inherent characteristics of phenolic compounds present challenges in devising a comprehensive extraction process (19). The polarity and characteristics of the solvent employed have an impact on the solubility of phenolic compounds (20).

The highest flavonoid content was observed in sun-dried ethanol extracts, 496.296 ± 3.70 mg/100 g CE (where CE stands for catechin equivalent), followed by shade-dried ethanol extracts at 451.852 ± 3.70 mg/100 g CE. Sun and shade-dried aqueous extracts showed minimum flavonoid content of 177.778 ± 3.70 mg/100 g CE and 159.259 ± 48.15 mg/100 g CE (Table 2). The flavonoid compositions of *Zizyphoxyphylla* exhibit similarities to those found in certain plants, such as *Artemisia campestris* and *A. arvensis* belonging to the same family as reported $\alpha\beta\gamma$

Table 2. Mean of flavonoids (leaves)

Solvent	Drying method		Mean
	Sun dry	Shade	
Ethanol	496.296 ± 3.70^A	451.852 ± 3.70^A	474.074 ± 24.57^A
Aqueous	177.778 ± 3.70^B	159.259 ± 48.15^B	168.519 ± 32.18^B
Mean	337.037 ± 174.49^A	305.556 ± 163.14^A	

Units= mg/100 g CE (CE stands for catechin equivalent). Significantly different mean values do not share a letter.

Sun-dried ethanol extract 3711.35 ± 2.70 μ g/100 g AAE (where AAE stands for ascorbic acid equivalent) exhibits a higher level of antioxidants compared to the shade-dried extract 1443.78 ± 2.70 μ g/100 g AAE. Likewise, the sun-dried aqueous extract 597.84 ± 7.15 μ g/100 g AAE demonstrates a significantly higher antioxidant value compared to the shade-dried aqueous extract 462.70 ± 2.70 μ g/100 g AAE (Table 3).

Ethanol extracts, whether from shade or sun-dried samples, display greater antioxidant activity when contrasted with both sun and shade-dried aqueous extracts.

Previous research conducted by (22, 23) demonstrated that the choice of solvent used for extraction significantly affects the antioxidant capacity of

Table 3. Mean of antioxidants (leaves)

Solvent	Drying method		Mean
	Sun dry	shade	
Ethanol	3711.35 ± 2.70 ^A	1443.78 ± 2.70 ^B	2577.57 ± 1242.00 ^A
Aqueous	597.84 ± 7.15 ^C	462.70 ± 2.70 ^D	530.27 ± 74.17 ^B
Mean	2154 ± 1705.35 ^A	953.24 ± 537.37 ^B	

Units= µg/100 g AA (AAE stand for ascorbic acid equivalent). Significantly different mean values do not share a letter.

the extracts. These findings align with the results obtained in the present study. The observed differences in antioxidant activity may be attributed to the varying levels of polyphenols in each solvent, as suggested by (24, 25). However, it is important to note that the quality of polyphenols, flavonoids and tannins also contributes to this variation. The utilization of solvents possessing varying polarity enables the extraction of a specific subset of antioxidants, hence influencing the estimation of antioxidant capacity (26).

Total phenolic contents in the fruit of *Z. oxyphylla* quantified as gallic acid equivalent are presented in Table 4. The total phenolic content of the fruit sample was measured to be 758.000 mg/100 g GAE for aqueous extracts and 739.333 mg/100 g GAE for ethanol extracts. Antioxidant content for fruit was measured to be 405.045 µg/100 g AAE for aqueous extracts and 329.369 µg/100 g AAE. The inherent characteristics of phenolic compounds present challenges in devising a comprehensive extraction process (19). Similar reports are on many bioactive compounds in the methanolic extract of *Artocarpus hirsutus* leaves (21). The polarity and characteristics of the solvent employed have an impact on the solubility of phenolic compounds (20).

Table 4. Means for TPC and antioxidants (Fruit)

Treatment	TPC	Total Antioxidants
Aqueous	758.000 ^A	405.045 ^A
Ethanol	739.333 ^B	329.369 ^B

Units= mg/100g GAE (GAE stands for Gallic acid equivalent) for TPC and µg/100g AA (AAE stand for ascorbic acid equivalent) for antioxidants. Means that do not share a letter are significantly different

Proximate analysis of fruit: The moisture content of *Z. oxyphylla* was measured to be 71.9333 ± 0.0208 % This moisture content found in *Z. oxyphylla* (11.9333 ± 0.0208 %) close to some common leafy vegetables like *Xanthosoma sagittifolium* and *Adansonia digitata* with 13.2 % and 9.50 % respectively (27). The fruit sample consisted of 8.2466 ± 0.0208 % fat; the findings of this study closely align with the fat content (28), who examined “nutrient value of North East Indian plant foods” and (29) with the topic “nutrient composition of less familiar leaves consumed by the tribals of Udaipur region.” Fat plays a vital role in facilitating the absorption of essential vitamins like A, B, E and K, making it a valuable energy source (30). Crude fiber % was found to be 5.03 ± 0.02 %. The findings from this study are corroborated by the results obtained in the research conducted earliest (31). Ash content was 5.6666 ± 0.1527 %, (32) considered the ash content, which ranged from 1.91 % to 8.73 %, to be rich. Results revealed that crude protein content was 2.0366 ± 0.0152 %. The present study benefits from the support of prior research conducted by (33-35), which validated the current outcomes. The NFE (Nitrogen-Free Extract) content of *Z. oxyphylla* was measured at 79.2333 ± 0.0208 % (Fig. 2), which is similar and can be compared to the carbohydrate content found in other fruits like *Morus alba*, *Mangifera indica*, *Prunus nepalensis*, *P. cerasoides* and *Terminalia chebulaz* which possess carbohydrate contents of 87.5 %, 84.0 %, 82.6 %, 84.1 % and 80.6 % respectively (32).




Fig. 2. Results of proximate analysis shown as %.

Table 5. Proximate analysis of *Z. oxyphylla* fruit

SAMPLE	Moisture %	Fat %	Fiber %	Ash %	Protein %	NFE %
<i>Ziziphus oxyphylla</i>	11.9333 ± 0.0208	8.2466 ± 0.02081	5.03 ± 0.02	5.6666 ± 0.1527	2.0366 ± 0.0152	79.233 ± 0.02081

Conclusion

In this study, total phenolic content, antioxidant activity of fruit and leaves, and proximate composition of fruit of *Ziziphus oxyphylla* were determined using aqueous and ethanol extracts. The leaves and fruits of this plant contain a high amount of total phenolic content, considerable total flavonoid content, antioxidant activity and high nutritional value. The results obtained from this study, which focused on the total phenolic contents and proximate compositions of *Z. oxyphylla*, can serve as a foundation for further investigations into its antioxidant capabilities. These research endeavours will help improve the National Food Composition Database in the country.

Acknowledgments

The manuscript is part of the M. Phil dissertation of the second author (Javaria Nargis). The authors are thankful to the Higher Education Commission of Pakistan (HEC) (Programme No. 20-4546/ R&D/NRPU/HEC/14/850) for financial, technical and practical support. The authors are also indebted to the laboratory of the Institute of Food Sciences and Nutrition (IFSC) for cooperation and provision of required facilities. The plant was collected from the study area (Soon Valley) and was deposited in the Herbarium of Department of Botany, University of Sargodha that was established with the financial support of ORIC Project No. UOS/ORIC/2016/43 University of Sargodha, Sargodha, Pakistan.

Authors' contributions

The manuscript is written by JN as M.Phil scholar. SR (as M.Phil supervisor) and MN (as M.Phil co-supervisor) helped in the tabulation of data. The rest of the authors AS, MK, GS, TT, UZ, ABS and NR helped in the finalization of the paper. All the authors approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None.

References

- Tanaka T. *Cyclopedia of edible plants of the World*. Keigaku Publishing Company; 1976.
- Facciola S. *Cornucopia: A source book of edible plants*. Kampong Publications; 1990.
- Moerman DE. *Native American ethnobotany*; 1998.
- Hou W-C, Lin R-D, Cheng K-T, Hung Y-T, Cho C-H, Chen C-H *et al*. Free radical-scavenging activity of Taiwanese native plants. *Phytomed*. 2003;10(2-3):170-75. <https://doi.org/10.1078/094471103321659898>
- Yanishlieva NV, Marinova E, Pokorný J. Natural antioxidants from herbs and spices. *Eur J Lipid Sci Technol*. 2006;108(9):776-93. <https://doi.org/10.1002/ejlt.200600127>
- Pietta P, Simonetti P, Mauri P. Antioxidant activity of selected medicinal plants. *J. of Agric Food Chemi*. 1998;46(11):4487-90. <https://doi.org/10.1021/jf980310p>
- Zheng W, Wang SY. Antioxidant activity and phenolic compounds in selected herbs. *J Agric Food Chem*. 2001;49(11):5165-70. <https://doi.org/10.1021/jf010697n>
- Silva E, Souza J, Rogez H, Rees J-F, Larondelle Y. Antioxidant activities and polyphenolic contents of fifteen selected plant species from the Amazonian region. *Food Chem*. 2007;101(3):1012-18. <https://doi.org/10.1016/j.foodchem.2006.02.055>
- Tawaha K, Alali FQ, Gharaibeh M, Mohammad M, El-Elimat T. Antioxidant activity and total phenolic content of selected Jordanian plant species. *Food Chem*. 2007;104(4):1372-78. <https://doi.org/10.1016/j.foodchem.2007.01.064>
- Balasundram N, Sundram K, Samman S. Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence and potential uses. *Food Chem*. 2006;99(1):191-203. <https://doi.org/10.1016/j.foodchem.2005.07.042>
- Sher H. Ethnoecological evaluation of some medicinal and aromatic plants of Kot Malakand agency, Pakistan. *Scientific Research and Essays*. 2011;6(10):2164-73. <https://doi.org/10.5897/SRE10.1114>
- Jan G, Khan MA, Gul F. Ethnomedicinal plants used against jaundice in Dir Kohistan valleys (NWFP), Pakistan. *Ethnobotanical Leaflets*. 2009;2009(8):7. <https://opensiuc.lib.siu.edu/ebl/vol2009/iss8/7>
- Khan MA, Khan SA, Qureshi MA, Ahmed G, Khan MA, Hussain M *et al*. Ethnobotany of some useful plants of Poonch valley Azad Kashmir. *J Med Plants Res*. 2011;5(26):6140-51. <https://doi.org/10.5897/JMPR11.045>
- Sher Z, Khan Z, Hussain F. Ethnobotanical studies of some plants of Chagharzai Valley, district Buner, Pakistan. *Pak J Bot*. 2011;43(3):1445-52. [http://www.pakbs.org/pjbot/PDFs/43\(3\)](http://www.pakbs.org/pjbot/PDFs/43(3))
- Vi S. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*. 1999;299:152-78. [https://doi.org/10.1016/S0076-6879\(99\)99017-1](https://doi.org/10.1016/S0076-6879(99)99017-1)
- Zhishen J, Mengcheng T, Jianming W. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem*. 1999;64(4):555-59. [https://doi.org/10.1016/S0308-8146\(98\)00102-2](https://doi.org/10.1016/S0308-8146(98)00102-2)
- Prieto P, Pineda M, Aguilar M. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Analytical Biochem*. 1999;269(2):337-41. <https://doi.org/10.1006/abio.1999.4019>
- AOAC. *Association of official analytical chemist. Official Methods of Analysis*. AOAC Gaithersburg, MD, USA; 2000.
- Vercauteren J, Chèze C, Triaud J. *Polyphenols 96: 18th International Conference on Polyphenols, Bordeaux (France)*. July 15-18, 1996: Editions Quae; 1998.
- Naczek M, Shahidi F. Extraction and analysis of phenolics in food. *J Chromatogr Sci A*. 2004;1054(1-2):95-111. <https://doi.org/10.1016/j.chroma.2004.08.059>
- Djeridane A, Yousfi M, Nadjemi B, Boutassouna D, Stocker P, Vidal N. Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. *Food Chem*. 2006;97(4):654-60. <https://doi.org/10.1016/j.foodchem.2005.04.028>
- Gülçin İ, Oktay M, Kireççi E, Küfrevioğlu Öİ. Screening of antioxidant and antimicrobial activities of Anise (*Pimpinella anisum* L.) seed extracts. *Food Chem*. 2005;97(1):8-12. [https://doi.org/10.1016/S0308-8146\(03\)00098-0](https://doi.org/10.1016/S0308-8146(03)00098-0)
- Tepe B, Daferera D, Sokmen A, Sokmen M, Polissiou M. Antimicrobial and antioxidant activities of the essential oil and various extracts of *Salvia tomentosa* Miller (Lamiaceae). *Food Chem*. 2005;90(3):333-40. <https://doi.org/10.1016/j.foodchem.2003.09.013>

24. Jayaprakasha G, Patil BS. *In vitro* evaluation of the antioxidant activities in fruit extracts from citron and blood orange. *Food Chem.* 2007;101(1):410-18. <https://doi.org/10.1016/j.foodchem.2005.12.038>
25. Hodzic Z, Pasalic H, Memisevic A, Srabovic M, Saletovic M, Poljakovic M. The influence of total phenols content on antioxidant capacity in the whole grain extracts. *Eur J Sci Res.* 2009;28(3):471-77.
26. Zhou K, Yu L. Effects of extraction solvent on wheat bran antioxidant activity estimation. *LWT-Food Science and Technology.* 2004;37(7):717-21. <https://doi.org/10.1016/j.lwt.2004.02.008>
27. Ladan M, Bilbis L, Lawal M. Nutrient composition of some green leafy vegetables consumed in'sokoto; 1996. <http://hdl.handle.net/123456789/259>
28. Longvah T. Nutrient value of North East India plant foods. *Nutrition News, NIN.* 2002;21.
29. Sankhala A, Sankha A, Bhatnagar B, Singh A. Nutrient composition of less familiar leaves consumed by the tribals of Udaipur region. *J Food Sci Technol.* 2005;42(5):446-48.
30. Wakili A, Abdullahi M, Madara M. Proximate composition of five commonly used horticultural products in Northern Nigeria. *Int J Curr Microbiol Appl Sci.* 2015;4(8):924-28. <http://www.ijcmas.com/vol-4-8/A.%20Wa>
31. Rathore M. Nutrient content of important fruit trees from arid zone of Rajasthan. *J Hortic.* 2009;1(7):103-08.
32. Seal T. Nutritional composition of wild edible fruits in Meghalaya state of India and their ethno-botanical importance. *J Bot.* 2011;6(2):58-67. <https://doi.org/10.3923/rjb.2011.58.67>
33. Ogle BM, Grivetti LE. Legacy of the chameleon: Edible wild plants in the kingdom of Swaziland, Southern Africa. A cultural, ecological, nutritional study. Part IV-Nutritional analysis and conclusions. *Ecology of Food and Nutrition.* 1985;17(1):41-64. <https://doi.org/10.1080/03670244.1985.9990881>
34. French B. Food composition tables for food plants in Papua New Guinea. West St Burnie, Tasmania; 2006.
35. Mahapatra AK, Mishra S, Basak UC, Panda PC. Nutrient analysis of some selected wild edible fruits of deciduous forests of India: An explorative study towards non conventional bio-nutrition. *Adv J Food Sci Technol.* 2012;4(1):15-21.
36. Patel J, Kumar GS Satyasai D. GC-MS analysis of bioactive components on the leaves extract of *Artocarpus hirsutus*: A potential folklore medicinal plant. *Int J Herb Med.* 2016;4(6):95-97.
37. Patel J, Kumar GS, Ahirwar K, Gupta MK, Singh SK, Chandel SS, Patel VK. Comparative analysis in hepatoprotective activity of crude extracts of important medicinal plants. *Research Journal of Pharmacy and Technology.* 2023;16(2):659-62. <https://doi.org/10.52711/0974-360X.2023.00112>