



RESEARCH COMMUNICATION

# Chemical components and antioxidant properties of acetone extracts of four different species of Zingiberaceae

Thi Hoi Huynh<sup>1</sup>, Gia Hoa Tran<sup>1</sup>, Ngoc Nam Trinh<sup>2</sup>, Quoc Hung Nguyen<sup>3</sup>, Thien Hoang Ho<sup>1</sup>, Le Pham Tan Quoc<sup>1</sup>, Nguyen Tuong An Huynh<sup>1</sup> & Hong Thien Van<sup>1\*</sup>

<sup>1</sup>Institute of Biotechnology and Food Technology, Industrial University of Ho Chi Minh City, No. 12 Nguyen Van Bao Street, Go Vap District, Ho Chi Minh City, Vietnam

<sup>2</sup>Office of Science Management and International Affairs, Industrial University of Ho Chi Minh City, No. 12 Nguyen Van Bao Street, Go Vap District, Ho Chi Minh City, Vietnam

<sup>3</sup>Centre of Analytical Services and Experimentation HCMC, Vietnam, 2 Nguyen Van Thu, Dakao Ward, District 1, Ho Chi Minh City, Vietnam

\*Email: [vanhongthien@iuh.edu.vn](mailto:vanhongthien@iuh.edu.vn)



## ARTICLE HISTORY

Received: 10 January 2024

Accepted: 24 January 2024

Available online

Version 1.0 : 27 March 2024



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Huynh TH, Tran GH, Trinh NN, Nguyen QH, Hoang Ho T, Quoc LPT, Huynh NTA, Van HT. Chemical components and antioxidant properties of acetone extracts of four different species of Zingiberaceae. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.3272>

## Abstract

In this report, the chemical components and antioxidant properties of the acetone extracts obtained from 4 Zingiberaceae species, including *Curcuma gracillima* Gagnep., *Curcuma pierreana* Gagnep., *Globba marantina* L. and *Amomum biflorum* Jack were investigated for the first time. A total of 142 chemical compounds were determined in 4 studied extracts using GC/MS techniques. Furthermore, the extract obtained from *C. gracillima* showed the strongest antioxidant effect with an IC<sub>50</sub> value of 1.76 mg/mL, followed by *G. marantina* (IC<sub>50</sub> = 2.45 mg/mL), *C. pierreana* (IC<sub>50</sub> = 2.97 mg/mL) and *A. biflorum* (IC<sub>50</sub> = 6.32 mg/mL).

## Keywords

Antioxidant; chemical compounds; GC/MS techniques; zingiberaceae

## Introduction

Zingiberaceae, the largest family in the order Zingiberales, includes around 1200 species belonging to 53 genera (1) being distributed mainly in subtropical and tropical zones, especially in Southeast Asia. Most members of this family are medicinal plants and are widely used in indigenous medicine and the pharmaceutical industry because of their good biological properties (2). In addition, the Zingiberaceae species are also used for dye and perfume (3), food, spices, ornamental plants and cosmetics, etc (4).

Vietnam is one of the countries with a high diversity of Zingiberaceae species. In 2000, Pham-Hoang recorded 24 genera and 113 species belonging to the Zingiberaceae family for the flora of Vietnam (5). In 2017, Nguyen in the book "Flora of Vietnam" reported 19 genera and 140 species of Zingiberaceae in Vietnam (6). Notably, a large number of plants belonging to this family have been discovered as new species for the flora of Vietnam by recent publications (7-11).

Binh Chau-Phuoc Buu is a large Nature Reserve with about 10451 ha located in the lowland dry forests in southern Vietnam (12). This Nature Reserve was established in 1996 and had about 732 vascular plants belonging to 113 genera and 63 families that were used as medicinal plants (12, 13). Recently, we have conducted some field trips to survey the biodiversity in Binh Chau-Phuoc Buu Nature Reserve, Ba Ria-Vung Tau province, and collected 4 species of Zingiberaceae family, including *Curcuma gracillima* Gagnep., *Curcuma pierreana* Gagnep., *Globba marantina* L. and *Amomum biflorum* Jack. Among 4 studied species, except

for *G. marantina*, an ethnomedicinal plant used in some Asian countries, the remaining three species are rare and they do not have any information about their use in traditional medicine. Accordingly, indigenous medicine in India has used *G. marantina* to treat some common diseases such as rheumatism, conjunctivitis, snake bite, cough, leucoderma, cold, asthma, etc (14, 15). Thus, in the current study, the chemical components and the antioxidant properties of the acetone extracts from these species were firstly reported.

## Materials and Methods

### Plant materials and taxonomy

The whole plants of 4 studied species, namely *C. gracillima*, *C. pierreana*, *G. marantina*, and *A. biflorum* were collected from Binh Chau-Phuoc Buu Nature Reserve from June to August 2023 by Mr. Van Son Le. The voucher specimens (Le VS 982-985) were deposited in the herbarium of this Nature Reserve. The process of collecting samples and processing specimens was carried out according to the instructions Kew Botanic Garden protocol (16). Comparative morphology was used to identify the scientific names of studied species based on the prior publications (5, 6).

### Extraction procedures

The whole plants of each species were dried at 50 °C and then ground into powder. 500 mL of acetone solution (Thermo Fisher Scientific, USA) was used to soak 100 g of sample powder for 72 h at room temperature. The soaking solution was filtered through the Whatman filter paper. The experiment was repeated two more times with the filter residue. The filtrate was evaporated on a vacuum evaporator at 45 °C and then the evaporation fluid was dried to completely remove acetone (17).

### Gas chromatography-mass spectrometry (GC/MS) analysis

The chemical components of the studied sample were determined using the gas chromatograph GC TRACE 1310 coupled with a single quadrupole mass spectrometer ISQ 7000 (Thermo Fisher Scientific Inc., Waltham, MA, USA) and the DB-5MS column (30 m x 0,25 mm x 0,25 μm) with a standard procedure (17, 18).


### Determination of the antioxidant activity of the extracts

The free radical scavenging property of the 4 studied extracts was performed by DPPH radical scavenging assay. The procedure was also performed (17, 19).

## Results

### Chemical compounds of acetone extracts from four studied species.

The chemical constituents of the acetone extracts isolated from 4 studied species are presented in Fig. 1 and Table 1. The major compounds of *G. marantina* extract were composed of n-hexadecanoic acid (13.42%), neophytadiene (7.35%), (*E*)-labda-8(17),12-diene-15,16-dial (6.65%), caryophyllene oxide (6.36%) and β-caryophyllene (5.07%). The *C. gracillima* extract was found to be rich in n-hexadecanoic acid (14.03%), β-caryophyllene (11.76%), (*E*)-labda-8(17),12-diene-15,16-dial (9.88%), caryophyllene oxide (8.81%), neophytadiene (8.78%) and linolenic acid (8.69%). The most abundant component of *C. pierreana* extract was n-hexadecanoic acid (12.07%), followed by (*E*)-labda-8(17),12-diene-15,16-dial (11.97%), cis-13-octadecenoic acid (6.86%), γ-bicyclohomofarnesal (6.11%), octadecanoic acid (5.34%), 9*E*,11*E*-octadecadienoic acid (4.97%) and caryophyllene oxide (3.78%) whereas (-)-alcanfor (14.12%), reynosin (8.74%), neophytadiene (8.31%) and n-hexadecanoic acid (6.75%), eucalyptol (4.24%) was the major compounds of the *A. biflorum* extract.



**Fig. 1.** The GC chromatogram of four studied extracts. TR5: *Globba marantina*, TR7: *Curcuma gracillima*, TR8: *Curcuma pierreana*, and TR10: *Amomum biflorum*.

**Table 1.** The relative percentage of chemical compounds of four studied extracts

No.	Compounds	The relative percentage (%)			
		TR5	TR7	TR8	TR10
1	2-Pentanone, 4-hydroxy-4-methyl-	2.50	1.27	9.70	4.30
2	Ethanone, 1-(3-methylenecyclopentyl)-	-	0.10	0.10	0.07
3	3-Methylcyclopentyl acetate	0.09	0.10	0.13	0.07
4	2-Heptanol, acetate	-	-	0.22	-
5	Benzene, (1-methylethyl)-	0.15	0.08	0.12	-
6	$\alpha$ -Pinene	-	0.14	0.23	-
7	Benzene, propyl	0.14	0.08	-	-
8	Camphene	-	-	0.09	0.25
9	Toluene, m-ethyl-	0.39	0.14	0.10	-
10	Benzene, 1,2,3-trimethyl-	0.17	0.19	-	0.12
11	Mesitylene	-	0.12	0.23	-
12	$\beta$ -Pinene	0.77	0.91	2.79	0.13
13	Benzene, 1,2,4-trimethyl-	0.49	0.06	-	-
14	o-Cymene	-	-	0.08	-
15	D-sylvestrene	-	-	0.13	-
16	Limonene	-	-	-	0.04
17	Eucalyptol	-	-	0.11	4.24
18	3-Carene	-	-	0.07	-
19	Pyrazine, tetramethyl-	-	-	0.09	-
20	2-Norbornanone	-	-	-	0.05
21	Linalool	0.25	-	-	0.24
22	rans-p-Mentha-2,8-dienol	-	-	-	0.12
23	$\alpha$ -Campholenal	-	-	0.16	-
24	1,7,7-Trimethylbicyclo[2.2.1]hept-5-en-2-ol	-	-	-	0.10
25	(1R)-(+)-Nopinone	0.13	0.09	0.47	-
26	(-)-Alcanfor	0.18	0.27	0.41	14.12
27	Camphene hydrate	-	-	-	0.24
28	Sabinone	-	-	0.52	0.45
29	Terpinen-4-ol	-	-	-	0.50
30	Myrtenal	0.20	0.14	0.76	0.69
31	Levoverbenone	-	-	-	0.30
32	Ethylmethylmaleimide	0.25	0.23	0.65	0.17
33	(-)-Carvone	-	-	-	0.15
34	p-Mentha-1,8-dien-3-one	-	-	-	0.16
35	Bornyl acetate	-	-	0.14	0.29
36	(-)-Neoclovene-(I), dihydro	0.20	-	-	-
37	Bicyclo[2.2.1]heptane-2-carboxylic acid, 3,3-dimethyl	-	-	-	0.30
38	Spiro[2.7]dec-4-ene, 1,1,5,6,6,9,9-heptamethyl-10-methylene	-	-	-	0.11
39	Bicyclo[4.1.0]hept-3-en-2-one, 4,7,7-trimethyl-	-	-	-	0.09
40	$\alpha$ -Cubebene	-	0.11	-	0.08
41	1,3,5,7-Tetramethyl-adamantane	0.19	-	-	-
42	Cypera-2,4-diene	0.39	-	-	-
43	Copaene	-	0.37	-	-
44	$\beta$ -elemene	0.35	-	-	0.35
45	9,9-Dimethyl-2-oxabicyclo[4.2.1]nonan-3-one	-	-	-	0.84
46	p-Cymene, 2,5-dimethoxy	-	-	0.19	-
47	$\alpha$ -Gurgujene	0.80	-	-	-
48	$\beta$ -Caryophyllene	5.07	11.76	1.08	-
49	Isocaryophyllene	-	-	-	0.43
50	Geranylacetone	-	-	-	0.25
51	(E)- $\beta$ -Famesene	0.14	-	-	-
52	Tetradecane, 2,6,10-trimethyl	0.62	-	-	-
53	Humulene	0.61	-	-	-
54	1,1,4,8-tetramethyl-cis,cis,4,7,10-cycloundecatriene	-	1.50	0.39	-
55	Benzene, 1-(1,5-dimethyl-4-hexenyl)-4-methyl	-	0.45	-	-
56	Aristolochene	-	-	-	0.81
57	$\beta$ -Selinene	0.51	-	-	1.36
58	$\alpha$ -Selinene	-	-	-	0.32
59	$\beta$ -Bisabolene	1.21	-	-	-
60	$\beta$ -Curcumene	-	0.94	-	-
61	Isogermacrene D	-	-	0.28	-
62	Cadina-1(10),4-diene	-	0.38	-	0.28
63	Guaia-10(14),11-diene	0.33	-	-	-
64	$\alpha$ -Guaiene	0.29	-	-	-

65	Dihydroactinolide	0.34	0.37	0.25	0.25
66	3,5,9-Trimethyl-deca-2,4,8-trien-1-ol	0.32	-	-	-
67	2(1H)-Naphthalenone, octahydro-4a,7,7-trimethyl-, cis	-	-	0.23	-
68	Dodecanoic acid	-	-	0.19	-
69	Nerolidol	-	0.31	0.45	0.37
70	Caryophyllene oxide	6.36	8.81	3.78	0.37
71	4,8,12-trimethyltrideca-1,3,7,11-tetraene	0.21	-	-	0.39
72	$\alpha$ -Costal	-	-	-	1.82
73	(-)-Spathulenol	-	-	-	1.00
74	Humulene-1,2-epoxide	0.64	1.03	0.31	-
75	Methyl (3-oxo-2-[(2Z)-2-pentenyl]cyclopentyl)acetate	2.15	-	-	-
76	Isospathuleno	-	-	-	0.66
77	Cyclolongifolene oxide, dehydro	-	0.39	-	-
78	Pogostole	-	-	0.50	-
79	Neointermedeo	-	-	-	1.06
80	Globulol	0.58	-	-	-
81	Mustakone	-	0.63	-	-
82	(+)- $\beta$ -Costol	-	-	-	1.02
83	Longifolenaldehyde	0.85	-	0.73	-
84	Pentadecanal	-	-	0.30	-
85	2-Phytene, isomer 1	-	-	0.43	-
86	Isovalencenyl formate	0.46	-	-	-
87	1,1,4,7-Tetramethyldecahydro-1H-cyclopropa[e]azulene-4,7-diol	-	-	-	0.63
88	Tetradecanoic acid	0.35	-	0.60	-
89	Aromadendrene oxide-(1)	-	-	-	0.89
90	3,7,11-Trimethyl-dodeca-2,4,6,10-tetraenal	-	-	-	0.76
91	trans-Methylisocosticate	-	-	-	1.20
92	Neophytadiene	7.35	8.78	1.36	8.31
93	$\gamma$ -Bicyclohomofarnesal	-	-	6.11	-
94	2-Pentadecanone, 6,10,14-trimethyl-	3.33	-	3.49	1.33
95	2-Hexadecen-1-ol, 3,7,11,15-tetramethyl	1.45	2.48	-	-
96	Valeric acid	-	-	-	1.29
97	Farnesyl acetone	-	-	-	1.41
98	2-Dodecen-1-yl(-)succinic anhydride	1.07	-	-	-
99	n-Hexadecanoic acid	13.42	14.03	12.07	6.75
100	(E)-15,16-Dinorlabda-8(17),11-dien-13-one	-	-	0.45	0.97
101	Z-(13,14-Epoxy)tetradec-11-en-1-ol acetate	0.64	-	-	-
102	trans-Geranylgeraniol	1.25	-	-	-
103	Shyobunol	-	-	0.68	-
104	Heptadecanoic acid	0.80	0.69	1.12	1.06
105	(E)-15,16-Bisnorlabda-8(17),12-diene-14-al	1.56	-	1.06	-
106	Naphthalene, decahydro-1,1,4a-trimethyl-5-(3-methylbutyl)-6-methylene-, (4aS,5S,8aS)	-	0.64	0.85	-
107	3,7,11,15-Tetramethyl-1-hexadecyn-3-ol	3.80	-	-	-
108	2,6-Dodecadien-1-ol, 3,7,11-trimethyl-, (Z,E)-	-	-	0.94	-
109	9E,11E-octadecadienoic acid	2.03	-	4.97	-
110	10E,12Z-octadecadienoic acid	-	-	-	1.97
111	Linolenic acid	1.81	8.69	-	-
112	cis-13-Octadecenoic acid	-	-	6.86	-
113	cis-Vaccenic acid	-	-	-	3.80
114	7-Hydroxy-6,9a-dimethyl-3-methylene-decahydro-azuleno[4,5-b]furan-2,9-dione	-	-	-	3.16
115	Octadecanoic acid	2.42	3.68	5.34	2.35
116	2,6-Di-tert-butylbenzoquinone	-	-	-	1.92
117	Androst-4-en-3-one, 17-methyl	-	0.93	-	-
118	7 $\beta$ -Hydroxydehydroepiandrosterone	-	-	1.35	-
119	Ambria	-	-	0.80	-
120	Reynosin	-	-	-	8.74
121	Eicosane	-	1.07	-	-
122	Ambrosin	-	-	-	2.52
123	Androst-5-en-3-one, 4,4-dimethyl	-	-	1.48	-
124	4,8,12,16-Tetramethylheptadecan-4-olide	0.97	-	1.06	-
125	(E)-labda-8(17),12-diene-15,16-dial	6.65	9.88	11.97	1.55
126	Androst-4-en-3-one, 17-(1-oxo-3-phenylpropoxy)-, (17 $\beta$ )-	1.12	-	-	-
127	Eicosanal	-	-	1.26	-
128	Pentacosane	1.43	2.10	-	-
129	Pentacosanal	-	-	0.97	1.68
130	2H-Pyran, 2-(7-heptadecynyloxy)tetrahydro	1.88	-	-	-
131	Pregn-20-yn-17-ol, (5 $\beta$ ,17 $\alpha$ )-	-	2.11	-	-
132	Villosin	1.40	1.14	3.81	-
133	1-Heptatriacotanol	1.81	6.71	-	-
134	Tetracosanoic acid	-	-	1.52	-
135	13-Docosenamido, (Z)-	-	-	1.12	0.77
136	Squalene	-	4.23	-	-
137	2-(3-Hydroxy-4-methoxyphenyl)-3,7-dimethoxy-4H-chromen-4-one	-	-	-	3.17
138	4',7-Dimethoxyflavonol	-	-	-	3.14
139	Cholesta-4,6-dien-3-ol, (3 $\beta$ )-	5.55	-	-	-
140	Tetratriacontane	-	-	-	1.45
141	Stigmasta-3,5-dien-7-one	3.11	-	-	-
142	$\gamma$ -Sitostenone	3.95	-	-	-
	Total	97.48	98.13	97.70	99.78

Note: TR5: *Globba marantina*, TR7: *Curcuma gracillima*, TR8: *Curcuma pierreana*, and TR10: *Amomum biflorum*

### Antioxidant activity of acetone extracts from four studied species

The DPPH radical scavenging agents of 4 studied extracts are presented in Table 2. Accordingly, the extract obtained from *C. gracillima* possessed the highest antioxidant property (IC<sub>50</sub> value of 1.76 mg/mL) compared to the other 3 extracts, including *G. marantina* (IC<sub>50</sub> = 2.45 mg/mL), *C. pierreana* (IC<sub>50</sub> = 2.97 mg/mL) and *A. biflorum* (IC<sub>50</sub> = 6.32 mg/mL).

### Discussion

The chemical compounds of the acetone extracts isolated from *G. marantina*, *C. gracillima*, *C. pierreana*, and *A. biflorum* were first presented in this study. However, their chemical constituents in the essential oils and other solvent extractions obtained from these species using GC/MS have been shown by prior studies. For instance, the chemical components of the essential oils (EOs) from *G. marantina* contained β-caryophyllene (19.30%), α-humulene (14.20%), (Z)-nerolidol (7.50%) and isoborneol (7.30%) as the main compounds (20) whereas the ethanolic extract of this species was found to be rich in pinocarvone (54.27%), β-caryophyllene (13.65%), terpineol (6.31%) and heptadecane (6.19%) (21). Furthermore, the EO isolated from *C. pierreana* was reported to contain camphene (18.82%), 3-carene (14.13%), β-caryophyllene (10.76%) and fenchyl acetate (10.60%) (22). The chemical compositions of the aerial part of the *C. gracillima* EOs were characterized by the predominance of caryophyllene (19.8%), caryophyllene oxide (13.7%), and 3-carene (9.3%) while the rhizome oil contained α-curcumin (18.9%), 3,4-dimethyl anisole (17.6%) and 2-bornanone (8.6%) as the main components (23). The chemical compounds of the EO obtained from *Amomum biflorum* were reported by the dominance of camphor (17.60%), α-bisabolol (16.00%), camphene (8.20%), and α-humulene (5.10%) (24).

Many bioactive compounds contained from 4 studied extracts in this study have been also found in Zingiberaceae species. Accordingly, (*E*)-labda-8(17),12-diene-15,16-dial was reported as a compound obtained from the extracts of various Zingiberaceae plants (25-28). This component has been described to have cytotoxic properties against some cancer cell lines in humans (29). It has also been shown that this compound was used as an agent to treat fungal infections, α-glucosidase inhibition, or as an antibiotic (30, 31). Also, (*E*)-labda-8(17),12-diene-15,16-dial isolated from *Curcuma amada* extract showed potent antibacterial activity against *Escherichia coli* (26). β-caryophyllene has also been shown as a bioactive component to have anticancer, antioxidant, antimicrobial,

re-epithelialization, and anti-inflammatory activities (32-35). Prior studies demonstrated that *n*-hexadecanoic acid also possessed cytotoxic (36), antibacterial and antioxidant (37), and anti-inflammatory (38) properties. Moreover, caryophyllene oxide has been announced to possess anti-inflammatory, analgesic, (39) antifungal (40), and cytotoxic (41) activities. The prior study showed that neophytadiene exhibited an anti-inflammatory effect in Sprague dawley rats and RAW 264.7 macrophages (42). This compound was also shown as an agent in neuropharmacological activity, a type of anticonvulsant, antidepressant, sedative, and anxiolytic (43).

### Conclusion

The present study firstly showed the chemical compositions and antioxidant activity of the acetone extracts isolated from 4 Zingiberaceae species in which the *C. gracillima* extract was found to be rich in *n*-hexadecanoic acid (14.03%), β-caryophyllene (11.76%), (*E*)-labda-8(17),12-diene-15,16-dial (9.88%) while the most abundant component of *C. pierreana* extract was *n*-hexadecanoic acid (12.07%), followed by (*E*)-labda-8(17),12-diene-15,16-dial (11.97%), *cis*-13-octadecenoic acid (6.86%). The major compounds of *G. marantina* extract were composed of *n*-hexadecanoic acid (13.42%), neophytadiene (7.35%), (*E*)-labda-8(17),12-diene-15,16-dial (6.65%) while *Amomum biflorum* contains (-)-alcanfor (14.12%), reynosin (8.74%), neophytadiene (8.31%) as the main compounds. The acetone extracts from *C. gracillima*, *G. marantina*, *C. pierreana*, and *A. biflorum* showed antioxidant activity with IC<sub>50</sub> values of 1.76, 2.45, 2.97, and 6.32 mg/mL respectively.

### Acknowledgements

The authors would like to thank the Board of Directors Institute of Biotechnology and Food Technology, Industrial University of Ho Chi Minh City for their support. We also wish to thank Mr. Van Son Le, a staff scientist at Binh Chau-Phuoc Buu Nature Reserve, who collected the samples for this research.

### Authors' contributions

This research was designed and conducted by THH. This author was also the person who wrote the manuscript. All authors searched literature and revised the manuscript as well as contributed many valuable options. HTV was a person who revised the final version of the manuscript as well as was responsible for working with the Journal.

**Table 2.** DPPH radical scavenging activity of four studied extracts

	<i>G. marantina</i>	<i>C. gracillima</i>	<i>C. pierreana</i>	<i>A. biflorum</i>	Vitamin C
IC <sub>50</sub> (mg/mL)	2.45	1.76	2.97	6.32	0.0075

## Compliance with ethical standards

**Declaration :** Authors do not have any conflict of interest to declare.

**Ethical issues:** None.

## References

- Kress WJ, Prince LM, Williams KJ. The phylogeny and a new classification of the gingers (Zingiberaceae): Evidence from molecular data. *Am J Bot.* 2002;89(10):1682-96. <https://doi.org/10.3732/ajb.89.10.1682>
- Van HT, Dam SM, Phan UTX, Nguyen TNA, Nguyen TBT, Tran TL *et al.* Chemical diversity of essential oils from aerial parts of eight species of Zingiberaceae family from Vietnam. *Acta Univ Agric Silvicae Mendelianae Brun.* 2022;70(4-5):273-81. <https://doi.org/10.11118/actaun.2022.020>
- Piyaporn S, Surapon S. Diversity, traditional uses and conservation status of Zingiberaceae in Udorn Thani Province, Thailand. *Biodiversitas J Biol Divers.* 2021;22(8):3083-97. <https://doi.org/10.13057/biodiv/d220801>
- Trimanto T, Dwiyantri D, Indriyani S. Morfologi, anatomi dan uji histokimia rimpang *Curcuma aeruginosa* Roxb., *Curcuma longa* L. DAN *Curcuma heyneana* Valetton dan Zijp. *Ber Biol.* 2018;17(2):2337-51. <https://doi.org/10.14203/beritabiologi.v17i2.3086>
- Pham HH, Araceae, In: Pham-hoang, H (Ed.). *Cây cỏ Việt Nam: An illustrated flora of Vietnam.* Youth Publishing House, Ho Chi Minh City. 2000.
- Nguyen QB. Zingiberaceae, Flora of Vietnam. Publisher of Science and Technology, Ha Noi; 2017.
- Ly NS, Hoang TS, Insiengmay O, Haevermans T, Newman MF. *Conamomum vietnamense* (Zingiberaceae), a new species from Tay Nguyen, Vietnam. *Phytotaxa.* 2022;531(2):129-35. <https://doi.org/10.11646/phytotaxa.531.2.5>
- Nguyen HT, Nguyen NA, Vervanov LA, Nguyen DD, Le CT. *Curcuma tuanii* (Zingiberaceae) a new species of subgenus *Ecomata* from Northern Vietnam based on morphological and molecular evidence. *Acta Bot Brasilica.* 2023;37. <https://doi.org/10.1590/1677-941x-abb-2023-0028>
- Nguyen DD, Le TA, Hoang QH, Le QT, Nguyen E. Two new taxa of *Curcuma* sect. *Ecomata* (Zingiberaceae: Zingiberaceae), from coastal Central Vietnam. *Biodiversitas J Biol Divers.* 2022;23(5):2512-19. <https://doi.org/10.13057/biodiv/d230531>
- Luu HT, Tran HD, Nguyen QD, Nguyen TQT, Tran G. *Conamomum odorum*, a new species of Zingiberaceae from central Vietnam. *Academia Journal of Biology.* 2019;41(3):55-59. <https://doi.org/10.15625/0866-7160/v41n3.13671>
- Ly NS, Leong-Škorničková J. *Amomum cristatissimum* (Zingiberaceae: Alpinieae), a new species with echinate fruits from central Vietnam. *Nord J Bot.* 2018;36(3):njb-01691. <https://doi.org/10.1111/njb.01691>
- Baltzerm C, Nguyen TD, Shore RG. Towards a vision for biodiversity conservation in the forests of the lower Mekong ecoregion complex. WWF Indochina/WWF US, Hanoi and Washington D.C, 2001.
- Bang TV, Long V, Duc HM. The mammal fauna of Binh Chau-Phuoc Buu Nature Reserve, Xuyen Moc District, Ba Ria-Vung Tau Province. *Proceeding 5th Natl Sci Conf Ecol Biol Resour.* Ha Noi; 2013.
- Saxena HO, Brahmam M. The Flora of Orissa. Vol-III. Bhubaneswar: Orissa Forest Development Corporation Ltd; 1995.
- Anonymous. *Reviews on Indian Medicinal Plants.* Vol-11. New Delhi: Indian Council of Medical Research; 2013.
- Bridson D, Forman L. *The Herbarium Handbook-Third Edition.* Royal Botanic Gardens. Royal Botanic Gardens, Kew, UK; 1999.
- Le HT, Luu TN, Nguyen HMT, Nguyen DHT, Le PTQ, Trinh NN *et al.* Antibacterial, antioxidant and cytotoxic activities of different fractions of acetone extract from flowers of *Dipterocarpus intricatus* Dyer (Dipterocarpaceae). *Plant Sci Today.* 2021;8(2):273-77. <https://doi.org/10.14719/pst.2021.8.2.1086>
- Bhutia S, Uriah T, Mohanty JP, Pal P, Kakoti BB. Physicochemical evaluation, GC-MS and antibacterial activity of essential oil of the fruit peels of *Citrus macroptera* Montr. of Meghalaya North East India. *WJPLS.* 2019;5(1):108-15.
- Sannigrahi S, Mazuder UK, Pal DK, Parida S, Jain S. Antioxidant potential of crude extract and different fractions of *Enhydra fluctuans* Lour. *Iranian J Pharmaceutical Res.* 2010;9(1):75-82.
- Menon AN, Dan M. Chemical composition of essential oils of *Globba* species from South India. *Journal of Essential Oil Research.* 2009;21(1):59-60. <https://doi.org/10.1080/10412905.2009.9700110>
- Nayak S, Jena AK, Mittal DK, Joshi D. GC-MS analysis of phytoconstituents of some wild Zingiberaceae plants methanolic rhizome extracts. *Res Plant Sci.* 2014;2(1):1-5. <http://doi.org/10.12691/plant-2-1-1>
- Van HT, Dam SM, Phan NTX, Nguyen HBN, Le TT, Nguyen TP, Huynh NTA. Chemical diversity of essential oils of rhizomes of six species of Zingiberaceae family. *J Arid Agric.* 2022;8:8-13. <https://doi.org/10.25081/jaa.2022.v8.7430>
- Van HT, Phan UTX, Doan VD, Le VS. Chemical Constituents of essential oil from rhizomes and aerial parts of *Curcuma gracillima*. *Chem Nat Compd.* 2021;57(3):569-71. <https://doi.org/10.1007/s10600-021-03419-6>
- Singthong C, Gagnon MJ, Legault J. Chemical composition and biological activity of the essential oil of *Amomum biflorum*. 2013;8(2):265-67. <https://doi.org/10.1177/1934578X1300800234>
- Van HT, Tran MD, Tran TKN, Nguyen TDH, Nguyen NA, Huynh NTA *et al.* Chemical profiles and antibacterial activity of acetone extract of two *Curcuma* species from Vietnam. *Plant Sci Today.* 2023;10(2):83-89. <https://doi.org/10.14719/pst.1993>
- Sheeja ADB, Nair MS. Facile isolation of (E)-labda-8(17),12-diene-15,16-dial from *Curcuma amada* and its conversion to other biologically active compounds. 2014;53:319-24.
- Chimnoi N, Pisutjaroenpong S, Ngwisara L, Dechtrirut D, Chokchaichamnankit D, Khunnawutmanotham N. Labdane diterpenes from the rhizomes of *Hedychium coronarium*. *Nat Prod Res.* 2008;22(14):1249-56. <https://doi.org/10.1080/14786410701726434>
- Peng W, Li P, Ling R, Wang Z, Feng X, Liu J *et al.* Diversity of volatile compounds in ten varieties of Zingiberaceae. *Molecules.* 2022;27(2):565. <https://doi.org/10.3390/molecules27020565>
- Sundram T, Serm L, Malek S, Annuar M, Khalid N. (E)-labda-8(17),12-diene-15,16-dial production from *Curcuma mangga* (Mango ginger) *in vitro* cultures. *Proc 4th Australas Metabolomics Symp Work.* Shah Alam, Malaysia; 2012.
- Awini T, Mediani A, Maulidiani, Leong SW, Faudzi SMM, Shaari K, Abas F. Phytochemical and bioactivity alterations of *Curcuma* species harvested at different growth stages by NMR-based metabolomics. *J Food Compos Anal.* 2019;77:66-76. <https://doi.org/10.1016/j.jfca.2019.01.004>
- Chakrabarty I, Vijayasekhar A, Rangan L. Therapeutic potential of labdane diterpene isolated from *Alpinia nigra*: Detailed hemato-compatibility and antimicrobial studies. *Nat Prod Res.* 2021;35(6):1000-04. <https://doi.org/10.1080/14786419.2019.1610756>
- Dahham SS, Tabana YM, Iqbal MA, Ahamed MBK, Ezzat MO, Majid ASA, Majid AMSA. The Anticancer, antioxidant and antimicrobial properties of the sesquiterpene  $\beta$ -caryophyllene

- from the essential oil of *Aquilaria crassna*. *Molecules*. 2015;20(7):11808-29. <https://doi.org/10.3390/molecules200711808>
33. Yoo HJ, Jwa SK. Efficacy of  $\beta$ -caryophyllene for periodontal disease related factors. *Arch Oral Biol*. 2019;100:113-18. <https://doi.org/10.1016/j.archoralbio.2019.02.015>
  34. Yoo HJ, Jwa SK. Inhibitory effects of  $\beta$ -caryophyllene on *Streptococcus mutans* biofilm. *Arch Oral Biol*. 2018;88:42-46. <https://doi.org/10.1016/j.archoralbio.2018.01.009>
  35. Gushiken LFS, Beserra FB, Hussni MF, Gonzaga MT, Ribeiro VP, Campos JCL *et al.* Beta-caryophyllene as an antioxidant, anti-inflammatory and re-epithelialization activities in a rat skin wound excision model. *Oxid Med Cell Longev*. 2022;9004014. <https://doi.org/10.1155/2022/9004014>
  36. Fattore E, Fanelli R. Palm oil and palmitic acid: A review on cardiovascular effects and carcinogenicity. *International Journal of Food Sciences and Nutrition*. 2013;64(5):648-59. <https://doi.org/10.3109/09637486.2013.768213>
  37. Ganesan T, Subban M, Leslee CDB, Kuppannan SB, Seedeivi P. Structural characterization of n-hexadecanoic acid from the leaves of *Ipomoea eriocarpa* and its antioxidant and antibacterial activities. *Biomass Convers Biorefinery*. 2022. <https://doi.org/10.1007/s13399-022-03576-w>
  38. Aparna V, Dileep KV, Mandal PK, Karthe P, Sadasivan C, Haridas M. Anti-inflammatory property of n-hexadecanoic acid: Structural evidence and kinetic assessment. *Chem Biol Drug Des*. 2012;80(3):434-39. <https://doi.org/10.1111/j.1747-0285.2012.01418.x>
  39. Chavan MJ, Wakte PS, Shinde DB. Analgesic and anti-inflammatory activity of Caryophyllene oxide from *Annona squamosa* L. bark. *Phytomedicine*. 2010;17(2):149-51. <https://doi.org/10.1016/j.phymed.2009.05.016>
  40. Yang D, Michel L, Chaumont JP, Millet-Clerc J. Use of caryophyllene oxide as an antifungal agent in an *in vitro* experimental model of onychomycosis. *Mycopathologia*. 1999;148(2):79-82. <https://doi.org/10.1023/A:1007178924408>
  41. Jun NJ, Mosaddik A, Moon JY, Jang KC, Lee DS, Ahn KS, Cho SK. Cytotoxic activity of  $\beta$ -caryophyllene oxide isolated from Jeju Guava (*Psidium cattleianum* Sabine) leaf. *Nat Prod*. 2011;5:242-46.
  42. Bhardwaj M, Sali VK, Mani S, Vasanthi HR. Neophytadiene from *Turbinaria ornata* Suppresses LPS-induced inflammatory response in RAW 264.7 macrophages and sprague Dawley Rats. *Inflammation*. 2020;43(3):937-50. <https://doi.org/10.1007/s10753-020-01179-z>
  43. Gonzalez-Rivera ML, Barragan-Galvez JC, Gasca-Martínez D, Hidalgo-Figueroa S, Isiordia-Espinoza M, Alonso-Castro AJ. *In vivo* neuropharmacological effects of neophytadiene. *Molecules*. 2023;28(8):3457. <https://doi.org/10.3390/>