



RESEARCH ARTICLE

Traditional knowledge of ethnomedicinal plants used by the Mishing community in Sivasagar District, Assam (India)

Priyakshi Buragohain¹, Barnali Das^{1,2}, Madhushmita Nath¹, Parag Jyoti Sarma¹, Mijing Boro¹, Shilpa Roy¹ & Namita Nath^{1*}

¹Department of Botany, Gauhati University, Gopinath Bordoloi Nagar, Guwahati-781014, Assam, India

²Department of Botany, Pragiyotish College, Santipur, Guwahati-09, Assam, India

*Email: nathnamita1@gauhati.ac.in



ARTICLE HISTORY

Received: 09 February 2024

Accepted: 22 May 2024

Available online

Version 1.0 : 07 June 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Buragohain P, Das B, Nath M, Sarma PJ, Boro M, Roy S, Nath N. Traditional knowledge of ethnomedicinal plants used by the Mishing community in Sivasagar District, Assam (India). *Plant Science Today* (Early Access). <https://doi.org/10.14719/pst.3360>

Abstract

Ethnobotany plays a significant role in traditional medicine practices. The present study aims to enlist the ethnomedicinal plants used by the Mishing community in Sivasagar district, Assam, India. The study was conducted from March 2021 to March 2022 to record the ethnomedicinally important plants used by the Mishing community. In the present study, 109 plant species were used in different ethnomedicinal practices which belong to 55 families and 101 genera. Out of 55 families, Asteraceae is dominant with the highest number of medicinal plants (8 species, 7.33%). In the survey, the symptomatic diseases category showed the highest agreement with an F_{IC} of 0.64%. The most commonly used plant parts for medicinal treatments were leaves (68 species, 62.38%). In comparison to other plant parts, the utilization of leaves causes less distress to the plant ensuring sustainability and its further conservation. It is concluded that the Mishing community of Sivasagar (Assam) District uses various medicinal plants to cure different diseases as their primary source of health care. This will be a significant contribution to the herbal and pharmaceutical industries for the welfare of mankind.

Keywords

Assam; ethnobotany; mishing; Sivasagar; traditional knowledge

Introduction

Traditional knowledge plays a crucial role in resource conservation, specifically of native plant species important for local communities (1-4). Since time immemorial, the local communities have been collecting ethnomedicinal plants from the wild in various parts of the world (5-12) and the information is passed from generation to generation (13). This information has to be conserved particularly in countries facing a high risk of biodiversity loss because of urbanization, migrations, deforestation, and natural disasters. In India, traditional knowledge is decreasing day by day at an alarming rate because of the above-mentioned factors. As a mega biodiversity hot spot, Northeast India is wealthy in endemic flora (14-18) and shelter to nearly 1,350 medicinal plants with great economic value that are utilized in different ethnomedicinal concoctions (19). Apart from being wealthy in floristic diversity, the region is also rich in diversified culture and traditional knowledge between 145 tribal communities (20). NE India is considered as one of the ecological hot spots of the globe and has plenty of medicinal plants familiar to the local people (18, 21-23). Assam, an important state of northeastern India falls under the Indo-Burma Global Biodiversity Hotspot (24). The Mishing tribes are especially found in the Dhemaji,

selected for data collection. During the process, importance was given to collecting data with a detailed account of every informant including their identity, address, qualifications, and tribal group. This was recorded before the collection of traditional knowledge-based information in the local language.

Plant Collection, Identification and Preservation

Plants were collected by the herbal practitioners of the Mishing community for proper recognition during the mature stage. Also, voucher specimens related to ethno-medicinal information were collected during the reproductive phase for proper identification. Collected specimens were first tagged with the local name during the survey; then they were identified by following relevant taxonomic literature such as Flora of Assam (29); Assam's Flora (30); A checklist of Angiosperms and Gymnosperms (31), also comparing with the deposited voucher specimens at GUBH (Gauhati University Botanical Herbarium). Digital databases like "IPNI: International Plant Name Index (<https://www.ipni.org>)" and "POWO: Plants of the World Online (<https://powo.science.kew.org>)" were used for checking the accepted scientific nomenclature of each specimen.

Statistical analysis

The collected data is represented orderly in tabular format. Demographic data contains the details about the informants; whereas data related to medicinal plants such as scientific name, family, vernacular name, use value, parts used, disease, and mode of application were tabulated together.

Determination of Use Value (UV)

By determining the use value the relative importance of each recommended medicinal plant was calculated (32, 33), for measuring the relative importance of plants used by local healers on a quantitative basis:

$$UV = \sum U_i/n$$

Here,

U_i = number of use reports cited by each informant for a given species

n = total number of informants

UV will be high for a plant when there are many use reports, and UV will approach zero (0) when there are few reports.

Determination of informants' consensus factor (F_{ic})

Usually, the informants' consensus factor (F_{ic}) is determined to find out the uniformity in the data given by the informants of the study area. By using the following formula F_{ic} was calculated (34-37).

$$F_{ic} = (N_{ur} - N_t) / N_{ur-1}$$

Here,

N_{ur} = number of use reports in a particular category of illness by informants

N_t = number of species of taxa that are used for the treatment of a particular disease category by informants of the study.

The informants' consensus factor value ranges from 0 to 1. When F_{ic} is higher or close to 1, it indicates higher reports about a plant species used by the informants in a specific ailment. When it is low or close to 0, it indicates disagreement by the informants about a plant used for a particular ailment.

Results and Discussion

Demography

In the Sivasagar District, Assam a total of 50 informants of the age group range from 30-79 years were interviewed (**Table 1**). Out of 50 informants, 18 were male (36%) and 32 were female (64%). From the present work, it was found that the average age of the informants was 59 years. Most of the informants were involved with other livelihood activities being farmers, social workers, teachers, shopkeepers, and housewives. The illiteracy rate was found to be 24% while the literacy rate at the primary level was 22%, the middle level was 20%, the secondary level was 18% and the university level was 16% (**Table 2; Fig. 2**).

Table 1. Distribution of ethnic informants based on age and sex

Age group	Male	Female	No. of Individuals	Percentage (%)
20-29	00	03	03	6
30-39	02	07	09	18
40-49	06	08	14	28
50-59	03	05	08	16
60-69	06	06	12	24
70-79	01	03	04	8
Total	18	32	50	100




Fig.2. Pie chart showing the informants percentage based on educational status

Table 2. Educational status of the informants

Education level	No. of Individuals	Percentage (%)
Illiterate	12	24
Primary	11	22
Middle	10	20
Secondary	09	18
University	08	16

An overview of medicinal plants

In the present study, 109 plant species were used in different ethnomedicinal practices which belong to 55 families and 101 genera. These were found to be used to heal certain human diseases which were grouped under 17 ICPC (International Classification of Primary Care) disease categories. It was found that the most documented ethnomedicinal plants were followed by herbs, shrubs, trees, lianas, and epiphytes (Fig. 3). The details on traditional knowledge accomplished by the Mishing people of Sivasagar District were arranged by scientific names together with their families, vernacular names and applications (Table 3). This is because of the availability of non-conventional herbs which are simple to plant in home gardens as compared to shrubs, trees, lianas, and epiphytes which take a longer period to grow. Due to this

reason herbs possess strong medicinal properties and more healing effects to combat illnesses (38-40). Most of these traditionally used plants are used by the Mishing tribe in their daily life for their livelihood and also to clear out serious/chronic health problems.




Fig.3. Percentage distribution of the medicinal plants habit wise

Table 3. Description of plants with use value, parts used, application, method of preparation, and route of administration collected from Sivasagar District, Assam, India

Sl. No.	Scientific name	Family	Common name	Use Value	Parts used	Application	Method of preparation	Route of administration
1	<i>Acmella ciliata</i> (Kunth) Cass.	Asteraceae	Huhoni-bon	0.14	Wh, Fl	Tongue bump, vision, fever, cough, headache, body pain, post-parturition pain	Raw, vegetable	Oral
2	<i>Acorus calamus</i> (L.)	Acoraceae	Bosh	0.14	Rh, L	Fever, stomach pain, dizziness	Raw, pill	External, oral
3	<i>Aegle marmelos</i> (L.) Correa	Rutaceae	Bael	0.02	Fr	Leukorrhoea	Juice	Oral
4	<i>Ageratum conyzoides</i> (L.)	Asteraceae	Gendhali-bon	0.08	L	Cut & wound,	Paste	External
5	<i>Allium cepa</i> (L.)	Amaryllidaceae	Piyanj	0.04	Bu	Diarrhoea (children), dizziness	Paste	External
6	<i>Allium sativum</i> (L.)	Amaryllidaceae	Nohoru	0.24	Bu	Fever, gall bladder stone, gastric, hypertension, post-parturition weakness, toothache	Raw, oil infusion, paste, juice, vegetable	External, oral
7	<i>Aloe vera</i> (L.) Burm. f.	Asphodelaceae	Sal-kunwori	0.06	L	Fever	Raw	External
8	<i>Alternanthera sessilis</i> (L.) R.Br. Ex DC.	Amaranthaceae	Mati-kanduri	0.04	L, St	Gastric, pneumonia	Vegetable, water infusion	Oral
9	<i>Amaranthus spinosus</i> L.	Amaranthaceae	Hati-khutura	0.12	L, St, R	Cut & wound, pinned, tuberculosis, leukorrhoea	Raw, juice	Oral, external
10	<i>Ananas comosus</i> (L.) Merr.	Bromeliaceae	Anaras, Mati-kothal	0.02	L	Roundworm	Paste	Oral
11	<i>Andrographis paniculata</i> (Burm.f.) Nees	Acanthaceae	Kalmegh	0.04	L	Fever	Water infusion	Oral
12	<i>Artocarpus heterophyllus</i> Lam.	Moraceae	Kothal	0.06	Fr, L	Diabetes, tonsillitis	Water infusion, smoke	Oral, external
13	<i>Azadirachta indica</i> A. Juss.	Meliaceae	Mahaneem	0.06	L	Itchiness, malaria, small-pox	Raw, water infusion	External, oral
14	<i>Bambusa tulda</i> Roxb.	Poaceae	Banh	0.08	L, B	Over menstrual bleeding, cuts & wounds, epiglottis enlargement	Juice, paste	Oral, external
15	<i>Basella alba</i> (L.)	Basellaceae	Puroisak	0.02	L, St	Piles	Pill	Oral
16	<i>Blumea lacera</i> (Burm. f.) DC.	Asteraceae	(Barbie in mishing)	0.02	L	Boil	Paste	External
17	<i>Bombax ceiba</i> L.	Malvaceae	Himolu (SinggiAppun)	0.02	L	Boil	Paste	External

18	<i>Cardiospermum halicacabum</i> L.	Sapindaceae	Kopalphuta	0.04	L	Gastric, and joint pain	Juice, paste	Oral, external
19	<i>Centella asiatica</i> (L.) Urb.	Apiaceae	Bor-manimuni	0.08	Wh, L	Leucorrhoea, dysentery, boil, pneumonia	Juice, paste	Oral, external
20	<i>Centipeda minima</i> (L.) A. Braun & Asch.	Asteraceae	Hasiyotibon	0.02	Wh, L	Waist pain	Juice	Oral
21	<i>Chrysopogon aciculatus</i> (Retz.) Trin.	Poaceae	Bonguti	0.04	St, Rh	Pneumonia, urinary infection	Pill, water infusion	Oral
22	<i>Cissampelos pareira</i> (L.)	Menispermaceae	Tubukilota	0.02	L	Sore throat	Paste	External
23	<i>Cissus quadrangularis</i> (L.)	Vitaceae	Harjuralota	0.04	Wh	Bone fracture, breakage	Paste	External
24	<i>Citrus × aurantiifolia</i> (Christm.) Swingle	Rutaceae	Kaji-nemu	0.10	L, Fr	Dizziness, itchiness, typhoid pneumonia, post-parturition pain, stomach scars	Juice, water infusion	Oral, external
25	<i>Citrus × limon</i> (L.) Osbeck	Rutaceae	Golnemu	0.12	L, Fr, Sp	Pneumonia, jaundice, dysentery, roundworm, diarrhea	Juice, paste, pill, water infusion	Oral
26	<i>Clerodendrum colebrookeanum</i> Walp.	Lamiaceae	Nephaphu	0.04	L	Hypertension	Juice, vegetable	Oral, external
27	<i>Coccinia grandis</i> (L.) Voigt	Cucurbitaceae	Belipoka	0.08	L, St	Dysentery, leucorrhoea, pneumonia	Water infusion, pill, juice	Oral
28	<i>Cocos nucifera</i> (L.)	Arecaceae	Narikol	0.18	Fr	Jaundice, fever, small-pox, dizziness, cuticle pain, heat rash	Juice	Oral, external
29	<i>Colocasia esculenta</i> (L.) Schott.	Araceae	Kola kosu, bonoriyako su	0.04	Rh, La	Cancer, epiglottis enlargement	Paste, raw	External
30	<i>Commelina diffusa</i> Burm.f.	Commelinaceae	Kona-himolu	0.02	L, St	Jaundice	Juice	Oral
31	<i>Coriandrum sativum</i> (L.)	Apiaceae	Dhaniya	0.02	S	Gall bladder	Water infusion	Oral
32	<i>Crinum asiaticum</i> (L.)	Amaryllidaceae	Bon-nohoru	0.02	Bu	Urinary problems (children)	Raw, smoked	External
33	<i>Curcuma caesia</i> Roxb.	Zingiberaceae	Kola halodhi	0.06	Rh	Stomach pain, gastric	Paste, raw	Oral
34	<i>Curcuma longa</i> (L.)	Zingiberaceae	Halodhi	0.1	Rh	Menstruation pain, dizziness, bone fracture, joint pain, memory booster	Raw, paste, pill, juice	Oral, external
35	<i>Cyathula prostrata</i> (L.) Blume	Amaranthaceae		0.02	L	Cut & wound	Paste	External
36	<i>Cynodon dactylon</i> (L.) Pers.	Poaceae	Dubori bon	0.1	Wh	Cut & wound, dysuria, urinary problem	Juice, paste	Oral, external
37	<i>Datura metel</i> (L.)	Solanaceae	Dhatura	0.02	L	Bone-breakage	Paste	External
38	<i>Dillenia indica</i> (L.)	Dilleniaceae	Ow-tenga (Sompapayur)	0.02	Fr	Boil	Paste	External
39	<i>Drymaria cordata</i> (L.) Willd. ex. Schult.	Caryophyllaceae	Lai-jabori	0.02	L	Sinusitis	Juice	External
40	<i>Eclipta prostrata</i> (L.)	Asteraceae	Keheraj	0.2	L, St	Pneumonia, jaundice, menstruation pain, over-menstrual bleeding, external scars, dysentery, stomach pain, post-parturition pain	Paste, juice, pill	External, oral
41	<i>Eleocharis dulcis</i> (Burm.f.) Trin. Ex Hensch.	Cyperaceae	Sensu	0.02	Co	Asthma	Juice	Oral
42	<i>Elettaria cardamomum</i> (L.) Maton	Zingiberaceae	Elachi	0.08	Fr	Cough, pneumonia, jaundice, gallbladder stone	Pill, raw	Oral

43	<i>Eleusine indica</i> (L.) Gaertn.	Poaceae	Bobosa bon	0.04	Wh, R	Pneumonia, scabies	Water infusion, raw	Oral, external
44	<i>Eryngium foetidum</i> L.	Apiaceae	Man-dhoniya	0.02	L	Indigestion	Juice	Oral
45	<i>Ficus carica</i> L.	Moraceae	Dimaru	0.06	L	Post-parturition weakness, post-parturition bleeding, menstruation pain	Vegetavble, juice	Oral
46	<i>Ficus hispida</i> L.f.	Moraceae	(Tapot)	0.02	L	Pneumonia	Water infusion	Oral
47	<i>Ficus lamponga</i> Miq.	Moraceae	(Takhot)	0.02	L	Ear pain	Smoke	External
48	<i>Foeniculum vulgare</i> Mill.	Apiaceae	Sauf	0.02	S	Gall bladder stone	Water infusion	Oral
49	<i>Guilandina bonduc</i> L.	Fabaceae	Letaguti	0.2	S	Cough, stomach pain, pneumonia	Water infusion, pill	Oral
50	<i>Hellenia speciosa</i> (J.Koeing) S.R.Dutta	Costaceae	Jam-lakhuti	0.06	Rh	Leukorrhoea	Juice	Oral
51	<i>Hibiscus rosa-sinensis</i> L.	Malvaceae	Jobaphool (Goshaniappun)	0.14	L, Fl	Fever, boil, small-pox	Paste	External
52	<i>Houttuynia cordata</i> Thunb.	Saururaceae	Masandari	0.06	L	Diarrhoea, gastric, dysentery	Raw, juice, vegetable, paste	Oral
53	<i>Hydrocotyle sibthorpioides</i> Lam.	Araliaceae	Horu-manimuni	0.18	Wh	Leukorrhoea, diarrhoea, pneumonia, dysentery, memory boosting	Raw, juice, vegetable, paste, pill	Oral
54	<i>Impatiens tripetala</i> Roxb. Ex DC.	Balsaminaceae	Demdeuka	0.04	L	Cancer, fever	Paste	External
55	<i>Jatropha curcas</i> L.	Euphorbiaceae	Bongali-enera	0.04	L, St	Diarrhoea, toothache	Paste, smoke	External
56	<i>Kalanchoe pinnata</i> (Lam.) Pers.	Crassulaceae	Dupor-tenga	0.04	L	Gall bladder stone, kidney stone	Juice, paste	Oral
57	<i>Lawsonia inermis</i> L.	Lythraceae	Jetuka	0.02	L	Cuticle pain	Paste	External
58	<i>Leucas aspera</i> (Willd.) Link	Lamiaceae	Durun bon	0.3	L	Boil, cough, stomach pain, gastric, cold, tonsillitis, sinusitis, eye pain, itchiness	Paste, water infusion, juice, vegetable, raw	External, oral
59	<i>Lippia alba</i> (Mill.) N.E.Br. Ex Britton & P. Wilson	Verbenaceae	(Pelutoying in mishing)	0.02	L	Roundworm	Vegetable	Oral
60	<i>Lygodium japonicum</i> (Thunb.) Sw.	Schizaeaceae	Kopou-dhekiya (Rakanmakat)	0.02	L	Menstruation pain	Pill	Oral
61	<i>Mesosphaerum suaveolens</i> (L.) Kuntze	Lamiaceae	Tokma	0.02	S	Urinary infection	Water infusion	Oral
62	<i>Mikania micrantha</i> Kunth	Asteraceae	Premlota	0.12	L	Dysentery, stomach pain, cut & wound	Paste, juice, pill	External, oral
63	<i>Mimosa pudica</i> L.	Fabaceae	Lajukilota	0.02	L	Pneumonia	Water infusion	Oral
64	<i>Miscanthus fuscus</i> (Roxb.) Benth	Poaceae	Nol, Khagori	0.02	L, St	Cancer	Paste	External
65	<i>Momordica charantia</i> L.	Cucurbitaceae	Tita-kerela	0.02	L	Nasal bleeding	Juice	External
66	<i>Morinda angustifolia</i> Roxb.	Rubiaceae	Akalbih (Aawmpul)	0.02	L	Boil	Paste	External
67	<i>Moringa oleifera</i> Lam.	Moringaceae	Sojina	0.02	B	Epilepsy	Water infusion	Oral
68	<i>Musa acuminata</i> Colla	Musaceae	Malbhogkol	0.04	Fr, Rh	Leukorrhoea, piles	Raw, juice	Oral
69	<i>Myristica fragrans</i> Houtt.	Myristicaceae	Jaiphal	0.06	S	Pneumonia, jaundice, gall bladder stone	Pill, water infusion	Oral
70	<i>Nicotiana tabacum</i> L.	Solanaceae	Mansadha	0.04	L	Toothache	Raw	External

71	<i>Nigella sativa</i> L.	Ranunculaceae	Kaljira	0.02	S	Gall bladder stone	Water infusion	Oral
72	<i>Nyctanthes arbor-tristis</i> L.	Oleaceae	Sewali phool	0.08	L, Fl	Fever, pneumonia, cough	Water infusion	Oral
73	<i>Ocimum tenuiflorum</i> L.	Lamiaceae	Tulashi	0.12	L	Cough, small-pox, ear pain	Paste, juice, pill, raw	Oral, external
74	<i>Oldenlandia corymbosa</i> L.	Rubiaceae	Bon-jaluk	0.14	L, St	Body pain, cough, stomach pain, pneumonia, dysentery	Vegetable, water infusion, pill	Oral
75	<i>Paederia foetida</i> L.	Rubiaceae	Bhedailota	0.16	L	Menstruation pain, boil, diarrhoea, gastric, post-parturition weakness, itchiness, pneumonia	Juice, paste, vegetable	Oral
76	<i>Persicaria chinensis</i> (L.) H.Gross	Polygonaceae	(Sibe-takkir)	0.02	L, St	Boil	Paste	External
77	<i>Phyla nodiflora</i> (L.) Greene	Verbenaceae	(Gakhirkuse rai)	0.04	L	Lactation deficiency, pneumonia	Juice	Oral
78	<i>Phyllanthus emblica</i> L.	Phyllanthaceae	Amlakhi	0.04	Fr	Liver weakness, hair fall	Juice, paste	Oral, external
79	<i>Phyllanthus niruri</i> L.	Phyllanthaceae	Bon-amlakhi	0.02	R	Pneumonia	Water infusion	Oral
80	<i>Physalis angulata</i> L.	Solanaceae	(Tumpet)	0.02	R	Pneumonia	Water infusion	Oral
81	<i>Piper betle</i> L.	Piperaceae	Paan	0.1	L	Cough; post-parturition vomiting, pain; pneumonia, stomach scars	Raw, paste, juice, water infusion	Oral
82	<i>Piper nigrum</i> L.	Piperaceae	Jaluk	0.6	Fr	Pneumonia, jaundice, over menstrual bleeding, gall bladder stone, cough, asthma, stomach pain, cold, menstruation pain, fever, headache, body pain, post-parturition weakness & pain, waist pain, stomach scars	Pill, juice, water infusion, vegetable	Oral
83	<i>Plumbago zeylanica</i> L.	Plumbaginaceae	Agasiat	0.08	L, R	Tuberculosis, hairy caterpillar bite, liver weakness	Oil infusion, paste, pill, vegetable	Oral, external
84	<i>Pogostemon benghalensis</i> (Burm.f.) Kuntz	Lamiaceae	Hukloti	0.02	L	Stomach pain	Vegetable	Oral
85	<i>Persicaria glabra</i> (Willd.) M.Gomez	Polygonaceae	Bihlogoni	0.08	L	Fever, cough, headache, body pain	Vegetable	Oral
86	<i>Portulaca oleracea</i> L.	Portulacaceae	Malbhogkhutura	0.08	Wh, L	Cough, asthma, jaundice	Juice, pill	Oral
87	<i>Potentilla indica</i> (Andrews) Th. Wolf	Rosaceae	Goru-khis (Bog-jer)	0.04	Rh	Tongue bump, throat ulcer	Paste, juice	Oral
88	<i>Pouzolzia zeylanica</i> (L.) Benn.	Urticaceae	Borali-bokua	0.02	L	Cut & wound	Paste	External
89	<i>Psidium guajava</i> L.	Myrtaceae	Madhuri	0.16	L	Pneumonia, stomach pain, cough, dysentery	Juice, pill, paste	Oral
90	<i>Punica granatum</i> L.	Lythraceae	Dalim	0.02	Pe	Epilepsy	Water infusion	Oral
91	<i>Rhynchosyilis retusa</i> (L.) Blume	Orchidaceae	Kopouphool	0.02	L	Ear pain	Juice	External
92	<i>Rubus alceifolius</i> Poir.	Rosaceae	Jetuli-poka	0.02	L	Menstruation pain	Juice	Oral
93	<i>Sapindus mukorossi</i> Gaertn.	Sapindaceae	Manisal	0.08	S	Pneumonia, cough	Water infusion, pill	Oral
94	<i>Scoparia dulcis</i> L.	Plantaginaceae	Bon-cheni	0.02	L	Boil	Paste	External
95	<i>Sida rhombifolia</i> L.	Malvaceae	Hun-boriyal	0.12	L, Fr, R	Pneumonia, jaundice, tongue bump	Pill, juice, raw	Oral
96	<i>Solanum violaceum</i> Ortega	Solanaceae	Horubhekuri	0.04	R	Asthma, pneumonia	Juice, pill	Oral
97	<i>Streblus asper</i> Lour.	Moraceae	Gowalsali, hora gos	0.06	L	Boil, dysentery, pneumonia	Paste	External
98	<i>Syzygium aromaticum</i> (L.) Merr. & L.M. Perry	Myrtaceae	Laung	0.18	Fl	Cough, pneumonia, jaundice, over menstrual bleeding, gall bladder stone, leucorrhoea, toothache, menstruation pain	Pill, paste, raw, vegetable, water infusion	Oral
99	<i>Tagetes erecta</i> L.	Asteraceae	Narjiphool	0.08	L	Cut & wound, boil	Paste	External
100	<i>Tamarindus indica</i> L.	Fabaceae	Teteli	0.04	Fr	Hypertension, dizziness	Raw, juice	Oral

101	<i>Terminalia arjuna</i> (Roxb.)	Combretaceae	Arjun gos	0.02	B	Menstruation pain	Juice	Oral
102	<i>Terminalia chebula</i> Retz.	Combretaceae	Hilikha	0.04	Fr	Liver weakness, jaundice	Juice, pill	Oral
103	<i>Tetrastigma leucostaphyllum</i> (Dennst.) Alston	Vitaceae	Nol-tenga	0.08	L	Boil, diabetes, dysentery, diarrhea	Paste, juice, vegetable	External, oral
104	<i>Tinospora cordifolia</i> (Willd.) Hook.f. & Thomson	Menispermaceae	Amarlota, hogunilota	0.02	B	Gall bladder stone	Water infusion	Oral
105	<i>Vitex negundo</i> L.	Lamiaceae	Pasatiya	0.14	L	Fever, dysentery, cough, headache, bodypain	Water infusion, juice, vegetable	Oral
106	<i>Volkameria inermis</i> L.	Lamiaceae	-	0.02	L	Dysentery	Juice	Oral
107	<i>Xanthium strumarium</i> L.	Asteraceae	Agoruwa	0.02	R	Cough	Pill	Oral
108	<i>Zanthoxylum nitidum</i> (Roxb.) DC.	Rutaceae	Tejmui	0.02	B	Toothache	Raw	External
109	<i>Zingiber officinale</i> Roscoe	Zingiberaceae	Ada	0.28	Rh	Pneumonia, jaundice, cough, stomach pain, cold, pneumonia, gastric, post-parturition weakness, toothache	Raw, juice, paste, pill, vegetable, water infusion	Oral

(Abbreviation: L-Leaf, Wh-Whole plant, St-Stem, B-Bark, Fr-Fruit, Fl-Flower, Pe-Peel, R-Root, Bu-Bulb, Co-Corm, S-Seed, Rh-Rhizome, La-latex)

Plant parts used and types of medication

The tribe has a deep native knowledge on the use of different plant parts and their therapeutic properties. Commonly used parts of plants were leaf, root, rhizome, flower, fruit, bulb, bark, stem, seed, and whole plant for different purposes in their daily life. The most commonly used plant parts for medicinal treatment were leaves (68 species, 62.38%). In comparison to other parts of plants, the use of leaves causes less distress to the plant thus ensuring sustainability and its further conservation (18). It is then followed by fruit (13 species, 11.92%), stems (11 species, 10.09%), rhizomes (9 species, 8.25%), Whole plant (8 species, 7.33%), root (7 species, 6.42%), seeds (7 species, 6.42%), bark (5 species, 4.58%), flowers (4 species, 3.66%), bulbs (3 species, 2.75%), corm (1 species, 0.91%), Latex (1 species, 0.91%), peel (1 species, 0.91%), Spine (1 species, 0.91%)(Fig. 4).

Out of 55 families, Asteraceae is dominant with the highest number of medicinal plants (8 species, 7.33%); followed by Lamiaceae (7 species, 6.42%); Poaceae, Solanaceae and Moraceae (5 species, 4.58%) (Table 4).

The medicinal plants that were used in different forms to treat various human diseases were plant juice (48 species, 44.03%) followed by plant paste (46 species, 42.20%), water infusion (31 species, 28.44%), pill (26 species, 24.52%), eaten raw (24 species, 22.01%), vegetable (18 species, 16.51%), smoke (4 species, 3.66%), oil infusion (2 species, 1.83%)(Fig. 5).




Fig. 4. Bar diagram showing the percentage of parts used of the medicinal plants

Quantitative data on the ethnomedicinal uses

Use value (UV)

In this study, the use value ranges from 0.6-0.02 (Table 3). The most frequently used species were *Piper nigrum* L. with a 0.6 use value and *Leucas aspera* (Willd.) Link with 0.3 use value; followed by *Zingiber officinale* Roscoe with a use value of 0.28. Various works have been done on the above-mentioned plants to evaluate their medicinal properties. *Piper nigrum* L. has medicinal properties such as antioxidant (41), Neuroprotective (42), Anti-inflammatory (43), Anticancer (44), Analgesic (43), hypoglycemic, and hypolipidemic (45) activities, etc. It is scientifically proven that *Leucas aspera* L. is a very potential source of traditional medicine as it shows Antioxidant (46), Antimicrobial (47), Anticancer (48), Anti-inflammatory (49), Hepatoprotective (50), Larvicidal (51), Renoprotective (52), Anthelmintic (53) activities, etc. *Zingiber officinale* Roscoe is also a very important herb in traditional medicinal practices. Works have been done on its medicinal properties such as Antiviral (54), Radioprotective (55), Anti-inflammatory (56), Anticancer (57), Antioxidant (58) activities, etc. (Fig. 6 and 7).

Informant consensus factor (F_{IC})

In ethnomedicinal studies, F_{IC} analysis gives a measure of accessibility for the given information of data collection (59). In this study, the medicinal plants used to cure various ailments in the Sivasagar District of Assam were classified into 17 ICPC (International Classification of Primary Care) (<https://www.who.int/standards/classifications/other-classification/international-classification-of-primary-care>)





Fig. 5. Bar diagram showing the forms of medication of the medicinal recipes

Table 4. Category wise distribution of various medicinal plant taxa in Sivasagar District

Sl. No.	Family	Number of genera	Percentage of genera	Number of species	Percentage of species
1	Asteraceae	8	8	8	7.34
2	Lamiaceae	7	7	7	6.42
3	Poaceae	5	5	5	4.59
4	Moraceae	3	3	5	4.59
5	Solanaceae	4	4	4	3.67
6	Apiaceae	4	4	4	3.67
7	Zingiberaceae	3	3	4	3.67
8	Rutaceae	3	3	4	3.67
9	Malvaceae	3	3	3	2.75
10	Amaranthaceae	3	3	3	2.75
11	Fabaceae	3	3	3	2.75
12	Rubiaceae	3	3	3	2.75
13	Amaryllidaceae	2	2	3	2.75
14	Sapindaceae	2	2	2	1.83
15	Menispermaceae	2	2	2	1.83
16	Lythraceae	2	2	2	1.83
17	Verbenaceae	2	2	2	1.83
18	Polygonaceae	2	2	2	1.83
19	Rosaceae	2	2	2	1.83
20	Myrtaceae	2	2	2	1.83
21	Vitaceae	2	2	2	1.83
22	Piperaceae	1	1	2	1.83
23	Cucurbitaceae	1	1	2	1.83
24	Phyllanthaceae	1	1	2	1.83
25	Combretaceae	1	1	2	1.83
26	Acoraceae	1	1	1	0.92
27	Other 28 families	28	28	28	25.69
Total	54	100	100%	109	100%

**Image 1-** A- Informant showing *Morinda angustifolia* Roxb., B- Informant showing *Plumbago zeylanica* L.**Image 2-** Some medicinal plants collected from study site: A- *Acmella ciliata* (Kunth) Cass., B- *Lippia alba* (Mill.) N.E.Br. Ex Britton & P. Wilson, C- *Eclipta prostrata* (L.) L., D- *Kalanchoe pinnata* (Lam.) Pers.

disease categories and the F_{ic} value of each disease category was calculated (**Table-5**). In the investigation, the symptomatic diseases category showed the highest agreement with a F_{ic} of 0.64%. It was followed by external injuries (0.54%), heart and cardiovascular system disorders and diabetes (0.50%), digestive system disorders (0.44%), gynecological problems (0.41%), pneumonia (0.40%), ENT (0.33%), respiratory & pulmonary system diseases (0.3%), hepatological & dermatological diseases (0.29%), musculoskeletal & nervous system disorders (0.21%), urinogenital & renal and miscellaneous disorders (0.2%), oral & dentistry (0.09%). The lowest agreement between the informants was documented in the responses related to cancer and piles both with 0% F_{ic} Value. Earlier different workers followed this F_{ic} value as an important tool to conduct respective ethnobotanical studies (60-64). These studies show the greatest level of agreement between the different ethnic tribes of the state of Assam having a rich traditional knowledge with diverse flora as well as fauna together with rich tradition and culture.

Conclusion

The current study presents a comprehensive dataset on the traditional medicinal knowledge and ethnobotanical practices of the Mishing tribe in Sivasagar District. A total of 109 plants associated with 17 different disease categories were documented. While the majority of these plants belong to various angiospermic families, one belongs to the pteridophytic family.

In treatment of various diseases, senior villagers and traditional healers gave more preferences to some medicinal plants, such as *Piper nigrum* L., and *Leucas aspera* (Willd.) Link, *Zingiber officinale* Roscoe, *Allium sativum* L., *Guilandina bonduc* L., *Eclipta prostrata* (L.) L., *Cocos nucifera* L., *Hydrocotyle sibthorpioides* Lam., *Syzygium aromaticum* (L.) Merr. & L.M.Perry, *Paederia foetida* L. and *Psidium guajava* L. The most commonly utilized plant parts for medicinal treatments were leaves (68 species, 62.38%). In comparison to other plant parts, the use of leaves causes less threat to the plant ensuring sustainability and its further conservation. In this survey, the symptomatic disease category showed the highest agreement with a F_{ic} of 0.64%. The lowest agreement between the informants was recorded in the responses related to cancer and piles both with a F_{ic} of 0%. These statistical analyses of the medicinal plants have validated their relative importance and capability towards curing different ailments. Therefore, plants having ethno medicinal properties can be chemically tested for proper recognition of bioactive compounds which can be utilized further for drug designing. This will be a significant contribution to the herbal and pharmaceutical industries for the welfare of mankind. This current investigation offers a fresh perspective aimed at raising awareness and establishing management strategies for both the ethnomedicinal plants and the floral diversity within Sivasagar District.

Table 5. Informant Consensus Factor (IFC) by disease category in Sivasagar District, Assam, India

Sl. No.	Disease category	Use Report (N_{ur})	No. of taxa (N_t)	F_{ic}
1	Symptomatic diseases [Fever, cold, headache, dizziness]	49	18	0.64
2	External injuries [Cut & wound, pinned, cuticle pain, external scars]	23	11	0.54
3	Heart & cardiovascular system disorders [Hypertension]	5	3	0.5
4	Diabetes	3	2	0.5
5	Digestive system disorders [Gastric, stomach ache, diarrhea, dysentery, indigestion, roundworm, stomach scars]	55	31	0.44
6	Gynecological problems [Menstruation pain, over-menstrual bleeding, leucorrhoea, post-parturition vomiting, bleeding, weakness, lactation deficiency]	35	21	0.41
7	Pneumonia	48	21	0.41
8	ENT diseases [Ulceration, sore throat, throat bumps, tonsillitis, sinusitis, ear pain, nasal bleeding]	13	9	0.33
9	Respiratory & pulmonary system diseases [Tuberculosis, cough, asthma]	31	22	0.3
10	Hepatological [Gall bladder, liver weakness, jaundice]	28	20	0.29
11	Dermatological diseases [Boil, itchiness, small-pox, scabies, heat rash]	25	18	0.29
12	Musculoskeletal & nervous system disorders [Bone fracture, joint pain, waist pain, body pain, epilepsy]	15	12	0.21
13	Urinogenital & renal [Dysuria, kidney-stone, urinary infection & problems]	6	5	0.2
14	Miscellaneous disorders [Vision, memory booster, hairy caterpillar bite, malaria]	6	5	0.2
15	Oral & dentistry problems [Tongue bumps, toothache, uvule enlargement]	12	11	0.09
16	Cancer	3	3	0
17	Piles	2	2	0

Acknowledgements

All the authors are grateful to the local informants and traditional healers of the Mishng community of Sivasagar district, Assam for providing valuable data and for their cooperation during the entire field survey. The authors are also grateful to Dr. Sauravjyoti Borah, Curator, GUBH for the permission to study the deposited voucher specimens. The authors are also thankful to the authority of the Dept. of Botany, Gauhati University for all necessary help during the study.

Authors' contributions

The fieldwork was conducted by PB. PB and BD calculated the main statistical data. NN and BD helped in the identification of the specimens. BD, MN, and PJS collectively wrote the manuscript. NN supervised the work; all authors read, corrected, and approved the manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None.

References

- Cox PA. Will tribal knowledge survive the millennium? *Science*. 2000;287:44–45. <https://dx.doi.org/10.1126/science.287.5450.44>
- Leonti M, Sticher O, Heinrich M. Medicinal plants of the Popoluca, México: organoleptic properties as indigenous selection criteria. *Journal of Ethnopharmacology*. 2002;81(3):307–315. [https://dx.doi.org/10.1016/s0378-8741\(02\)00078-8](https://dx.doi.org/10.1016/s0378-8741(02)00078-8)
- Leonti M. The future is written: impact of scripts on the cognition, selection, knowledge, and transmission of medicinal plant use and its implications for ethnobotany and ethnopharmacology. *Journal of Ethnopharmacology*. 2011;134(3):542–555. <https://dx.doi.org/10.1016/j.jep.2011.01.017>
- Kayani S, Ahmad M, Sultana S, Shinwari ZK, Zafar M. Ethnobotany of medicinal plants among the communities of alpine and sub-alpine regions of Pakistan. *Journal of Ethnopharmacology*. 2015;164:186–202. <https://dx.doi.org/10.1016/j.jep.2015.02.004>
- Rana VS, Sharma S, Rana N, Kumar V, Sharma U, Modgill V, Prasad H. Underutilized fruit crops in North-Western Himalayan region under changing climatic scenario. *Genetic Resources and Crop Evolution*. 2023;70(1):37–69. <https://doi.org/10.1007/s10722-022-01470-y>
- Matu EN, Staden JV. Antibacterial and anti-inflammatory activities of some plants used for medicinal purposes in Kenya. *Journal of Ethnopharmacology*. 2003;87(1):35–41. [https://dx.doi.org/10.1016/s0378-8741\(03\)00107-7](https://dx.doi.org/10.1016/s0378-8741(03)00107-7)
- Mall B, Gauchan DP, Chhetri RB. An ethnobotanical study of medicinal plants used by ethnic people in Parbat District of western Nepal. *Journal of Ethnopharmacology*. 2015;165:13–17. <https://dx.doi.org/10.1016/j.jep.2014.12.057>
- Pasquini MW, Mendoza JS, Sanchez-Ospina C. Traditional Food Plant knowledge and Use in Three Afro-Descendant Communities in the Colombian Caribbean Coast: Part I Generational Differences. *Economic Botany*. 2018;72(3):278–294. <https://dx.doi.org/10.1007/s12231-018-9422-6>
- Phumthum M, Balslev H. Use of Medicinal Plants among Thai Ethnic Groups: A Comparison. *Economic Botany*. 2018;73(1):64–75. <https://dx.doi.org/10.1007/s12231-018-9428-0>
- Tomasini S, Theilade I. Local Knowledge of Past and Present Uses of Medicinal Plants in Prespa National Park, Albania. *Economic Botany*. 2019;73:217–232. [10.1007/s12231-019-09454-3](https://doi.org/10.1007/s12231-019-09454-3). <https://doi.org/10.1007/s12231-019-09454-3>
- Dixit S, Tiwari S. Investigation of anti-diabetic plants used among the ethnic communities of Kanpur division, India. *Journal of Ethnopharmacology*. 2020;253:112639. <https://dx.doi.org/10.1016/j.jep.2020.112639>
- Qamariah N, Mulia DS, Fakhri D. Indigenous Knowledge of Medicinal Plants by Dayak Community in Mandomai Village, Central Kalimantan, Indonesia. *Pharmacognosy Journal*. 2020;12(2):386–390. <https://dx.doi.org/10.5530/pj.2020.12.60>
- Tabuti JRS, Dhillon SS, Lye KA. Traditional medicine in Bulamogi Country, Uganda: its practitioners, users and viability. *Journal of Ethnopharmacology*. 2003;85(1):119–129. [https://dx.doi.org/10.1016/s0378-8741\(02\)00378-1](https://dx.doi.org/10.1016/s0378-8741(02)00378-1)
- Mao AA, Hynniewta TM. Floristic diversity of North East India. *Journal Assam Science Society*. 2000;41(4):255–266
- Sajeng AL, Rout J, Nath M. Traditional tribal knowledge and status of some rare and endemic medicinal plants of North Cachar Hills District of Assam, northeast India. *Ethnobotanical leaflets*. 2008;12:261–275
- Barbhuiya AR, Sharma GD, Arunachalam A, Deb S. Diversity and conservation of medicinal plants in Barak Valley, Northeast India. *Indian Journal of Traditional Knowledge*. 2009;8(2):169–175
- Mao AA, Hynniewta TM, Sanjappa M. Plant wealth of Northeast India with reference to Ethnobotany. *Indian Journal of Traditional Knowledge*. 2009;8(1):96–103
- Panmei R, Gajurel PR, Singh B. Ethnobotany of medicinal plants used by the Zeliangrong ethnic group of Manipur, Northeast India. *Journal of Ethnopharmacology*. 2019;235:164–182. <https://dx.doi.org/10.1016/j.jep.2019.02.009>
- Dutta BK, Dutta PK. Potential of ethnobotanical studies in northeast India: an overview. *Indian Journal of Traditional Knowledge*. 2005;4(1):7–14
- Ali ANMI, Das I. Tribal situation in northeast India. *Studies of Tribes and Tribals*. 2003;1(2):141–148. <https://dx.doi.org/10.1080/0972639X.2003.11886492>
- Asati BS, Yadav DS. Diversity of horticultural crops in northeastern regions. *Envis Bulletin: Himalayan Ecology*. 2004;12(2):1–10
- Debbarma M, Pala NA, Kumar M, Bussmann RW. Traditional knowledge of medicinal plants in tribes of Tripura in Northeast, India. *African Journal of Traditional Complementary and Alternative Medicines*. 2017;14(4):156–168. <https://dx.doi.org/10.21010/ajtcam.v14i4.19>
- Lanusunep AT, Jamir AN, Longkumer SI, Jamir NS. Traditional knowledge of herbal medicines practiced by Ao- Naga tribe in Nagaland, India. *Pleione*. 2018;12(1):11–17. <https://dx.doi.org/10.26679/Pleione.30.6.2018.011-017>
- Mittermeier RA, Turner WR, Larsen FW, Brooks TM, Gascon C. Global biodiversity conservation: the critical role of hotspots. *Biodiversity Hotspots*. 2011. London Springer Publishers. 3–22 pp. https://doi.org/10.1007/978-3-642-20992-5_1
- Teklehaymanot T, Giday M. Ethnobotanical study of medicinal plants used by people in Zegie Peninsula, Northwestern Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2007;3:1–12. <https://dx.doi.org/10.1186/1746-4269-3-12>
- Das RJ, Pathak K. Use of indigenous plants in traditional health care systems by Mishng tribe of Dikhowmukh, Sivasagar District, Assam. *International Journal of Herbal Medicine*. 2013;1(3):50–57

27. Panging SM, Sharma S. Studies on the ethno medicinal and traditional healing practices among Mishing community of Desangmukh Gaon Panchayat, Sivasagar District of Assam, India. *Journal of Medicinal Plants Studies*. 2017;5(54):193-196
28. Jain SK. *A Manual of Ethnobotany*. Jodhpur, India. Scientific Publisher. 1987.
29. Kanjilal UN, Kanjilal PC, Das A, De RN. *Flora of Assam*. Government of Assam Publication. 1940.
30. Chowdhury, S. *Assam's Flora Present Status of Vascular Plants*. Assam Science Technology and Environment Council, Guwahati. 2005.
31. Barooah C, Ahmed I. *Plant Diversity of Assam (A checklist of Angiosperms and Gymnosperms)*, ASTEC. Bigyan Bhawan, Guwahati, Assam. 2014.
32. Phillips O, Gentry AH, Reynel C, Wilkin P, Durand BCG. Quantitative ethnobotany and Amazonian conservation. *Conservation Biology*. 1994;8(1):225-248. <https://doi.org/10.1046/j.1523-1739.1994.08010225.x>
33. Zenderland J, Hart R, Bussmann RW, Zambrana NYP, Sikharulidze S. The Use of Use Value: Quantifying Importance in Ethnobotany. *Economic Botany*. 2019;73(1):1-11. <https://dx.doi.org/10.1007/s12231-019-09480-1>
34. Trotter RT, Logan MH. Informant census: a new approach for identifying potentially effective medicinal plants. *Plants in Indigenous Medicine and Diet*. 1986;91:112. <https://doi.org/10.4324/9781315060385-6>
35. Henrich M, Ankli A, Frei B, Weimann C, Sticher O. Medicinal plants in Mexico: Healer's consensus and Cultural importance. *Social Science and Medicine*. 1998;47(11):1859-1871. [https://dx.doi.org/10.1016/s0277-9536\(98\)00181-6](https://dx.doi.org/10.1016/s0277-9536(98)00181-6)
36. Singh GA, Kumar A, Tewari DD. An ethnobotanical survey of medicinal plants used in Terai forest of western Nepal. *Journal of Ethnobiology and Ethnomedicine*. 2012;8:19. <https://dx.doi.org/10.1186/1746-4269-8-19>
37. Bhat P, Hedge GR, Hedge G, Mulgund GS. Ethnomedicinal plants to cure skin diseases—an account of the traditional knowledge in the coastal parts of central Western Ghats, Karnataka, India. *Journal of Ethnopharmacology*. 2013;151:493-502. <https://dx.doi.org/10.1016/j.jep.2013.10.062>
38. Abbas Z, Khan SM, Khan SW, Abbasi AM. Medicinal plants used by inhabitants of the Shigar Valley, Baltistan region of Karakorum range-Pakistan. *Journal of Ethnobiology and Ethnomedicine*. 2017;13:53. <http://dx.doi.org/10.1186/s13002-017-0172-9>
39. Chekole G. Ethnobotanical study of medicinal plants used against human ailments in Gubalafto District, Northern Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2017;13:55. <https://dx.doi.org/10.1186/s13002-017-0182-7>
40. Umair M, Altaf M, Abbasi AM. An ethnobotanical survey of indigenous medicinal plants in Hafizabad district, Punjab Pakistan. *PLoS One*. 2017;12(6):e0177912. <https://dx.doi.org/10.1371/journal.pone.0177912>
41. Kapoor IP, Singh B, Singh G, Heluani CS De, Lampasona MP De, Catalan CA. Chemistry and in vitro antioxidant activity of volatile oil and oleoresins of black pepper (*Piper nigrum*). *Journal of Agricultural and Food Chemistry*. 2009;57(12):5358-64. <https://doi.org/10.1021/jf900642x>
42. Hritcu L, Noumedem JA, Cioanca O, Hancianu M, Postu P, Mihasan M. Anxiolytic and antidepressant profile of the methanolic extract of *Piper nigrum* fruits in beta-amyloid (1-42) rat model of Alzheimer's disease. *Behavioral and Brain Functions*. 2015;11:13. <https://doi.org/10.1186/s12993-015-0059-7>
43. Tasleem F, Azhar I, Ali SN, Perveen S, Mahmood ZA. Analgesic and anti-inflammatory activities of *Piper nigrum* L. *Asian Pacific Journal of Tropical Medicine*. 2014;7S1:S461-S8. [https://doi.org/10.1016/S1995-7645\(14\)60275-3](https://doi.org/10.1016/S1995-7645(14)60275-3)
44. Ee GC, Lim CM, Lim CK, Rahmani M, Shaari K, Bong CF. Alkaloids from *Piper sarmentosum* and *Piper nigrum*. *Natural Product Research*. 2009;23(15):1416-23. <https://doi.org/10.1080/14786410902757998>
45. Kaleem M, Sheema H, Sarma D, Bano B. Protective effects of *Piper nigrum* and *Vinca rosea* in alloxan induced diabetic rats. *Indian Journal of Physiology and Pharmacology*. 2005;49:65-71.
46. Eleazu CO, Eleazu KC. Physico-chemical properties and antioxidative potentials of 6 new varieties of ginger (*Zingiber officinale*). *American Journal of Food Technology*. 2012;7(4):214-221. <https://doi.org/10.3923/ajft.2012.214.221>
47. Sunilson JAJ, Suraj R, Rejitha G, Anandarajagopa K. In vitro antibacterial evaluation of *Zingiber officinale*, *Curcuma longa* and *Alpinia galangal* extracts as natural foods preservatives. *American Journal of Food Technology*. 2009; 4(5):192-200. <https://doi.org/10.3923/ajft.2009.192.200>
48. Katiyar SK, Agarwal R, Mukhtar H. Inhibition of tumor promotion in SENCAR mouse skin by ethanol extract of *Zingiber officinale* rhizome. *Cancer Research*. 1996; 56(5):1023-1030.
49. Jung HW, Yoon CH, Park KM, Han HS, Park YK. Hexane fraction of *Zingiberis Rhizoma Crudus* extract inhibits the production of nitric oxide and proinflammatory cytokines in LPS-stimulated BV2 microglial cells via the NF-kappaB pathway. *Food Chemistry and Toxicology*. 2009; 47(6):1190-1197. <https://doi.org/10.1016/j.fct.2009.02.012>
50. Khaki AA, Khaki A. Antioxidant effect of ginger to prevents lead-induced liver tissue apoptosis in rat. *Journal of Medicinal Plants Research*. 2010;4(14):1492-1495. 10.5897/JMPR09.397.
51. Lin RJ, Chen CY, Chung LY, Yen CM. Larvicidal activities of ginger (*Zingiber officinale*) against *Angiostrongylus cantonensis*. *Acta Tropica*. 2010; 115(1- 2): 69-76. <https://doi.org/10.1016/j.actatropica.2009.12.007>
52. Kuhad A, Tirkey N, Pilkhwal S, Chopra K. 6-Gingerol prevents cisplatin-induced acute renal failure in rats. *BioFactors*. 2006;26(3):189-200. 10.1002/biof.5520260304. <https://doi.org/10.1002/biof.5520260304>
53. Dubey RD, Verma S, Rane D, Wani VK, Pandey AK, Paroha S. Comparative studies of anthelmintic activity of *Zingiber officinale* and *Cassia tora*. *International Journal of Chemistry and Pharmaceutical Sciences*. 2010;1:1-4
54. Chang JS, Wang KC, Yeh CF, Shieh DE, Chiang LC. Fresh ginger (*Zingiber officinale*) has anti-viral activity against human respiratory syncytial virus in human respiratory tract cell lines. *Journal of Ethnopharmacology*. 2013;145(1):146-151. <https://doi.org/10.1016/j.jep.2012.10.043>
55. Baliga MS, Haniadka R, Pereira MM, Thilakchand KR, Rao S, Arora R. Radioprotective effects of *Z. officinale* Roscoe (ginger): past, present and future. *Food & Function*. 2012;3(7):714-723. 10.1039/c2fo10225k. <https://doi.org/10.1039/c2fo10225k>
56. Mao QQ, Xu XY, Cao SY, Gan RY, Corke H, Beta T, Li HB. Bioactive compounds and bioactivities of ginger (*Zingiber officinale* Roscoe). *Foods*. 2019;8(185):1-21. <https://doi.org/10.3390/foods8060185>
57. Manju V, Nalini N. Chemoprotective efficacy of ginger, a naturally occurring anticarcinogen during the initiation, postinitiation stages of 1,2 dimethylhydrazine induced colon cancer. *Clin Chim Acta*. 2005;358(1-2):60-67. <https://doi.org/10.1016/j.cccn.2005.02.018>
58. Kabuto H, Nishizawa M, Tada M, Higashio C, Shishibori T, Kohno M. Zingerone [4-(4-hydroxy-3-methoxyphenyl)-2-butanone] prevents 6-hydroxydopamine depression in mouse striatum and increases superoxide scavenging activity in Serum. *Neurochemical Research*. 2005;30(3):325-332. <https://doi.org/10.1007/s11064-005-2606-3>

59. Malla B, Chhetri RB. Indigenous knowledge on medicinal non-timber forest products (NTFP) in Parbat district of Nepal. *Indo-Global Research Journal of Pharmaceutical Sciences*. 2012;2(2):213–225. <https://doi.org/10.35652/IGJPS.2012.26>.
60. Inta A, Trisonthi P, Trisonthi C. Analysis of Traditional knowledge in medicinal plants used by Yuan in Thailand. *Journal of Ethnopharmacology*. 2013; 149: 344–351. <https://dx.doi.org/10.1016/j.jep.2013.06.047>
61. Singh H, Husain T, Agnihotri P, Pande PC, Khatoon S. An ethnobotanical study of medicinal plants used in sacred groves of Kumaon Himalaya, Uttarakhand, India. *Journal of Ethnopharmacology*. 2014;154:98–108. <https://dx.doi.org/10.1016/j.jep.2014.03.026>
62. Mall B, Gauchan DP, Chhetri RB. An ethnobotanical study of medicinal plants used by ethnic people in Parbat District of western Nepal. *Journal of Ethnopharmacology*. 2015;165:13–17. <https://dx.doi.org/10.1016/j.jep.2014.12.057>
63. Boro M, Das B, Boro KK, Nath M, Buragohain P, Roy S, Sarma PJ, Kalita S, Nath N. Quantitative ethnobotany of medicinal plants used by the Bodo Community of Baksa District, Assam, India. *Biodiversitas: Journal of Biological Diversity*. 2023 ;24(6): 3169–3182. <https://doi.org/10.13057/biodiv/d240610>
64. Das B, Tissopi G, Devi N, Nath N. Study on folk remedies using medicinal plants by Karbi tribe of West Karbi Anglong District, Assam, India. *Indian Journal of Traditional Knowledge (IJTK)*. 2023; 22(3):576–86. <https://doi.org/HYPERLINK> "http://dx.doi.org/10.56042/ijtk.v22i3.57