



REVIEW ARTICLE

Understanding the advances in Sorghum grain quality improvement: An overview

Anamika Dhar¹, B. Meena Kumari^{2*}, D. Kavithamani², N. Manikanda Boopathi³ & P. Meenakshi⁴

¹Department of Genetics and Plant Breeding, TNAU, Coimbatore - 641 003

²Department of Millets, Centre for Plant Breeding and Genetics, TNAU, Coimbatore - 641 003

³Department of Plant Biotechnology, Centre for Plant Molecular Biology and Biotechnology, TNAU, Coimbatore - 641 003

⁴Department of Plant Molecular Biology and Bioinformatics, Centre for Plant Molecular Biology and Biotechnology, TNAU-Coimbatore - 641 003

*Email: meenakumari.b@tnau.ac.in



ARTICLE HISTORY

Received: 14 March 2024

Accepted: 01 May 2024

Available online

Version 1.0 : 16 June 2024

Version 2.0 : 01 July 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Dhar A, Kumari BM, Kavithamani D, Boopathi NM, Meenakshi P. Understanding the advances in Sorghum grain quality improvement: An overview. Plant Science Today. 2024; 11(3): 61-78. <https://doi.org/10.14719/pst.3527>

Abstract

Sorghum, a crucial cereal crop with versatile applications, is increasingly recognized for its grain quality attributes. The nutritional and biochemical diversity within sorghum, encompassing elements such as iron (Fe), zinc (Zn), proteins, starch, dietary fibers, and β -carotene, plays a pivotal role in enhancing the quality across diverse sorghum accessions. Breeding programs offer a promising avenue for further improvement in these traits. Additionally, sorghum features a spectrum of phenolic compounds, including tannins and flavonoids, influencing both pigmentation and potential health benefits. The antioxidative properties of these compounds underscore their critical role in promoting health and mitigating oxidative stress. The significance of sorghum is shaped by genetic factors, environmental conditions, ripening stages, and varietal distinctions, highlighting the complex interplay between grain structure, genetics, and nutritional content. As the global demand for diverse, nutritionally rich food sources continues to rise, this review provides insights aimed at deepening our understanding of sorghum's potential as a staple crop with substantial nutritional and health-promoting attributes.

Keywords

Sorghum grain; phenolic compounds; kafirin; flavonoids; quantitative trait loci; antioxidant activity

Introduction

Sorghum (*Sorghum bicolor* (L.) Moench), a cereal crop of the Poaceae family serves as an indispensable dietary staple for over 500 million individuals in Africa and Asia (1). It was originated in sub-Saharan Africa, domesticated, and later spread to India and China (2). It is a nutrient-rich, annual diploid crop species ($2n=2x=20$) and is one of the most significant cereal crops in the world and ranks 5th in terms of production behind maize, rice, wheat, and barley (3). The USA is the top sorghum-producing country contributing 14% whereas India contributes 7 % of the total world production. The major sorghum-producing states in India are Maharashtra, Karnataka, Tamil Nadu, Rajasthan, and Madhya Pradesh (4). As the majority of the human population depends on rice and wheat as their staple food, sorghum being the crop adapted to the dry land conditions can be used to optimize the utilization of dry land and increase the farmers' income. Also, being the C4 crop, sorghum is climate resilient and can be grown in harsh environmental conditions including temperature and water constraints. It is diversified into four main categories: grain sorghum, sweet sorghum, forage sorghum and biomass sorghum, each featuring unique genetic variations, heights, traits and applications (5). Traditionally, sorghum

has been utilized in various forms, including whole meal flour and milled fractions for dishes like 'rotis' and gruel. In some instances, sorghum is employed for puffing and creating special dishes, albeit to a limited extent. Sorghum malt is globally used, notably by the Bantu people in Africa where they have a traditional drink known as kafir beer (6). Although the current regional system introduces hybrids, these are less favored for human consumption. In contrast, local and improved landraces are preferred for household consumption because of their superior grain quality, characterized by the bold, white appearance and a sweeter taste (7).

Sorghum shares many of the same essential components as millet and maize, including starch, fat, protein, and non-starch polysaccharides. It is a good source of micro and macronutrients, carotenoids, and polyphenols, as well as bioactive nutrients like vitamin B and fat-soluble vitamins D, E, and K (8). Sorghum has distinct and variety-dependent nutritional qualities. It is gluten-free, has resistant starch and thus can be consumed by people with celiac disease (9). It is well known for having very low digestion of starch and protein and has the potential for controlling obesity and weight. Compared to other cereals, sorghum has high and diverse phenolic compounds including all types with the prevalence of simple phenolic acids, flavonoids, and tannins (10). Identifying the genes regulating the composition of sorghum grains would help modify grain texture and quality to suit current end-use markets and promote the development of new products. This overview delves into the intricate details of sorghum's carbohydrate profile, emphasizing the pivotal role of starch, its granular morphology, and the genetic variations that categorize sorghum grains into waxy and non-waxy types. Beyond carbohydrates, the exploration encompasses the fiber landscape of sorghum, encompassing insoluble and soluble fibers, including notable polysaccharides like β -D-glucans and phytoglycogens. It also extends to the bioactive compounds present in sorghum, which confer the grains with potential health benefits to humans like lowering oxidative stress, anti-cancerous activity, anti-inflammatory action, etc (11). The association studies, combining ability and variability among the diverse sorghum genotypes result in the identification of promising lines with enhanced nutrient contents. Marker trait association as well as the Quantitative Trait Loci (QTLs) was identified in various sorghum accessions for their quality traits. Therefore, this review aims to provide a holistic understanding of sorghum's grain quality, its importance, nutritional composition and therapeutic properties offering insights into its potential benefits for human health by improving its quality attributes through different breeding approaches for alleviating malnutrition in developing countries.

1. Sorghum classification and distribution

Sorghum, belonging to the Andropogoneae tribe within the grass family (Poaceae), encompasses approximately 25 species and 5 subgenera. Certain species within this genus have been cultivated for human consumption, utilized as animal fodder, or exploited for industrial purposes. Notably, *Sorghum bicolor* (L.) Moench is a widely cultivated species

globally. The remaining species are primarily found in the wild or as weeds, exhibiting varying levels of interspecific compatibility, ranging from 0 % to 100 % (12). The five subgenera identified within Sorghum are Chaetosorghum, Heterosorghum, Parasorghum, Stiposorghum, and Eusorghum. Among these, Eusorghum, which includes the cultivated species, *S. bicolor*, has been further classified into 5 major races (bicolor, durra, kafir, guinea, and caudatum) and 10 intermediate races (Fig. 1) represent hybrid combinations based on panicle characteristics (13). Indian sorghum varieties mainly include the Durra, Caudatum, Guinea, and Guinea-Kafir types, with a smaller amount of bicolor being grown (14). Durra is the primary cultivated sorghum race in India which is also found in Eastern Africa with large, globular grains. High-quality flours can be derived from caudatum, durra, and kafir races, contingent on specific cultivars. The guinea margaritifera subrace yields excellent grain, which is cooked like rice in Western Africa. In Mali, couscous made with guinea gambicum is highly valued. The white grain forms of *Guinea guineense* are ground into flour, while red grain forms are predominantly used in beer production. *Guinea roxburghii* serves as pop-sorghum in India and high-quality grain in southern Africa (15). In the United States, grain sorghum production is dominated by the Kafir-Caudatum variety. In Nigeria, the Kaura sorghum is classified as Durra-Caudatum, while Zerzeras and Hegaris are considered part of the Caudatum category. In Sudan, the Feterita sorghum ranges across different classifications from Guinea-Caudatum and Caudatum to Durra-Caudatum. Varieties such as broomcorns, sorgos, and Sudan grass are categorized under the bicolor type.

2. Sorghum grain structure, appearance, and its genetics

The bran layer that comprises of pericarp and testa, the endosperm, and the germ makes up the sorghum grain (Fig. 2) and their relative proportion varies depending upon variety and environment. Generally, sorghum contains 84.2 % endosperm, 9.4 % germ, and 6.5 % pericarp (16). The sorghum bran generally contains non-starch polysaccharides, a wide range of phenolic compounds, such as flavonoids, phenolic acids, and condensed tannins, as well as certain vitamins, such as carotenoids. Proteins, carbohydrates, and a few vitamins, including vitamin B complex, make up the majority of the endosperm. The germ fraction is primarily made up of lipids and proteins, but it also contains a lot of minerals and vitamins, particularly fat-soluble and complex B vitamins (17, 18). Some sorghum types may also contain a pigmented testa, which is situated between the pericarp and endosperm (17, 19). Genetic and environmental variables mostly determine sorghum quality and appearance. Its caryopsis color and general appearance are influenced by several genetically interrelated elements mentioned in Table 1. Color and thickness of the pericarp, the presence of pigmented testa, and endosperm color all have a major impact on appearance (19). Thus, sorghum can be categorized into 5 types: white, yellow, red, brown and black based on color, phenolic profile and genotypes (Fig. 3).

The 2 genes R and Y are responsible for controlling pericarp color. The combination of these genes expresses the white (R_{yy} or rryy), lemon yellow (rrY₁), or red (R_Y)




Fig. 1. Sorghum classification.




Fig. 2. Sorghum plant and grain structure

Table 1. Genes and their inheritance patterns associated with sorghum grain traits

Traits	Genes governing the traits	Inheritance	References
Pericarp color	Y, R	rryy/R_yy : white/colorless	(20)
		rrY_ : yellow	(21)
		R_Y_ : red/black	(22)
Testa color	B ₁ ,B ₂	B ₁ B ₂ : pigmented testa (brown/purple), B ₁ b ₂ b ₂ / b ₁ B ₁ B ₂ _ / b ₁ b ₁ b ₂ b ₂ : non-pigmented testa	(17)
Pericarp thickness	Z	Z_ : thin pericarp, ZZ : thick pericarp	(17)
Color intensity	I (Intensifier gene, only for red and yellow pericarp)	I_ : increased intensity, ii : lighter colors	(127)

**Fig. 3.** Sorghum grain color diversity. A : red grains, B: light red/orange grains, C: white grains, D: yellow grains, E: black grains.

color. White sorghum has low total phenolic content and tannin and 3-deoxyanthocyanidin levels are extremely low or non-existent (20). Yellow sorghum is greater in total phenolic content and higher in flavanones than white sorghum (21). Red sorghum has a significant amount of phenolic chemicals and condensed tannins. Black sorghum is essentially a unique form of red sorghum at the genetic level, where the pericarp undergoes a transformation to black under the influence of solar radiation during maturation. The pericarp of black sorghum contains a high concentration of phenolic compounds, notably 3-deoxyanthocyanidin (22). The presence of a pigmented testa layer is governed by B₁ and B₂ genes. Condensed tannins are found in sorghums classed as type III (also known as brown or bird-resistant sorghum) that have pigmented testa (B₁B₂) and a dominant spreader (S₁) gene. Pericarp thickness is controlled by the Z gene ranging from 8 to 160 mm (23), and varies within an individual kernel. Sorghum carrying homozygous recessive (rr) genes have thick mesocarp. The presence of a thick pericarp typically

masks the color of the testa and endosperm, resulting in kernels that appear chalky due to the existence of small starch granules in the pericarp (23). The Wx gene impacts endosperm starch composition. The dominant Wx allele balances amylose and amylopectin, while the recessive wx allele creates a waxy endosperm rich in amylopectin (24). The genetics of important sorghum grain quality traits such as grain size, grain shape, and grain lustre were studied. While grain size and grain shape are governed by dominant genes that are polygenic in nature, the grain lustre trait is under the control of complementary recessive genes (25).

3. Nutritional Composition and Functional Attributes of Sorghum Grain

Carbohydrates are the most abundant component of sorghum grain, followed by proteins, lipids, fiber, dietary fiber, and ash (26). On average, 100 g of the grain consists of approximately 72.1 g carbohydrates, 12.4 g water, 10.6 g proteins, 6.7 g fibers, and 3.5 g fats, providing an estimated energy of around 1,377 kJ (27).

3.1 Polysaccharides: Starch, fibers, and their significance

Sorghum is rich in starch, primarily stored in the endosperm, with granule morphology varying between floury and corneous regions, as observed (28). The granules in the corneous endosperm are smaller and possess a more angular shape, while those in the floury endosperm are larger and exhibit a spherical form. Starch consists of amylose and amylopectin and their proportion affects the functional and nutritional attributes of sorghum starch and starch-derived products. Amylose plays a role in starch thermal characteristics like gelatinization and pasting (29). Foods high in amylose have been demonstrated to have a lower glycemic index, which promotes various health benefits, including improved diabetes and obesity control (30). Corneous endosperm starch had less amylose than floury endosperm starch (31). Genetic diversity in amylose concentration is evident, with levels ranging from 24 to 33 % in 95 Zimbabwean sorghum landraces and 16.1 % to 55.8 % in 55 different genotypes (32, 33). The majority of sorghum grains contain 20-30 % amylose, which is common for normal cereal grains (28). Sorghum grains are categorized as waxy and non-waxy, with waxy grains containing a higher amount of amylopectin and minimal amylose (up to 5 %) and exhibiting distinctive characteristics like higher starch digestibility, more susceptible to enzymatic degradation, etc (34, 35). Moreover, Waxy sorghum starch has quick cooking, high peak viscosity, and low stability during cooking. It resists gel formation and retrogradation (36).

Apart from being a source of carbohydrates, sorghum is rich in fibers, both insoluble (cellulose and lignin) and soluble (arabinoxylans, β -glucans), that are located in the pericarp and endosperm cell walls and account for 6 to 15 g per 100 g of grain (37) with significant phenolic acids like p-coumaric and ferulic acids (38). The β -D-glucan content in sorghum grain varies from 0.12 % (39) to \approx 0.09 % with Glucan A and B identified as water-soluble glucans (40). A few varieties, such as sweet sorghum, contain water-soluble polysaccharides called phytoglycogens, with the mutant gene "su" being homologous to sweet corn (41). The average chain length of sorghum phytoglycogen is 9.4–13.4 and the content can reach up to 13.2 % of the grain.

3.2 Sorghum Protein Composition: Distribution, Classes and Nutritional Implications

Sorghum protein distribution reveals that approximately 80 %, 16 %, and 3 % are found in the endosperm, germ, and pericarp respectively (42). The major protein fractions in sorghum include prolamin proteins, particularly kafirins, and non-prolamin proteins like globulins, glutelins, and albumins. Prolamin proteins, primarily situated in the endosperm's protein bodies, constitute the majority of the protein fraction, with nitrogen fertilizer significantly increasing prolamin levels in sorghum and millets (43). Sorghum exhibits greater hardness compared to other food grains, attributed to its elevated prolamin concentration ranging from 3.6 % to 5.1 % and lysine content ranging from 1.06 to 3.64 (44). Kafirins are categorized into α -, β -, γ - and δ -kafirins and makeup 48 % to 70 % of sorghum grain's total protein content (45), exhibiting high polymerization and resistance to enzymatic digestion in the digestive tract (46). The corneous endosperm has higher

kafirin levels, while the opaque endosperm contains more albumins and globulins, with equal glutelin concentration in both forms (47). Glutelins, the second-largest high molecular protein fraction located in the endosperm's protein matrix (48). Sorghum proteins are lysine-deficient but rich in glutamic acid, leucine and proline. Cultivars from breeding programs exhibit 52 % to 115 % higher lysine compared to conventional varieties (42). Earlier studies reported crude protein levels of 8 % to 12.5 %. At the same time, a recent examination of 19 sorghum genotypes showed significant variation, emphasizing the impact of landrace selection in developing biofortified sorghum, with high-protein genotypes PYPS 2 and PYPS 13 emerging from landraces PSLRC 2 and PSLRC 21 (49).

3.3 Carotenoids in Sorghum: Endosperm Color and Composition

Carotenoids are the characteristic yellow or orange pigments present in various seeds, leaves, and plant materials. In sorghum, particularly in the endosperm, show a significant association with grain color. Sorghum is broadly categorized into white endosperm, with low carotenoid levels, and yellow endosperm, with elevated carotenoid levels, primarily lutein and zeaxanthin (50). Yellow endosperm sorghum demonstrates higher carotenoid content, with zeaxanthin identified as the predominant carotenoid in most varieties and β -carotene exclusively found in yellow endosperm genotypes (51). Lutein and zeaxanthin are highlighted as the most prevalent carotenoids in grain sorghum, although at concentrations lower than those in yellow maize (52, 53). Enhanced bioavailability of provitamin A, specifically β -carotene, has been documented in certain transgenic sorghum lines, with 5 Quantitative Trait Loci identified on chromosomes 1, 2, and 10. Yellow endosperm sorghum exhibits higher levels of lutein, zeaxanthin, and β -carotene compared to white endosperm types, with crossbreeding resulting in inbreds with intermediate carotenoid values (54). A study underscored zeaxanthin as the predominant carotenoid in most sorghum varieties, with β -carotene or provitamin A activity exclusively identified in yellow endosperm genotypes (51). Other carotenoids present in smaller proportions include α -cryptoxanthin, β -cryptoxanthin, and α -carotenes (52, 53). A study investigated carotenoid and pro-vitamin A content in sorghum accessions with different pericarp pigments using LC-MS (55). Among the 5 identified carotenoids, 3 (α -carotene, β -carotene and β -cryptoxanthin) are vitamin A precursors. Sorghum with yellow pericarp exhibited the highest total carotenoid content, while brown or yellow pericarp varieties had the highest β -carotene. White pericarp sorghum had the lowest carotenoid levels. Pro-vitamin A content was highest in yellow pericarp, followed by brown and white pericarp varieties. The study suggests that breeding sorghum with specific pericarp pigments may help to prevent vitamin A deficiency in regions where sorghum is a staple crop.

3.4 Mineral Composition of Sorghum Grains: Fe and Zn Variability and Genetic Insights

Sorghum grains have been extensively studied for their iron (Fe) and zinc (Zn) content due to their essential role in

human nutrition. Research has consistently shown significant variability in the mineral content of sorghum grains among different varieties investigated the genetic and environmental impact on iron, zinc, and phytate in food sorghum, emphasizing the importance of understanding the factors influencing the availability of these minerals in sorghum varieties. Fe ranged from 30 to 113 mg/kg, Zn from 11 to 44 mg/kg, and phytate from 0.4 to 3.5 % and Fe and Zn did not consistently link to genetics but varied across locations, indicating environmental impact (56). Additionally, there are substantial differences in mineral content among different sorghum varieties, with greater Fe and Zn in red and black varieties compared to white (57). A study was conducted in a total of 19 sorghum genotypes derived from 11 superior landraces, where significant variations in the concentration of total and available Fe and Zn were found. Most sorghum genotypes demonstrated elevated levels of iron (ranging from 14.21 to 28.41 mg/100 g) and zinc (ranging from 4.81 to 8.16 mg/100 g). Thus, this study identified high Fe and Zn-containing landraces which underscores the potential of utilizing them for biofortification purposes.

Moreover, quantitative trait loci (QTLs) and candidate genes associated with high grain Fe and Zn concentrations in sorghum, provide insights into the genetic basis of mineral accumulation in sorghum grains (58). Twenty-three sorghum landrace accessions representing 5 distinct color types: white, buff, red, brown, and purple collected from 5 different countries, revealed significant variability among the accessions in terms of Fe and Zn levels, whereas no significant differences were observed in the Mn and Na contents (59). The concentration of Fe ranged from 25 µg/g to 42 µg/g, with the highest recorded concentration being 42 µg/g and the lowest at 25 µg/g. Zn concentrations exhibited variation within the range of 18 µg/g to 30 µg/g. In an assessment of the zinc (Zn) and iron (Fe) content in 29 accessions, specifically selected for their elevated mineral levels, as part of the screening process involving over 2200 sorghum lines, the iron content ranged from 26 to 60 mg/kg, while the zinc content varied between 21 and 57 mg/kg (60).

4. Antinutritional Factors in Sorghum: Implications for Nutrient Accessibility and Bioavailability

In addition to examining the nutritional composition, it is crucial to consider the accessibility of nutrients for consumers, particularly given the variation in naturally occurring antinutritional factors that diminish the nutritional value of sorghum compared to wheat or maize (61). These factors include phytates, enzyme inhibitors, condensed tannins, and others, collectively compromising the digestibility of the grain. Phytic acids or phytates form complexes with essential dietary minerals like calcium, zinc, iron, and magnesium, rendering them less available for absorption (62). This process involves the chelation of positively charged minerals, such as iron and zinc, resulting in insoluble complexes and reduced mineral absorption. Besides, the phytate forms complexes with proteins, slowing the digestion rate and impeding their bioavailability (63).

Primarily serving as a reservoir of phosphorus and inositol, phytates in sorghum are predominantly located in the aleurone layer (phytin bodies or aleurone grains) and to a lesser extent, in the germ. Abrasive decortication of sorghum can remove 40 to 50 % of phytate and total phosphorus. The levels of phytate-phosphorus in 30 sorghum varieties ranged from 0.17 % to 0.38 %, constituting 80 % to 87 % of the total phosphorus in the kernel (64). Some low phytic acid-containing landraces, such as Malkhed-1, Nalwar-2, and the variety Phule Maulee, have been identified (65). Because of these anti-nutrient interactions and sorghum's low bioavailability in comparison to other grains (66), maximizing the potential of nutrient-rich landraces and genotypes such as PSLRC-2, PSLRC-3, PSLRC-9, PYPS 2 and PYPS 13 by reducing anti-nutritive factors should be a priority, particularly in Asia and Sub-Saharan Africa, where malnutrition and nutritional deficiencies are common in women and children (49).

5. Phenolic Compounds in Sorghum: Diversity, Distribution and Health Implications

Phenolic compounds, which contain at least one aromatic ring and one or more hydroxyl groups, are categorized into soluble compounds like flavonoids, quinones, and phenylpropanoids within the plant cell vacuole. They also include insoluble compounds such as lignins, condensed tannins, and hydroxycinnamic acid, which are attached to the cell wall (67). These compounds contribute significantly to the antioxidant properties of sorghum, with potential health benefits linked to chronic disease prevention such as breast and colon cancer as well as diabetes (68). The composition of phenolic compounds in sorghum is influenced by factors such as variety, genotype, pericarp color, and grain testa pigmentation (69). The range of the phenolic compounds estimated in various sorghum accessions is mentioned in Table 2.

Phenolic acids, constituents of the grain's endosperm, pericarp, and testa exist in both free and bound and are abundant in sorghum grains, with ferulic acid being the most prevalent in the bound form (70, 71). Phenolic acids, the simplest yet abundant phenolic compounds present in all sorghum grains, exhibit a total concentration ranging from 445 to 2850 µg/g (72). Notable phenolic acids reported in sorghum include gallic, vanillic, protocatechuic, cinnamic, p-coumaric, p-hydroxybenzoic, syringic, ferulic, caffeic, and sinapic acids (73- 75). Studies on sorghum genetic resources reveal a wide range of total polyphenol content, with ferulic acid consistently exhibiting the lowest content among examined phenolic compounds (76). Higher total phenol levels in specific sorghum genotypes under salinity stress suggest a potential adaptive mechanism for mitigating sub-cellular damage during stress conditions (77). Variability in Total Phenolic Content (TPC) is observed among sorghum accessions from different countries (78). Genotypes with red and brown pericarp show elevated levels of total proanthocyanidin and total phenolic content, contributing to heightened antioxidant activities (79). Sorghum polyphenol extracts have antioxidant properties and show

Table 2. Total phenol content (TPC), Total flavonoid content (TFC), and Total tannin content (TTC) ranges in different sorghum accessions

Sorghum source	Extract solvent	TPC Range (mg GAE/g)	TFC Range	TTC Range	References
Moroccan ecotypes	Methanol	125.86 - 314.91	4.53 - 14.04 mg GAE/g	-	(128)
Kenyan varieties	Methanol	7.9 - 29.6	-	-	(109)
Australian varieties	Methanol	0.24 - 11.50	-	-	(129)
Senegalese varieties	Methanol	0.22 - 0.57	-	-	(130)
Genetic resources collected from 5 countries	Methanol	16.33 - 231.21	16.46 - 67.71 mg QE/g	-	(76)
Accessions from 15 regions	Methanol	9.97 - 333.58	10.35 - 252.67 mg QE/g	-	(78)
Breeding lines	Methanol	0.29 - 10.38	-	0.68 - 434.22 mg CE/g	(131)
Black sorghum lines	Methanol	5 - 20	-	-	(22)
Black sorghum hybrids	Methanol	9 - 17	-	14 - 19 mg CE/g	(132)
Korean sorghum landraces	Methanol	1.17 - 10.23	-	0.12 - 428.95 mg CE/g	(132)
Sorghum landraces	Methanol	1.37 - 9.20	-	25.18 - 408.84 mg CE/g	(133)
Sorghum genotypes		173.68 - 1040.73	-	1.13 - 12.4 mg/100 g	(79)

inhibition of the activity of cholesterol biosynthesis-related enzymes (80). Biologically active compounds linked to sorghum polyphenols have been shown to lower blood LDL (low-density lipoprotein) cholesterol levels and elevate HDL (high-density lipoprotein) cholesterol, demonstrating significant hypocholesterolemic effects. Sorghum's cholesterol-lowering impact has been recognized compared to cereals like wheat, oats and millet as the sorghum diet reduced the absorption of cholesterol in the intestine (81). In a study, it was found that sorghum extracts reduce cholesterol levels by influencing the expression of proteins involved in hepatic cholesterol metabolism (82).

Tannins, a diverse group of polyphenolic polymers, exhibit varying molecular weights and complexity and are associated with numerous health benefits, including antioxidant properties, anticancer activities, and cardioprotective effects (83). Despite their positive health effects, tannins also possess anti-nutritional properties, hindering protein digestibility and iron absorption through complex formation (84). Influenced by phenol-protein ratios, tannins form soluble and insoluble complexes with proteins and impact carbohydrates in sorghum, including hemicellulose, cellulose, starch, and pectin (85). Sorghum predominantly contains condensed tannins also known as proanthocyanidins, with hydrolyzable tannins found in various plants demonstrating diverse biological capacities (86, 87). Sorghum tannins, mainly condensed, vary significantly among varieties, with black testa varieties having the highest content (88). Sorghum varieties are classified into three types based on tannin content, with Type I having low levels, Type II moderate levels, and Type III high levels. Type I sorghums, characterized by recessive B_1 and/or B_2 genes ($B_1b_2b_2$, $b_1b_1B_2$, or $b_1b_1b_2b_2$), Type II sorghums possess dominant B_1 and B_2 genes but a homozygous recessive S gene (B_1B_2ss) and Type III

sorghums, with dominant B_1 , B_2 and S genes (B_1B_2S) (17, 89).

Sorghum grains harbor condensed or non-hydrolyzable tannins with varied degrees of polymerization degrees, including dimers, oligomers, and polymers of catechins, exhibiting variations in flavanol interlinkages, phenolic hydroxylation patterns and the configuration of the hydroxylated C-ring C3 center of the flavan-3-ol building block (90). The primary unit of condensed tannins is (epi)catechin, accompanied by significant units like (epi)galocatechin, (epi)galocatechin gallate, and (epi)afzelechin (91). Specifically, procyanidins, a type of condensed tannin in sorghum grains, act as potent radical scavengers, significantly enhancing antioxidant capacity. Variability in condensed tannin levels is observed among sorghum bran varieties, with coarse sorghum bran exhibiting the highest content (179.77 mg/100 g) and fine sorghum bran having the lowest content (172.38 mg/100 g). This variability may be attributed to factors such as ripening stage, varietal differences, climatic conditions, or their interplay (92).

Flavonoids, primarily concentrated in sorghum bran, exhibit diverse types and concentrations influenced by factors such as pericarp color, thickness, and the presence of pigmented testa (93). Representing the most abundant and diverse class of phenolic compounds in sorghum, flavonoids, with recognized health benefits including antioxidant, anticancer, anti-inflammatory, and gastroprotective properties, exist in both free (soluble) and bound forms (94). Specific flavonoid compounds identified in sorghum include flavan-4-ols, flavanones, flavonols, and dihydroflavonols, with notable content in red and black sorghums (22, 89, 95). Sorghum is a primary source of rare flavonoids known as 3-deoxyanthocyanins, contributing to red-to-black pigmentation in various plant

parts. Particularly, black sorghum grains exhibit elevated levels of these compounds in the bran, ranging from 4 to 16 mg/g (96). Significant variability in total flavonoid content among sorghum landraces underscores their diversity. The diversity in flavonoid levels among sorghum landraces may result not only from genetic differences but also from variations in collection sites, indicating an influence of environmental factors on flavonoid concentration, as observed in the study conducted (97). The flavonoid concentrations in 8 sorghum genotypes revealed significant variations influenced more by genotype than environmental factors (98). In black sorghum, four 3-deoxy anthocyanidin compounds, including apigeninidin (AP), luteolinidin (LUT), 7-methoxyapigeninidin (7-MeO-AP) and 5-methoxyluteolinidin (5-MeO-LUT), exhibited diversity in locations and genotypes. Flavones, yellow-colored flavonoids present in sorghum grain with varying concentrations influenced by factors such as pericarp color and thickness, with red and yellow pericarp sorghums often associated with elevated flavone levels (21, 99). A study conducted on 8 sorghum varieties in Austria aimed to assess their antioxidant content and activity. Flavonoids were detected in quantities ranging from 27.03 $\mu\text{g/g}$ (Kalatur variety) to 87.52 $\mu\text{g/g}$ (Huggo variety) (100). On another note, sorghum flavonoids are pivotal for resisting biotic stress. Various diet-based feeding experiments have underscored the importance of testing flavonoids and phenolics against challenging herbivores, implicating these compounds in host plant resistance (101). Notably, research indicates that flavonoids in sorghum adversely affect Fall armyworm larval growth and survival, induce changes in the peritrophic membrane of the larval midgut and mitigate herbivory damage (102, 103) The role of sorghum flavonoids in resisting corn leaf aphids (CLA) *Rhopalosiphum maidis* was studied and how the sorghum flavonoids, controlled by the yellow seed1 (*y1*) gene, repel CLA (106). Sorghum with functional *y1* alleles accumulated flavonoids, deterring CLA, while those lacking *y1* alleles attracted more aphids. Flavonoids from functional *y1* plants increased CLA mortality compared to those from plants lacking *y1* alleles or controls. This underscores the significance of *y1*-regulated flavonoids in sorghum's defense against CLA.

6. Sorghum Phenolics: Antioxidant, Anticancer and Antidiabetic Properties

Sorghum and its co-products have gained recognition as valuable sources of nutraceuticals and functional foods, owing to their high content of phenolic compounds, some of the most significant naturally occurring antioxidants and chemopreventive agents (104, 105). The antioxidant activity of sorghum phenolics, assessed through various widely-used *in vitro* assays such as Oxygen radical absorbance capacity (ORAC), Ferric reducing antioxidant power (FRAP), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) and 1,1-diphenyl-2-picrylhydrazyl (DPPH), contributes significantly to the associated health benefits (106). However, it is important to acknowledge

the limitations of these assays in directly translating to *in vivo* antioxidant capacity, considering the complexities of the physiological environment like pH, temperature, bioavailability, and metabolism. Sorghum phenolic compounds act as natural antioxidants, protecting biomolecules from oxidative damage caused by reactive oxygen species, making sorghum grains a valuable source of readily available antioxidants through dietary intake (107). Pigmented sorghum varieties, in particular, exhibit significantly higher antioxidant activity compared to red wheat (108). Recent studies have highlighted strong correlations between phenolic compounds and antioxidant activities in sorghum, with flavonoids and tannins demonstrating significant associations with FRAP and total antioxidant activity (97). Beyond direct antioxidant effects, sorghum phenolics, including 3-deoxyanthocyanidins and condensed tannins, offer additional health benefits through mechanisms such as induction of endogenous detoxifying enzymes and acting as free radical "sinks" to mitigate oxidative stress (11, 109, 110). Sorghum, known for its rich content of specific compounds, including phenolic extracts, demonstrates potential anticancer properties as evidenced by *in vitro* and animal studies, showcasing the ability to reduce and inhibit cancer cell proliferation in the colon, liver, and esophagus (111). Flavones in sorghum, exhibiting estrogenic effects, contribute to *in vitro* anticancer effects, while 3-deoxyanthocyanins in black sorghum show similar properties. The presence of condensed tannins in sorghum, particularly in brown sorghum bran, may play a crucial role in cancer prevention by inhibiting aromatase associated with breast cancer, thereby impeding the formation of cancer growth stimuli (112). Sorghum bran, recognized for its abundance of phenolic substances, is also beneficial for individuals at risk of diabetes due to its low glycemic index (113). In studies investigating the antidiabetic properties of sorghum phenol, sorghum phenolic extract significantly reduced plasma glucose concentrations in diabetic rats, comparable to the effectiveness of the antidiabetic drug glibenclamide (114, 115). Additionally, tannin-rich polyphenolic extracts from sorghum grain demonstrated inhibitory activity against α -amylase and α -glucosidase enzymes, with *in vivo* studies showing significant hypoglycemic activity in diabetic rats, reducing various metabolic parameters (114, 116).

7. Breeding for improving the sorghum grain quality

Sorghum, renowned for its substantial genetic diversity, serves as a valuable resource for breeding programs, particularly in the context of enhancing nutritional profiles. A comprehensive analysis of 265 sorghum accessions over 3 years, revealed substantial variation in grain protein (8.1-18.8 %), fat (1.0-4.3 %), and starch (61.7-71.1 %) content (117). The study identified high heritability for protein, followed by fat and starch, highlighting significant genetic contributions to trait variation. Specific races exhibited distinct compositions, with durras/bicolor-durras from Ethiopia/India having the highest protein and fat, while kafirs from the USA/India/South Africa displayed the lowest protein and the highest starch content. These

findings not only offer new genetic sources for enhancing sorghum nutrition but also provide insights into the genetic architecture influencing grain composition variation.

A total of 108 sorghum landraces from Tigray, Ethiopia, along with 2 improved varieties, Melkam and Dekeba, were analyzed for genetic diversity and associations in grain yield and nutritional traits. Significant variations were observed among genotypes in all studied traits. The average starch, protein, ash, zinc, iron, calcium and magnesium contents were 65.1 %, 10.3 %, 1.9 %, 35.3 ppm, 36.8 ppm, 243 ppm and 1096.8 ppm respectively. LR14, LR65, LR10, LR23 and LR38 exhibited the highest values for these traits. Strong associations were found among traits and ten promising landraces were selected based on Genotype x yield x trait (GYT) analysis for potential use in breeding programs in Tigray and similar environments (118).

A study conducted a study employing three sets of full diallel crosses with 15 parents to investigate the inheritance patterns of grain iron (Fe) and zinc (Zn) concentrations in sorghum (119). It was found that the genetic regulation of grain zinc (Zn) concentrations, primarily governed by additive gene effects, suggests the efficacy of progeny selection for Zn enhancement. Conversely, grain iron (Fe) concentrations were influenced predominantly by non-additive gene effects, emphasizing the potential of heterosis breeding in addition to progeny selection for increased Fe levels. A study further contributed by developing 12 new hybrids, revealing the

importance of non-additive gene action in improving nutritional traits, with some crosses outperforming parents for both Fe and Zn concentrations without yield penalties (120).

Additionally, sorghum's adaptability to diverse environments contributes to its significant polyphenol diversity. This diversity presents opportunities for biofortification and crop enhancement, integrating desirable traits into elite varieties, but requires quantitative phenotyping to identify alleles responsible for quantitative variation in grain polyphenols (121). An average total polyphenol concentration of 7.00 mg GAE/g, with mean proanthocyanidin and 3-deoxyanthocyanidin concentrations at 7.73 mg CE/g and 27.40 abs/mL/g respectively was earlier reported (122), showcasing the potential for diverse polyphenolic content in sorghum grains. More advances have been achieved in breeding sorghum varieties (ATx3363 and BTx3363 germplasms) with elevated levels of 3-deoxyanthocyanidins in the grain pericarp while maintaining satisfactory grain yield (123). Moreover, different studies related to the correlation between quality traits of sorghum are mentioned in Table 3.

8. QTL mapping, Association mapping and GWAS in Sorghum for quality traits

Sorghum, a versatile crop used for food, animal feed, biofuels and more, exhibits diverse quality traits influenced by complex genetic architectures. Association mapping and Genome-Wide Association Studies (GWAS) offer powerful tools to unravel the genetic basis of these traits, aiding in

Table 3. Pearson's correlation between different quality traits (phenolic compounds, macro and the micronutrients, Estimated glycemic index) in sorghum

Trait 1	Trait 2	Correlation	References
Total phenols	Proanthocyanidins	Strong positive correlation	
Total phenols	3-deoxyanthocyanidins	Weak positive correlation	
Grain weight	Proanthocyanidins	No significant correlation	(122)
Grain weight	3-deoxyanthocyanidins	No significant correlation	
Grain weight	Total phenols	Small negative correlation	
Phosphorus	Magnesium	Strong positive correlation	
Phosphorus	Potassium	Weak positive correlation	
Phosphorus	Protein	Weak positive correlation	
Iron	Zinc	Significant positive correlation	
Iron	Manganese	Significant positive correlation	
Phosphorus	Calcium	No significant correlation	(59)
Phosphorus	Iron	Significant positive correlation	
Calcium	Sodium	Significant positive correlation	
Protein	Magnesium	Significant positive correlation	
Zinc	Magnesium	Significant positive correlation	
Zinc	Phosphorus	Significant positive correlation	
Manganese	Zinc	Significant positive correlation	
Protein	Zinc	Significant positive correlation	(134)
Iron	Zinc	Strong positive correlation	
Total starch	Estimated glycemic index	Significant positive correlation	
Non-resistant starch	Estimated glycemic index	Significant positive correlation	
β- glucan	Estimated glycemic index	Significant negative correlation	(135)
Flavonoids, phenolic compounds, antioxidant activity	Estimated glycemic index	Significant negative correlation	

Table 4. Quantitative trait loci (QTLs) identified for different nutritional and therapeutic traits

Traits	QTLs/Gene	Chromosome number	Population studied	References
Starch	QGSTR1.1	1	BTx623/Rio	(136)
	QPRCN3.1	3		
	QGSTR3.4, QGSTR3.2 QGSTR3.3	3	RTx430/Sureno	(137)
	QGSTR1.2, QGSTR1.3	1	ICRISAT mini core collection, landraces, Chinese cultivars	(138)
	QGSTR3.5	3		
	QGSTR4.3	4		
	QGSTR4.2	4	Diversity panel	(126)
	QGSTR3.1	3	Diverse set	(125)
Proteins	QPRCN1.1	1	BTx623/Rio	(136)
	QPRCN1.4	1		
	QPRCN2.15, QPRCN2.20, QPRCN2.16, QPRCN2.29, QPRCN2.4, QPRCN2.17, QPRCN2.11, QPRCN2.10, QPRCN2.5, QPRCN2.2, QPRCN2.12, QPRCN2.6, QPRCN2.35, QPRCN2.32, QPRCN2.7, QPRCN2.3, QPRCN2.13, QPRCN2.24, QPRCN2.14, QPRCN2.1, QPRCN2.8, QPRCN2.9, QPRCN2.33, QPRCN2.18, QPRCN2.19, QPRCN2.25, QPRCN2.30, QPRCN2.21, QPRCN2.22, QPRCN2.28, QPRCN2.31, QPRCN2.27, QPRCN2.26, QPRCN2.23	2	Sorghum Association Panel (SAP)	(139)
	QPRCN9.1	9	RTx430/Sureno	(137)
	QPRCN10.2, QPRCN10.3, QPRCN10.1	10		
	QPRCN9.2	9		
	QPRCN5.1	5	Diversity Panel	(140)
	QPRCN2.36, QPRCN2.38, QPRCN2.37, QPRCN2.39	2		

Carotenoids	QGRCC2.2, QGRCC2.3, QGRCC2.4, QGRCC2.5	2		
	QGRCC3.2, QGRCC3.5, QGRCC3.4, QGRCC3.3	3		
	QGRCC4.2	4		
	QGRCC5.1, QGRCC5.2	5		
	QGRCC6.1, QGRCC6.2 QGRCC6.3, QGRCC6.29 QGRCC6.4, QGRCC6.7, QGRCC6.9, QGRCC6.8, QGRCC6.6, QGRCC6.10, QGRCC6.11, QGRCC6.31, QGRCC6.12, QGRCC6.13, QGRCC6.14, QGRCC6.15, QGRCC6.16, QGRCC6.17, QGRCC6.30, QGRCC6.18, QGRCC6.19, QGRCC6.20, QGRCC6.21, QGRCC6.22, QGRCC6.23, QGRCC6.24, QGRCC6.25, QGRCC6.26, QGRCC6.27, QGRCC6.28	6	Sorghum Association Panel (SAP)	(141)
	QGRCC7.3, QGRCC7.4	7		
	QGRCC1.1, QGRCC1.2, QGRCC1.3, QGRCC1.4, QGRCC1.5	1		
	QGRCC4.1	4		
	QGRCC6.1	6	KS115/Macia	(50)
	QGRCC7.1, QGRCC7.2	7		
	QGRCC10.1, QGRCC10.2, QGRCC10.3, QGRCC10.5, QGRCC10.6	10		

Polyphenol	QPCGA1.2, QPCGA1.1	1				
	QPCGA2.1, QPCGA2.2	2				
	QPCGA4.11, QPCGA4.12, QPCGA4.14, QPCGA4.15, QPCGA4.8, QPCGA4.6, QPCGA4.16, QPCGA4.13, QPCGA4.7, QPCGA4.1, QPCGA4.10, QPCGA4.2, QPCGA4.5, QPCGA4.9, QPCGA4.3, QPCGA4.4	4		Sorghum Association Panel (SAP)	(117)	
	QPCGA5.2, QPCGA5.3, QPCGA5.1	5				
	QPCGA7.3, QPCGA7.4, QPCGA7.1, QPCGA7.2, QPCGA7.5					
	QPCGA8.1	8				
	QPCGA7.7, QPCGA7.6	7				
	QPCGA6.1	6				
	QPCGA4.32, QPCGA4.31, QPCGA4.25, QPCGA4.27, QPCGA4.29, QPCGA4.28, QPCGA4.30, QPCGA4.26, QPCGA4.24, QPCGA4.22, QPCGA4.20, QPCGA4.23, QPCGA4.21, QPCGA4.17, QPCGA4.19, QPCGA4.18	4		Diverse landraces and <i>S. bicolor</i> x <i>S. halepense</i> RIL	(142)	
	QPCGA1.4, QPCGA1.5, QPCGA1.3	1				
	Tannins	QTANN4.13, QTANN4.14, QTANN4.15, QTANN4.16, QTANN4.17, QTANN4.18	4		NAM (BTx623 x Niu Shen Zui, Hong Ke Zi & Kaoliang) - JLM	(143)
		QTANN2.4	2			
		QTANN10.1	10			
		QTANN1.1	1			
		QTANN2.2, QTANN2.3	2			
		QTANN3.2, QTANN3.3	3			
		QTANN4.10, QTANN4.11, QTANN4.12	4		Diversity panel	(126)
		QTANN5.3	5			
		QTANN7.4	7			
		QTANN8.1	8			
QTANN9.2		9				
QTANN2.1		2				
QTANN3.1		3				
QTANN4.9, QTANN4.7, QTANN4.2, QTANN4.6, QTANN4.8, QTANN4.4, QTANN4.3, QTANN4.5		4		Sorghum Association Panel (SAP)	(117)	
QTANN5.1, QTANN5.2		5				
QTANN7.3, QTANN7.1, QTANN7.2		7				
QTANN9.1		9				

Source: Sorghum QTL Atlas | OZ Sorghum (aussorgm.org.au) (124)

targeted breeding and improvement. QTLs for different quality traits have been reported and recorded in the Sorghum QTL Atlas (<http://aussorgm.org.au/sorghum-qtl-atlas/>) (124) and the identified QTLs for some traits are also mentioned in Table 4.

A study on candidate gene association mapping in a diverse set of 300 sorghum accessions to evaluate marker-trait associations related to 10-grain quality traits, using 1290 single nucleotide polymorphisms (SNPs) separated the panel into 5 subpopulations (125). Variations were observed in kernel hardness, acid detergent fiber and total digestible nutrients among the subpopulations. Following the model testing, association analysis between 333 SNPs in candidate genes and/or loci and grain quality traits revealed eight significant marker-trait associations (Table 5).

Genome-wide association study in genotypes of grain sorghum identified significant SNPs associated with tannins, starch and amino acids, revealing 14, 15 and 711 genetic loci respectively (126). Notably, SNPs for tannins colocalized with known loci, while those for starch linked to the sucrose phosphate synthase gene. Homologs of *opaque1* and *opaque2* genes associated with amino acids were also identified. Promising candidate genes were mapped to metabolic pathways and biosynthetic pathways for aspartate and branched-chain amino acids were reconstructed. Such a study provides valuable insights into the genetic basis of grain quality traits in sorghum, serving as a foundation for further validation studies.

Conclusion and Future Prospects

Sorghum being a climate-resilient crop, is raising significant attention as a future grain option due to its agricultural benefits in the face of climate change as well as the beneficial effects of its bioactive compounds on human health. This comprehensive review consolidates the current knowledge on the advances in sorghum grain quality improvement, covering various aspects such as grain structure, appearance, genetics, nutritional composition, functional attributes and the diverse components contributing to sorghum's nutritional value. The exploration of polysaccharides, proteins, carotenoids, minerals, phenolic compounds and antinutritional factors

provides a thorough understanding of the complex interplay influencing sorghum grain quality. Specifically, sorghum stands out for its abundance and diversity of phenolic compounds, unique among cereal grains. Considerable research has explored sorghum's phenolic profile, with most studies concentrating on overall content (such as total phenolic, flavonoids or condensed tannins) using colorimetric methods, rather than individual phenolics. The review also highlighted the multifaceted properties of sorghum phenolics, emphasizing their antioxidant, anticancer and antidiabetic potential. However, the majority of evidence comes from *in vitro* studies, indicating a need for additional *in vivo* and potentially clinical research to validate these health advantages, along with exploring interactions between bioactive compounds and their underlying mechanisms. Integrating association mapping and genome-wide association studies (GWAS) in sorghum provides valuable insights into the genetic basis of quality traits, paving the way for targeted breeding strategies. The identification of quantitative trait loci (QTLs) associated with nutritional and therapeutic traits adds a crucial dimension to sorghum breeding efforts. Continued research in this field will likely unveil further opportunities for optimizing sorghum grain quality and harnessing its full potential for the global food and nutritional security of people from developing and underdeveloped countries.

Acknowledgements

I would like to express my sincere gratitude to the Tamil Nadu Agricultural University Library for the invaluable digital assistance during the research and writing of this review paper. The extensive online resources, databases, and support services provided by the library significantly enhanced the quality and depth of the information presented in this work. I am grateful for the unwavering commitment of the library staff in ensuring a conducive digital environment for academic exploration and knowledge acquisition.

Authors' contributions

Anamika Dhar collected the literature and wrote the article. B. Meena Kumari contributed by suggesting the idea for writing the review and compilation of the content.

Table 5. Association of the eight significant Single nucleotide polymorphisms (SNPs) in a candidate gene/locus with the quality traits

Marker	Locus/Gene	Chromosome	Trait	R _{LR} ² (%) value
SB00116.3	SSIa	10	Kernel hardness	8
SB00214.1	pSB1700	3	Kernel hardness	10
SB00214.2	pSB1700	3	Kernel hardness	10
SB00156.1	pSB0289	3	Ca content	6
SB00054.1	PRC1044	6	Ca content	6
SB00068.1	pSB0140	6	P content	5
SB00086.1	pSB1120	3	Starch	9
SB00115.3	SSIb	4	Starch	10

Source: Sukumaran *et al.* (125)

D. Kavithamani helped in summarizing and revising the manuscript. N. Manikanda Boopathi helped in editing and summarizing the manuscript. P. Meenakshi provided information related to the biochemical aspects.

All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None.

References

- Kumar AA. Botany, taxonomy and breeding. The Sorghum Genome. 2016;27-45. https://doi.org/10.1007/978-3-319-47789-3_2
- De Wet J. Special paper: Systematics and evolution of Sorghum sect. Sorghum (Gramineae). American Journal of Botany. 1978;65(4):477-84. <https://doi.org/10.1002/j.15372197.1978.tb06096.x>
- Food and Agriculture Organization Corporate Statistical Database [Internet]. FAO; 2021. Available from: <http://www.fao.org/faostat/en/#data/QC>.
- International production assessment division [Internet]. USDA; 2023. Available from: <https://ipad.fas.usda.gov/cropexplorer/cropview/commodityView.aspx?cropid=0459200>.
- Mullet J, Morishige D, McCormick R, Truong S, Hilley J, McKinley B *et al*. Energy Sorghum—A genetic model for the design of C4 grass bioenergy crops. Journal of Experimental Botany. 2014;65(13):3479-89. <https://doi.org/10.1093/jxb/eru229>
- ME, Tonapi V, Bhat P, Varanavasiappan S, Seetharama N. Indian Sorghum landraces and their protection. 2015.
- Nagaraj N, Basavaraj G, Rao PP, Bantilan C, Halder S. Sorghum and pearl millet economy of India: Future outlook and options. Economic and Political Weekly. 2013;74-81.
- Przybylska-Balcerk A, Frankowski J, Stuper-Szablewska K. Bioactive compounds in sorghum. European Food Research and Technology. 2019;245:1075-80. <https://doi.org/10.1007/s00217-018-3207-0>
- Taylor JR, Schober TJ, Bean SR. Novel food and non-food uses for sorghum and millets. Journal of Cereal Science. 2006;44(3):252-71. <https://doi.org/10.1016/j.jcs.2006.06.009>
- Shen S, Huang R, Li C, Wu W, Chen H, Shi J *et al*. Phenolic compositions and antioxidant activities differ significantly among sorghum grains with different applications. Molecules. 2018;23(5):1203. <https://doi.org/10.3390/molecules23051203>
- González-Montilla FM, Chávez-Santoscoy RA, Gutiérrez-Urbe JA, Serna-Saldivar SO. Isolation and identification of phase II enzyme inducers obtained from black Shawaya sorghum [*Sorghum bicolor* (L.) Moench] Bran. Journal of Cereal Science. 2012;55(2):126-31. <https://doi.org/10.1016/j.jcs.2011.10.009>
- Venkateswaran K, Elangovan M, Sivaraj N. Origin, domestication and diffusion of *Sorghum bicolor*. Breeding Sorghum for diverse end uses. Elsevier. 2019; p. 15-31. <https://doi.org/10.1016/B978-0-08-101879-8.00002-4>
- Harlan JR, de Wet JM. A simplified classification of cultivated sorghum 1. Crop Science. 1972;12(2):172-76. <https://doi.org/10.2135/cropsci1972.0011183X001200020005x>
- Prabhakar, Madhusudhana R, Aruna C. Sorghum breeding. Fundamentals of Field Crop Breeding. Springer. 2022; p. 367-447. https://doi.org/10.1007/978-981-16-9257-4_7
- Deu M, Hamon P. The genetic organisation of sorghum; 1994.
- Serna-Saldivar S. Structure and chemistry of sorghum and millets. Sorghum and Millets: Chemistry and Technology. 1995;69-124. <https://doi.org/10.1016/B978-0-340-61457-0/50006-1>
- Earp C, McDonough C, Rooney L. Microscopy of pericarp development in the caryopsis of *Sorghum bicolor* (L.) Moench. Journal of Cereal Science. 2004;39(1):21-27. [https://doi.org/10.1016/S0733-5210\(03\)00060-2](https://doi.org/10.1016/S0733-5210(03)00060-2)
- Slavin J. Whole grains and human health. Nutrition Research Reviews. 2004;17(1):99-110. <https://doi.org/10.1079/NRR200374>
- Waniska RD, Rooney LW. Sorghum grain quality for increased utilization. Sorghum and Millets Diseases. 2002;327-35.
- Awika JM, Rooney LW. Sorghum phytochemicals and their potential impact on human health. Phytochemistry. 2004;65(9):1199-221. <https://doi.org/10.1016/j.phytochem.2004.04.001>
- Dykes L, Peterson GC, Rooney WL, Rooney LW. Flavonoid composition of lemon-yellow sorghum genotypes. Food Chemistry. 2011;128(1):173-79. <https://doi.org/10.1016/j.foodchem.2011.03.020>
- Dykes L, Rooney WL, Rooney LW. Evaluation of phenolics and antioxidant activity of black sorghum hybrids. Journal of Cereal Science. 2013;58(2):278-83. <https://doi.org/10.1016/j.jcs.2013.06.006>
- Earp C, Rooney L. Scanning electron microscopy of the pericarp and testa of several sorghum varieties. Food Structure. 1982;1(2):3.
- Nagaraja T, Parveen SG, Aruna C, Hariprasanna K, Singh S, Singh AK *et al*. Millets and pseudocereals: A treasure for climate resilient agriculture ensuring food and nutrition security. Indian Journal of Genetics and Plant Breeding. 2024;84(01):1-37. <https://doi.org/10.31742/ISGPB.84.1.1>
- Audilakshmi S, Aruna C. Genetic analysis of physical grain quality characters in sorghum. The Journal of Agricultural Science. 2005;143(4):267-73. <https://doi.org/10.1017/S0021859605005368>
- Cabrera-Ramírez A, Luzardo-Ocampo I, Ramírez-Jiménez A, Morales-Sánchez E, Campos-Vega R, Gaytán-Martínez M. Effect of the nixtamalization process on the protein bio accessibility of white and red sorghum flours during *in vitro* gastrointestinal digestion. Food Research International. 2020;134:109234. <https://doi.org/10.1016/j.foodres.2020.109234>
- USDA. National nutrient database for standard reference legacy release: Full report (all nutrients) 20067, sorghum grain. [Internet]; 2019. Available from: <https://ndb.nal.usda.gov/ndb/foods/show/20067?n1=%7BQv%3D1%7D&mp>
- Zhu F. Structure, physicochemical properties, modifications and uses of sorghum starch. Comprehensive Reviews in Food Science and Food Safety. 2014;13(4):597-610. <https://doi.org/10.1111/1541-4337.12070>
- Sasaki T, Yasui T, Matsuki J. Effect of amylose content on gelatinization, retrogradation and pasting properties of starches from waxy and nonwaxy wheat and their F1 seeds. Cereal Chemistry. 2000;77(1):58-63. <https://doi.org/10.1094/CCHEM.2000.77.1.58>
- Behall KM, Scholfield DJ. Food amylose content affects postprandial glucose and insulin responses. Cereal Chemistry. 2005;82(6):654-59. <https://doi.org/10.1094/CC-82-0654>
- Cagampang G, Kirleis A. Properties of starches isolated from sorghum flourey and corneous endosperm. Starch-Stärke. 1985;37(8):253-57. <https://doi.org/10.1002/star.19850370802>
- Beta T, Corke H, Rooney LW, Taylor JRN. Starch properties as affected by sorghum grain chemistry. Journal of the Science of Food and Agriculture. 2001;81(2):245-51. [https://doi.org/10.1002/1097-0010\(20010115\)81:2<245::AID-JSFA805>3.0.CO;2-S](https://doi.org/10.1002/1097-0010(20010115)81:2<245::AID-JSFA805>3.0.CO;2-S)
- Hill H, Lee LS, Henry RJ. Variation in sorghum starch synthesis genes associated with differences in starch phenotype. Food Chemistry. 2012;131(1):175-83. <https://doi.org/10.1016/j.foodchem.2011.08.057>

34. Wester T, Gramlich S, Britton R, Stock R. Effect of grain sorghum hybrid on *in vitro* rate of starch disappearance and finishing performance of ruminants. *Journal of Animal Science*. 1992;70(9):2866-76. <https://doi.org/10.2527/1992.7092866x>
35. Sullins R, Rooney L. Microscopic evaluation of the digestibility of sorghum lines that differ in endosperm characteristics. 1974.
36. Sang Y, Bean S, Seib PA, Pedersen J, Shi Y-C. Structure and functional properties of sorghum starches differing in amylose content. *Journal of Agricultural and Food Chemistry*. 2008;56(15):6680-85. <https://doi.org/10.1021/jf800577x>
37. Taylor JR, Emmambux MN. Developments in our understanding of sorghum polysaccharides and their health benefits. *Cereal Chemistry*. 2010;87(4):263-71. <https://doi.org/10.1094/CCHEM-87-4-0263>
38. Verbruggen MA, Beldman G, Voragen AG. Enzymic degradation of sorghum glucuronarabinoylans leading to tentative structures. *Carbohydrate Research*. 1998;306(1-2):275-82. [https://doi.org/10.1016/S0008-6215\(97\)10065-9](https://doi.org/10.1016/S0008-6215(97)10065-9)
39. Niba LL, Hoffman J. Resistant starch and β -glucan levels in grain sorghum (*Sorghum bicolor* M.) are influenced by soaking and autoclaving. *Food Chemistry*. 2003;81(1):113-18. [https://doi.org/10.1016/S0308-8146\(02\)00386-2](https://doi.org/10.1016/S0308-8146(02)00386-2)
40. Ramesh HP, Tharanathan RN. Structural characteristics of a mixed linkage β -D-glucan from sorghum (*Sorghum bicolor*). *Carbohydrate Research*. 1998;308(1-2):239-43. [https://doi.org/10.1016/S0008-6215\(98\)00064-0](https://doi.org/10.1016/S0008-6215(98)00064-0)
41. Boyer CD, Liu K-C. Starch and water-soluble polysaccharides from sugary endosperm of sorghum. *Phytochemistry*. 1983;22(11):2513-15. [https://doi.org/10.1016/0031-9422\(83\)80151-4](https://doi.org/10.1016/0031-9422(83)80151-4)
42. Taylor J, Schüssler L. The protein compositions of the different anatomical parts of sorghum grain. *Journal of Cereal Science*. 1986;4(4):361-69. [https://doi.org/10.1016/S0733-5210\(86\)80040-6](https://doi.org/10.1016/S0733-5210(86)80040-6)
43. Rooney LW, Serna-Saldivar SO. Sorghum. *Handbook of Cereal Science and Technology, Revised and Expanded*: CRC Press. 2000; p. 149-75.
44. Proietti I, Frazzoli C, Mantovani A editors. Exploiting nutritional value of staple foods in the world's semi-arid areas: Risks, benefits, challenges and opportunities of sorghum. *Healthcare*;2015. <https://doi.org/10.3390/healthcare3020172>
45. Bean S, Ioerger B, Smith B, Blackwell D. Sorghum protein structure and chemistry: Implications for nutrition and functionality. *Advances in Cereal Science: Implications to Food Processing and Health Promotion*. ACS Publications. 2011; p. 131-47.
46. Taylor J, Bean SR, Ioerger BP, Taylor JR. Preferential binding of sorghum tannins with γ -kafirin and the influence of tannin binding on kafirin digestibility and biodegradation. *Journal of Cereal Science*. 2007;46(1):22-31. <https://doi.org/10.1016/j.jcs.2006.11.001>
47. Watterson J, Shull J, Kirleis A. Quantitation of α -, β -, and γ -kafirins in vitreous and opaque endosperm of Sorghum bicolor. 1993.
48. Gebremariam MM, Zarnkow M, Becker T. Teff (*Eragrostis tef*) as a raw material for malting, brewing and manufacturing of gluten-free foods and beverages: A review. *Journal of Food Science and Technology*. 2014;51:2881-95. <https://doi.org/10.1007/s13197-012-0745-5>
49. Kumar MVN, Ramya V, Maheshwaramma S, Ganapathy KN, Govindaraj M, Kavitha K, et al. Exploiting Indian landraces to develop biofortified grain sorghum with high protein and minerals. *Frontiers in Nutrition*. 2023;10. <https://doi.org/10.3389%2Ffnut.2023.1228422>
50. Fernandez MGS, Hamblin MT, Li L, Rooney WL, Tuinstra MR, Kresovich S. Quantitative trait loci analysis of endosperm color and carotenoid content in sorghum grain. *Crop Science*. 2008;48(5):1732-43. <https://doi.org/10.2135/cropsci2007.12.0684>
51. de Morais Cardoso L, Pinheiro SS, da Silva LL, de Menezes CB, de Carvalho CWP, Tardin FD et al. Tocochromanols and carotenoids in sorghum (*Sorghum bicolor* L.): Diversity and stability to the heat treatment. *Food Chemistry*. 2015;172:900-08. <https://doi.org/10.1016/j.foodchem.2014.09.117>
52. Kean EG, Ejeta G, Hamaker BR, Ferruzzi MG. Characterization of carotenoid pigments in mature and developing kernels of selected yellow-endosperm sorghum varieties. *Journal of Agricultural and Food Chemistry*. 2007;55(7):2619-26. <https://doi.org/10.1021/jf062939v>
53. Kean EG, Bordenave N, Ejeta G, Hamaker BR, Ferruzzi MG. Carotenoid bioaccessibility from whole grain and decorticated yellow endosperm sorghum porridge. *Journal of Cereal Science*. 2011;54(3):450-59. <https://doi.org/10.1016/j.jcs.2011.08.010>
54. Lipkie TE, De Moura FF, Zhao Z-Y, Albertsen MC, Che P, Glassman K et al. Bioaccessibility of carotenoids from transgenic provitamin A biofortified sorghum. *Journal of Agricultural and Food Chemistry*. 2013;61(24):5764-71. <https://doi.org/10.1021/jf305361s>
55. Shen Y. Sorghum pericarp pigments are associated with the contents of carotenoids and provitamin A: Kansas State University; 2016.
56. Kayodé AP, Linnemann AR, Hounhouigan JD, Nout MJ, van Boekel MA. Genetic and environmental impact on iron, zinc and phytate in food sorghum grown in Benin. *Journal of Agricultural and Food Chemistry*. 2006;54(1):256-62. <https://doi.org/10.1021/jf0521404>
57. Pontieri P, Troisi J, Calcagnile M, Bean SR, Tilley M, Aramouni F et al. Chemical composition, fatty acid and mineral content of food-grade white, red and black sorghum varieties grown in the mediterranean environment. *Foods*. 2022;11(3):436. <https://doi.org/10.3390/foods11030436>
58. Kotla A, Phuke R, Hariprasanna K, Mehtre SP, Rathore A, Gorthy S et al. Identification of QTLs and candidate genes for high grain Fe and Zn concentration in sorghum [*Sorghum bicolor* (L.) Moench]. *Journal of Cereal Science*. 2019;90:102850. <https://doi.org/10.1016/j.jcs.2019.102850>
59. Motlhaodi T, Bryngelsson T, Chite S, Fatih M, Ortiz R, Geleta M. Nutritional variation in sorghum [*Sorghum bicolor* (L.) Moench] accessions from Southern Africa revealed by protein and mineral composition. *Journal of Cereal Science*. 2018;83:123-29. <https://doi.org/10.1016/j.jcs.2018.08.010>
60. Ashok Kumar A, Reddy BV, Ramaiah B, Sahrawat KL, Pfeiffer WH. Genetic variability and character association for grain iron and zinc contents in sorghum germplasm accessions and commercial cultivars. *European Journal of Plant Science and Biotechnology*. 2012;6(1):1-5.
61. Bean S, Wilson J, Moreau R, Galant A, Awika J, Kaufman R et al. Structure and composition of the sorghum grain. *Sorghum: A State of the Art and Future Perspectives*. 2019;58:173-214. <https://doi.org/10.2134/agronmonogr58.c9>
62. Lasztity R. Phytic acid in cereal technology. *Advances in Cereal Science and Technology*. 1990;10:309-71.
63. Reddy N, Pierson MD, Sathe SK, Salunkhe D. Phytates in cereals and legumes: CRC Press;1989.
64. Doherty C, Rooney L, Faubion J editors. Phytin content of sorghum and sorghum products. *Proc Int Symp On Sorghum Grain Quality (LW Rooney and DS Murty eds) Int Crops Res Inst Semi-arid Tropics (ICRISAT), Patancheru, AP India; 1981*.
65. Badigannavar A, Girish G, Ganapathi T. Genetic variability for phytic acid and inorganic phosphorous in Indian Sorghum (*Sorghum bicolor*) landraces. *Electronic Journal of Plant Breeding*. 2014;5(3):451-58.
66. Soetan K, Oyewole O. The need for adequate processing to reduce the anti-nutritional factors in plants used as human foods and animal feeds: A review. *African Journal of Food Science*. 2009;3(9):223-32.
67. Jahromi SG. Extraction techniques of phenolic compounds from

- plants. *Plant Physiological Aspects of Phenolic Compounds*. 2019;1-18.
68. Wu G, Bornman JF, Bennett SJ, Clarke MW, Fang Z, Johnson SK. Individual polyphenolic profiles and antioxidant activity in sorghum grains are influenced by very low and high solar UV radiation and genotype. *Journal of Cereal Science*. 2017;77:17-23. <https://doi.org/10.1016/j.jcs.2017.07.014>
 69. Salazar-López NJ, Gonzalez-Aguilar G, Rouzaud-Sandez O, Robles-Sanchez M. Technologies applied to sorghum (*Sorghum bicolor* L. Moench): Changes in phenolic compounds and antioxidant capacity. *Food Science and Technology*. 2018;38:369-82. <https://doi.org/10.1590/fst.16017>
 70. Chiremba C, Taylor JR, Rooney LW, Beta T. Phenolic acid content of sorghum and maize cultivars varying in hardness. *Food Chemistry*. 2012;134(1):81-88. <https://doi.org/10.1016/j.foodchem.2012.02.067>
 71. Yang L, Allred KF, Geera B, Allred CD, Awika JM. Sorghum phenolics demonstrate estrogenic action and induce apoptosis in nonmalignant colonocytes. *Nutrition and Cancer*. 2012;64(3):419-27. <https://doi.org/10.1080/01635581.2012.657333>
 72. Girard AL, Awika JM. Sorghum polyphenols and other bioactive components as functional and health promoting food ingredients. *Journal of Cereal Science*. 2018;84:112-24. <https://doi.org/10.1016/j.jcs.2018.10.009>
 73. Xiong Y, Zhang P, Warner RD, Fang Z. Sorghum grain: From genotype, nutrition and phenolic profile to its health benefits and food applications. *Comprehensive Reviews in Food Science and Food Safety*. 2019;18(6):2025-46. <https://doi.org/10.1111/1541-4337.12506>
 74. Althwab S, Carr TP, Weller CL, Dweikat IM, Schlegel V. Advances in grain sorghum and its co-products as a human health promoting dietary system. *Food Research International*. 2015;77:349-59. <https://doi.org/10.1016/j.foodres.2015.08.011>
 75. Vanamala JK, Massey AR, Pinnamaneni SR, Reddivari L, Reardon KF. Grain and sweet sorghum (*Sorghum bicolor* L. Moench) serves as a novel source of bioactive compounds for human health. *Critical Reviews in Food Science and Nutrition*. 2018;58(17):2867-81. <https://doi.org/10.1080/10408398.2017.1344186>
 76. Seo JW, Ham DY, Lee JG, Kim NY, Kim MJ, Yu CY *et al*. Antioxidant activity, phenolic content and antioxidant gene expression in genetic resources of sorghum collected from Australia, Former Soviet Union, USA, Sudan and Guadeloupe. *Agronomy*. 2023;13(7):1698. <https://doi.org/10.3390/agronomy13071698>
 77. Punia H, Tokas J, Bhadu S, Mohanty AK, Rawat P, Malik A *et al*. Proteome dynamics and transcriptome profiling in sorghum [*Sorghum bicolor* (L.) Moench] under salt stress. *3 Biotech*. 2020;10:1-10. <https://doi.org/10.1007/s13205-020-02392-1>
 78. Ghimire B-K, Seo J-W, Yu C-Y, Kim S-H, Chung I-M. Comparative study on seed characteristics, antioxidant activity and total phenolic and flavonoid contents in accessions of *Sorghum bicolor* (L.) Moench. *Molecules*. 2021;26(13):3964. <https://doi.org/10.3390/molecules26133964>
 79. Punia H, Tokas J, Malik A, Satpal, Sangwan S. Characterization of phenolic compounds and antioxidant activity in sorghum [*Sorghum bicolor* (L.) Moench] grains. *Cereal Research Communications*. 2021;49:343-53. <https://doi.org/10.1007/s42976-020-00118-w>
 80. Kim J, Noh SK, Woo K-S, Seo M-C. Sorghum extract lowers lymphatic absorption of trans fat and cholesterol in rats. *Journal of the Korean Society of Food Science and Nutrition*. 2016;45(6):783-88. <https://doi.org/10.3746/jkfn.2016.45.6.783>
 81. Liu H, Huang L, Pei X. Effects of sorghum rice and black rice on genes associated with cholesterol metabolism in hypercholesterolemic mice liver and intestine. *Food Science and Nutrition*. 2021;9(1):217-29. <https://doi.org/10.1002/fsn3.1986>
 82. Kim E, Kim S, Park Y. Sorghum extract exerts cholesterol-lowering effects through the regulation of hepatic cholesterol metabolism in hypercholesterolemic mice. *International Journal of Food Sciences and Nutrition*. 2015;66(3):308-13. <https://doi.org/10.3109/09637486.2014.1000839>
 83. Queiroz VAV, da Silva Aguiar A, de Menezes CB, de Carvalho CWP, Paiva CL, Fonseca PC *et al*. A low calorie and nutritive sorghum powdered drink mix: Influence of tannin on the sensorial and functional properties. *Journal of Cereal Science*. 2018;79:43-49. <https://doi.org/10.1016/j.jcs.2017.10.001>
 84. Iyabo O, Ibiyinka O, Abimbola Deola O. Comparative study of nutritional, functional and antinutritional properties of white *Sorghum bicolor* (Sorghum) and *Pennisetum glaucum* (Pearl Millet). *International Journal of Engineering Technologies and Management Research*. 2018;5(3):151-58.
 85. Parnian F, Taghizadeh A, Paya H, Nobari BB. *In vitro* fermentation response to alkaline treated sorghum grain. *Journal of Bioscience & Biotechnology*. 2014;3(1).
 86. de Oliveira KG, Queiroz VAV, de Almeida Carlos L, de Morais Cardoso L, Pinheiro- Sant'Ana HM, Anunciação PC *et al*. Effect of the storage time and temperature on phenolic compounds of sorghum grain and flour. *Food Chemistry*. 2017;216:390-98. <https://doi.org/10.1016/j.foodchem.2016.08.047>
 87. Aguilera CN, Aguilera-Carbo A, Robledo A, Ventura J, Belmares R, Martinez D *et al*. Production of antioxidant nutraceuticals by solid-state cultures of pomegranate (*Punica granatum*) peel and creosote bush (*Larrea tridentata*) leaves. *Food Technology and Biotechnology*. 2008;46(2):218.
 88. Vila-Real C, Pimenta-Martins A, Maina N, Gomes A, Pinto E. Nutritional value of indigenous whole grain cereals millet and sorghum. *Nutrition and Food Science International Journal*. 2017;4(1).
 89. Dykes L, Rooney LW. Sorghum and millet phenols and antioxidants. *Journal of Cereal Science*. 2006;44(3):236-51. <https://doi.org/10.1016/j.jcs.2006.06.007>
 90. Daglia M. Polyphenols as antimicrobial agents. *Current Opinion in Biotechnology*. 2012;23(2):174-81. <https://doi.org/10.1016/j.copbio.2011.08.007>
 91. Jiang Y, Zhang H, Qi X, Wu G. Structural characterization and antioxidant activity of condensed tannins fractionated from sorghum grain. *Journal of Cereal Science*. 2020;92:102918. <https://doi.org/10.1016/j.jcs.2020.102918>
 92. Ahmad F, Pasha I, Saeed M, Asgher M. Biochemical profiling of Pakistani sorghum and millet varieties with special reference to anthocyanins and condensed tannins. *International Journal of Food Properties*. 2018;21(1):1586-97. <https://doi.org/10.1080/10942912.2018.1502198>
 93. Awika JM, McDonough CM, Rooney LW. Decorticating sorghum to concentrate healthy phytochemicals. *Journal of Agricultural and Food Chemistry*. 2005;53(16):6230-34. <https://doi.org/10.1021/jf0510384>
 94. Chandrasekara A, Shahidi F. Content of insoluble bound phenolics in millets and their contribution to antioxidant capacity. *Journal of Agricultural and Food Chemistry*. 2010;58(11):6706-14. <https://doi.org/10.1021/jf100868b>
 95. Khoddami A, Truong HH, Liu SY, Roberts TH, Selle PH. Concentrations of specific phenolic compounds in six red sorghums influence nutrient utilisation in broiler chickens. *Animal Feed Science and Technology*. 2015;210:190-99. <https://doi.org/10.1016/j.anifeeds.2015.09.029>
 96. Awika JM, Rooney LW, Waniska RD. Properties of 3-deoxyanthocyanins from sorghum. *Journal of Agricultural and Food Chemistry*. 2004;52(14):4388-94. <https://doi.org/10.1021/jf049653f>
 97. Semere T, Tsehaye Y, Tareke L, Westengen OT, Fjellheim S. Nutritional and antinutritional potentials of sorghum: A comparative study among different sorghum landraces of Tigray,

- Northern Ethiopia. *Agriculture*. 2023;13(9):1753. <https://doi.org/10.3390/agriculture13091753>
98. Taleon V, Dykes L, Rooney W, Rooney L. Effect of genotype and environment on flavonoid concentration and profile of black sorghum grains. *Journal of Cereal Science*. 2012;56(2):470-75. <https://doi.org/10.1016/j.jcs.2012.05.001>
 99. Dykes L, Seitz LM, Rooney WL, Rooney LW. Flavonoid composition of red sorghum genotypes. *Food Chemistry*. 2009;116(1):313-17. <https://doi.org/10.1016/j.foodchem.2009.02.052>
 100. Speranza S, Knechtl R, Witlaczel R, Schönlechner R. Reversed-phase HPLC characterization and quantification and antioxidant capacity of the phenolic acids and flavonoids extracted from eight varieties of sorghum grown in Austria. *Frontiers in Plant Science*. 2021;12:769151. <https://doi.org/10.3389/fpls.2021.769151>
 101. Riddick EW. Potential of quercetin to reduce herbivory without disrupting natural enemies and pollinators. *Agriculture*. 2021;11(6):476. <https://doi.org/10.3390/agriculture11060476>
 102. Chatterjee D, Lesko T, Peiffer M, Elango D, Beuzelin J, Felton GW *et al.* Sorghum and maize flavonoids are detrimental to growth and survival of fall armyworm *Spodoptera frugiperda*. *Journal of Pest Science*. 2023;96(4):1551-67. <https://doi.org/10.3390/agriculture11060476>
 103. Kariyat RR, Gaffoor I, Sattar S, Dixon CW, Frock N, Moen J *et al.* Sorghum 3-deoxyanthocyanidin flavonoids confer resistance against corn leaf aphid. *Journal of Chemical Ecology*. 2019;45:502-14. <https://doi.org/10.1007/s10886-019-01062-8>
 104. Lopez-Contreras JJ, Zavala-Garcia F, Urias-Orona V, Martinez-Avila GCG, Rojas R, Guillermo N-M. Chromatic, phenolic and antioxidant properties of *Sorghum bicolor* genotypes. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*. 2015;43(2):366-70. <https://doi.org/10.15835/nbha4329949>
 105. Shahidi F, Peng H. Bioaccessibility and bioavailability of phenolic compounds. *Journal of Food Bioactives*. 2018;4:11-68. <https://doi.org/10.31665/JFB.2018.4162>
 106. Granato D, Shahidi F, Wrolstad R, Kilmartin P, Melton LD, Hidalgo FJ *et al.* Antioxidant activity, total phenolics and flavonoids contents: Should we ban *in vitro* screening methods? *Food Chemistry*. 2018;264:471-75. <https://doi.org/10.1016/j.foodchem.2018.04.012>
 107. Farida K, Messaoud B, Nouredine B, Bernard W. Free radical scavenging activity correlated with phenolic.
 108. Awika JM, Rooney LW, Waniska RD. Anthocyanins from black sorghum and their antioxidant properties. *Food Chemistry*. 2005;90(1-2):293-301. <https://doi.org/10.1016/j.foodchem.2004.03.058>
 109. Awika JM, Yang L, Browning JD, Faraj A. Comparative antioxidant, antiproliferative and phase II enzyme inducing potential of sorghum (*Sorghum bicolor*) varieties. *LWT-Food Science and Technology*. 2009;42(6):1041-46. <https://doi.org/10.1016/j.lwt.2009.02.003>
 110. Tian Y, Zou B, Li C-m, Yang J, Xu S-f, Hagerman AE. High molecular weight persimmon tannin is a potent antioxidant both *ex vivo* and *in vivo*. *Food Research International*. 2012;45(1):26-30. <https://doi.org/10.1016/j.foodres.2011.10.005>
 111. Cox S, Noronha L, Herald T, Bean S, Lee S-H, Perumal R *et al.* Evaluation of ethanol- based extraction conditions of sorghum bran bioactive compounds with downstream anti- proliferative properties in human cancer cells. *Heliyon*. 2019;5(5). <https://doi.org/10.1016/j.heliyon.2019.e01589>
 112. Hargrove JL, Greenspan P, Hartle DK, Dowd C. Inhibition of aromatase and α -amylase by flavonoids and proanthocyanidins from *Sorghum bicolor* bran extracts. *Journal of Medicinal Food*. 2011;14(7-8):799-807. <https://doi.org/10.1089/jmf.2010.0143>
 113. Moraes ÉA, da Silva Marineli R, Lenquiste SA, Queiroz VAV, Camargo RL, Borck PC *et al.* Whole sorghum flour improves glucose tolerance, insulin resistance and preserved pancreatic islets function in obesity diet-induced rats. *Journal of Functional Foods*. 2018;45:530-40. <https://doi.org/10.1016/j.jff.2017.03.047>
 114. Chung I-M, Kim E-H, Yeo M-A, Kim S-J, Seo MC, Moon H-I. Antidiabetic effects of three Korean sorghum phenolic extracts in normal and streptozotocin-induced diabetic rats. *Food Research International*. 2011;44(1):127-32. <https://doi.org/10.1016/j.foodres.2010.10.051>
 115. Kim J, Park Y. Anti-diabetic effect of sorghum extract on hepatic gluconeogenesis of streptozotocin-induced diabetic rats. *Nutrition & Metabolism*. 2012;9:1-7. <https://doi.org/10.1186/1743-7075-9-106>
 116. Links MR, Taylor J, Kruger MC, Taylor JR. Sorghum condensed tannins encapsulated in kafirin microparticles as a nutraceutical for inhibition of amylases during digestion to attenuate hyperglycaemia. *Journal of Functional Foods*. 2015;12:55-63. <https://doi.org/10.1016/j.jff.2014.11.003>
 117. Rhodes D, Gadgil P, Perumal R, Tesso T, Herald TJ. Natural variation and genome-wide association study of antioxidants in a diverse sorghum collection. *Cereal Chemistry*. 2017;94(2):190-98. <https://doi.org/10.1094/CCHEM-03-16-0075-R>
 118. Welderufael S, Abay F, Ayana A, Amede T. Genetic diversity, correlation and genotype \times yield \times trait (GYT) analysis of grain yield and nutritional quality traits in sorghum (*Sorghum bicolor* [L.] Moench) genotypes in Tigray, Northern Ethiopia. *Discover Agriculture*. 2024;2(1):4. <https://doi.org/10.1007/s44279-024-00016-3>
 119. Kumar AA, Reddy BV, Ramaiah B, Sahrawat KL, Pfeiffer WH. Gene effects and heterosis for grain iron and zinc concentration in sorghum [*Sorghum bicolor* (L.) Moench]. *Field Crops Research*. 2013;146:86-95. <https://doi.org/10.1016/j.fcr.2013.03.001>
 120. Gaddameedi A, Phuke RM, Polavarapu KKB, Gorthy S, Subhasini V, Jagannathan J *et al.* Heterosis and combining ability for grain Fe and Zn concentration and agronomic traits in sorghum [*Sorghum bicolor* (L.) Moench]. *Journal of King Saud University-Science*. 2020;32(7):2989-94. <https://doi.org/10.1016/j.jksus.2020.08.003>
 121. Flint-Garcia SA. Genetics and consequences of crop domestication. *Journal of Agricultural and Food Chemistry*. 2013;61(35):8267-76. <https://doi.org/10.1021/jf305511d>
 122. Rhodes DH, Hoffmann Jr L, Rooney WL, Ramu P, Morris GP, Kresovich S. Genome-wide association study of grain polyphenol concentrations in global sorghum [*Sorghum bicolor* (L.) Moench] germplasm. *Journal of Agricultural and Food Chemistry*. 2014;62(45):10916-27. <https://doi.org/10.1021/jf503651t>
 123. Rooney WL, Portillo O, Hayes C. Registration of ATx3363 and BTx3363 black sorghum germplasm. *Journal of Plant Registrations*. 2013;7(3):342-46. <https://doi.org/10.3198/jpr2013.01.0006crg>
 124. Mace E, Innes D, Hunt C, Wang X, Tao Y, Baxter J *et al.* The sorghum QTL Atlas: A powerful tool for trait dissection, comparative genomics and crop improvement. *Theoretical and Applied Genetics*. 2019;132:751-66. <https://doi.org/10.1007/s00122-018-3212-5>
 125. Sukumaran S, Xiang W, Bean SR, Pedersen JF, Kresovich S, Tuinstra MR *et al.* Association mapping for grain quality in a diverse sorghum collection. *The Plant Genome*. 2012;5(3). <https://doi.org/10.3835/plantgenome2012.07.0016>
 126. Kimani W, Zhang L-M, Wu X-Y, Hao H-Q, Jing H-C. Genome-wide association study reveals that different pathways contribute to grain quality variation in sorghum (*Sorghum bicolor*). *BMC Genomics*. 2020;21:1-19. <https://doi.org/10.1186/s12864-020-6538-8>
 127. Rooney LW, Waniska RD. Sorghum food and industrial utilization. *Sorghum: Origin, History, Technology and Production*. 2000;689-729. https://doi.org/10.1007/1-4020-0613-6_8400
 128. Bouargalne Y, Ben Mrid R, Bouchmaa N, Zouaoui Z, Benmrid B, Kchikich A *et al.* Genetic diversity for agromorphological traits, phytochemical profile and antioxidant activity in Moroccan

- sorghum ecotypes. *Scientific Reports*. 2022;12(1):5895. <https://doi.org/10.1038/s41598-022-09810-9>
129. Rao S, Santhakumar AB, Chinkwo KA, Wu G, Johnson SK, Blanchard CL. Characterization of phenolic compounds and antioxidant activity in sorghum grains. *Journal of Cereal Science*. 2018;84:103-11. <https://doi.org/10.1016/j.jcs.2018.07.013>
130. Ba K, Tine E, Destain J, Cisse N, Thonart P. Comparative study of phenolic compounds, antioxidant power of various Senegalese sorghum cultivars and amyolytic enzymes of their malt. *Biotechnologie, Agronomie, Société et Environnement*. 2010;14(1):131-39.
131. Lee S, Choi Y-M, Shin M-J, Yoon H, Wang X, Lee Y *et al*. Exploring the potentials of sorghum genotypes: A comprehensive study on nutritional qualities, functional metabolites and antioxidant capacities. *Frontiers in Nutrition*. 2023;10. <https://doi.org/10.3389%2Ffnut.2023.1238729>
132. Lee S, Choi Y-M, Shin M-J, Yoon H, Wang X, Lee Y *et al*. Agromorphological and biochemical characterization of Korean sorghum (*Sorghum bicolor* (L.) Moench) landraces. *Agronomy*. 2022;12(11):2898. <https://doi.org/10.3390/agronomy12112898>
133. Desta KT, Choi Y-M, Shin M-J, Yoon H, Wang X, Lee Y *et al*. Comprehensive evaluation of nutritional components, bioactive metabolites and antioxidant activities in diverse sorghum (*Sorghum bicolor* (L.) Moench) landraces. *Food Research International*. 2023;173:113390. <https://doi.org/10.1016/j.foodres.2023.113390>
134. Nguni D, Geleta M, Hofvander P, Fatih M, Bryngelsson T. Comparative genetic diversity and nutritional quality variation among some important Southern African sorghum accessions [*Sorghum bicolor* (L.) Moench]. *Australian Journal of Crop Science*. 2012;6(1):56-64.
135. Moraes ÉA, da Silva Marineli R, Lenquiste SA, Steel CJ, de Menezes CB, Queiroz VAV *et al*. Sorghum flour fractions: Correlations among polysaccharides, phenolic compounds, antioxidant activity and glycemic index. *Food chemistry*. 2015;180:116-23. <https://doi.org/10.1016/j.foodchem.2015.02.023>
136. Murray SC, Sharma A, Rooney WL, Klein PE, Mullet JE, Mitchell SE *et al*. Genetic improvement of sorghum as a biofuel feedstock: I. QTL for stem sugar and grain non-structural carbohydrates. *Crop Science*. 2008;48(6):2165-79. <https://doi.org/10.2135/cropsci2008.01.0016>
137. Patil NY, Pugh NA, Klein RR, Martinez HS, Martinez RS, Rodriguez-Herrera R *et al*. Heritability and quantitative trait loci of composition and structural characteristics in sorghum grain. *Journal of Crop Improvement*. 2019;33(1):1-24. <https://doi.org/10.1080/15427528.2018.1536006>
138. Chen B-R, WANG C-y, Ping W, ZHU Z-x, Ning X, SHI G-s *et al*. Genome-wide association study for starch content and constitution in sorghum (*Sorghum bicolor* (L.) Moench). *Journal of Integrative Agriculture*. 2019;18(11):2446-56. [https://doi.org/10.1016/S2095-3119\(19\)62631-6](https://doi.org/10.1016/S2095-3119(19)62631-6)
139. Rhodes DH, Hoffmann L, Rooney WL, Herald TJ, Bean S, Boyles R *et al*. Genetic architecture of kernel composition in global sorghum germplasm. *BMC Genomics*. 2017;18(1):1-8. <https://doi.org/10.1186/s12864-016-3403-x>
140. Li J, Tang W, Zhang Y-W, Chen K-N, Wang C, Liu Y *et al*. Genome-wide association studies for five forage quality-related traits in sorghum (*Sorghum bicolor* L.). *Frontiers in Plant Science*. 2018;9:1146. <https://doi.org/10.3389/fpls.2018.01146>
141. Cruet-Burgos C, Cox S, Ioerger BP, Perumal R, Hu Z, Herald TJ *et al*. Advancing provitamin A biofortification in sorghum: Genome-wide association studies of grain carotenoids in global germplasm. *The Plant Genome*. 2020;13(1):e20013. <https://doi.org/10.1002/tpg2.20013>
142. Habyarimana E, Dall'Agata M, De Franceschi P, Baloch FS. Genome-wide association mapping of total antioxidant capacity, phenols, tannins and flavonoids in a panel of *Sorghum bicolor* and *S. bicolor* × *S. halepense* populations using multi-locus models. *PLoS One*. 2019;14(12):e0225979. <https://doi.org/10.1371/journal.pone.0225979>