



RESEARCH COMMUNICATION

Phytochemistry, antibacterial and antioxidant properties from *Adinandra hongiaoensis* and *Polyspora huongiana* (Theaceae)

Tan Viet Pham¹, Thi Diep Nguyen¹, Thi Thanh Truc Ngo¹, Hanh Thi Dieu Nguyen¹, Ngoc An Nguyen¹, Quoc Hung Nguyen², Thanh Tho Le², Van Hop Nguyen^{3,4} & Hong Thien Van^{1*}

¹Institute of Biotechnology and Food Technology, Industrial University of Ho Chi Minh City, No. 12 Nguyen Van Bao Street, Go Vap district, Ho Chi Minh City 710 000, Vietnam

²Center of Analytical Services and Experimentation HCMC, Vietnam, No.2 Nguyen Van Thu, Dakao Ward, District 1, Ho Chi Minh City 710 000, Vietnam

³Faculty of Natural Resources and Environment, Vietnam National University of Forestry-Dongnai Campus, Trang Bom Town, Trang Bom District, Dong Nai Province, Vietnam

⁴College of Forestry, Fujian Agriculture and Forestry University, Fuzhou City, Fujian Province, China

*Email: vanhongthien@iuh.edu.vn



ARTICLE HISTORY

Received: 10 May 2024

Accepted: 17 July 2024

Available online

Version 1.0 : 29 October 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Pham TV, Nguyen TD, Ngo TTT, Nguyen HTD, Nguyen NA, Nguyen OH, Le TT, Nguyen VH, Van HT. Phytochemistry, antibacterial and antioxidant properties from *Adinandra hongiaoensis* and *Polyspora huongiana* (Theaceae). Plant Science Today (Early Access). <https://doi.org/10.14719/pst.3677>

Abstract

Adinandra hongiaoensis and *Polyspora huongiana* are the rare and native species found in Vietnam. Firstly, the present studies demonstrated the chemical components, antioxidant and antibacterial effects of the acetone extracts from these plants. Here, a total of 54 constituents are identified in the leaf extracts of 2 studied species using the gas chromatography/mass spectrometry (GC/MS) method. The *P. huongiana* extract displayed activity against 3 *Staphylococcus* strains, including *S. saprophyticus*, *S. aureus* ATCC 29213 and *S. aureus* ATCC 25923, while the *A. hongiaoensis* extract was not active against any studied bacteria. In addition, *A. hongiaoensis* and *P. huongiana* extracts was found effective against the DPPH radical scavenger with IC₅₀ values of 213.31 and 47.60 µg/mL, respectively. The results of this report are expected to provide a foundation for further research aimed at applying these species in the development of pharmaceutical products in the future.

Keywords

Adinandra hongiaoensis; *Polyspora huongiana*; GC/MS; antibacterial; antioxidant properties

Introduction

Theaceae includes about 480 species and 9 genera mainly found in tropical and subtropical regions of East and Southeast Asia (1-5). In Vietnam, about 80 species and 11 genera have been recorded (6-9). *Adinandra* and *Polyspora* are the 2 genera belonging to the Theaceae family. The first genus comprises over 100 species widely distributed across China, India, Bangladesh, Japan, tropical Africa and Southeast Asian countries and about 12 species in Vietnam (7, 10, 11). Whereas, the later one includes about 40 species mainly found in East and Southeast Asia (5), of which 13 species recorded in Vietnam (5-7, 9, 12-20). The members of *Adinandra* genus have been reported to possess so many bioactive compounds as well as the biological activities, including analgesic, antitumor, anti-toxicity and antibacterial. In addition, the extracts from leaves of some *Adinandra* plants have been consumed as beverage for hundreds of years (21-26). Meanwhile, because of their beautiful tree shape, especially, blooming in winter, the *Polyspora* plants have been reported to have the ornamental value and they, thus, are usually used as street or shade

trees in gardens (27-29). Moreover, studies have also demonstrated the medicinal value of *Polyspora* plants, highlighting their natural antioxidant effects (30, 31) and cytotoxic properties (32-34).

Adinandra hongiaoensis Son & L.V.Dung and *Polyspora huongiana* Orel, Curry & Luu are 2 new species belonging to Theaceae family which have been recently described from Vietnam (11, 16). The type specimens of the first species were collected from Bidoup Nui Ba National Park, Lam Dong province, Vietnam. The morphological features of this species were characterized by having: trees up to 10 m tall, cordate leaf blade base, short pedicel length and pubescent petal pubescence (11). Similarly, the later plant was also found in Bidoup Nui Ba National Park. It is distinguished from other *Polyspora* species by its narrowly elliptic mature leaves, stamens arranged in three concentric circles, filaments basally joined to the inner petal whorl, and a densely tomentose ovary (16). So far, both species have been known as the rare species and only found in the type locations (11, 16). The present study, thus, demonstrated the chemical components, antioxidant and antibacterial activities of the acetone extracts from *A. hongiaoensis* and *P. huongiana* for the first time.

Materials and Methods

Bacterial strains

Eight bacterial strains were used to identify the antibacterial activity of the studied species, including 4 Gram-negative strains (*Salmonella typhimurium* ATCC 13311, *Escherichia coli* ATCC 25922, *Shigella flexneri* ATCC 9199, *Enterobacter hormaechei* ATCC 700323) and 4 Gram-positive strains (*Staphylococcus saprophyticus* BAA750, *Staphylococcus aureus* ATCC 29213, *Bacillus cereus* ATCC 13883, *S. aureus* ATCC 25923).

Plant materials

The leaves of *Adinandra hongiaoensis* and *Polyspora huongiana* were collected from Bidoup Nui Ba National Park, Lam Dong province, Vietnam, at the coordinates of 12°10'59"N and 111°27'49"E; 12°11'00"N and 111°27'47"E, respectively. The vouchered specimens of *A. hongiaoensis* (BDNP24012021) and *P. huongiana* (BDNP25012021) were deposited in the Herbarium of Faculty of Natural Resources and Environment, Vietnam National University of Forestry at Dongnai.

Extraction procedures

The fresh samples were dried at 50 °C and ground into fine powder. 100 g of powder was soaked in 500 mL of 99% acetone (Thermo Fisher Scientific, USA) for 72 h and then the extract was filtered using Whatman paper. The residue material was subjected to 2 additional acetone extractions and the final product was obtained by pooling all the filtrates. Solvent removal was performed at 45 °C using a vacuum evaporator, resulting in the brown extracts (26).

Gas chromatography/mass spectrometry (GC/MS)

Chemical components in different extracts were first analyzed using Thermo Scientific™ ISQ™ 7000 GC-MS

(Thermo Fisher Scientific, USA) system with the stationary phase Agilent DB-5MS column. Running parameters were set as previously described (25). The chemical compositions were determined based on the NIST 2017 library.

Antibacterial activity

Agar disk diffusion method was used to determine the extract's antibacterial effect as recommended by CLSI (35). Bacterial cultures of 0.5 McFarland standard equivalent were spread on sterile MHA plates and different acetone extracts diluted with DMSO 15 % into concentrations of 100, 150 and 200 mg/mL were subsequently added to the standard sterile disc. The plates were incubated at 35 ± 2 °C for about 18 h for measurement of the zone of inhibition. Gentamicin disc (10 µg, Nam Khoa BioTek, Vietnam) was used as positive control, whereas DMSO 15 % was used as negative control. The zone of inhibition of studied sample was calculated by the formula

$$A = D - d \quad (\text{Eqn. 1})$$

Where, D is a zone of inhibition; d is the diameter of disc (6 mm). The experiment was done in triplicate and data were presented as mean ± standard deviation (SD). Fisher's least significant difference (LSD) and one-way analysis of variance (ANOVA) were employed for statistical analysis.

Antioxidant activity

Antioxidant effect was determined by 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay followed with minor modifications (25). The extract was completely dissolved in 99.8% methanol (Thermo Fisher Scientific, USA) to obtain different concentrations. Then, 0.3 mL was mixed with 3.7 mL of 0.1 mM DPPH. The mixture was incubated for 30 min in the dark at room temperature. The absorbance was subsequently measured at 517 nm using a UV-Vis spectrophotometer (Genesys 20, USA). The antioxidant effect of the extract was determined by IC₅₀ value in comparison with ascorbic acid, which was used as a control. Antioxidant activity was calculated as:

$$\% \text{ DPPH} = \frac{A_0 - A_i}{A_0} \times 100 \quad (\text{Eqn. 2})$$

Where, A₀ and A_i are the absorbance of the blank (DPPH solution) and the sample (DPPH solution and extract sample) respectively.

Statistical analysis

The experiments were carried out 3 times and the results were reported as the mean ± standard deviation (SD). The data were analyzed using a one-way analysis of variance (ANOVA) to compare different groups, with Fisher's least significant difference (LSD) procedure (p<0.05) used for multiple comparisons. Statistical analysis was done with Statgraphics Centurion XV (version 15.1.02, Statgraphics Technologies, Inc., USA).

Results

Chemical compositions from two studied extracts

The chemical compositions of the acetone extracts from the leaf of *A. hongiaoensis* and *P. huongiana* were shown in Table 1 and Fig. 1. Accordingly, the *A. hongiaoensis* extract was found to be rich in linolenic acid (16.40 %), neophytadiene (9.76 %), γ -sitosterol (9.73 %), phytol (9.51 %), 2-pentanone, 4-hydroxy-4-methyl- (7.47 %), l-(+)-ascorbic acid 2,6-dihexadecanoate (7.24 %), tetracosan-10-yl acetate (5.31 %), labda-8(20)-13-diene-15,19-dioic acid, (E)- (4.89 %). The *P. huongiana* extract was characterized by the predominance of 1,2,3-benzenetriol (18.02 %), neophytadiene (13.40 %), 3,7,11,15-tetramethyl-2-hexadecen-1-ol (12.94 %), vitamin E (8.81 %), squalene (5.72 %), benzoic acid, 4-hydroxy- (4.94 %).

Antibacterial properties of acetone extracts from two Theaceae species

The antibacterial effect of the acetone extracts from 2 studied plants were conducted using agar disc diffusion method. Overall, the *P. huongiana* extract was found to be effective against 3 out of 8 bacterial strains, including *B. cereus*, *S. aureus* ATCC 29213 and *S. aureus* ATCC 25923, while no active agent was found in the *A. hongiaoensis* extract (Table 2). At dose of 100 mg/mL, the *P. huongiana* extract showed a weaker effect than those at dose of 150 and 200 mg/mL, whereas there was no significant difference in the zone of inhibition between the concentrations of 150 and 200 mg/mL (Table 2).

Antioxidant properties of acetone extracts from two Theaceae species

The antioxidant effect of acetone extract from *A. hongiaoensis* and *P. huongiana* were investigated using the DPPH radical scavenging test with IC₅₀ value of 213.31 and 47.60 μ g/mL, respectively (Table 3).

Discussion

The chemical components and biological properties of the different solvent extracts isolated from other Adinandra species from Vietnam have been reported by prior reports (22-26). For instance, the methanol extract of Adinandra poilanei stem showed 5 compounds, including tyrosol; scopoletin; lupeol; 2,6-dimethoxy-1,4-benzoquinone and 2 β -hydroxypomolic acid (26). The chemical compositions and biological properties of the ethanol extract and its fractions, ethyl acetate and dichloromethane, isolated from Adinandra glischroloma stem were also investigated. Accordingly, the ethanol extract contained some chemical groups such as flavonoids, polyphenolic and coumarins (22). Moreover, the dichloromethane fraction from this species was found to be effective against *B. subtilis* and *L. plantarum*. The antioxidant activities of the dichloromethane, ethyl acetate fractions and ethanol extract were quantified with EC₅₀ values of 86.7, 88.5 and 111.2 μ g/mL, respectively. The ethanol extract also possessed effect against gastric, lung and breast cancer cell lines with IC₅₀ values of 59.1, 63.23 and 67.48 μ g/mL, respectively (22). Similarly, the ethanol extract and its

Table 1. The chemical components of acetone extracts from *A. hongiaoensis* (AHO) and *P. huongiana* (PHU).

RT	Compounds	AHO (%)	PHU (%)
2.45	2-Pentanone, 4-hydroxy-4-methyl-	7.47	4.00
4.62	3-Hexenoic acid, (E)-	0.21	-
4.67	cis-1,2-Dihydrocatechol	-	0.27
5.84	Cyclohexene, 4-ethenyl-1,4-dimethyl	-	0.09
5.92	Eucalyptol	-	0.19
7.22	Alpha-l-rhamnopyranose	-	0.70
7.24	Phenethylamine, 3-benzyloxy-2-fluoro- α -hydroxy	0.09	-
7.44	Paromomycin	0.09	-
8.22	12,15-Octadecadiynoic acid, methyl ester	0.04	-
8.56	5-Hydroxymethylfurfural	-	2.77
9.50	10-Heptadecen-8-ynoic acid, methyl ester, (E)-	0.09	-
9.87	Pyrogallol	-	18.02
10.02	Melezitose	0.32	2.27
10.16	Vanillin lactoside	0.05	-
10.20	Octahydrochromen-2-one	0.11	-
10.85	Benzoic acid, 4-hydroxy-	-	4.49
11.19	cis-Vaccenic acid	0.12	-
11.51	1,3-Benzenediol, 4-propyl-	0.34	-
11.57	Z-(13,14-Epoxy)tetradec-11-en-1-ol acetate	0.25	-
11.82	Benzenepropanol, 4-hydroxy-3-methoxy-	-	0.24
12.00	7-Oxo-2-oxa-7-thiatricyclo[4.4.0.0(3,8)]decan-4-ol	-	0.73
12.14	(E)-2,6-Dimethoxy-4-(prop-1-en-1-yl)phenol	0.31	-
12.37	(E)-4-(3-Hydroxyprop-1-en-1-yl)-2-methoxyphenol	0.73	-
12.42	Tetradecanoic acid	0.30	-
12.88	Neophytadiene	9.76	13.40
12.90	1-Hexadecanol, 2-methyl-	0.28	-
13.11	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	3.61	12.94
13.49	n-Hexadecanoic acid	-	2.29
13.51	l-(+)-Ascorbic acid 2,6-dihexadecanoate	7.24	-
13.70	trans-Sinapyl alcohol	0.61	-
14.27	Phytol	9.51	3.45
14.40	Linolenic acid	16.40	2.12
14.48	Octadecanoic acid	0.97	-
14.88	9-Hexadecenoic acid	0.32	-
15.26	Ethyl iso-allocholate	0.23	-
15.80	Kauren-19-oic acid	1.10	-
16.07	17-Pentatriacontene	3.70	-
16.19	Palmitin, 2-mono-	1.15	0.46
16.30	1-Heptatriacontanol	0.32	-
16.67	17-Pentatriacontene	0.96	-
17.37	Tetracosan-10-yl acetate	5.31	-
17.41	Linolenin, 1-mono-	-	0.31
18.05	4,6-Androstadien-3 β -ol-17-one, acetate	0.98	-
18.38	Squalene	2.84	5.72
19.23	Octadecane, 3-ethyl-5-(2-ethylbutyl)-	-	0.67
19.28	Labda-8(20)-13-diene-15,19-dioic acid, (E)-	4.89	-
21.12	Ethyl iso-allocholate	-	0.36
22.15	17-Pentatriacontene	0.93	-
22.52	Vitamin E	-	8.81
25.03	Stigmasterol	2.06	5.56
26.09	γ -Sitosterol	9.73	-
26.91	Stigmast-7-en-3-ol, (3 β ,5 α)-	-	2.52
27.37	α -Amyrin	-	5.47
28.07	dl- α -Tocopherol	3.14	-
	Total	96.56	97.85

Table 2. The antibacterial activity of acetone extracts from *A. hongiaoensis* and *P. huongiana*.

Bacterial strains	Concentration (mg/mL)			Gentamicine	Negative control
	100	150	200		
<i>E. coli</i>	-	-	-	13.5 ± 0.87	-
<i>S. typhimurium</i>	-	-	-	17.67 ± 0.58	-
<i>S. flexneri</i>	-	-	-	14.67 ± 1.15	-
<i>E. hormachei</i>	-	-	-	15.33 ± 1.53	-
<i>B. cereus</i>	-	-	-	13.17 ± 0.76	-
<i>S. saprophyticus</i>	-	6.33 ± 1.53 ^a	5.67 ± 0.76 ^a	20.67 ± 0.58 ^b	-
<i>S. aureus</i> ATCC 29213	3.25 ± 0.25 ^a	6.17 ± 1.26 ^b	5.17 ± 0.29 ^b	18.33 ± 1.15 ^c	-
<i>S. aureus</i> ATCC 25923	3.75 ± 0.43 ^a	7.40 ± 1.15 ^b	6.83 ± 0.76 ^b	18.50 ± 0.87 ^c	-


Different superscript lower-case letters in the same row denote significant differences ($p < 0.05$). - not active.

Table 3. The antioxidant activity of acetone extracts from *A. hongiaoensis* and *P. huongiana*.

	<i>A. hongiaoensis</i>	<i>P. huongiana</i>	Ascorbic Acid
IC ₅₀ (µg/mL)	213.31	47.60	2.41

fractions, dichloromethane and ethyl acetate fractions, from the *Adinandra megaphylla* stem were also reported to consist of some bioactive compounds and biological activity. As a result, the ethanol extract possessed polyphenolic and coumarins compounds (24). The dichloromethane and ethyl acetate fractions of this species had an inhibitory effect on *B. subtilis*, *L. plantarum* and *E. coli*. The ethyl acetate fraction and ethanol extract were found effective against the DPPH radical scavenger with EC₅₀ values of 32.0 and 28.9 µg/mL, respectively. The ethanol extract also showed cytotoxic effects against several cell lines, including lung, gastric, and breast, with IC₅₀ values of 43.52, 56.54, and 58.24 µg/mL, respectively (24).

A study showed a new triterpene compound, lupan-3 β ,20-dihydroxy-28-carbaldehyde and 11 known compounds from the methanol extracts of the stem obtained from *Adinandra hainanensis* grown in Vietnam (23). Furthermore, 10 out of 12 compounds, including a new compound, isolated from this species were also found to possessed the cytotoxicity (KB, Hep-G2, LU, MCF-7 cell lines) and α -glucosidase inhibitory effects (23). The ethanol extract from the leaf of *Adinandra bockiana* collected from Vietnam also contained some chemical groups like tannin, polyphenol and coumarin (21). Also, ethanol extract and its fractions, dichloromethane and ethyl acetate, isolated from *A. bockiana* displayed activity against *B. subtilis*, *L. plantarum* and *S. macescens* as well as effective against DPPH radical scavenger with EC₅₀ values of 0.2, 0.47 and 5.7 µg/mL, respectively. The ethanol extract of this plant also showed cytotoxic activities against breast, lung, gastric cell lines with IC₅₀ values of

**Fig. 1.** The GC chromatogram of acetone extracts from *A. hongiaoensis* (AHO) and *P. huongiana* (PHU).

43.15, 48.73 and 58.62 $\mu\text{g}/\text{mL}$, respectively (21).

Previous studies have also provided insights into the chemical constituents of *Gordonia chrysandra*, a synonym for *Polyspora chrysandra*. For instance, 2 new flavanonol glycosides, chrysandrosides A and B, were obtained from *P. chrysandra* roots (36), whereas a new phenolic glycoside, chrysandroside A was also isolated from the stem of this species (37). A study demonstrated the polyphenolic compounds and antioxidant properties from the fruit extract of *Gordonia axillaris*, a synonym species of *Polyspora axillaris* (30). Accordingly, the fruit extract of this plant contained some polyphenolic constituents, including rutin, gallic acid, protocatechuic acid, epicatechin, 2-hydrocinnamic acid, p-coumaric acid, quercetin, epicatechin gallate and ferulic acid. The ABTS radical scavenging effect of the extract was 525.05 ± 14.34 μmol Trolox/g DW (30).

Conclusion

In this study, the phytochemical properties, antibacterial and antioxidant activities of the acetone extracts isolated from 2 Theaceae species, *Adinandra hongiaoensis* and *Polyspora huongiana* were investigated for the first time. Along with the applicability of many species of Theaceae family, hopefully, the result of this study may serve as a foundation for future research, exploring the use of these species in pharmaceutical and other fields.

Authors' contributions

TVP and HTV participated in the design of the study and performed the experiments and statistical analysis. TDN, TTTN, HTDN, NAN, QHN, TTL and VHN performed experiments and handled the research data. HTV and TVP drafted the manuscript and resolved all the queries of editors and reviewers. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None.

References

- Keng H. Comparative morphological studies in Theaceae. University of California Press.1962.
- Prince LM, Parks CR. Phylogenetic relationships of Theaceae inferred from chloroplast DNA sequence data. *Am J Bot.* 2001; 88(12):2309-20. <https://doi.org/10.2307/3558391>.
- Prince LM. A brief nomenclatural review of genera and tribes in Theaceae. *Aliso.* 2007; 24(1):105-21. <https://doi.org/10.5642/aliso.20072401.08>.
- Stevens PF, Dressler S, Weitzman AL. Flowering plants: Dicotyledons (Celastrales, Oxidales, Rosales, Cornales, Ericales), in the families of and genera of flowering plants, Kubitzki K, Editor. Springer-Verlag, Berlin, Germany. 2004.
- Ming TL, Bartholomew B. Theaceae, in *Flora of China*, Wu ZY, Raven PH, Hong DY, Editors. Beijing: Science Press; St. Louis: Missouri Botanical Garden Press. 2007.
- Gagnepain F. Ternstroemiaceae, in supplément a la flore générale de l'Indo-Chine, Lecomte MH, Editor. Paris: Muséum National D'Histoire Naturelle. 1943.
- Pham HH. Theaceae, in *Cây cỏ Việt Nam: An illustrated flora of Vietnam*, Pham HH, Editor. Tre Publishing House, Ho Chi Minh City, Vietnam. 1999.
- Nguyen HH. Họ Chè - Theaceae, in checklist of plant species of Vietnam, Nguyen TB, Editor. Agricultural Publishing House, Hanoi, Vietnam. 2003.
- Nguyen HH. Thực vật chí Việt Nam (Flora of Vietnam) Vol.19 - Họ Chè (Theaceae D. Don). Publishing House for Science and Technology, Hanoi, Vietnam. 2017.
- Nguyen TB. A list of all plant species in Vietnam. Agricultural Publishing House, Hanoi, Vietnam. 2003.
- Hoang TS, Luong VD. *Adinandra hongiaoensis* (Theaceae), a new species from Lam Dong, Vietnam. *J Jpn Bot.* 2014; 89:331-34. <https://doi.org/10.13140/2.1.2395.3924>.
- Chun WY. Materials for the flora of Hainan (II). *J Syst Evol.* 1963; 8(4):341. Available from: <https://www.jse.ac.cn/EN/Y1963/V8/I4/341>.
- Ming TL. *Gordonia*. FL. Yunnan. Vol. 8. 1997. <https://doi.org/10.1179/014703797788763544>
- Chang HT. *Gordonia*. Fl. Reipubl. Popularis Sin. Vol. 49(3): Science Press, Beijing. 1998.
- Bartholomew B, Ming TL. New combinations in Chinese *Polyspora* (Theaceae). *Novon.* 2005; 15(2):264-66. Available from: <https://www.jstor.org/stable/3393331>.
- Orel G, Wilson PG, Curry AS, Luu HT. *Polyspora huongiana* sp. nov. (Theaceae) from Vietnam and notes on related species. *Nord J Bot.* 2012; 30(1):47-52. <https://doi.org/10.1111/j.1756-1051.2011.01330.x>.
- Orel G, Wilson P, Curry A, Luu HT. *Camellia oconoriana* (Theaceae), a new species from Vietnam. *Edinb J Bot.* 2013; 70 (3):439-47. <https://doi.org/10.1017/S0960428613000103>.
- Luu HT, Nguyen QD, Nguyen TQT. *Camellia sonthaiensis* (Theaceae), a new species from Vietnam. *Ann Bot Fennici.* 2015; 52:289-95. <https://doi.org/10.5735/085.052.0502>.
- Luong VD, Le NHN. *Camellia ninhii*-A new yellow *Camellia* species from Vietnam. *Int Camellia J.* 2016; 48:117. <https://doi.org/10.1097/JNN.00000000000023520>.
- Rivers MC. *Polyspora balansae*. The IUCN Red List of Threatened Species 2018: e.T62079054A62079057. 2018 Accessed on 04 April 2024. Available from: <https://www.iucnredlist.org/>.
- Nguyen QH, Nguyen LTN, Nguyen NTT, Sy TD, Chu MH. Thành phần hóa học và hoạt tính sinh học của cao chiết từ lá loài Dương đồng bích (*Adinandra bockiana* E. Pritz. ex Diels). *VN J Sci Technol.* 2022; 64(12):34-38. <https://doi.org/10.31276/VJST>.
- Pho HTT, Tran DD, Nguyen NTT, Tu TQ, Nguyen QH, Chu MH. Thành phần hóa học và hoạt tính sinh học của cao chiết từ thân của loài sum lông. *TNU J Sci Technol.* 2022; 227(5):232-39. <https://doi.org/10.34238/tnu-jst.5561>.
- Vu OTK, Tran GH, Bui TH, Diep TLP, Nguyen TVA, Le NT. Chemical constituents and biological activity of the stems of *Adinandra hainanensis* Hayata. *Rec Nat Prod.* 2022; 16(4):346-52. <https://doi.org/10.25135/rnp.282.2108.2158>.
- Nguyen LTN, Nguyen QH. Thành phần hóa học và hoạt tính sinh học của cao chiết từ thân của loài sum lá lớn thu tại tỉnh Lào Cai, Việt Nam. *TNU J Sci Technol.* 2021; 226(10):55-61. <https://doi.org/10.34238/tnu-jst.4458>.
- Huynh TH, Tran GH, Trinh NN, Nguyen QH, Ho TH, Le TQP, et al. Chemical components and antioxidant property of acetone

- extracts of four different species of Zingiberaceae. *Plant Sci Today*. 2024; 11(2): In press. <https://doi.org/10.14719/pst.3272>.
26. Vu OTK, Bui TH, Dinh NT, Le NT. Isolation and structure determination of some compounds from *Adinandra poilanei* Gagnep. *VN J Sci Technol*. 2021;63(7). [https://doi.org/10.31276/VJST.63\(7\).22-25](https://doi.org/10.31276/VJST.63(7).22-25).
 27. Fan ZF, Han L, Ma, CL. Research advances of *Polyspora* sweet (Theaceae). *Guihaia*. 2021; 41(10):1755-66 (in Chinese). <https://doi.org/10.11931/guihaia.gxzw202010042>
 28. Fan ZF, Qian SJ, Zhang YH, Ma CL. Characterization of the complete chloroplast genome of *Polyspora tiantangensis* (Theaceae), an endemic and endangered species in south western China. *Mitochondrial DNA Part B: Resources*. 2021; 6(3):814-15. <https://doi.org/10.1080/23802359.2021.1884013>
 29. Ma CL, Li J, Bai Q, Huang XX, Cheng XM. Distribution and utilization of the indigenous tree species of genus *Polyspora* in Yunnan province. *Heilongjiang Agricultural Sciences*. 201; 55:78-80 (in Chinese). <https://doi.org/10.11942/j.issn1002-2767.2015.05.0078>
 30. Li Y, Cao SY, Lin SJ, Zhang JR, Gan RY, Li HB. Polyphenolic profile and antioxidant capacity of extracts from *Gordonia axillaris* fruits. *Antioxidants*. 2019; 8(6):150. <https://doi.org/10.3390/antiox8060150>
 31. Li Y, Li S, Lin SJ, Zhang JJ, Zhao CN, Li HB. Microwave assisted extraction of natural antioxidants from the exotic *Gordonia axillaris* fruit: Optimization and identification of phenolic compounds. *Molecules*. 2017; 22(9):1481. <https://doi.org/10.3390/molecules22091481>
 32. Fu HZ. Studies on the chemical constituents and bioactivities of *Gordonia chrysandra* and *Gordonia kwangsiensis*. Chinese Academy of Medical Sciences & Peking Union Medical College (in Chinese). 2012.
 33. Tang J. Studies on chemical constituents and bioactivities of *Polyspora longicarpa*: Synthesis and bioactivities of carbazole alkaloid Claulansine F (in Chinese). Chinese Academy of Medical Sciences & Peking Union Medical College. 2013.
 34. Xu L, Ren Q, Wan KH, Fu HZ. Chemical constituents from stem of *Gordonia chrysandra*. *Journal of Chinese Medicinal Materials*. 2019; 42(9):2045-48 (in Chinese). <https://doi.org/10.13863/j.issn1001-4454.2019.09.015>
 35. CLSI. Methods for antimicrobial dilution and disk susceptibility testing of infrequently isolated or fastidious bacteria. 3rd ed. CLSI Guideline M45. Wayne, PA. Clinical and Laboratory Standard Institute. 2016.
 36. Wang K, Yang JZ, Zuo L, Zhang DM. Two new flavanonol glycosides from *Gordonia chrysandra*. *Chinese Chem Lett*. 2008; 19(1):61-64. <https://doi.org/10.1016/j.ccllet.2007.10.033>
 37. Fang WU. A new phenolic glycoside from stems of *Gordonia chrysandra*. *Chinese Traditional and Herbal Drugs*. 2019; 24: 2997-000.