



RESEARCH ARTICLE

Phytochemical screening and antioxidant properties of alcoholic extract and antibacterial activity of *Rosmarinus officinalis* L., Leaves

Thekra Sideeq Al-Tayawi*, Esraa Mohammed Ade1 & Farah Hazem Omer

Department of Clinical Laboratory Sciences, College of Pharmacy, University of Mosul, Mosul 41001, Iraq

*Email: thekra.siddeq@uomosul.edu.iq



ARTICLE HISTORY

Received: 11 May 2024
Accepted: 08 June 2024

Available online
Version 1.0 : 24 July 2024
Version 2.0 : 25 July 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Al-Tayawi TS, Ade EM, Omer FH. Phytochemical screening and antioxidant properties of alcoholic extract and antibacterial activity of *Rosmarinus officinalis* L., Leaves. Plant Science Today. 2024; 11(3): 383-391. <https://doi.org/10.14719/pst.3897>

Abstract

Antibiotic resistance is increasing due to the increased use of antibiotics. The emergence of new antibacterial drugs with fewer side effects than antibiotics is an issue. The medicinal plant rosemary is widely used in traditional medicine. In this study, an ethanol extract from the leaves of this plant was tested against the most common hospital-acquired infections. The following study aimed to ascertain the phytochemicals, antibacterial, and antioxidant activities of commercial rosemary flavonoids in leaf extract used on major hospital pathogens through disc diffusion and MIC tests. The *Rosmarinus officinalis* plant was used in this experimental study to evaluate its antimicrobial efficacy against pathogens by disc diffusion and MIC tests. The leaves of this plant were extracted in ethanol at concentrations of 200, 100, 50, and 25 mg/ml and evaluated for their antimicrobial effect against several pathogenic strains. In addition, the optical density in the spectrophotometer (620 nm) was used to calculate the minimum inhibitory concentration (MIC) test. In this investigation, *Pseudomonas aeruginosa*, *Enterobacter faecalis*, *Staphylococcus aureus*, and *Escherichia coli* were all sensitive to the effects of the ethanolic extract of rosemary leaves at a concentration of 200 mg/mL. The minimum inhibitory concentration (MIC) of *E. coli* and *P. aeruginosa* in this extract changed from 25 mg/ml to 100 mg/ml concentration, while the other types of bacteria required only 200 and 100 mg/ml. These findings imply that high concentrations of rosemary extract inhibited all species, including *E. coli*, *S. aureus*, *P. Aeruginosa*, and *E. faecalis*.

Keywords

Rosmarinus officinalis; antimicrobial activity; plant extracts; flavonoids

Introduction

Rosemary, biologically known as *Rosmarinus officinalis* L, is a perennial woody plant native to Asia and the Mediterranean region. It has fragrant, needle-like, evergreen leaves and white, blue, purple, or pink flowers. The leaves of an aromatic evergreen plant called rosemary resemble hemlock needles (1). The leaves are added to stuffing and roasted lamb, chicken, pork, turkey, and beef dishes as a spice and condiment. Rosemary plant is a good source of iron, calcium, glutathione, enzymes, phenolic compounds, vitamins B-6 and E and C, and antioxidant substances, in addition to having a pleasant flavor in culinary preparations (2).

Since ancient times, it has been praised for its therapeutic qualities. According to published studies, "rosemary crude ethanol extract (RO)" may

be an effective antioxidant, anti-inflammatory and anti-tumor agent. In recent years, plant extracts have been created and used as antioxidants in foods. In addition to their antioxidant activity, these extracts may have antimicrobial activity against Gram-positive and Gram-negative bacteria, as they include a wide variety of phenolic compounds, including abietane diterpenoids, carnosol, and ursolic acid (3, 4).

Natural agents are currently being encouraged to replace synthetic antioxidants due to the development of bacteria resistance to antibiotics and the adverse effects of synthetic antioxidants. So, rosemary has long been used as an example in traditional medicine and cosmetics (5, 6). In addition to its therapeutic use, *R. officinalis* essential oil is significant due to its potent antibacterial, cytotoxic, antioxidant, antichloristic, and chemo-preventive qualities (7, 8).

Although it is evident that rosemary extracts are bioactive, nothing is known about their antibacterial capabilities. Despite their antibacterial properties, rosemary extracts' strong taste has kept them from being utilized extensively in food (9).

the other hand, microorganisms can create bioactive substances with cytotoxic, antifungal, and antibacterial properties through their metabolic pathways. All drugs or chemicals are referred to as general antimicrobial agents, which can kill or inhibit the growth of bacteria. The specific toxicity of an antimicrobial agent varies. Some have similar effects on all cell types and act in a relatively non-selective manner. Chemotherapy for infectious diseases often benefits greatly from antimicrobials with selective toxicity. Antimicrobial drugs affect bacteria to varying degrees. They have been found to have many targets within the thin microbial cell, and when all of these targets are disrupted, a microbicidal effect occurs (10, 11).

Interest in phytochemicals as a new source of organic antioxidants, antimicrobials, and hospital infection treatments is on the rise in developed countries. Pathogens are often attacked by synthetic chemicals, but sadly, this indiscriminate use of commercial antibiotics has led to the development of antibiotic resistance in many microorganisms. In addition, these drugs may have negative side effects such as immunosuppression and allergic reactions (12, 13). Therefore, the use of plant extracts is safer for the environment and human health.

Pathogens of the Enterobacteriaceae family are Gram-negative and opportunistic and include *E. faecalis* and *E. coli*. *S. aureus* is also common in hospitals as a Gram-positive bacterium. It regards one of the biggest problems plaguing people who use medications or people whose lifestyles are somehow linked and associated with hospitals is hospital-associated infections (14, 15). One of the most prevalent microorganisms causing infections in hospitalized patients is *P. aeruginosa*. In this work, we investigated the response of *P. aeruginosa* and other bacterial strains to concentrated extracts of rosemary extract. Enterobacteriaceae have recently become an important pathogen of intranasal infections (16). To

identify the phytochemical, antibacterial, and antioxidant activities of commercial rosemary flavonoids for use against Gram-negative bacteria (*P. aeruginosa*, *E. faecium*, *E. coli*) and Gram-positive bacteria (*S. aureus*).

Materials and Methods

This study was conducted at the Laboratory of the Clinical & laboratories Science Department, College of Pharmacy, University of Mosul, in 2019.

Samples Collection:

Leaves of rosemary were collected and identified in cooperation with the Herbarium and Herbal Research Center of Mosul. The test organisms *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Enterobacter faecalis* were obtained from the Biology Department- College of the Science/University of Mosul Iraq.

Preparation of Plant extracts:

The method of Golshani and Sharifzadeh (17) was adopted for the preparation of plant extracts with little modifications. Briefly, the fresh leaves of *R. officinalis* were collected and dried in a place not exposed to sunlight. Then, the dried leaves were ground into a powder, and 25 g of the plant powder was placed in 150 ml of ethanol absolute and stirred for 24 h at room temperature. Whatman No. 1 filter paper was used to filter the mixture.

Phytochemical screening:

The evaluation and the screening of rosemary extract were done using a conventional approach as used by (18, 19).

Alkaloid:

A total of 1 ml of Mayer's reagent was added to a test tube containing a small amount of sample (mercuric potassium iodide). Alkaloid was present because a yellow-cream precipitate was formed.

Terpenoids testing:

After adding two drops of chloroform and 1 ml of strong hydrochloric acid to a small sample and heating for two minutes, a reddish-brown color appeared, indicating the presence of terpenoids.

Phenol testing:

A total of 2 ml of ferric chloride solution (5%) was added to a small portion of the sample and allowed to incubate at room temperature for 5 minutes. The presence of phenol is indicated by the production of a black, blue hue.

Tannin testing:

A few drops of natural lead acetate solution (10%) were added to a small amount of the sample to show the presence of tannins by producing a yellow ppt precipitate.

Carbohydrates testing:

Molisch test: The sulfuric acid-treated extract was carefully added along the side of the test tube, where it formed a violent ring indicating the presence of carbohydrates (20).

Saponins testing: A little sample was obtained, and 2-3 drops of distilled water were added before shaking thoroughly. The formation of foam shows the presence of saponin.

Coumarin glycosides testing:

The presence of coumarin was detected by the appearance of a yellow precipitate after the introduction of a few drops of the extract into diluted HNO_3 (21).

Test for flavonoids:

FeCl_3 solution was added to the sample solution, and the onset of a tingling, greenish-black color signifies the presence of flavonoids.

Quinine test:

Quinine can be detected by adding ml of sodium hydroxide solution to a tiny sample and waiting two minutes. Quinine will then turn reddish-brown.

Protein testing:

A little sample was obtained and placed in a test tube. A drop of concentrated HNO_3 was then added. Reddish color formation revealed the presence of the protein.

Steroid testing through Salkowski reaction:

A small sample was mixed with 1 ml of concentrated H_2SO_4 and 1 ml of chloroform, stirred thoroughly, and left for 5 minutes. When the lower layer appears red, it indicates the presence of steroids.

Extraction of Flavonoid:

The dried leaves of rosemary were ground to a powder, and 25 g of the plant powder was placed in 150 ml of 100% ethanol and stirred for 24h at room temperature. Whatman No. 1 filter paper was used to filter the mixture. With the remaining residue, the above procedure was performed again, and 150 ml of absolute ethanol and the two filtrates were properly combined. The filtrate was then mixed with 100 ml of 1% lead acetate solution for 4 hours to ensure complete precipitation, after which the filtrate was collected. The filtrate was mixed with 250 ml of acetone and 30 ml of hydrochloric acid. The entire mixed solution was stored at 4°C. The 100 ml liquid sample was completely evaporated each time using a rotary evaporator while operating under reduced pressure to produce a dark brown precipitate, which was then used in various scientific experiments (22).

Chemical screening of Flavonoids:

The presence of flavonoids and phenolic compounds was checked by adding 5 ml of plant extract to a few drops of 1% potassium hydroxide (KOH) and 1% ferric chloride ($\text{FeCl}_3 \cdot 2\text{H}_2\text{O}$) solutions, respectively, at a time. The formation of the yellow and blue-green colors showed the presence of flavonoids and phenolic chemicals in the plant extracts, respectively (23).

Determining the net flavonoid content in the extracts

A drop of glacial acetic acid, 1.0 ml of aluminum trichloride in ethanol (20 mg/ml), and 1 ml of rosemary extract were mixed to detect the flavonoid content. The mixture was then diluted with ethanol to a volume of 25 ml to quantify

the overall flavonoid concentration. After 40 minutes, the absorbance at 415 nm was measured using (a spectrophotometer). 1.0 ml of rosemary extract, one drop of glacial acetic acid, and 25 ml of ethanol were used to prepare the blank. Under the same conditions, the absorption of a standard (quercetin) solution (0.5 mg/ml) of ethanol was determined.

The following equation was used to determine the total amount of flavonoids present in the rosemary extract as quercetin equivalents (QE):

$$X=(A.mg)/(Aq.m) \text{ (Eq.1)}$$

Where: X is the number of flavonoids (mg/mg rosemary extract in QE), A is the amount of Rosemary extract absorbed, Aq is the number of standard quercetin solutions absorbed, m solution is the number of pomegranate flowers (mg) in the extract, and mq in rosemary s is the amount of quercetin in the solution (mg).

Effect of extract on DPPH radical scavenging

The ability of plant extracts to scavenge free radicals was assessed using the following method. Plant extracts were added to a known volume of test tubes (50-150 L), completed with a known volume (1.0 mL) by D.W., and then (1.0 mL) of DPPH solution (0.2 mM of ethanol) was added to each test tube. After that, the tubes were left at room temperature for 30 minutes. The same procedure was used to make a control but without the use of plant extracts. As a positive control, an ascorbic acid solution (0.03% w/v) was used. The absorbance (A) of this solution was determined at 517 nm using a Jenway 6300 spectrophotometer. The following formula was used to determine the percentage inhibition of DPPH radicals in this product. The percentage I am equal to $[(Ac-As)/Ac]$ (24).

Identification of the compounds:

FTIR spectrophotometer was used to evaluate the plant extract. Japan's SIMADZU, Model No. 8400S.

Bacterial Culture:

The rosemary flavonoid extracts were then stored in a refrigerator, covered with aluminum foil to protect them from the effects of light. Flavonoid extracts were administered and dissolved in 5% dimethyl sulfoxide at doses of 25, 50, 100, and 200 mg/ml (DMSO). The inhibitory activity was then evaluated using the disc diffusion technique and MIC test.

Evaluation of the rosemary flavonoid extract's inhibitory activity:

Mueller-Hinton (MH) agar is used to maintain bacterial cultures. It was made according to the instructions provided by the manufacturer. The surfaces of agar plates previously infected with a standard number (1.5105) of microorganisms were covered with sterile filter paper sheets of uniform size (6 mm) that had been soaked with various concentrations, which were 25, 50, 100, and 200 mg/ml of RO alcohol extract. The incubation period of the plates was of one day at room temperature of 37°C. At the end of the incubation, the plates were checked for the

presence of an inhibition zone (IZ) around the plate that would prevent bacterial growth.

MIC test:

The antimicrobial activity of the extracts was observed after six hours, and this was determined by the minimum inhibitory concentration of each target bacteria individually. Tubes containing nutrient broth enriched with different amounts (10-500 L) of extracts were filled with 100 L of 10^6 cells/ml. After 24 h at 37°C, sample readings were compared to negative controls by measuring the optical density (620 nm) in a spectrophotometer, and the results were recorded (media without bacteria), with samples without turbidity classified as MIC. Each test was performed three times, and the mean value was then displayed (25).

Antibiotic resistance:

Antibiotic plates were placed on inoculation plates at 37°C for 24 h. Following this, bacterial strains suspended in nutrient broth were inoculated onto the surface of each nutrient agar plate. Chloramphenicol (C), ceftriaxone (CRO), ciprofloxacin (CIP), and amoxicillin (AM) were utilized in this study.

Bacterial cultures and preparation of rosemary extracts:

The leaves were powdered after sun-drying. The leaves weighing up to 50 g were added to a sterilized Erlenmeyer flask. The next stage was the addition of 250 ml of 100% ethanol to dissolve the components of the herbs. The alcohol and herbal powder were placed in a conical flask and shaken for 48 hours until the solvent started to act at a temperature of 40°C. The solvent was then removed using the rotary method. Finally, the rosemary extract was stored in sterilized refrigerator-safe plates and covered with aluminum foil to protect it from light. The extracts were provided in 5% dimethyl sulfoxide at strengths of 25, 50, 100, and 200 mg/ml. Inhibitory activity was then evaluated using disc diffusion techniques and MIC testing.

Results

As shown in Table 1 and Fig. 1, the chemical analysis of rosemary leaf extracts revealed the presence of different concentrations of active substances and other

Table 1. Results of phytochemical analysis Results of phytochemical screening of *R. officinalis* leaf extracts.

Phytoconstituents test	Observations	Result
Alkaloids	formation of yellow	+
Terpenoids	Formation reddish colour	+
Phenols	Observed black-bluish precipitate	+
Tannins	Observed yellow precipitate	+
Saponins	No Formation foam	-
Coumarine glycoside	Observed yellow precipitate	+
Flavonoids	Formation greenish blue	+
Quinine	Formation reddish brown	+
Steroids	Formation red colour	+
Protein	No Formation of reddish colour	-
Carbohydrate	No Formation purple ring	-

(+) = positive reaction; (-) = negative reaction.

unidentified components.

Flavonoid Content

In Table 2, the total flavonoids of rosemary ethanolic extract are given as quercetin equivalents (quercetin/g extract). The results showed that the total flavonoid

Table 2. Total solid and flavonoid content of *R. officinalis* leaves

Alcoholic <i>R. officinalis</i> leaves	Total solids (%)	Total flavonoids (mg/g)
	36	8.34



Fig. 1. Results of phytochemical Screening *R. officinalis* leaves extracts

content of the ethanolic extract was 8.34 mg QUE/g dry weight.

Antioxidant Study:

DPPH radical scavenging activity

The radical scavenging potential of different concentrations of rosemary leaf extract varied from 75% concentration at 62.5 ppm to 76% at 125 ppm and 79% at 187 ppm, depending on the concentration. Table 3 summarizes the results of scavenging DPPH activity by

Table 3. Scavenging activity of ethanolic extract of rosemary leaves against DPPH.

Treatment	Conc. (ppm)	Scavenging activity (%)
Alcoholic rosemary leaves	62.5	75
	125	76
	187	79.3
Ascorbic acid	23	81.96




Fig. 2. Discoloration of DPPH under the influence of *R. officinalis* leaves extract at different concentrations.

flavonoids of rosemary leaves ethanolic extract and compares them with the standard antioxidant ascorbic acid, as shown in (Fig. 2).

Fourier Transform Infrared Spectroscopy (FTIR) Assay

The solutions were analyzed individually using an FT-IR spectrophotometer (SHIMADZU, Japan, model 8400S).




Fig. 3. The result of FTIR Spectroscopy

Table 4. The functional groups of the purified flavonoid compound from the IR-spectrum

Wave no.2	Band shape	Band	Functional group
3329	Band	OH	Alcohol
2927	sharp	-CH2-	Aliphatic stretch
1446-1517	sharp	-C=C-	Stretching of olefinic
1684	sharp	C=O	Stretching C=O of -CHO Group
1270	sharp	C-O	C-O Stretching

According to the FT-IR analysis (Fig. 3), the molecule includes multifunctional groups such as phenolic -CH₂-, -C=C-, C=O, etc. (Table 4).

According to the data, quinoline alkaloids, phytol, eucalyptol, camphor, and verbenone were the most prevalent substances. Other substances, such as borneol flavor, borneol acetate flavor, fragrance, heliotrope, geraniol, and lidocaine, were found in lower concentrations.

Activities of referred antibiotics:

Table 5. Antimicrobial activities of the suggested antibiotics against tested microorganisms represented as resistant (R), susceptible (S), and intermediate (I).

Bacterial strains	Antibiotics			
	C*	CRO*	CIP*	AM*
<i>Staphylococcus aureus</i>	R	R	R	R
<i>Escherichia coli</i>	S	R	R	S
<i>Pseudomonas aeruginosa</i>	I	S	S	I
<i>Enterobacter faecalis</i>	I	R	S	S

*Chloramphenicol (C), Ceftriaxone (CRO) Ciprofloxacin (CIP), Amikacin (AM)



Fig. 4. Inhibition zones (mm) of antibiotics against 1- *E. faecalis*, 2- *P. aeruginosa*, 3- *E. coli*, 4- *S. aureus*, respectively.

The bactericidal activities of the suggested antibiotics were also conducted on four types of bacteria that were taken into consideration in this study. (Table 5) (Fig. 4)

Antibacterial activity of rosemary extracts on various microorganisms:

In this investigation, the presence of IZ in millimeters was

Table 6. Antimicrobial activities of RO alcoholic extract against tested microorganisms represented as inhibition zone diameter (mm)

Concentration of Extracts mg/ml	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. aureus</i>	<i>E. faecalis</i>
200	19mm	21mm	10mm	10mm
100	13mm	15	8	8
50	10	11	-	-
25	8	8	-	-




Fig. 5. Inhibition zones (mm) of rosemary ethanolic extract against : 1- *S. aureus*, 2- *E. coli*, 3- *P. aeruginosa* and *E. faecalis*, respectively.

used to assess the antimicrobial activity of phenolic components of rosemary leaf extracts against the tested types. (Table 6) (Fig. 5)

Minimum inhibitory concentration (MIC)

Table 7. MIC values of alcoholic rosemary extract determined with agar diffusion method.

Bacterial species	MIC
<i>Pseudomonas aeruginosa</i>	0.25
<i>Escherichia coli</i>	0.125
<i>Enterobacter faecalis</i>	0.5
<i>Staphylococcus aureus</i>	0.5

As shown in Table 7, the ethanolic extract of rosemary leaves showed MICs ranging from 0.125 to 0.5 mg/ml against the indicated bacteria.

Discussion

The main causes of food quality deterioration and shelf life reduction are bacterial infection and lipid oxidation. Therefore, controlling bacterial cross-contamination and delaying lipid oxidation is of critical importance to food processors (26). Although the rosemary plant is a rich source of phenolic compounds with strong antioxidant and antimicrobial properties, little is known about its antimicrobial effects. Furthermore, there is some evidence that minor elements have an important role in antimicrobial activity, perhaps through synergistic effects with other elements. In this investigation, the flavonoids found to be most effective against bacterial species were rosmarinic acid and carnosic acid (27). Their unique antioxidant function may be related to their antibacterial effect. It was found that ethanolic plant extracts showed broad-spectrum activity against Gram-positive and Gram-negative bacteria (28).

The results of the study showed that phenolic flavonoids and substituted benzene hydrocarbons were the more readily available components of rosemary extracts. GC-MS analysis of rosemary led to the identification of more than 20 compounds by comparing the recorded mass spectra with those in a computer library, although the most active substances were eucalyptol, camphor, lobster, lobster acetate, and verbenone (29). The quinoline alkaloids found in rosemary extracts are of high medicinal value; in addition to their recognized antimicrobial properties, they are frequently used in the food and soft drink industries. Phytol had the largest percentage, and it was used as a mouthwash and preservative. Stearic acid, geraniol, and geraniene were the substances with the lowest percentages (0.64%) at

(0.71%), respectively (30, 31). These findings are consistent with a related study that found that rosemary extract included higher levels of camphor, lobster, lobster acetate, and eucalyptol. These variations may be due to seasonal variations, plant conditions, and extraction techniques.

E. coli was sensitive to some drugs but resistant to ciprofloxacin and ceftriaxone, while *S. aureus* showed resistance to all prescribed antibiotics. On the other hand, *P. aeruginosa* showed some susceptibility to other antibiotics. These findings support the notion that *E. coli* strains resistant to some antibiotics become vulnerable to alcoholic extracts of *R. officinalis* plants (32).

The flavonoids in the rosemary extract were active against Gram-positive (*S. aureus*) and Gram-negative (*E. coli*, *P. aeruginosa*, and *E. faecalis*) pathogenic microorganisms at concentrations 200, 100, 50, and 25 mg/ml ranging from 8 to 20 mm (Table 6). The zone of inhibition increased with the increasing concentration of methanolic extract. Furthermore, the results showed that rosemary extract affected *P. aeruginosa*, contrary to the previous investigations (33, 34). In addition, *P. aeruginosa* and *E. coli* were more sensitive to the concentration of 200 mg/ml. The results of the study showed that there were significant differences in the sensitivity of the tested bacteria to ethanol extracts. In other words, Gram-negative bacteria such as *E. coli* and *P. aeruginosa* showed more sensitivity than Gram-positive bacteria such as *S. aureus*.

Their growth was stopped in 25 mg/ml of extract. These findings are inconsistent with those who claim that Gram-positive bacteria are the most sensitive compared to other bacterial species. According to those who found that the changes in antimicrobial activity could be caused by changes in the bacterial cell wall composition, the extract and its components were more active against the bacterial species (35). In addition, the presence of an inhibition zone does not always indicate that a compound is inactive. For example, non-polar substances may not disperse into the culture medium.

The MIC of rosemary extract is considered to be one of the lowest levels in inhibiting the growth of bacterial cells. The data suggest that higher extract concentrations are needed to inhibit microorganisms. *E. coli* can be inhibited with 25 mg/ml of extract, but *S. aureus* requires 100 mg/ml. The use of plants to treat diseases, including infectious diseases, has been widely used, and data in the literature, as well as our findings, reveal the great potential of phototherapy, although they have not been fully investigated (36, 37). Therefore, more research is needed to find new compounds. Once extracted, they should be tested for toxicity in vivo bioassays before being used in new therapeutic approaches, as has been demonstrated for different plant extracts.

The development of antibiotic resistance has been of particular concern and is currently one of the main therapeutic issues. Since rosemary extracts have been shown to have antibacterial effects against various

bacteria, they are recommended for treating bacteria-causing infections. Further research into medicinal plants is recommended as potential alternatives to chemical drugs, which, although beneficial, have negative side effects (38).

The European Food Safety Authority (EFSA) has examined the safety of rosemary extracts. It has been concluded that estimates of excessive intake of rosemary alcohol and rosemary acid are 0.09 (elderly) to 0.81 (adolescents) mg/kg per day. Rosemary extracts are now added to food and beverages in the EU in amounts up to 400 mg/kg (as the sum of carnosic acid and carnosol).

Conclusion

In conclusion, it can be said that *Rosmarinus officinalis* L. leaf extract has inhibitory effects on different types of bacteria and can be used in the pharmaceutical industry to create new synthetic drugs to treat infectious diseases brought about by food pathogens of these bacterial species, such as *E. coli*, *P. aeruginosa*, etc., which may cause food spoilage and subsequently human diseases.

These findings imply that plant extracts belonging to *R. officinalis* might be employed as natural antibacterial agents in medical settings. Further study will be needed to determine the active component in each extract and to investigate its phytopharmaceutical properties.

Overall, the prevalence of drug-resistant strains has grown over the past several decades due to the widespread use of antibiotics and other antimicrobial treatments, especially in the treatment of infections. Due to the harmful effects and high levels of toxicity of most manufactured remedies, there has become a great urgent need to create alternative materials such as herbal extracts or their active ingredients in treatments.

Acknowledgements

The authors wish to thank the Department of Microbiology, College of Science, University of Mosul, for providing bacterial strains, and the Department of Clinical Laboratory Sciences, College of Pharmacy, University of Mosul, for providing the necessary facilities to carry out the present investigation.

Authors' contributions

TA carried out the Conception and design, Analysis and interpretation of the data, and Drafting of the article, while EA performed the critical revision of the article for important intellectual content and provision of study materials. FO conceived the study and participated in its design and coordination. All authors contributed to the study design, performed the statistical analysis, and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None.

References

- Oussaid A, Azzouzi M, Mansour AI, Azouagh M, Koudad M, Oussaid A. Assessment of the Chemical/biological activities of extracts and essential oil of *Rosmarinus officinalis* L. from the Oriental region of Morocco. Moroccan Journal of Chemistry. 2020; 8 (3):8-3. <https://doi.org/10.48317/IMIST.PRSM/morjchem-v8i3.20151>
- Chetouani M, Mzabri I, Amar A, Boukroute A, Kouddane N, Berrichi A, Morphological-physiological and biochemical responses of Rosemary (*Rosmarinus officinalis*) to salt stress, Materials Today: Proceedings 13 (2019) 752-761. <https://doi.org/10.1016/j.matpr.2019.04.037>
- Jeba Priya GR, Anburaj R, Kanmani M, Singh CR. *In vitro* antibacterial activity and phytochemical analysis of ethanolic leaf extract of *Rosmarinus officinalis* L. International Journal of Innovative Engineering and Management Research. 2020;3 (09):12.
- Aziz E, Batool R, Akhtar W, Shahzad T, Malik A, Shah MA, Iqbal S, Rauf A, Zengin G, Bouyahya A, Rebezov M, Dutta N, Khan MU, Khayrullin M, Babaeva M, Goncharov A, Shariati MA, Thiruvengadam M. Rosemary species: a review of phytochemicals, bioactivities and industrial applications. South African Journal of Botany. 2022; 151:1-18. <https://doi.org/10.1016/j.sajb.2021.09.026>
- Bendahou M, Benabdellah M, Hammouti B. A study of rosemary oil as a green corrosion inhibitor for steel in 2M H₃PO₄. Pigm Resin Tech. 2006; 95-100. <https://doi.org/10.1108/03699420610652386>
- Cocan I, Alexa E, Danciu C, Radulov I, Galuscan A, Obistioiu D, Morvay AA, Sumalan RM, Poiana MA, Pop G, Dehelean CA. Phytochemical screening and biological activity of Lamiaceae family plant extracts. Experimental and Therapeutic Medicine. 2018;15(2):1863-1870. <https://doi.org/10.3892/etm.2017.5640>
- Celiktas OY, Kocabas EH, Bedir E, Sukan FV, Ozek T, Baser KH. Antimicrobial activities of methanol extracts and essential oils of *Rosmarinus officinalis*, depending on location and seasonal variations. Food Chemistry. 2007;100(2):553-559. <https://doi.org/10.1016/j.foodchem.2005.10.011>
- Farouk S, Al-Amri SM, Exogenous melatonin-mediated modulation of arsenic tolerance with improved accretion of secondary metabolite production, activating antioxidant capacity and improved chloroplast ultrastructure in rosemary herb. Ecotoxicology and Environmental Safety. 180 (2019) 333-347. <https://doi.org/10.1016/j.ecoenv.2019.05.021>
- Sushanthi S, Prabakar J, Indiran MA, Kumar RP, Doraikannan S, Leelavathi L, Kumar SR. Assessment of cytotoxic and antioxidant effects of *Rosmarinus officinalis* and fluoride-containing dentifrice-an *in vitro* study. Research Journal of Pharmacy and Technology. 2021;14(7):3779-3784. <https://doi.org/10.52711/0974-360X.2021.00654>.
- Mushtaq W, Naim A, Naeem S. *In vitro* antibacterial activity of *Rosmarinus officinalis* L. and *Murraya koenigii* L. against multi-drug resistant *staphylococcus species*. Int J Biol Biotechnol. 2019; 16(4):1005-10.
- Adris AA, Soultan AA, Bellail AA, Ibrahim FA. Antioxidant and Antimicrobial Activities of *Rosmarinus officinalis* L. Growing Naturally in El-Jabal El-Akhdar Province–Libya and its effect on keeping quality of cold serola dumereri Fillets. Journal of Food and Dairy Sciences. 2019; 10(2):23-30. <https://doi.org/10.21608/jfds.2019.36149>
- Kamli MR, Sharaf AA, Sabir JS, Rather IA. Phytochemical Screening of *Rosmarinus officinalis* L. as a potential anticholinesterase and antioxidant–medicinal plant for cognitive decline disorders. Plants. 2022; 11(4):514. <https://doi.org/10.3390/plants11040514>
- Paudel S, Mishra N, Agarwal R. Phytochemicals as immunomodulatory molecules in cancer therapeutics. pharmaceuticals. 2023, 16(12): 1652. <https://doi.org/10.3390/ph16121652>
- Charpentier T, Boisard S, Le Ray AM, Bréard D, Chabrier A, Esselin H, Guilet D, Ripoll C, Richomme P. A descriptive chemical composition of concentrated bud macerates through an optimized SPE-HPLC-UV-MS2 method—application to *Alnus glutinosa*, *Ribes nigrum*, *Rosa canina*, *Rosmarinus officinalis*, and *Tilia tomentosa*. Plants. 2022; 11(2):144. <https://doi.org/10.3390/plants11020144>
- Behbahani BA, Yazdi FT, Shahidi F, Noorbakhsh H, Vasiee A, Alghooneh A. Phytochemical analysis and antibacterial activities extracts of mangrove leaf against the growth of some pathogenic bacteria. Microbial pathogenesis.2018;114:225-32. <https://doi.org/10.1016/j.micpath.2017.12.004>
- Amel AA, Ayat AA, Wafaa MA, Samar MS, Hind HA. Antimicrobial and antioxidant activities and phytochemical analysis of *Rosmarinus officinalis* L. Pod and *Thymus vulgaris* L. leaf ethanolic extracts on *Escherichia coli* urinary isolates. International Journal of Microbiology. 2023;4171547:1-7 pages. <https://doi.org/10.1155/2023/4171547>
- Golshan Z, Sharifzadeh A. Evaluation of antibacterial activity of alcoholic extract of rosemary leaves against pathogenic strains. Zahedan Journal of Research in Medical Sciences. 2014; 16(3): 12-15.
- Kumar A, Singh S, Mahour K, Vihan VS, Gurura K . Phytochemical analysis of some Indigenous plants potent against Ectoparasite. Asian Journal of Experimental Biological Sciences. 2011; 2(3): 506-509.
- Das BK, Al-Amin MM, Russel SM, Kabir S, Bhattacharjee R, Hannan JMA. Phytochemical screening and evaluation of analgesic activity of *Oroxylum indicum*. Indian J Pharm Sci. 2014; 76(6): 571-575. PMID: PMC4293694
- Manilal A, Sabu KR, Shewangizaw M, Aklilu A, Seid M, Merdekios B, Tsegaye B. *In vitro* antibacterial activity of medicinal plants against biofilm-forming methicillin-resistant *Staphylococcus aureus*: efficacy of *Moringa stenopetala* and *Rosmarinus officinalis* extracts. Heliyon. 2020;6(1):e 03303. <https://doi.org/10.1016/j.heliyon.2020.e03303>
- Nefzi K, Ben Jemaa M, Baraket M, Dakhlaoui S, Msaada K, Nasr Z. *In Vitro* Antioxidant, Antibacterial and Mechanisms of Action of Ethanolic Extracts of Five Tunisian Plants against Bacteria. Applied Sciences. 2022 ; 12(10):5038. <https://doi.org/10.3390/app12105038>
- Milena T, Vasil A, Zvezdelina Y, Donika I, Toncho D. Selectivity of Current Extraction Techniques for Flavonoids from Plant Materials. Processes. 2020; 8(10), 1222; <https://doi.org/10.3390/pr8101222>
- Trease GE, Evans WC. *Pharmacognosy*. 15th Ed. London: Saunders Publishers; 2002. pp. 42–44.
- Foti MC, Daquino C, Geraci C. Electron-transfer reaction of cinnamic acids and their methyl esters with the DPPH radical in alcoholic solutions. J Org Chem. 2004;69(7):2309-14. <https://doi.org/10.1021/jo035758q>
- Althobaiti F. Antibacterial Activity And Genotoxicity Effect of Ethanolic Leaves Extract of *Rosmarinus Officinalis*. Journal of

- Pure and Applied Microbiology. 2020; 14(3):2165-72. <https://doi.org/10.22207/JPAM.14.3.57>
26. Maliki I, Moussaoui AE, Ramdani M, EL Badaoui K. Phytochemical screening and the antioxidant, antibacterial and antifungal activities of aqueous extracts from the leaves of *Salvia officinalis* planted in Morocco. Moroccan Journal of Chemistry. 2021; 9 (2):9-2. <http://doi.org/10.48317/IMIST.PRSM/morjchem-v9i2.24840>
 27. Zaïri A, Nouir S, Khalifa MA, Ouni B, Haddad H, Khélifa A, Trabelsi M. Phytochemical analysis and assessment of biological properties of essential oils obtained from thyme and *Rosmarinus species*. Current Pharmaceutical Biotechnology. 2020;21(5):414-24. <https://doi.org/10.2174/1389201020666191019124630>
 28. Christopoulou SD, Androutsopoulou C, Hahalis P, Kotsalou C, Vantarakis A, Lamari FN. Rosemary extract and essential oil as drink ingredients: An evaluation of their chemical composition, genotoxicity, antimicrobial, antiviral, and antioxidant properties. Foods.2021 Dec 18;10(12):3143. <https://doi.org/10.3390/foods10123143>
 29. Ielciu I, Niculae M, Pall E, Barbălată C, Tomuță I, Olah NK, Burtescu RF, Benedec D, Oniga I, Hanganu D. Antiproliferative and antimicrobial effects of *Rosmarinus officinalis* L. loaded liposomes. Molecules. 2022;27(13):3988. <https://doi.org/10.3390/molecules27133988>
 30. Nieto G, Ros G, Castillo J. Antioxidant and antimicrobial properties of rosemary (*Rosmarinus officinalis*, L.): A review. Medicines.2018;5(3):98. <https://doi.org/10.3390/medicines5030098>
 31. Jeevalatha A, Kalaimathi RV, Basha AN, Kandeepan C, Ramya S, Loganathan T, Jayakumararaj R. Profile of bioactive compounds in *Rosmarinus officinalis*. Journal of Drug Delivery and Therapeutics. 2022 Jan 15;12(1):114-22. <https://doi.org/10.22270/jddt.v12i1.5189>
 32. Fazeli-Nasab B, Shahraki-Mojahed L, Dahmardeh N. Evaluation of antimicrobial activity of essential oil and ethanolic extract of 10 medicinal plants on *Rathayibacter tritici* and *Xanthomonas translucens*. Plant Biotechnology Persa.2022 Jun 10;4(1):10-7. <https://doi.org/10.52547/pbp.4.1.3>
 33. Janampa E, Perez H, Mauricio F, Alvéitez-Temoche D, Mauricio-Vilchez C, Gallo W, Mayta-Tovalino F. Antibacterial activity of ethanol extract of the leaves of *Rosmarinus officinalis* L. at different concentrations versus *Streptococcus mutans*: An *In vitro* comparative study. European Journal of General Dentistry.2020 Sep;9(03):152-6. https://doi.org/10.4103/ejgd.ejgd_148_20.
 34. Hussein Rajaa A. Extraction and Identification of A Flavonoid compound from Oak Plant *Quercus infectoria* (Olivier) and study of its Antibacterial Activity, *in vitro.*, QMJ. 2013; 13:80-94. <https://doi.org/10.28922/qmj.2013.9.16.80-94>.
 35. Malek A, M Sadaka MW, Hamo S, Al-Mahbashi HM. Evaluation of antidiabetic activity of *rosmarinus officinalis* var. prostratus growing in syria in alloxan diabetic rats. Current Bioactive Compounds. 2021 Feb 1;17(2):187-93. <https://doi.org/10.2174/1573407216999200426235739>.
 36. Abdulbary M, Shakir MR, Sadik AM. Detection about the anti-inflammatory effect in alcoholic extract of *Rosmarinus officinalis in vitro*. Journal of Pharmaceutical Sciences and Research. 2019; 11 (4):1531-3.
 37. Chaul LT, Alves VF, de Sá S, Oliveira LD, de Oliveira Silva E, de Sousa Fiuza T, Faustino LM, Torres IM, dos Santos PA, da Conceição EC, de Paula JR. Antimicrobial Activity an Physicochemical characterization of extracts and fractions of *Rosmarinus officinalis* and *Origanum vulgare*. Revista Fronteiras: Journal of Social, Technological and Environmental Science. 2022; 11(1). <https://doi.org/10.21664/2238-8869.2022v11i1.p8-30>.
 38. Divya S, Arivoli S, Tennyson S. Phytochemical analysis and *in vitro* Antioxidant and antibacterial activity of the chloroform leaf extract of *Abelmoschus manihot* (L.) Medik (Malvaceae). Research Journal of Pharmacy and Technology. 2021;14(9):4719-26. <https://doi.org/10.52711/0974-360X.2021.00821>.