



RESEARCH ARTICLE

Chemical composition and antioxidant activity of *Physalis angulata* L. (Solanaceae) and its effect on blood clotting and biofilm formation of some wound bacterial isolates

Sara H. Shaker*, Ashwaq T. Hameed

Department of Biology, University of Anbar, 31008, Iraq

*Email: sar21w4008@uoanbar.edu.iq



ARTICLE HISTORY

Received: 30 May 2024
Accepted: 05 June 2024

Available online
Version 1.0 : 12 August 2024
Version 2.0 : 13 August 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Shaker SH, Hameed AT. Chemical composition and antioxidant activity of *Physalis angulata* L. (Solanaceae) and its effect on blood clotting and biofilm formation of some wound bacterial isolates. Plant Science Today. 2024; 11(3): 583-591. <https://doi.org/10.14719/pst.3898>

Abstract

Physalis angulata a member of the Solanaceae family, is a plant recognized for its significant nutritional and medicinal properties. Widely distributed across tropical and subtropical regions, this plant has garnered attention for its diverse biological activities. The ethanol extract from the plant is being used in this study to investigate its biological and antioxidant properties. The ability of the extract to reduce free radicals was also tested using the compound phenyl-1-picrylhydrazyl). The DPPH method was used because it is a modern color method and is considered the most common and easy to apply, in addition to its speed and superior sensitivity in testing antioxidants for samples of plant extracts, and the values were compared with the value of ascorbic acid IC₅₀. The study also looks at the extract's ability to prevent the formation of biofilms by different pathogens, including *E. coli* and Staphylococcus strains that were isolated from community labs and Ramadi Teaching Hospital. The ethanol extract demonstrated a noteworthy ability to inhibit the formation of cellular membranes in staphylococcal bacteria and *E. coli*. The efficacy of the extract at different concentrations (150, 200, 250 mg/mL) was also assessed, revealing that a concentration of 250 mg/mL resulted in significant inhibition rates (74.3% and 61.45% for *E. coli*). Furthermore, the study explores the impact of ethanol extract on blood coagulation. At high concentrations (150, 200, 250 mg/mL), the extract exhibited properties akin to aspirin and warfarin, extending the clotting times in both total and partial coagulation tests (Prothrombin time, Partial thromboplastin time) at lower concentrations (100, 50, 10 mg/mL), the extract appeared to promote blood clot formation, suggesting a concentration-dependent effect on hemostasis. This dual action positions *P. angulata* as a potential therapeutic agent, warranting further investigation into its biomedical applications.

Keywords

Antioxidant activity; biofilm formation; blood coagulation management; chemical composition; *Physalis angulata* L.

Introduction

The plant *Physalis angulata* belongs to the Solanaceae family and contains more than 120 species that may be perennial or herbaceous. It is a dense herbaceous annual plant up to about 50cm tall. It has bell-like flowers of a yellow color, and its fruits resemble a balloon hanging down, and the fruits are edible. It grows in areas where the weather is temperate and warm. This plant is distributed worldwide in tropical, subtropical, and coastal regions such as South America. It is known by different names such as kamabo, wild

tomato, winter cherry, golden raspberry, etc (1). The plant also contains compounds such as alkaloids, glycosides, terpenoids, saponins, flavonoids, tannins, and withanolides (2); it has immunomodulatory and immunostimulating activity, anti-tumor, anti-inflammatory and anti-malarial activity and has a significant effect on microbes (3). It is used in Japan as an antipyretic. It is used to treat many ailments and diseases of the intestines and gastrointestinal tract, such as ulcers, boils, wounds, etc. Studies have proved that this *Physalis angulata* plant offers a lot of therapeutic substances, such as antiallergic, asthmatic, anti-Leishmania, anti-malarial, and immunomodulatory activity (4).

Withanolides are steroids found in *Physalis*. *Physalis* is a great source of Vitamin C, Beta Carotene, and Vitamin K, which support wound healing and a healthy immune system, and Withanolides can help combat inflammatory diseases (5). Medicinal plants have protective properties resulting from their antioxidant activities and are anti-hypercholesterolemic and anti-cardiovascular diseases (6). Flavonoids are anticoagulants that have a role in vasodilation and have antioxidant activity (7), protecting against blood clots. *E. coli* causes many diseases. Diseases, including septicemia and bacteremia, are caused by the possession of many virulence factors, which are common in humans and animals (8). *S. epidermidis* is widely present and is negative for the blood coagulation test. It is an opportunistic pathogenic bacterium that causes diseases. *P. angulata*, also known as the "Ciplukan" plant, was discovered to have antimicrobial properties. Many studies have proven its ability to prevent the growth of bacteria and fungi. Essential oils extracted from various parts of the plant be effective against a range of microorganisms, including *Candida albicans*, *C. stellate*, *C. torulopsis*, *K. pneumoniae*, *B. subtilis*, *S. aureus*, and *Pseudomonas aeruginosa* (9,10). Physalin B, a compound found in the plant, has been specifically linked to antimicrobial activity against *Staphylococcus aureus* and other bacteria. (11,12). These results support the traditional use of *Physalis angulata* in treating of pathological conditions. Consequently, the objective was to investigate the chemical composition, antioxidant capacity, and their impact on bacteria, growth, and some blood characteristics, such as their influence on the clotting factors par and part.

Materials and Methods

Plant Collection

The plant material was collected from Al Khaldiya city in Ramadi, Iraq, in June 2023. The plant was identified by the author. Fruits and leaves were dried at room temperature and stored until use.

Preparation of Plant Extract

Fruits and leaves were separated, carefully cleaned, and air-dried. They were processed using an electric grinder to a fine powder once they were totally dry. The leaves and fruits were macerated separately in a heated process to create the extract. Every component was dried in an electric oven several times. After weighing 50 grams of the

powdered plant material (leaves and fruits combined) from the *Physalis angulata* plant

150 ml of 70% concentrated ethanol was added. The powdered plant material (fruits and leaves) was cooked in a water bath at 50 - 80 degrees Celsius temperature while macerated in the solvent in a glass flask. After that, the flasks were placed in a shaking incubator. The mixture was filtered through cheesecloth several times, and filtered with gauze several times. After that, the fragrance was taken, and the precipitate was discarded and dried by placing it in Petri dishes inside the oven. Then, the skimming process was carried out to obtain the powder extract formed (13).

Diphenyl picrylhydrazyl Scavenging Activity (Change from violet to yellow color)

The scavenging activity of the leaf and fruit extracts was evaluated using the DPPH (2,2-diphenyl-1-picrylhydrazyl) assay to determine their free radical scavenging activity as natural antioxidants. The antioxidant activity was measured according to the method described in (14); 250 mg of the plant extract was dissolved in 50 ml of ethanol to obtain a 5 mg/ml concentration. The volume was adjusted to 10 ml with ethanol to achieve a final concentration of 0.5 mg/ml. A standard solution of ascorbic acid was prepared by dissolving 100 mg of ascorbic acid in 25 ml of distilled water, resulting in a final concentration of 10 mg/ml. This served as the stock solution from which 0.1, 0.2, and 0.3 mg/ml solutions were prepared. Sixty microliters of the standard ascorbic acid solution and the plant extract (as a control) were placed in separate tubes. Then, 3 ml of DPPH solution was added to each tube and vortexed thoroughly. The tubes were kept in the dark for 30 minutes, after which the absorbance was measured at a wavelength of 517 nm compared to the DPPH control.

The DPPH free radical scavenging activity (%) was calculated using the following equation:

$$\text{DPPH Activity (\%)} = \frac{\text{Absorbance of control} - \text{Absorbance of sample}}{\text{Absorbance of control}} \times 100$$

Effect of the Ethanolic Extract on Blood Coagulation

Five healthy people in the 20–40 age range had blood samples drawn from their venous blood. The samples were collected and centrifuged for 15 minutes at 2000 rpm. After separation, the extract was added to the plasma at a 1:1 ratio. To prepare these concentrations of 10, 50, 100, 150, 200, and 250 mg/ml, the stock solution, 500 mg/ml, was made by diluting distilled water; to enable the plasma and extract to combine, the samples were incubated in a rotary mixer for five minutes at 37°C. After that, they were prepared to detect intrinsic and extrinsic coagulation factors. A plasma sample was retained from each sample to serve as a reference and point of comparison (15).

PT and PTT Assays

A glass tube is filled with 100 microliters of plasma, and the obtained extract concentration (10, 50, 100, 150, 200, 250 mg/ml). It is then incubated at 37°C for five minutes. After

adding 200 microliters of PT solution, the timer is set to begin immediately. Seconds are used to measure the clotting time. 100 microliters of extract-treated plasma and 100 microliters of PTT solution at 37°C are added for the PTT test. The clotting time is expressed in seconds, and the timer is set to begin immediately. Every produced concentration of plasma treated with the leaf ethanolic extract, as well as every sample, goes through this process again.

Antibiofilm Activity of Plant Extract against Wound Bacteria

A stock solution is prepared from the powder extract powder by dissolving 5 g of it in 25 ml of dimethyl sulfoxide solution (DMSO) and placing it in the rotary mixer to mix it well. Serial concentrations of 150, 200, and 250 mg/ml were prepared (16), and tubes were prepared containing the culture medium to which concentrations of the ethanolic extract of the leaves of the *P. angulata* plant were added. The bacterial suspension was inoculated into a medium made from the heart and brain marrow. These tubes were incubated for 24 hours at a temperature of 37°C, and then 200 microliters of the culture were added to etch the microtiter plate with three replicates in rows. Verticality for each type of bacterial isolates under study, which is considered the control treatment, then add 200 microliters of the culture of bacteria growing in the culture medium containing the extract, then incubate at a temperature of 37°C for 24 hours.

After the end of the incubation period, the holes are washed with physiological saline water to get rid of non-adherent bacterial cells; then, they are dyed with crystal violet dye at a concentration of 1% for 5 minutes. Then, the holes are washed three times to remove the dye and left to dry at room temperature (17).

To extract the dye from the adherent bacteria, 160 microliters of glacial acetic acid were added. The adherent bacterial cells' and the dye's optical density (OD) were measured using an ELISA reader at a wavelength of 360 nm to ascertain the isolates' effectiveness in producing biofilms. This was compared to the bacterial cells treated with the plant extract to calculate the percentage inhibition of bacterial adhesion using the following equation. (18):

Percentage inhibition of biofilm (%) = $1 - \text{OD}(\text{treated}) / \text{OD}(\text{control}) * 100\%$

Statistical Analysis

The data were analyzed using a one-way, completely randomized design (C.R.D.). The differences between means were tested using the L.S.D. test at a significant level of 0.05. The data were analyzed using SPSS statistical software, and Excel was used to present the results (19).

Results

Chemical Composition of the Plant

Physalis angulata fruits and leaves have been subjected to a phytochemical analysis, which has revealed the presence of several secondary metabolites, including alkaloids,

glycosides, flavonoids, tannins, and phenols (4). Additionally, it includes phytol, an unsaturated branched-chain terpene that is thought to be an effective sterilizing agent because most studies have shown that it may stop the growth of a variety of microorganisms, including fungus and bacteria (20), Table 2.

Antioxidant Activity

The results of the DPPH assay of the fruit and leaf extracts of *P. angulata* showed significant radical scavenging in their ability to capture free radicals. Ascorbic acid was used as a standard for comparing values. It has been mentioned by (21) that plant-derived antioxidants such as tannins, phenolic acids, flavones, flavonoids, and other compounds are considered natural scavengers of free radicals that protect living cells. The ethanolic extracts of the leaves and fruits of *P. angulata* had good ability for both assays.

The ethanol extract of leaves showed radical scavenging activity of 93% over a standard value for ascorbic acid of 88% in DPPH assays, and the ethanol extract of fruits showed radical scavenging activity of 60% over a standard value for ascorbic acid with 88% in DPPH




Fig. 1 Antioxidant activity of the fruit and leaf extract of the *P. angulata* plant. DPPH radical scavenging ability.

assays as shown in (Fig. 1)

All results are presented as mean. Studies by (22) have confirmed that the antioxidant activity of the ethanolic extract can be attributed to the plant's content of phenols and flavonoids, which are known for their effectiveness as antioxidants. These compounds are among the most effective in displacing free radicals by donating hydrogen atoms through their hydroxyl groups.

Effect of the ethanolic extract of *P. angulata* leaves on blood clotting factors.

The findings demonstrated that at doses of 250, 200, and 150 mg/ml, the ethanolic extract of *Physalis angulata* leaves significantly extended both the whole and partial clotting times ($p < 0.05$). At these concentrations, the measurements varied from 138-29, 130-36, and 150-41 seconds, in that order. However, the extract shortened the clotting time to 10, 50, and 100 mg/ml doses. At these concentrations, the measurements varied from 23-19, 23-18, and 18-15 seconds, in that order. The treated samples' total clotting time (PT - Prothrombin time) test findings,




Fig 2. Effect of leaf extract on prothrombin total clotting time

Table 1 : Percentages of inhibition of the formation of the life membrane of the species of wound bacteria using the ethanol extract of the leaves of the harankash plant at a concentration of 150, 200, 250) mg/mL.

| Bacteria isolates | Biofilm Inhibition % | concentration of 150 mg\ml | Control |
|-----------------------|----------------------|----------------------------|----------|
| <i>S. aureus</i> | 50.3 | 0.081 | 0.163 |
| <i>S. epidermidis</i> | 74.3 | 0.065 | 0.253 |
| <i>E. coli</i> | 61.45 | 0.074 | 0.192 |
| <i>P. aeruginosa</i> | 45.92 | 0.073 | 0.135 |
| Mean | 57.9925 | 0.07325 | 0.18575 |
| SD | 12.68606 | 0.006551 | 0.050513 |
| SE | 6.343032 | 0.003276 | 0.025257 |

| Bacteria isolates | Biofilm Inhibition % | concentration of 200 mg\ml | Control |
|-----------------------|----------------------|----------------------------|---------|
| <i>S. aureus</i> | 61.96 | 0.062 | 0.163 |
| <i>S. epidermidis</i> | 75.89 | 0.061 | 0.253 |
| <i>E. coli</i> | 63.54 | 0.070 | 0.192 |
| <i>P. aeruginosa</i> | 48.89 | 0.069 | 0.135 |
| Mean | 62.570 | 0.066 | 0.186 |
| SD | 11.044 | 0.005 | 0.051 |
| SE | 5.522 | 0.002 | 0.025 |

| Bacteria isolates | Biofilm Inhibition % | concentration of 250 mg\ml | Control |
|-----------------------|----------------------|----------------------------|---------|
| <i>S. aureus</i> | 61.96 | 0.062 | 0.163 |
| <i>S. epidermidis</i> | 76.28 | 0.06 | 0.253 |
| <i>E. coli</i> | 59.89 | 0.077 | 0.192 |
| <i>P. aeruginosa</i> | 56.29 | 0.059 | 0.135 |
| Mean | 63.605 | 0.065 | 0.186 |
| SD | 8.769 | 0.008 | 0.051 |
| SE | 4.384 | 0.004 | 0.025 |

which varied from 16 to 14 seconds, were noticeably longer than the control samples (Fig. 2&3 and Table 1).

The partial clotting time (PTT- Partial thromboplastin time) test findings indicated that the clotting time was prolonged at the high doses (150, 200, and 250 mg/ml). The values varied from (43-172), (47-195), and (56-180) seconds, in that order. In comparison to the control sample, which yielded a clotting time ranging from (40-32) seconds at a level of P < 0.05, the values at the doses of (10, 50, and 100) mg/ml ranged from (52-31)




Fig 3. Effect of the ethanolic extract of the leaves on the factors of total clotting time (PT).




Fig 4. Average effect of leaf extract on thromboplastin partial clotting












Fig 5: Effect of the ethanolic extract of the leaves on the factors of partial coagulation time (PTT).





seconds, (100-35) seconds, and (59-31) seconds, respectively (Fig 4 & 5 and Table 2).

Consuming a large number of medicinal plants has been demonstrated in previous research to lower the chance of getting various illnesses in general and cardiovascular disorders in specific. They also mentioned the anticoagulant properties of alcoholic and watery plant extracts (23) and that plants are recognized for their high protein and carbohydrate content, both of which have been demonstrated to have biological effects on blood coagulation. Isolated from a variety of plant species, sugars, including pectin and hemicellulose, have demonstrated anti-thrombin and anti-thrombosis properties (24).

According to earlier research, medicinal herbs have an anticoagulant effect that helps cure illnesses involving venous and arterial thrombosis.

Table 2: Phytochemical analysis

| Retention Time (Min) | Molecular Formula | Molecular Mass g/mol | Peak Area (%) | Compound Name | Structure |
|----------------------|--|----------------------|---------------|---|--|
| 49.849 | C ₁₈ H ₃₆ O | 268.4780 | 3.51 | Hexahydrofarnesyl acetone |  |
| 51.072 | C ₂₀ H ₄₀ O | 296.5315 | 2.055 | 3,7,11,15-Tetramethyl-2-hexadecen-1-ol |  |
| 58.896 | C ₁₈ H ₃₆ O ₂ | 284.4772 | 5.908 | Octadecanoic acid |  |
| 59.848 | C ₁₈ H ₃₂ O ₂ | 280.4464 | 2.673 | 9,12-Octadecadienoic acid (Z,Z)- |  |
| 57.095 | C ₂₀ H ₄₀ O | 296.5315 | 10.192 | Phytol |  |
| 62.796 | C ₂₁ H ₃₈ O ₄ | 354.5247 | 0.281 | 9,12-Octadecadienoic acid (Z,Z)-, 2,3-dihydroxypropyl ester |  |
| 59.848 | C ₁₈ H ₃₂ O ₂ | 280.4464 | 2.673 | 9,12-Octadecadienoic acid (Z,Z)- |  |
| 62.906 | C ₂₁ H ₄₀ O ₃ | 340.5419 | 0.370 | Oleic acid, 3-hydroxypropyl ester |  |

| | | | | | |
|--------|---|----------|-------|--|--|
| 64.095 | C ₂₁ H ₄₀ O ₃ | 340.5415 | 0.605 | Glycidyl stearate |  |
| 70.654 | C ₃₀ H ₄₄ O ₂ | 436.6709 | 0.180 | Anthiaergosatin-5,7,9,22-tetraen, 3-acetoxy- |  |
| 73.695 | C ₂₉ H ₅₀ O ₂ | 430.7075 | 0.716 | α-Tocopherol |  |
| 58.998 | C ₂₃ H ₃₉ NO ₂ | 361.5627 | 0.747 | 3-[(1,5-Dimethyl-hexylamino)-methyl]-5,8a-dimethyl-3a,5,6,7,8,8a,9,9a-octahydro-3H-naphtho[2,3-b]furan-2-one |  |

One such study was conducted by (25) to evaluate the anticoagulant effect of some medicinal plant extracts. The results of the study confirmed that these extracts have an anticoagulant effect.

Anti-biofilm activity of the leaf extract

The results of the study showed that the production of biofilm by the bacterial isolates (*Pseudomonas aeruginosa*, *Escherichia coli*, *Staphylococcus epidermidis*, and *Staphylococcus aureus*) used in the experiment gave different readings depending on the absorption of the bacterial isolates used, as shown in (Table 3). The decrease in the absorption values of the bacterial isolates after treatment with the plant extract indicated that the extract could inhibit the adhesion of bacteria in the wells in the assay plate (26).

The results also showed that the leaf extract of *P. angulata* had an effective inhibiting biofilm formation at concentrations of (150, 200, 250) mg/ml. The percentage inhibition of biofilm formation for *S. aureus*, *S. epidermidis*, *E. coli*, and *P. aeruginosa* after treatment with a concentration of (150) mg/ml were (45.92, 61.45, 74.03, 50.3) %, respectively. However, at a concentration of (200)

mg/ml, the inhibitory percentages for biofilm formation were as follows: (48.89, 63.54, 75.89, 61.96) %. The percentages at a concentration of (250) mg/ml were (56.29, 59.89, 76.28, and 61.96) % for the types of bacterial isolates, respectively.

The study showed that the effectiveness of the extract was at its best at a concentration of (250) mg/ml and that the highest rate of inhibition of biofilm formation was due to *S. epidermidis* at the three concentrations used in the experiment, as shown in Fig. 6.

According to research by (27), bacterial isolates can produce biofilms, which are dependent on the pH (ranging from 5-7.4), temperature (25-37°C), and nutrients that are present in the bacterial environment. Additionally, the study demonstrated a connection between bacteria's pathogenicity and their capacity to produce biofilms, which are the source of several diseases. Plant extracts rich in active chemical components, such as hexadecenoic acid, pentadecanoic acid, phytol, stearic, and oleic acid, are extremely suppressive against numerous bacteria, according to a study by (28) at the Indian University's Department of Biotechnology. Research by (29-32) also

Table 3: Statistical analysis of total blood clotting factor PT

| cons. | Control (mg/mL) | 250 (mg/mL) | 200 (mg/mL) | 150 (mg/mL) | 100 (mg/mL) | 50 (mg/mL) | 10 (mg/mL) |
|----------|--------------------|----------------|----------------|----------------|----------------|---------------|---------------|
| SD | 0.894 | 5.683 | 12.818 | 6.189 | 2.966 | 1.871 | 1.140 |
| SE | 0.400 | 2.542 | 5.732 | 2.768 | 1.327 | 0.837 | 0.510 |
| LSD 5% | | | | 7.73 | | | |
| P- value | | | | 0.000** | | | |




Fig 6. Biofilm formation by wound bacterial isolates using *P. angulata* leaf extract.

Table 4. Statistical analysis of partial blood coagulation factor

| cons. | Control | 250 (mg/mL) | 200 (mg/mL) | 150 (mg/mL) | 100 (mg/mL) | 50 (mg/mL) | 10 (mg/mL) |
|---------|---------|-------------|-------------|-------------|-------------|------------|------------|
| SD | 3.90 | 13.37 | 43.40 | 11.74 | 8.50 | 6.58 | 10.89 |
| SE | 1.74 | 5.98 | 19.41 | 5.25 | 3.80 | 2.94 | 4.87 |
| LSD 5% | | | | 24.24 | | | |
| P-value | | | | 0.000** | | | |

Table 5. Inhibition of membrane biogenesis represents % represent results

| Bacterial type | Extract concentrations (mg/ml) | | |
|---------------------------|--------------------------------|-------------|-------------|
| | 150 | 200 | 250 |
| <i>Staph. Aureus</i> | 50.3± 0.34 | 61.96± 1.35 | 61.96± 0.31 |
| <i>Staph. Epidermidis</i> | 74.3± 0.29 | 75.89± 0.06 | 76.82± 0.02 |
| <i>E. coli</i> | 61.45± 0.24 | 63.54± 0.11 | 59.89± 0.22 |
| <i>P. aeruginosa</i> | 45.92± 0.54 | 48.89± 0.32 | 56.29± 0.06 |

Note: (mean ± SD)

demonstrated the antibacterial properties of the polyphenols found in plants, *P. angulata* has a good amount of anti-inflammatory polyphenol, which is one of the natural antioxidants and has various pharmaceutical importance.

Conclusion

With the leaves having a proportion of 93% and the fruits 60%, *P. angulata*'s fruits and leaves demonstrated strong antioxidant activity and the capacity to displace free radicals. By employing the MTP (Micro titration plate) technique, the leaf extract may prevent the growth of biofilms of various bacterial species, including *S. epidermidis*, *S. aureus*, *E. coli*, and *P. aeruginosa*. It can also influence total and partial clotting factors at 10, 50, and 100 mg/mL doses. It is possible to thin blood using large

dosages (150, 200, or 250 mg/mL). After more investigation, it may be utilized as a blood thinner that helps treat blood clotting disorders.

Acknowledgements

We are grateful to the College of Education for Women, Department of Life Sciences, to all the authors of the various sources that were consulted in the research, and to all who provided information. This research is supported by the College of Education for Women, Anbar University, and we appreciate the support.

Authors' contributions

SH performed the experiments, wrote the paper draft, and AT corrected the paper draft, MA: Supervised the extraction experiment, MJ supervised the experiment of the anti-biofilm activity, and MB Supervised the experiment Coagulation tests and analysis of data by OH. All authors agree to be accountable for all aspects of the paper. All authors read the manuscript and approved it.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest

Ethical issues: None

References

- Kasali FM, Tusiimire J, Kadima JN, Tolo CU, Weisheit A, Agaba AG. Ethnotherapeutic uses and phytochemical composition of *Physalis peruviana* L.: an overview. *Sci World J.* 2021; 2021. <https://doi.org/10.1155/2021/5212348>
- Mohammed Hasan ZY. Phytochemical investigation and antioxidant activity of total phenols in the aerial parts of some Asteraceae family wild plants grown in western of Iraq. *Syst Rev Pharm.* 2020; 11(1).
- Ferreira LM dos SL, Vale AE do, Souza AJ de, Leite KB, Sacramento C, Moreno MLV, et al. Anatomical and phytochemical characterization of *Physalis angulata* L.: A plant with therapeutic potential. 2019; 11(2):171-177. https://doi.org/10.4103/pr.pr_97_18
- de Oliveira AM, Malunga LN, Perussello CA, Beta T, Ribani RH. Phenolic acids from fruits of *Physalis angulata* L. in two stages of maturation. *South African J Bot.* 2020; 131:448-53. <https://doi.org/10.1016/j.sajb.2020.02.029>
- Al Rajab ATH, Khalaf HY. The activity of ethanolic extract of *Capsella bursa* on some biochemical parameters of mice. *Biochem Cell Arch.* 2019; 19(1).
- Ramakrishna Pillai J, Wali AF, Menezes GA, Rehman MU, Wani TA, Arafah A, et al. Chemical composition analysis, cytotoxic, antimicrobial and antioxidant activities of *Physalis angulata* L.: A comparative study of leaves and fruit. *Molecules.* 2022; 27(5):1480. <https://doi.org/10.3390/molecules27051480>
- Shenstone E, Lippman Z, Van Eck J. A review of nutritional properties and health benefits of *Physalis* species. *Plant Foods Hum Nutr.* 2020;75:316-25. <https://doi.org/10.1007/s11130-020-00821-3>
- Hu S, Li X, Yin X, Li R, Zhang R, Zang J, et al. Species-specific identification of *Pseudomonas* based on 16S–23S rRNA gene internal transcribed spacer (ITS) and its combined application with next-generation sequencing. *BMC Microbiol.* 2022;22(1):188. <https://doi.org/10.1186/s12866-022-02607-w>
- Sambyal SS, Sharma P, Shrivastava D. Anti-biofilm activity of selected plant essential oils against *Pseudomonas aeruginosa* and *Staphylococcus aureus*. *Int J Curr Microbiol Appl Sci.* 2017;6(3). <https://doi.org/10.20546/ijcmas.2017.603.051>
- Farhan HN, Hameed AT, Aobad HM. The biological activity of some *Pseudomonas* sp. isolates on the growth of three plant pathogenic fungi under incubator conditions. *Adv Environ Biol.* 2010;4(1):53-7.
- Osho A, Adetunji T, Fayemi SO, Moronkola DP. Antimicrobial activity of essential oils of *Physalis angulata* L. *African J Tradit Complement Altern Med.* 2010;7(4). <https://doi.org/10.4314/ajtcam.v7i4.56696>
- Silva MTG, Simas SM, Batista TGFM, Cardarelli P, Tomassini TCB. Studies on antimicrobial activity, *in vitro*, of *Physalis angulata* L. (Solanaceae) fraction and physalin B bringing out the importance of assay determination. *Mem Inst Oswaldo Cruz.* 2005;100:779-82. <https://doi.org/10.1590/S0074-02762005000700018>
- Jilbhkate YJ, Awachat AP, Lohiya RT, Umekar MJ, Hemke AT, Gupta KR. Extraction: An important tool in the pharmaceutical field. *Int J Sci Res Arch.* 2023;10(1):555-68. <https://doi.org/10.30574/ijrsra.2023.10.1.0768>
- Adámez JD, Samino EG, Sánchez EV, González-Gómez D. *In vitro* estimation of the antibacterial activity and antioxidant capacity of aqueous extracts from grape-seeds (*Vitis vinifera* L.). *Food Control.* 2012;24(1–2):136-41. <https://doi.org/10.1016/j.foodcont.2011.09.016>
- Gunendren M, Noordin SS, Muggundha R, Nozlana AS. Effect of *Ocimum sanctum* (Tulsi) aqueous leaf extract on prothrombin time (PT), activated partial thromboplastin time (APTT), and thrombin time (TT) of human plasma. *J Biomed Clin Sci.* 2017;2(1).
- Rutter WC, Burgess DR, Burgess DS. Increasing incidence of multidrug resistance among cystic fibrosis respiratory bacterial isolates. *Microb drug Resist.* 2017;23(1):51-5. <https://doi.org/10.1089/mdr.2016.0048>
- Dahal RH, Chaudhary DK. Microbial infections and antimicrobial resistance in Nepal: current trends and recommendations. *Open Microbiol J.* 2018;12:230. <https://doi.org/10.2174/1874285801812010230>
- Gudiña EJ, Rocha V, Teixeira JA, Rodrigues LR. Antimicrobial and antiadhesive properties of a biosurfactant isolated from *Lactobacillus paracasei* ssp. *paracasei* A20. *Lett Appl Microbiol.* 2010;50(4):419-24. <https://doi.org/10.1111/j.1472-765X.2010.02818.x>
- Steel RGD, Torrie JH. Principles and procedures of statistics. *Princ Proced Stat.* 1960;
- Chou MY, Andersen TB, Mechan Llontop ME, Beculheimer N, Sow A, Moreno N, et al. Terpenes modulate bacterial and fungal growth and sorghum rhizobiome communities. *Microbiol Spectr.* 2023;11(5):e01332-23. <https://doi.org/10.1128/spectrum.01332-23>
- Amer AA, Mohammed RS, Hussein Y, Ali ASM, Khalil AA. Development of *Lepidium sativum* extracts/PVA electrospun nanofibers as wound healing dressing. *ACS omega.* 2022;7(24):20683-95. <https://doi.org/10.1021/acsomega.2c00912>
- Lee CY, Sharma A, Semenya J, Anamoah C, Chapman KN, Barone V. Computational study of ortho-substituent effects on antioxidant activities of phenolic dendritic antioxidants. *Antioxidants.* 2020;9(3):189. <https://doi.org/10.3390/antiox9030189>
- Omar G, Abdallah L, Barakat A, Othman R, Bourinee H. Eficácia hemostática *in vitro* de extratos aquosos, metanólicos e etanólicos de três espécies de plantas medicinais na Palestina. *Brazilian J Biol.* 2019;80:763-8. <https://doi.org/10.1590/1519-6984.219186>
- Mengome LE, Voxeur A, Akue JP, Lerouge P, Ngouamizokou A, Moutsimbi RM. Analysis of toxicity and hemostatic properties of polysaccharides from plants endemic to Gabon. *Br J Pharmacol Toxicol.* 2014;5(6):186-93. <https://doi.org/10.19026/bjpt.5.5185>
- Alabdallat NG, Dukhyil AAA Bin. *In vitro* anticoagulant effect of methanolic extracts of *Artemisia herba-alba*, *Achillea fragrantissima* and *Citrullus colocynthis* grown in Saudi Arabia. 2021;
- Stiefel P, Rosenberg U, Schneider J, Mauerhofer S, Maniura-Weber K, Ren Q. Is biofilm removal properly assessed? Comparison of different quantification methods in a 96-well plate system. *Appl Microbiol Biotechnol.* 2016;100:4135-45. <https://doi.org/10.1007/s00253-016-7396-9>

27. Namasivayam SKR, Vivek JM. Screening of quorum sensing (qs) modulatory effect of medicinal plant extracts against quorum sensing mediated virulence factors of human pathogenic gram-negative bacteria. *Int J Pharm Phytochem Res.* 2016;8:263.
28. Álvarez-Martínez FJ, Barrajón-Catalán E, Encinar JA, Rodríguez-Díaz JC, Micol V. Antimicrobial capacity of plant polyphenols against gram-positive bacteria: A comprehensive review. *Curr Med Chem.* 2020;27(15):2576-606. <https://doi.org/10.2174/0929867325666181008115650>
29. Jassim RA, Hamid AT. Molecular Characterization of Selected Genera of the (Apiaceae) Family using SSR Molecular Markers. *Indian J Forensic Med Toxicol.* 2020;14(4).
30. Mohammed IH, Hameed AT, Salman HF. Phytochemical and biological of *Anthemis nobilis* (Asteraceae family) a native herb of Iraq. *Sys Rev Pharm.* 2020;11(2):458-61.
31. Ushie OA, Neji PA, Abeng FE, Okpashi VE, Baba NH, Azuaga TI. Phytochemical Screening and Antimicrobial Activities of Chloroform and Ethyl Acetate Extracts of *Physalis angulata*. *J Chem Soc Niger.* 2019;44(6).
32. Al-Joboury KR, Aliwy SA. Survey With Revised Checklist of Compositae in The Herbarium of Iraq Natural History Research Center and Museum. *Bull Iraq Nat Hist Museum.* 2023;17(3):375-407. <https://doi.org/10.26842/binhm.7.2023.17.3.0375>