



RESEARCH COMMUNICATION

# Sole and sequential application of herbicides and botanicals for weed management in summer sesame

Swastika Hota<sup>1</sup>, Subhprada Dash<sup>1\*</sup> & Utpal B. Joshi<sup>1</sup>

<sup>1</sup>Department of Agronomy, Faculty of Agricultural Sciences, Siksha 'O' Anusandhan, Deemed to be University, Bhubaneswar-751029, Odisha, India

\*Email: [subhpradadash@soa.ac.in](mailto:subhpradadash@soa.ac.in)

 OPEN ACCESS

## ARTICLE HISTORY

Received: 17 June 2024  
Accepted: 17 September 2024  
Available online  
Version 1.0 : 23 October 2024  
Version 2.0 : 24 October 2024



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Hota S, Dash S, Joshi UB. Sole and sequential application of herbicides and botanicals for weed management in summer sesame. Plant Science Today. 2024; 11(4): 1022-1027. <https://doi.org/10.14719/pst.4122>

## Abstract

Weed infestation poses a significant challenge in sesame cultivation, leading to yield losses and decreased economic returns for farmers. The present study comprehensively analyses various weed control approaches, including allelopathic interaction with integrating herbicides and hand-weeding practices. The experiment was conducted at the Agriculture Farm, SOA, Odisha, during the summer of 2023 with 10 weed management treatments in RBD with 3 replications. The objective was to study different weed management practices, including the sole and sequential application of herbicides and botanicals, to control diverse weeds and enhance summer sesame growth, yield and profitability. The dominance of *Digitaria sanguinalis* (L.) Scop., *Echinochloa colona* (L.) Link. and *Poa annua* L. among grassy weeds and *Melochia corchorifolia* L., *Cleome viscosa* L. and *Cassia tora* L. among the broadleaved weeds were evident in summer sesame. The execution of different treatments showed that integrated use of aqueous leaf extract of *Parthenium hysterophorus* at 5 % concentration, followed by quizalofop-p-ethyl at 50 g/ha, was found on par with hand-weeding and brought down the weed count and biomass, which in turn increased the growth of summer sesame. The maximum yield of (793 kg seed/ha and 3035 kg stalk/ha) was registered under twice hand weeding. It was comparable with the spraying of aqueous leaf extract of *Parthenium hysterophorus* at 5 % concentration followed by (*fb*) quizalofop-p-ethyl at 50 g/ha (772 kg seed/ha and 2968 kg stalk/ha). Economic analysis of sesame revealed that the maximum return (Rs. 37454/- per ha) and return/rupee(2.28) were recorded with aqueous leaf extract of *P. hysterophorus* at 5 % concentration *fb* quizalofop-p-ethyl at 50 g/ha treatment.

## Keywords

Allelopathy; botanicals; integrated weed management; nerium; parthenium

## Introduction

Sesame (*Sesamum indicum* L.) is prominent amongst the oilseeds for its high-quality oil and rich nutritional offerings. It is an annual herbaceous crop which belongs to the family Pedaliaceae. The flower is axillary and pedicellate and the fruit is known as a capsule. Sesame holds a wealth of nutritional benefits. These tiny seeds are the powerhouse of essential nutrients, boasting high concentrations of protein, healthy fats, dietary fibre and an array of fat-soluble vitamins and amino acids. It is also rich in minerals such as calcium and iron (1). Globally, the sesame is grown in an 11743 thousand (000) ha area with a production of 6016 thousand (000) M

tons and 512 kg/ha productivity. India contributed around 1730 Thousand (000) ha in area, with a production and productivity of 746 Thousand M tons and 431 kg/ha in 2018 respectively. India produces nearly 12.4 % of sesame cultivation (2). Weed infestation in sesame cultivation can lead to substantial yield losses, hinder efficient harvesting and compromise the quality of the produce. Sesame is often cultivated in diverse agro-climatic conditions, infested with varying weed species. To achieve the required productivity, weed control is essential. If weeds are not effectively managed, they can severely affect the yield of crops by competing with sesame plants for resources such as nutrients, space, water and sunlight.

*Cynodon dactylon* (L.) Pers., *Echinochloa colona* (L.) Link., *Digitaria sanguinalis* (L.) Scop., *Dactyloctenium aegyptium* (L.) P. Beauv. Willd., *Cyperus rotundus* L., *Trianthema portulacastrum* L., *Spilanthes acmella* L., *Amaranthus viridis* L., *Phyllanthus niruri* L., *Commelina benghalensis* L., *Digera arvensis* Forsk. and *Malvastrum coromandalianum* Garcke. are commonly found as predominant weeds in sesame cultivation (3-5). Sesame experienced a significant reduction in seed yield, ranging from 50 % to 75 %, because of intense weed competition (6). Weed management is not only about protecting the crop yield but also about optimizing input costs. As the demand for this oilseed crop continues to rise, the challenges posed by weed infestations have become increasingly evident. Excessive reliance on chemical herbicides poses environmental risks in pursuing productive sesame cultivation. It leads to a shift in weed flora and the emergence of herbicide-resistant weed populations (7). Relying solely on a single method can have several drawbacks. At the same time, manual weeding has been a traditional practice in sesame cultivation, which requires a significant amount of labour and incurs substantial costs for farmers (8). Manual weeding is dependent on weather conditions. Wet conditions can delay and make manual weeding difficult, allowing weeds to establish and compete with the crop and keep the weed in an advantageous position (9).

Allelopathy is a biological phenomenon where certain plants release biochemicals, known as allelochemicals, into the environment. These allelochemicals can potentially influence crops' growth, development and germination, either inhibiting or promoting their growth (10). In *Parthenium hysterophorus*, the allelochemicals named sesquiterpene lactones and parthenin have herbicidal properties, which can impact the growth and development of weeds (11). Similarly, the flowering plant *Nerium oleander* L. is well-known for its flavonoid and glycoside compounds, among which oleandrine is the most prominent glycoside, which can be used as botanicals to control weeds at different concentrations (12). The aqueous extract of *P. hysterophorus* significantly reduced the germination of *Digitaria sanguinalis* (L.) Scop. and *Eleusine indica* (L.) Gaertn. (13). Similarly, the application of dry leaf extracts of *Nerium oleander* at different concentrations effectively affected the initiation and growth of *Cyperus rotundus* L.

(14). In addition to their herbicidal activities, the allelopathic potential of *P. hysterophorus* and *N. oleander* has been exploited for control of fungal and bacterial diseases (15-17). These findings may shed light on using allelopathic approaches to an overall integrated weed management (IWM) strategy to manage weeds sustainably. Therefore, a field trial was conducted to investigate the competent IWM approach to manage weeds and enhance sesame's growth, yield and profitability during the summer season.

## Materials and Methods

### a. Experimental site:

The investigation was executed during the summer of 2023 at the Agriculture Farm, Binjhagiri, Chhatabar, Odisha, positioned at coordinates 20°15' N latitude, 85°40' E longitude and 58 m altitude above MSL. The soil of the experimental field exhibited a clayey texture, was deep and featured a slight slope from west to east, contributing to adequate drainage. It was characterized by acidic nature with a pH of 5.87. The soil content showed lower levels of soil organic carbon (0.36 %) as analyzed by the volumetric weight combustion method and available N (204.5 kg/ha) by alkaline permanganate method, while being moderate in both available P<sub>2</sub>O<sub>5</sub> (21.27 kg/ha) and K<sub>2</sub>O (131.1 kg/ha) by Bray's method no. 1 and flame photometer method respectively.

### b. Treatments and crop management:

A randomised block design framed the field experiment consisting of 10 treatments with 3 replications. Sesame seeds of the 'Smarak' variety were line sown with a 6 kg/ha seed rate at 30 cm x 10 cm spacing in the north-south direction. The treatments consist of solely applying pendimethalin, quizalofop-p-ethyl, *Parthenium hysterophorus* and *Nerium oleander* leaf extract at 5 and 10 % concentration. Further botanicals and herbicides were used for sequential application and compared with 2 hand weeding and weedy check. Pendimethalin, the aqueous leaf extract of *P. hysterophorus* and *N. oleander* were applied one day after sowing (DAS), whereas quizalofop-p-ethyl was applied at 20 DAS. The crop was supplied with nutrients of 40-20-20 kg of N-P<sub>2</sub>O<sub>5</sub>-K<sub>2</sub>O respectively, per ha, using urea, SSP and MOP. The complete dose of phosphorus, potassium and 50 % nitrogen was applied as a basal before the sowing. The remaining dose of N was top-dressed at 21 DAS under sufficient soil moisture.

### c. Data collection and observation taken:

The observations on plant growth characteristics, including plant height and dry matter accumulation (DMA), were measured at the 40 DAS. The weed count and biomass were also documented at 40 DAS using two randomly selected quadrats, each measuring 50 cm x 50 cm within the sampling area of each plot. Weed samples were cut close to the ground level, sun-dried and oven-dried at 65 ± 5 °C until a constant weight was obtained to record biomass. The yield was measured from each plot's

net plot area (5 m x 4 m) and recorded in kg/ha. The rate of increase in dry weight per unit of ground area per unit of time calculated crop growth rate (CGR). Weed control efficiency was computed using the dry weight of weeds. It was worked out from a reduction in weed dry weight due to the weed control method over the unweeded check. The weed index was a gain in crop yield due to weed control as a percentage of yield from weed-free crops. The significance of different sources of variation was tested by the "Error Mean Square Method" of Fisher Snedecor's "F" test at probability level 0.05. The analysis of variance method was followed for analyzing the experimental data statistically (18). The standard error of means (SEM  $\pm$ ) and the value of critical differences (C.D.) between means have been provided.

## Results and Discussion

During the investigation, 8 weed species *viz.* *Digitaria sanguinalis* (L.) Scop., *Echinochloa colona* (L.) Link. and *Poa annua* L. and *Dactyloctenium aegyptium* (L.) P. Beauv. Willd. among the grasses; *Melochia corchorifolia* L., *Cleome viscosa* L. and *Cassia tora* L. among the broadleaved and *Cyperus iria* L. only sedge, were predominantly found in the experimental field. Similar grassy weeds were documented in sesame cultivation (4). The least density and weed biomass were observed in plots with twice hand weeding ( $T_9$ ) (13 No./m<sup>2</sup> and 2.77 g/m<sup>2</sup>). Among the integrated practices, the aqueous leaf extract of *P. hysterophorus* at 5 % concentration, *fb* quizalofop-p-ethyl at 50 g/ha ( $T_8$ ) (18 No./m<sup>2</sup> and 4.53 g/m<sup>2</sup>) exhibited lower weed density and biomass (Table 1). This was because of the reduction of the first flush of weeds by the *Parthenium* extract and later, weeds were controlled by the herbicide. Aqueous extract of *Parthenium hysterophorus* at 40 % resulted in a potent allelopathic interaction inhibiting germination, seedling growth and dry matter accumulation of *T. populnea* (19). Similarly, applying *P. hysterophorus* extracts remarkably reduced germination alongside the initiation of *Cyperus iria*, i.e., coleoptile and radicle length. They also found that the increase in the

concentration of *P. hysterophorus* extract was exhibited in lower biomass and the *Cyperus iria* leaf area (20). The weed biomass was effectively reduced under the integrated application of aqueous leaf extract of *N. oleander* at 5 % conc. *fb* quizalofop-p-ethyl was significantly at par with the sole application of quizalofop-p-ethyl. Integrated weed management approach, including quizalofop, effectively lowered the weed biomass (21, 22). Apart from  $T_8$  and  $T_9$  treatments, the total weed density was reduced under the integrated application of aqueous leaf extract of *Nerium oleander* at 5 % concentration *fb* quizalofop-p-ethyl at 50 g/ha, which was closely followed by the sole application of quizalofop-p-ethyl at 50 g/ha, pendimethalin at 750 g/ha and aqueous leaf extract of *Parthenium hysterophorus* at 10 % concentration treatments in sesame during summer season (Table 1). Application of quizalofop-p-ethyl at 50 g/ha decreased the grassy weed population (7.33 no/0.25 m<sup>2</sup>) (23). The application of pendimethalin at 1 kg/ha controlled most of the annual grasses and broad-leaved weeds during the germination of weed seeds. Still, pendimethalin at 750 g/ha failed to control *Commelina benghalensis* more effectively (24).

Applying twice hand weeding had 95.29 % weed control efficiency (WCE) among weed management treatments - *P. hysterophorus* at 5 % conc. *fb* quizalofop-p-ethyl recorded the maximum WCE (92.32 %) followed by *Nerium oleander* at 5 % at 1 DAS *fb* quizalofop-p-ethyl (80.41 %) (Fig. 1). The use of quizalofop-p-ethyl at 50 g/ha had weed control efficiency of 47.79 % (23). The maximum dry matter and CGR growth were registered under two-hand weeding, which was closely followed by aqueous leaf extract of *Parthenium hysterophorus* at 5 % *fb* quizalofop-p-ethyl at 50 g/ha (Table 1). The integrated weed management treatments exhibited the maximum dry matter accumulation of sesame, which was on par with the two-hand weeding (25)-however, the sole application of quizalofop-p-ethyl registered maximum growth parameters. In the present study, the application of both *Parthenium* and *Nerium* aqueous leaf extract at 10 %

**Table 1.** Weed and crop growth under various weed management treatments at 40 DAS.

Treatments	Weed growth		Crop growth		
	Population (no./m <sup>2</sup> )	Biomass (g/m <sup>2</sup> )	Plant height (cm)	DMA (g/m <sup>2</sup> )	CGR (g/m <sup>2</sup> /day)
T <sub>1</sub> - Pendimethalin at 750 g/ha	6.09 (36.6)*d	3.60 (12.4)d	54.18ab	117.2bc	6.03b
T <sub>2</sub> - Quizalofop-p-ethyl at 50 g/ha	6.02 (35.7)d	3.56 (12.1)d	55.00ab	126.1b	6.07b
T <sub>3</sub> - ALEN at 5 %	8.15 (66.0)b	5.60 (30.8)b	48.11bc	102.4cd	5.19bc
T <sub>4</sub> - ALEN at 10 %	7.07 (49.6)c	4.65 (21.1)c	45.08c	88.8d	4.62cd
T <sub>5</sub> - ALEP at 5 %	6.95 (48.0)c	4.79 (22.5)c	48.36bc	103.7cd	5.38bc
T <sub>6</sub> - ALEP at 10 %	6.39 (40.3)cd	4.39 (18.8)c	45.95c	91.5d	5.09bc
T <sub>7</sub> - T <sub>3</sub> <i>fb</i> T <sub>2</sub>	5.78 (33.1)d	3.45 (11.5)d	54.29ab	123.1b	5.90b
T <sub>8</sub> - T <sub>5</sub> <i>fb</i> T <sub>2</sub>	4.29 (18.0)e	2.23 (4.5)e	55.92a	145.4a	7.35a
T <sub>9</sub> - Two HW	3.67 (13.0)e	1.80 (2.7)e	56.51a	147.0a	7.42a
T <sub>10</sub> -Weedy check	10.03 (100.3)a	7.70 (58.9)a	34.94d	46.7e	3.46d
CD(P=0.05)	0.69	0.45	7.48	16.93	1.24

\*The bracketed value is original, and the square root was transformed before analysis. Treatments with similar letters are not significantly different.

WCE: Weed control efficiency, DMA: Dry matter accumulation, CGR: Crop growth rate, ALEN: Aqueous leaf extract of *Nerium oleander*, ALEP: Aqueous leaf extract of *Parthenium hysterophorus*, H.W.: Hand weeding


showed a phytotoxic effect and a significant decrement in the plant height of sesame. An increment in the concentration of *P. hysterophorus* extract notably hampered gingly's initiation and seedling length (26).

The highest productivity (793 kg seed/ha and 3035 kg stalk/ha) was observed in hand-weeded plots, followed by plots treated with the *P. hysterophorus* leaf extract at 5 % concentration *fb* quizalofop (772 kg seed/ha and 2968 kg stalk/ha) (Table 2). The application of pre-emergence herbicide followed by quizalofop-p-ethyl documented significantly higher seed yield (1002 kg/ha) and stalk yield (3374 kg/ha) (5). In the present study, the subsequent best treatment in registering higher levels of seed and stalk yield was the sole application of quizalofop-p-ethyl at 50 g/ha at 15 DAS (727 kg seed/ha and 2629 kg stalk/ha<sup>-1</sup>) which was at par with the aqueous leaf extract of *Nerium oleander* at 5 % concentration at 1 DAS *fb* quizalofop-p-ethyl at 50 g/ha at 15 DAS (709 kg seed/ha and 2626 kg stalk/ha) and lone application of pendimethalin at 750 g/ha<sup>-1</sup> at 1 DAS (692 kg seed/ha and 2533 kg stalk/ha) (Table 2). The infestation of grassy weeds was higher than broadleaved and sedge weeds, which was effectively controlled by the sole application of quizalofop-p-ethyl. The post emergence application of quizalofop significantly improved the seed yield and plant growth parameters by effectively managing the weed population (27). Applying

pendimethalin at 750 g/ha produced a much greater seed yield (5.463 q/ha) and caused low injury in sesame plants (28). The harvest index of sesame did not indicate any variation by the imposition of different weed management practices and remained relatively similar in all the treatments.

The weedy check plot registered % maximum weed index (W.I.) of 81.70 %. The lowest W.I. was observed in 2 hand weeding treatments followed by the integrated application of *Parthenium hysterophorus* at 5 % concentration *fb* quizalofop (2.64 %) and sole application of quizalofop-p-ethyl (8.25 %) (Fig. 2).


The higher net return of Rs. 37454/ha and return/rupee (2.28) were recorded with the application of aqueous leaf extract of *Parthenium hysterophorus* at 5 % concentration *fb* quizalofop-p-ethyl at 50 g/ha. It was followed by applying quizalofop-p-ethyl with net return and return/rupee of Rs. 35107/ha and 2.27 respectively (Table 2). The net returns as well as the B:C ratio, were higher with the treatment of weed-free check, found to be significantly at par with the integrated application involving quizalofop as a postemergence herbicide (29). The lowest net return was noticed in weedy plots (Table 2). The lowest return values are documented in the weedy check plot due to the inability of the sesame crop to perform under higher weed competition (30).



**Fig. 1.** Effect of weed management on weed control efficiency (%) of summer sesame at 40 DAS.

T<sub>1</sub>- Pendimethalin at 750 g/ha, T<sub>2</sub>- Quizalofop-p-ethyl at 50 g/ha, T<sub>3</sub>- ALEN at 5 %, T<sub>4</sub>- ALEN at 10 %, T<sub>5</sub>- ALEP at 5 %, T<sub>6</sub>- ALEP at 10 %, T<sub>7</sub>- T<sub>3</sub>*fb* T<sub>2</sub>, T<sub>8</sub>- T<sub>5</sub>*fb* T<sub>2</sub>, T<sub>9</sub>- Two HW, T<sub>10</sub>-Weedy check.

\* ALEN: Aqueous leaf extract of *Nerium oleander*, ALEP: Aqueous leaf extract of *Parthenium hysterophorus*, H.W.: Hand weeding



**Fig. 2.** Weed index (%) of sesame under different weed management treatments.

T<sub>1</sub>- Pendimethalin at 750 g/ha, T<sub>2</sub>- Quizalofop-p-ethyl at 50 g/ha, T<sub>3</sub>- ALEN at 5 %, T<sub>4</sub>- ALEN at 10 %, T<sub>5</sub>- ALEP at 5 %, T<sub>6</sub>- ALEP at 10 %, T<sub>7</sub>- T<sub>3</sub>*fb* T<sub>2</sub>, T<sub>8</sub>- T<sub>5</sub>*fb* T<sub>2</sub>, T<sub>9</sub>- Two HW, T<sub>10</sub>-Weedy check.

\* ALEN: Aqueous leaf extract of *Nerium oleander*, ALEP: Aqueous leaf extract of *Parthenium hysterophorus*, H.W.: Hand weeding

**Table 2.** Effect of weed management on yield and economics of sesame.

Treatments	Yield (kg/ha)		Harvest Index (%)	Economics	
	Seed	Stalk		Net return (Rs./ha)	Return/ Rupee invested
T <sub>1</sub> - Pendimethalin at 750 g/ha	692.12c	2533.76cd	21.58a	30669	2.05
T <sub>2</sub> - Quizalofop-p-ethyl at 50 g/ha	727.88bc	2629.51bc	21.76a	35107	2.27
T <sub>3</sub> - ALEN at 5 %	610.30de	2179.82de	21.89a	25704	1.95
T <sub>4</sub> - ALEN at 10 %	510.61f	1846.18e	21.80a	16496	1.60
T <sub>5</sub> - ALEP at 5 %	629.70d	2195.58de	22.30a	27379	2.01
T <sub>6</sub> - ALEP at 10 %	562.42ef	2051.94e	21.57a	20970	1.76
T <sub>7</sub> - T <sub>3</sub> <i>fb</i> T <sub>2</sub>	709.88c	2626.79bc	21.29a	32053	2.10
T <sub>8</sub> - T <sub>5</sub> <i>fb</i> T <sub>2</sub>	772.42ab	2968.30ab	20.68a	37454	2.28
T <sub>9</sub> - Two HW	793.33a	3035.27a	20.74a	16009	1.30
T <sub>10</sub> -Weedy check	145.15g	599.51f	19.39a	-12961	0.49
CD (P=0.05)	62.35	374.17	NS	-	-

Treatments with similar letters are not significantly different.

WI: Weed index, ALEN: Aqueous leaf extract of *Nerium oleander*, ALEP: Aqueous leaf extract of *Parthenium hysterophorus*, H.W.: Hand weeding



## Conclusion

The aqueous leaf extract of *Parthenium hysterophorus* at 5 % concentration fb quizalofop-p-ethyl at 50 g/ha significantly reduced the weed densities and biomass. The seed and stalk yield of the treatment were 772 and 2968 kg/ha respectively. Economic analysis of sesame revealed that the maximum return (Rs. 37454/- per ha) and return/rupee (2.28) was recorded with the application of *Parthenium hysterophorus* extract at 5 % concentration at 1 DAS fb quizalofop-p-ethyl at 50 g/ha at 20 DAS. Thus, integration of aqueous leaf extract of *Parthenium hysterophorus* at 5 % concentration at 1 DAS sequenced with the early postemergence application of quizalofop-p-ethyl at 50 g/ha proved to be a viable option for managing the weed infestation and has the potential to strengthen the growth, yield and profit in summer sesame. Future research should delve deeply into the role of botanicals in weed management, as their potential benefits are not yet fully understood. Detailed studies are needed to assess their effectiveness, mechanisms of action and any possible side effects or interactions with crops and soil ecosystems. By exploring various plant extracts and their properties, researchers can uncover novel solutions for controlling weed growth, improving crop yields and promoting ecological balance.

## Acknowledgements

The authors express their gratitude to Siksha' O' Anusandhan, Deemed to be University, for providing the facilities for conducting the experiment and Lab-analysis.

## Authors' contributions

SH carried out the field experiment and collected and analyzed the data. SD conceptualized and supervised the experiment and prepared the manuscript. UJ participated in the data collection and drafted the manuscript. All authors read and approved the final manuscript.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interest to declare.

**Ethical issues:** None

## References

- Wei P, Zhao F, Wang Z, Wang Q, Chai X, Hou G, Meng Q. Sesame (*Sesamum indicum* L.): A comprehensive review of nutritional value, phytochemical composition, health benefits, development of food and industrial applications. *Nutr.* 2022;14. <https://doi.org/10.3390/nu14194079>.
- Food and Agriculture Organization Statistical Databases (FAOSTAT). Available online: <http://faostat.fao.org/>.
- Joshi N, Joshi S, Sharma JK, Shekhawat HS, Shukla UN. Efficacy of sequential application of pre- and postemergence herbicides for weed management in sesame. *Ind J Weed Sci.* 2022;54(3): 279-82.
- Fatima A, Duary B. Integrated use of herbicide and straw mulch in suppressing the weed species in summer sesame (*Sesamum indicum*). *Ind J Agric Sci.* 2020;90(11):92-97.
- Ambika M, Sundari A. Weed management in irrigated sesame (*Sesamum indicum* L.). *World Sci News (An Int Sci J).* 2019;131(1):272-78.
- Mruthul T, Halepyati AS, Chittapur BM. Chemical weed management in sesame (*Sesamum indicum* L.). *Karnataka J Agric Sci.* 2015;28(2):151-54.
- Hazarika JR, Deka AM, Borah B, Gogoi B, Gogoi A, Kalita B, et al. Weedflora shift as affected by cropping systems. *Int J Res Agron.* 2024;SP-7(8):415-20. <https://doi.org/10.33545/2618060X.2024.v7.i8Sf.1285>.
- Zimbere E, Mashingaidze N, Muzemu S, Rugare JT. Growth and tolerance of sesame (*Sesamum indicum* L.) varieties to pre and postemergence graminicides. *Adv Agric.* 2022;2022:1-7. <https://doi.org/10.1155/2022/7439281>.
- Ramesh K, Matloob A, Aslam F, Florentine SK, Chauhan BS. Weeds in a changing climate: Vulnerabilities, consequences and implications for future weed management. *Front Plant Sci.* 2017;8:95.
- Erida G, Ichsan CN, Syamsuddin, Kurniawanc T, Khan IH, Javaid A. Potential of secondary metabolites of *Ageratum conyzoides* L. in weed management: A review. *Allelopathy J.* 2023;58(1):23-40.
- Pandey DK. Allelochemicals from *Parthenium* for water hyacinth control. *Indian J Weed Sci.* 2015;47(3):321-28.
- Mojarad AK, Majd A, Fahimi H. Allopathic effects of *Nerium oleander* L. on growth and anatomy structure of *Hordeum vulgare* (monocotyledon) and *Vicia sativa* (dicotyledon) seedlings. *Adv Environ Biol.* 2013;1:766-72.
- Bashar HMK, Juraimi AS, Ahmad, Hamdani MS, Uddin MK, Asib N, Anwar MP. Evaluation of allelopathic effects of *Parthenium hysterophorus* L. methanolic extracts on some selected plants and weeds. *Plos One.* 2023;18(1):e0280159.
- Begam UJ, Srikrishnah S, Sutharsan S. Allelopathic suppressive effects of *Thevetia peruviana* L. and *Nerium oleander* L. leaf extracts on germination and seedling growth of *Cyperus rotundus* L. *AGRIEAST: J Agric Sci.* 2020;14(2):43-51.
- Bajwa R, Shafique S, Khaliq A, Javaid A, Shafique S. Fungal biological control potential of aqueous extract of *Parthenium hysterophorus* L. In: Proceedings of 4<sup>th</sup> National Conference of Plant Pathology. Arid Agriculture University, Rawalpindi; 14-16 October, 2003:89-96.
- Saeed HM, Ferdosi MFH, Khan IH, Javaid A, Sultan MW. Antibacterial activity and GC-MS analysis of white flowers extract of *Nerium oleander* L. *Int J of Biol Biotechnol.* 2023;20(1):163-68.
- Saeed HM, Ferdosi MFH, Javaid A, Khan IH, Sattar T. Antibacterial activity and phytochemical profile of flower extract of pink oleander. *Int J Biol Biotechnol.* 2023;20(2):353-58.
- Gomez KA, Gomez AA. Statistical procedure for agricultural research. John Wiley and sons, London. 2003;139-67, 204-07.
- Thirupathi P, Senthivelu M, Prabha AC. Allelopathic effect of Lantana and *Parthenium* on germination and growth of *Thespesia* tree species. *Ind J Weed Sci.* 2020;52(4):400-04.
- Motmainna M, Juraimi AS, Uddin MK, Asib NB, Islam AKMM, Hasan M. Bioherbicidal properties of *Parthenium hysterophorus*, *Cleome ruidosperma* and *Borreria alata* extracts on selected crop and weed species. *Agron.* 2021;11:643.
- Mathukia RK, Sagarka BK, Jadav CN. Integrated weed management in summer sesame. *Indian J Weed Sci.* 2015;47(2):150-52.

22. Sujithra P, Hemalatha M, Joseph M, Sobhana E. Effect of different weed management practices on growth and yield of rainfed sesame (*Sesamum indicum* L.) under vertisol. Madras Agric J. 2019;106(3):1.
23. Mane SV, Kanade VM, Shendage GB, Sarawale PP, Shetye VN. Weed management in sesamum (*Sesamum indicum* L.) grown under coastal region of Maharashtra. J Indian Soc Coast Agric Res. 2017;35(1):31-33.
24. Babu KS, Subramanyam D, Sumathi V, Umamahesh V. Weed management in sesame with sequential application of herbicides. Indian J Weed Sci. 2016;48(4):455-57.
25. Venu M, Reddy GS, Mahesh N, Devi MU. Effect of integrated weed management practices on growth and yield of sesame (*Sesamum indicum* L.). Biol Forum-Int J. 2022;14(2a):271-75.
26. Bhuvaneshwari K, Muniyappan C, Priyadharshini D, Sathyapriya P, Rathika S, Ramesh T. Allelopathic effect of *Parthenium hysterophorus* L. on germination and seedling growth of different field crops. J Pharm Innov. 2019;8(7):257-61.
27. Gupta S, Kushwah SS. Postemergence herbicides for weed control in sesame. Ind J Weed Sci. 2016;48(1):97-98.
28. Singh R, Ghosh D, Dubey RP, Singh VP. Weed control in sesame with pre-emergence herbicides. Ind J Weed Sci. 2018;50(1):91-93.
29. Mane SV, Kanade VM, Mhashelkar AU. Effect of mechanical and chemical weed control measures on yield and economics of sesame. Int J Curr Microbiol Appl Sci. 2018;6:2979-83.
30. Nisha Bhadauria, KS Yadav, RL Rajput, VB Singh. Integrated weed management in sesame. Indian J Weed Sci. 2012;44(4):235-237.